



Irreducible modules of modular Lie superalgebras and super version of the first Kac–Weisfeiler conjecture

Bin Shu

Abstract. Suppose $\mathfrak{g} = \mathfrak{g}_0 + \mathfrak{g}_1$ is a finite-dimensional restricted Lie superalgebra over an algebraically closed field \mathbf{k} of characteristic $p > 2$. In this article, we propose a conjecture for maximal dimensions of irreducible modules over the universal enveloping algebra $U(\mathfrak{g})$ of \mathfrak{g} , as a super generalization of the celebrated first Kac–Weisfeiler conjecture. It is demonstrated that the conjecture holds for all basic classical Lie superalgebras and all completely solvable restricted Lie superalgebras. In this process, we investigate irreducible representations of solvable Lie superalgebras.

1 Lie superalgebras in characteristic p

Since the works [1–3, 6, 13, 21] etc. on irreducible representations of algebraic supergroups in odd characteristic, especially Wang–Zhao’s work [29] focusing on irreducible representations of basic classical Lie superalgebras, the study of irreducible representations of finite-dimensional restricted Lie superalgebras in odd characteristic has found big progress. For instance, see [16, 30, 33, 39–43] etc. for determination of irreducible modules of classical Lie superalgebras; see [14, 22–24, 28, 32, 34–37] etc. for determination of irreducible modules of Cartan-type Lie superalgebras; and see [16, 23, 30, 34, 39, 40, 42] etc. for dimensions or character formulas of irreducible modules. Nevertheless, their irreducible modules are not well-understood. The purpose of this article is to propose a formulation of maximal dimensions of their irreducible modules. In particular, we thoroughly investigate irreducible representations of finite-dimensional solvable Lie superalgebras.

Throughout the paper, the notions of vector spaces (resp. modules and subalgebras) mean vector superspaces (resp. super-modules and super-subalgebras). For simplicity, we will often omit the adjunct word “super.” All vector spaces are defined over \mathbf{k} which is an algebraically closed field of characteristic $p > 2$. For superspace $V = V_0 + V_1$, we will mention the super-dimension of V which means

Received by the editors May 10, 2023; revised December 5, 2023; accepted December 6, 2023.
 Published online on Cambridge Core December 11, 2023.

This work is partially supported by the National Natural Science Foundation of China (Grant Nos. 12071136, 11771279, and 12271345), supported in part by Science and Technology Commission of Shanghai Municipality (Grant No. 22DZ2229014).

AMS subject classification: 17B50, 17B20, 17B30.

Keywords: Modular Lie superalgebras, maximal dimensions of irreducible modules, the first Kac–Weisfeiler conjecture.



$\dim V = (\dim V_0 | \dim V_1)$, in the meanwhile, we mention the dimension of V which means $\dim_{\mathbf{k}} V := \dim V_0 + \dim V_1$. As usual, we denote by V^* the linear dual space of V . Throughout the paper, all Lie (super)algebras are finite-dimensional unless other statements.

1.1 Restricted Lie superalgebras

A Lie superalgebra $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$ is called a restricted one if \mathfrak{g}_0 is a restricted Lie algebra and \mathfrak{g}_1 is a restricted module of \mathfrak{g}_0 , more precisely, there exists a p -mapping $[p] : \mathfrak{g}_0 \rightarrow \mathfrak{g}_0$ satisfying:

- (a) $(kx)^{[p]} = k^p x^{[p]}$ for all $k \in \mathbf{k}$ and $x \in \mathfrak{g}_0$,
- (b) $[x^{[p]}, y] = (\text{ad} x)^p(y)$ for all $x \in \mathfrak{g}_0$ and $y \in \mathfrak{g}$,
- (c) $(x + y)^{[p]} = x^{[p]} + y^{[p]} + \sum_{i=1}^{p-1} s_i(x, y)$ for all $x, y \in \mathfrak{g}_0$, where $(\text{ad}(x \otimes t + y \otimes 1))^{p-1}(x \otimes 1) = \sum_{i=1}^{p-1} s_i(x, y) \otimes t^{i-1} \in \mathfrak{g}_0 \otimes_{\mathbf{k}} \mathbf{k}[t]$. Here, $\mathbf{k}[t]$ denotes the polynomial ring over \mathbf{k} with indeterminant t .

With emphasis on the p -mapping $[p]$, we sometimes denote the restricted Lie algebra \mathfrak{g}_0 by $(\mathfrak{g}_0, [p])$. One can refer to [10, Section V.7] or [26, Chapter 2] for more details on restricted Lie algebras and restricted modules.

Denote by $\mathcal{Z}(\mathfrak{g})$, the center of $U(\mathfrak{g})$, i.e., $\mathcal{Z}(\mathfrak{g}) := \{u \in U(\mathfrak{g}) \mid \text{ad} x(u) = 0 \ \forall x \in \mathfrak{g}\}$. For a restricted Lie superalgebra \mathfrak{g} , the p -center \mathcal{Z}_0 of $U(\mathfrak{g}_0)$ which is defined to be the subalgebra generated by $\{x^p - x^{[p]} \mid x \in \mathfrak{g}_0\}$, lies in \mathcal{Z} . Fix a basis $\{x_1, \dots, x_s\}$ of \mathfrak{g}_0 and a basis $\{y_1, \dots, y_t\}$ of \mathfrak{g}_1 . Set $\xi_i = x_i^p - x_i^{[p]}$, $i = 1, \dots, s$. The p -center \mathcal{Z}_0 is a polynomial ring $\mathbf{k}[\xi_1, \dots, \xi_s]$ generated by ξ_1, \dots, ξ_s (see, for example, [29, Section 2.3]).

By the PBW theorem, one easily knows that the enveloping superalgebra $U(\mathfrak{g})$ is a free module over \mathcal{Z}_0 with basis

$$x_1^{a_1} \dots x_s^{a_s} y_1^{b_1} \dots y_t^{b_t}, 0 \leq a_i \leq p-1, b_j \in \{0, 1\} \text{ for } i = 1, \dots, s, j = 1, \dots, t$$

(see, for example, [29, Section 2.3]).

1.2 Reduced enveloping algebras of restricted Lie superalgebras

Suppose V is an irreducible $U(\mathfrak{g})$ -module. By the above argument, for any $x \in \mathfrak{g}_0$, $x^p - x^{[p]}$ lies in the center of $U(\mathfrak{g})$. By definition, $x^p - x^{[p]}$ acts on V as an even linear transformation for $x \in \mathfrak{g}_0$. Schur's lemma entails that each $x^p - x^{[p]}$ for $x \in \mathfrak{g}_0$ acts on V by scalar $\chi(x)^p$ for some $\chi \in \mathfrak{g}_0^*$. Such χ is called the p -character of V . Suppose $\chi \in \mathfrak{g}_0^*$ is given, which is naturally regarded in \mathfrak{g}^* by trivial extension. Denote by I_χ , the ideal of $U(\mathfrak{g})$ generated by the even central elements $x^p - x^{[p]} - \chi(x)^p$ with x running over \mathfrak{g}_0 . More generally, we can say that a $U(\mathfrak{g})$ -module M is a χ -reduced module for any given $\chi \in \mathfrak{g}_0^*$ if for any $x \in \mathfrak{g}_0$, $x^p - x^{[p]}$ acts by the scalar $\chi(x)^p$. All χ -reduced modules for any given $\chi \in \mathfrak{g}_0^*$ constitute a full subcategory of the $U(\mathfrak{g})$ -module category. The quotient algebra $U_\chi(\mathfrak{g}) := U(\mathfrak{g})/I_\chi$ is called the reduced enveloping superalgebra of p -character χ . Then the χ -reduced module category of \mathfrak{g} coincides with the $U_\chi(\mathfrak{g})$ -module category. If \mathfrak{h} is a restricted Lie subalgebra of \mathfrak{g} , we often make use of $\chi|_{\mathfrak{h}_0}$ for $\chi \in \mathfrak{g}_0^*$ when we consider the $U_\chi(\mathfrak{g})$ -module category and

its objects induced from \mathfrak{h} -modules. By abuse of notations, we will simply write $\chi|_{\mathfrak{h}_0}$ as χ .

By the PBW theorem, the superalgebra $U_\chi(\mathfrak{g})$ has a basis

$$x_1^{a_1} \dots x_s^{a_s} y_1^{b_1} \dots y_t^{b_t}, 0 \leq a_i \leq p-1; b_j \in \{0, 1\} \text{ for } i = 1, \dots, s; j = 1, \dots, t,$$

and $\dim U_\chi(\mathfrak{g}) = p^{\dim \mathfrak{g}_0} 2^{\dim \mathfrak{g}_1}$.

The reduced enveloping algebra corresponding to $\chi = 0$ is $U_0(\mathfrak{g})$. We call it the restricted enveloping algebra of \mathfrak{g} . The modules of $U_0(\mathfrak{g})$ are called restricted modules of \mathfrak{g} . The following observation is clear.

Lemma 1.1 *The dimensions of irreducible modules of a restricted Lie algebra \mathfrak{g} are not greater than $p^{\dim \mathfrak{g}_0} 2^{\dim \mathfrak{g}_1}$.*

1.3 Minimal p -envelopes of finite-dimensional Lie superalgebras

Any finite-dimensional Lie superalgebra can be embedded in a finite-dimensional restricted Lie superalgebra (see the appendix section). Let $\mathfrak{g} = \mathfrak{g}_0 + \mathfrak{g}_1$ be any given Lie superalgebra. There is a minimal finite-dimensional restricted Lie superalgebra \mathfrak{g}_p such that $\mathfrak{g}_p = (\mathfrak{g}_0)_p + \mathfrak{g}_1$ is a p -envelope of \mathfrak{g} , and $(\mathfrak{g}_0)_p$ a p -envelope of \mathfrak{g}_0 (see Lemma A.3 in the appendix section). Then, one can still show that dimensions of all irreducible modules of \mathfrak{g} has unified upper-bound, by considering its minimal p -envelope. There is a natural question.

Question 1.2 What is the maximal dimension for irreducible modules over \mathfrak{g} ?

With aim at the above question, the purpose of the present paper is to formulate the maximal irreducible dimensions for finite-dimensional restricted Lie superalgebras over \mathbf{k} , as a conjecture (see Conjecture 2.3). This conjecture is regarded a super version of the plausible first Kac–Weisfeiler conjecture (see Remark 2.4(2), or [12, 31]). The progress of the work on the first Kac–Weisfeiler conjecture can be learnt from [15, 18].

The main body of the text is devoted to the verification of the super first Kac–Weisfeiler conjecture in the case of basic classical Lie superalgebras and complete solvable Lie superalgebras.

2 Maximal dimensions of irreducible modules for a finite-dimensional restricted Lie superalgebra

Keep the notations and assumption as above. In particular, $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$ is a finite-dimensional restricted Lie superalgebra over \mathbf{k} . For any given $\chi \in \mathfrak{g}_0^*$, consider the bilinear form B_χ on \mathfrak{g} with regarding $\chi \in \mathfrak{g}^*$ by trivial extension

$$B_\chi : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbf{k}, (X, Y) \mapsto \chi([X, Y]).$$

Set $\ker(B_\chi) = \{X \in \mathfrak{g} \mid B_\chi(X, \mathfrak{g}) = 0\}$. Generally, \mathfrak{g}^* can be regarded a \mathfrak{g} -module via defining for $X \in \mathfrak{g}_{|X|}, f \in \mathfrak{g}_{|f|}^*, X.f : \mathfrak{g} \rightarrow \mathbf{k}$ with $(X.f)(Y) = -(-1)^{|X||f|} f([X, Y])$, $\forall Y \in \mathfrak{g}$. Here, $|X|$ and $|f|$ denote the parities of the \mathbb{Z}_2 -homogeneous element $X \in \mathfrak{g}$

and $f \in \mathfrak{g}^*$, respectively. So we can define the centralizer of χ in \mathfrak{g} , which is denoted by \mathfrak{z}^χ . By definition,

$$\mathfrak{z}^\chi = \{X \in \mathfrak{g} \mid X \cdot \chi = 0, \text{ equivalently, } \chi([X, \mathfrak{g}]) = 0\}.$$

Then this \mathfrak{z}^χ is exactly equal to $\ker(B_\chi)$. Furthermore, with B_χ , we may define bilinear forms on the spaces $\tilde{\mathfrak{g}} := \mathfrak{g}/\mathfrak{z}^\chi$, $\tilde{\mathfrak{g}}_0 := \mathfrak{g}_0/\mathfrak{z}_0^\chi$ and $\tilde{\mathfrak{g}}_1 := \mathfrak{g}_1/\mathfrak{z}_1^\chi$, respectively. By abuse of notations, those bilinear forms are still denoted by B_χ .

In the following arguments, we need some conventions and notations. Let $\lceil a \rceil$ denote the greatest integer lower bound of a for a real number $a \in \mathbb{R}$, and $\lfloor a \rfloor$ denote the least integer upper bound of a .

Lemma 2.1 *The following statements hold.*

- (1) *The centralizer $\mathfrak{z}^\chi = \mathfrak{z}_0^\chi + \mathfrak{z}_1^\chi$ is a restricted subalgebra of \mathfrak{g} if \mathfrak{g} itself is a restricted Lie superalgebra.*
- (2) *B_χ is a nondegenerate skew-symmetric bilinear form on $\tilde{\mathfrak{g}}_0$, and a nondegenerate skew-symmetric bilinear form on $\tilde{\mathfrak{g}}_1$. Consequently, $\dim(\mathfrak{g}_0 - \mathfrak{z}_0^\chi)$ is even.*
- (3) *Any maximal isotropic space in \mathfrak{g}_0 with respect to B_χ has dimension $\frac{\dim \mathfrak{g}_0 + \dim \mathfrak{z}_0^\chi}{2}$.*
- (4) *Any maximal isotropic space in \mathfrak{g}_1 with respect to B_χ has dimension $\frac{\dim \mathfrak{g}_1 + \dim \mathfrak{z}_1^\chi}{2}$ if $\dim \mathfrak{g}_1 - \dim \mathfrak{z}_1^\chi$ is even, and has dimension $\frac{\dim \mathfrak{g}_1 + \dim \mathfrak{z}_1^\chi - 1}{2}$ if $\dim \mathfrak{g}_1 - \dim \mathfrak{z}_1^\chi$ is odd.*

Proof The parts (1) and (2) directly follows from the definition. As to (3), we first note that \mathfrak{z}^χ is an isotropic subspace of \mathfrak{g} with respect to B_χ . From the part (2), it follows that a maximal isotropic subspace \tilde{V} of $\tilde{\mathfrak{g}}_0$ has dimension $\frac{\dim \mathfrak{g}_0 - \dim \mathfrak{z}_0^\chi}{2}$. So naturally, the preimage space of \tilde{V} in V which contains \mathfrak{z}^χ is a maximal isotropic subspace of \mathfrak{g}_0 . This maximal isotropic subspace has dimension $\frac{\dim \mathfrak{g}_0 + \dim \mathfrak{z}_0^\chi}{2}$.

As to the part (4), from the part (2), again it follows that a maximal isotropic subspace \tilde{W} of $\tilde{\mathfrak{g}}_1$ has dimension $\lceil \frac{\dim \mathfrak{g}_1 - \dim \mathfrak{z}_1^\chi}{2} \rceil$. By the same reason, the preimage space of \tilde{W} of \mathfrak{g}_1 which contains \mathfrak{z}^χ is a maximal isotropic subspace of \mathfrak{g}_1 . Consequently, this maximal isotropic subspace has dimension $\lceil \frac{\dim \mathfrak{g}_1 + \dim \mathfrak{z}_1^\chi}{2} \rceil$.

The proof is completed. ■

Remark 2.2 With the notations $\lceil a \rceil$ and $\lfloor a \rfloor$ for $a \in \mathbb{R}$, Lemma 2.1(4) becomes that the maximal isotropic space with respect to B_χ in \mathfrak{g}_1 has dimension $\lceil \frac{\dim \mathfrak{g}_1 + \dim \mathfrak{z}_1^\chi}{2} \rceil$. Set

$$d(\mathfrak{g}, \chi) = \left(\frac{\dim \mathfrak{g}_0 + \dim \mathfrak{z}_0^\chi}{2} \parallel \frac{\dim \mathfrak{g}_1 + \dim \mathfrak{z}_1^\chi}{2} \right).$$

This $d(\mathfrak{g}, \chi)$ is the maximal super-dimension of the isotropy subspaces of \mathfrak{g} with respect to B_χ . Set $i(\mathfrak{g}, \chi) = \underline{\dim} \mathfrak{g} - d(\mathfrak{g}, \chi)$. Then

$$i(\mathfrak{g}, \chi) = \left(\frac{\dim \mathfrak{g}_0 - \dim \mathfrak{z}_0^\chi}{2} \parallel \frac{\dim \mathfrak{g}_1 - \dim \mathfrak{z}_1^\chi}{2} \right).$$

2.1 The set $D(\mathfrak{g}, \chi)$ of degraded subalgebras associated with χ

We regard $\chi \in \mathfrak{g}_0^*$ as a linear function on \mathfrak{g}^* by trivial extension. Associated with χ , we say that a subalgebra \mathfrak{h} is degraded if $\dim \mathfrak{h} = d(\mathfrak{g}, \chi)$ and $\chi(\mathfrak{h}^{(1)}) = 0$. Here and further, $L^{(1)}$ for a Lie (super)algebra L denotes the derived subalgebra of L , i.e., $L^{(1)} = [L, L]$. Obviously, such subalgebras contain \mathfrak{z}^χ if they exist. In this case, they are further restricted subalgebras whenever \mathfrak{g} is a restricted Lie superalgebra.

Denote by $D(\mathfrak{g}, \chi)$, the set of all degraded subalgebras \mathfrak{h} of \mathfrak{g} .

For the simplicity of arguments, we say that a pair of nonnegative integers $(a|b)$ is a super-datum. Call a and b its even entry and odd entry, respectively. For $\chi \in \mathfrak{g}_0^* \subset \mathfrak{g}^*$, we set

$$\begin{aligned} b_0^\chi &= \dim \mathfrak{g}_0 - \dim \mathfrak{z}_0^\chi, \\ b_1^\chi &= \dim \mathfrak{g}_1 - \dim \mathfrak{z}_1^\chi. \end{aligned}$$

Correspondingly, $i(\mathfrak{g}, \chi) = (\frac{b_0^\chi}{2} \parallel \lfloor \frac{b_1^\chi}{2} \rfloor)$. Also set

$$\mathcal{M}(\mathfrak{g}) = \max_{\chi \in \mathfrak{g}_0^*} p^{\frac{b_0^\chi}{2}} 2^{\lfloor \frac{b_1^\chi}{2} \rfloor}.$$

Conjecture 2.3 *Let \mathfrak{g} be a finite-dimensional restricted Lie superalgebra over \mathbf{k} . The maximal dimension of irreducible \mathfrak{g} -modules is $\mathcal{M}(\mathfrak{g})$.*

Remark 2.4 (1) Clearly, by definition, $\mathcal{M}(\mathfrak{g})$ can be expressed as $p^{\frac{b_0}{2}} 2^{\lfloor \frac{b_1}{2} \rfloor}$ for some nonnegative integers b_0 and b_1 . In general, such b_0 and b_1 are not necessarily unique. However, we will see that in many cases, $b_0 = \max_{\chi \in \mathfrak{g}_0^*} b_0^\chi$ and $b_1 = \max_{\chi \in \mathfrak{g}_0^*} b_1^\chi$, which are unique.

(2) The formulation in the above conjecture becomes the first Kac–Weisfeiler conjecture when $\mathfrak{g}_1 = 0$, i.e., a finite-dimensional restricted Lie algebra \mathfrak{g}_0 is regarded a restricted Lie superalgebra with the odd part being zero.

(3) This conjecture is a super version of the first Kac–Weisfeiler conjecture on irreducible modules of restricted Lie algebras (see [12]).¹ For the latter, the study has been in a great progress, but the question is still open (see [15, 18]). There are remarkable works (see [12, 17, 29]) concerning another (the second) Kac–Weisfeiler conjecture on irreducible modules of Lie algebras of reductive groups in prime characteristic and its super version. Some related progress can be found in [9, 30, 39, 40].

3 Irreducible modules of basic classical Lie superalgebras

In this section, we suppose \mathfrak{g} is a basic classical Lie superalgebra over \mathbf{k} . Then $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$ with even part being a reductive Lie algebra. As to classical Lie superalgebras of type P and Q , Conjecture 2.3 was very recently confirmed by taking quite different and nontrivial arguments (see [19]).

¹There is some counterexample against the first Kac–Weisfeiler conjecture for nonrestricted Lie algebras (see [27]).

3.1 Basic classical Lie superalgebras

We list basic classical Lie superalgebras and their even parts over \mathbf{k} with the restriction on p (see, for example, [11, 29]). The restriction on p could be relaxed, but we always assume this restriction on p in this section). The most important feature is that each basic classical Lie superalgebra listed below admits a nondegenerate even supersymmetric bilinear form.

3.1.1

Basic classical Lie superalgebra \mathfrak{g}	\mathfrak{g}_0	Characteristic of \mathbf{k}
$\mathfrak{gl}(m n)$	$\mathfrak{gl}(m) \oplus \mathfrak{gl}(n)$	$p > 2$
$\mathfrak{sl}(m n)$	$\mathfrak{sl}(m) \oplus \mathfrak{sl}(n) \oplus \mathbf{k}$	$p > 2, p \nmid (m - n)$
$\mathfrak{osp}(m n)$	$\mathfrak{so}(m) \oplus \mathfrak{sp}(n)$	$p > 2$
$F(4)$	$\mathfrak{sl}(2) \oplus \mathfrak{so}(7)$	$p > 15$
$G(3)$	$\mathfrak{sl}(2) \oplus G_2$	$p > 15$
$D(2, 1, \alpha)$	$\mathfrak{sl}(2) \oplus \mathfrak{sl}(2) \oplus \mathfrak{sl}(2)$	$p > 3$

For Lie superalgebra \mathfrak{g} in the list, there is an algebraic supergroup G with $\text{Lie}(G) = \mathfrak{g}$ satisfying:

- (1) G has a purely-even subgroup scheme G_{ev} which is an ordinary connected reductive algebraic group with $\text{Lie}(G_{\text{ev}}) = \mathfrak{g}_0$.
- (2) There is a well-defined action of G_{ev} on \mathfrak{g} , giving rise to the adjoint action of \mathfrak{g}_0 .

The above algebraic supergroup are usually called basic classical supergroups, which can be constructed as Chevalley supergroups (see [7, 8]). Generally, for an algebraic supergroup G with $\mathfrak{g} = \text{Lie}(G)$, \mathfrak{g} does not determine G . Instead, the theory of supergroups shows that the pair $(G_{\text{ev}}, \mathfrak{g})$ determines G (see, for example, [4, Chapter 7]). The pair $(G_{\text{ev}}, \mathfrak{g})$ is called a super Harish–Chandra pair (*ibid.*). More precisely, the category of algebraic supergroups is equivalent to the category of super Harish–Chandra pairs (*ibid.*).

One easily knows that $\mathfrak{g} = \text{Lie}(G)$ for an algebraic supergroup G is a restricted Lie superalgebra (cf. [21, Lemma 2.2] or [25]).

Let \mathfrak{g} be a given basic classical Lie superalgebra. We fix a Cartan subalgebra \mathfrak{h} in \mathfrak{g}_0 . Denote by Φ , the root system associate with \mathfrak{h} . Then $\Phi = \Phi_0 \cup \Phi_1$, where Φ_0 stands for the set of even roots, and Φ_1 for the set of odd roots. Fix a triangular decomposition $\mathfrak{g} = \mathfrak{n}^- \oplus \mathfrak{h} \oplus \mathfrak{n}^+$ which is equivalent to say, fix a positive root system Φ^+ , or to say fix a simple root system Δ . Here, \mathfrak{n}^\pm stand the Lie subalgebras of positive and negative root vectors, respectively. Furthermore, $\Phi^- = -\Phi^+$, and $\Phi^\pm = \Phi_1^\pm \cup \Phi_0^\pm$. Moreover, without loss of generality we can assume $\chi(\mathfrak{n}_0^+) = 0$ for any $\chi \in \mathfrak{g}_0^*$, up to G_{ev} -conjugation. Let $\mathfrak{b} = \mathfrak{h} \oplus \mathfrak{n}^+$. Then any $\lambda \in \mathfrak{h}^*$ defines a one-dimensional $U_\chi(\mathfrak{h})$ -module \mathbf{k}_λ as long as λ satisfies $\lambda(H)^p - \lambda(H^{[p]}) = \chi(H)^p$ for all $H \in \mathfrak{h}$. Set

$$\Lambda(\chi) = \{\lambda \in \mathfrak{h}^* \mid \lambda(H)^p - \lambda(H^{[p]}) = \chi(H)^p \quad \forall H \in \mathfrak{h}\},$$

which clearly contains $p^{\dim \mathfrak{h}}$ elements.

The one-dimensional space \mathbf{k}_λ can be regarded a $U_\chi(\mathfrak{b})$ -module with trivial \mathfrak{n}^+ -action, because of $\chi(\mathfrak{n}_0^+) = 0$. Then we define an induced module (called a baby Verma module)

$$Z_\chi(\lambda) := U_\chi(\mathfrak{g}) \otimes_{U_\chi(\mathfrak{b})} \mathbf{k}_\lambda.$$

Obviously, $\dim Z_\chi(\lambda) = p^{\dim \mathfrak{n}_0^-} 2^{\dim \mathfrak{n}_1^-}$. The following fact is clear.

Lemma 3.1 *Any irreducible $U_\chi(\mathfrak{g})$ -module has dimension not bigger than $\dim Z_\chi(\lambda)$.*

The proof is standard. We give an account on it. By the above arguments, we only need to consider the case $\chi(\mathfrak{n}^+) = 0$. In this case, for any irreducible $U_\chi(\mathfrak{g})$ -module V , \mathfrak{n}^+ acts nilpotently on V . Hence, V admits one-dimensional $U_\chi(\mathfrak{b})$ -module \mathbf{k}_λ with \mathfrak{n}^+ -trivial action, and $U_\chi(\mathfrak{h})$ -action by some function $\lambda \in \Lambda(\chi)$. So V coincides with $U_\chi(\mathfrak{n}^-)\mathbf{k}_\lambda$ which is isomorphic to an irreducible quotient of $Z_\chi(\lambda)$. The lemma follows.

Recall that $\mathfrak{g}_0 = \text{Lie}(G_{\text{ev}})$, and any element $X \in \mathfrak{g}_0$ admits Jordan–Chevalley decomposition $X = X_s + X_n$ with X_s being semisimple and X_n nilpotent. Note that for a basic classical Lie algebra \mathfrak{g} listed in 3.1.1, there is a nondegenerate G_0 -invariant symmetric bilinear form (\cdot, \cdot) on \mathfrak{g}_0 . Hence, there is a G -equivariant isomorphism between \mathfrak{g}_0 and \mathfrak{g}_0^* . Consequently, for any $\chi \in \mathfrak{g}_0^*$, there exists a unique X such that $\chi = (X, -)$, which leads to the Jordan–Chevalley decomposition $\chi = \chi_s + \chi_n$ with $\chi_s = (X_s, -)$ and $\chi_n = (X_n, -)$. Furthermore, there exists $g \in G_{\text{ev}}$ such that $(g \cdot \chi_s)(\mathfrak{n}_0^\pm) = 0$, and $(g \cdot \chi_n)(\mathfrak{b}_0) = 0$, where the action of g means the coadjoint action. We say that χ is semisimple if $\chi = \chi_s$, and that χ is nilpotent if $\chi = \chi_n$. For simplicity, we always assume that $\chi = \chi_s + \chi_n$ with $\chi_s(\mathfrak{n}_0^\pm) = 0$, and $\chi_n(\mathfrak{b}_0) = 0$ in the following because the coadjoint action gives rise to an isomorphism between $U_\chi(\mathfrak{g})$ and $U_{g \cdot \chi}(\mathfrak{g})$.

For a semisimple p -character $\chi \in \mathfrak{g}_0^*$, we say that χ is regular semisimple if $\chi(H_\alpha)$ are nonzero for all α from Φ , where H_α is the Cartan toral element corresponding to α . By Lei Zhao's result, we have the following theorem.

Theorem 3.2 [42, Theorems 4.6 and 4.7] *Suppose χ is regular semisimple, and $\lambda \in \Lambda_\chi$. Then $Z_\chi(\lambda)$ is irreducible.*

For a regular semisimple p -character $\chi \in \mathfrak{g}_0^*$, from the definition, it follows that $3^\chi = \mathfrak{h}$. Hence, $(\lfloor \frac{b_0^\chi}{2} \rfloor \mid \lfloor \frac{b_1^\chi}{2} \rfloor) = (\dim \mathfrak{n}_0^- \mid \dim \mathfrak{n}_1^-)$. Correspondingly, $\dim Z_\chi(\lambda) = p^{\lfloor \frac{b_0^\chi}{2} \rfloor} 2^{\lfloor \frac{b_1^\chi}{2} \rfloor}$, which coincides with $p^{\frac{b_0}{2}} 2^{\lfloor \frac{b_1}{2} \rfloor}$. On the other hand, Wang–Zhao's theorem concerning Kac–Weisfeiler property (see [29, Theorem 4.3]) says that for any $\chi \in \mathfrak{g}_0^*$ and any irreducible $U_\chi(\mathfrak{g})$ -module V , $\dim V$ is divisible by $p^{\frac{b_0^\chi}{2}} 2^{\lfloor \frac{b_1^\chi}{2} \rfloor}$. Hence, we have

$$\dim V \leq \dim Z_\chi(\lambda) = p^{\frac{b_0^\chi}{2}} 2^{\lfloor \frac{b_1^\chi}{2} \rfloor}.$$

Combining the above with Lemma 3.1 and Theorem 3.2, we finally have the following result.

Corollary 3.3 Suppose \mathfrak{g} is a basic classical Lie superalgebra over \mathbf{k} . Then the maximal dimension of irreducible $U(\mathfrak{g})$ -modules is exactly $\mathcal{M}(\mathfrak{g})$. Moreover, $\mathcal{M}(\mathfrak{g})$ can be precisely described as $p^{\frac{b_0}{2}} 2^{\lfloor \frac{b_1}{2} \rfloor}$ for $b_0 = \max\{b_0^X | \chi \in \mathfrak{g}_0^*\}$ and $b_1 = \max\{b_1^X | \chi \in \mathfrak{g}_0^*\}$.

Consequently, the statement of Conjecture 2.3 is true for basic classical Lie superalgebras.

4 Irreducible modules of solvable Lie superalgebras

In the next two sections, we will study irreducible representations of finite-dimensional solvable Lie superalgebras over \mathbf{k} , by exploiting the arguments for ordinary solvable Lie algebras (see [20, 31] or [26, Chapter 5]). Keep the notations and assumptions as before. In particular, for a Lie (super)algebra L , we denote by $L^{(1)}$ the derived subalgebra of L , i.e., $L^{(1)} = [L, L]$.

4.1 Basic properties on solvable Lie superalgebras

The following results are important for the later arguments.

Lemma 4.1 Let $\mathfrak{g} = \mathfrak{g}_0 + \mathfrak{g}_1$ is a finite-dimensional Lie superalgebra over \mathbf{k} .

- (1) Suppose V is an irreducible module of \mathfrak{g} . If all elements of $[\mathfrak{g}, \mathfrak{g}]$ act nilpotently on V , then V is one-dimensional.
- (2) Suppose additionally, \mathfrak{g} is solvable and non-abelian, then the center $C(\mathfrak{g}) := \{X \in \mathfrak{g} \mid \text{ad}(X)(\mathfrak{g}) = 0\}$ does not contain all abelian ideals of \mathfrak{g} .

Proof (1) It is an obvious fact.

(2) We prove this statement by reductio ad absurdum. Suppose $C(\mathfrak{g})$ contains all abelian ideals. We intend to deduce a contradiction.

Note that $C(\mathfrak{g})$ is an ideal of \mathfrak{g} . Consider the natural surjective homomorphism of Lie superalgebras $\bar{\cdot}: \mathfrak{g} \rightarrow \mathfrak{g}/C(\mathfrak{g})$. Of course, $\bar{\mathfrak{g}} = \mathfrak{g}/C(\mathfrak{g})$. By assumption, $\bar{\mathfrak{g}} \neq 0$ which is still a solvable Lie superalgebra. Take a minimal ideal $I = I_0 + I_1 \triangleleft \mathfrak{g}$ containing $C(\mathfrak{g})$ properly. The solvableness of \mathfrak{g} yields that I properly contains $I^{(1)}$. So $I^{(1)} \subset C(\mathfrak{g})$. Under the assumption of $C(L)$ containing all abelian ideals, it follows that $I^{(1)} \neq 0$. Consequently, $I_0 \neq 0$. Moreover, there exists a linear function $\lambda \in C(\mathfrak{g})^*$ such that $\lambda|_{I^{(1)}} \neq 0$. Note that $\bar{I} \neq 0$ and it becomes an irreducible \mathfrak{g} -modules. This irreducible representation of \mathfrak{g} on \bar{I} is denoted by (ρ, \bar{I}) . The linear function λ gives rise to a bilinear form Λ on \bar{I} by defining $\Lambda: \bar{I} \times \bar{I} \rightarrow \mathbf{k}$ via $\Lambda(\bar{v}_1, \bar{v}_2) = \lambda([\bar{v}_1, \bar{v}_2])$ for any $v_1, v_2 \in I$. Then it is easily checked that Λ satisfies \mathfrak{g}_0 -invariant property in the sense that

$$(4.1) \quad \Lambda(X.\bar{v}_1, \bar{v}_2) + \Lambda(\bar{v}_1, X.\bar{v}_2) = 0 \text{ for } X \in \mathfrak{g}_0, \bar{v}_i \in \bar{I}, (i = 1, 2),$$

where $X.\bar{v}_i = \overline{[X, v_i]}$. As $\lambda \neq 0$, we can take \bar{v}, \bar{w} such that $\Lambda(\bar{v}, \bar{w}) \neq 0$.

We claim $\rho(\mathfrak{g}) = 0$, which means \mathfrak{g} acts trivially on \bar{I} . Otherwise, if $\rho(\mathfrak{g})$ is nonzero, then there exists a nonzero abelian ideal J because $\rho(\mathfrak{g})$ is solvable. For any given nonzero $Z \in J_0$, consider the action of Z^p on \bar{I} . Note that for any $Y \in \rho(\mathfrak{g})$, $[Z^p, Y] =$

$\text{ad}(Z)^p Y \in J^{(1)} = 0$. By Schur's lemma, there exists $\alpha(Z) \in \mathbf{k}$ such that $Z^p \cdot \bar{v} = \alpha(Z) \bar{v}$ for any $Z^p \bar{v} = \alpha(Z) \bar{v}$. Comparing with (4.1), we have

$$\alpha(Z) \Lambda(\bar{v}, \bar{w}) = \alpha(Z) \Lambda(Z^p \cdot \bar{v}, \bar{w}) = \Lambda(\bar{v}, Z^p \bar{w}) = -\alpha(Z) \Lambda(\bar{v}, \bar{w}).$$

Hence, $\alpha(Z) = 0$ for all $Z \in J_0$. Thus, J_0 acts nilpotently on \bar{I} , naturally so does J_1 . But \bar{I} is irreducible, which implies that J acts trivially on \bar{I} . This is to say $J = 0$, a contraction.

By the arguments, it is already deduced that $\rho(\mathfrak{g}) = 0$ and the irreducible \mathfrak{g} -module \bar{I} must be one-dimensional. This means, $I = \mathbf{k}v + C(\mathfrak{g})$ for $v \in \mathfrak{g}$, which leads to a contradiction $I^{(1)} = 0$. The proof is completed. ■

4.2 Induced modules for solvable restricted Lie superalgebras

Now, we begin to investigate irreducible modules of solvable restricted Lie superalgebras. In the following two subsections, we assume that $\mathfrak{g} = \mathfrak{g}_0 + \mathfrak{g}_1$ is a finite-dimensional solvable restricted Lie superalgebra, and $\chi \in \mathfrak{g}_0^*$ which will be often regarded a linear function on \mathfrak{g} by trivial extension. At first, Lemma 4.1(1) implies the following conclusion.

Lemma 4.2 *Let $\mathfrak{g} = \mathfrak{g}_0 + \mathfrak{g}_1$ be a finite-dimensional solvable restricted Lie superalgebra, and $\chi \in \mathfrak{g}_0^*$. If $\mathfrak{g}^{(1)}$ is nilpotent, and $\chi(\mathfrak{g}^{(1)}) = 0$, then any irreducible $U_\chi(\mathfrak{g})$ -module is one-dimensional.*

Next, let $\rho : \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ be a finite-dimensional representation of $U_\chi(\mathfrak{g})$. Suppose I is an ideal of \mathfrak{g} such that $I^{(1)} \subset \ker \rho$ and $[\mathfrak{g}, I] \not\subset \ker \rho$. So there is an irreducible I -submodule in V which is one-dimensional. This yields that

$$(4.2) \quad V_{I,\chi} := \{v \in V \mid X \cdot v = \chi(X)v \ \forall X \in I\}$$

is nonzero. Consider $I^\chi := \{X \in \mathfrak{g} \mid \chi([X, I]) = 0\}$. It is easily shown that I^χ is a restricted subalgebra of \mathfrak{g} containing I . Clearly, I^χ is still solvable, and $V_{I,\chi}$ becomes a module of $U_\chi(I^\chi)$ from which we will consider an induced module of $U_\chi(\mathfrak{g})$.

We claim that I^χ is a proper subalgebra of \mathfrak{g} . We will show this by reductio ad absurdum. If $\mathfrak{g} = I^\chi$, then $V_{I,\chi}$ is a $U_\chi(\mathfrak{g})$ -submodule and therefore coincides with V , which means $\rho|_{[\mathfrak{g}, I]} = \chi|_{[\mathfrak{g}, I]} \text{id}_V = 0$. Thus $[\mathfrak{g}, I] \subset \ker \rho$, which contradicts the condition of I . So we can take a cobasis $\{e_1, \dots, e_s\}$ of I_0^χ in \mathfrak{g}_0 for $s = \dim \mathfrak{g}_0 - \dim I_0^\chi$, and a cobasis $\{f_1, \dots, f_t\}$ of I_1^χ in \mathfrak{g}_1 for $t = \dim \mathfrak{g}_1 - \dim I_1^\chi$, which means

$$\mathfrak{g}_0 = I_0^\chi \oplus \bigoplus_{i=1}^s \mathbf{k}e_i \quad \text{and} \quad \mathfrak{g}_1 = I_1^\chi \oplus \bigoplus_{i=1}^t \mathbf{k}f_i.$$

Now, we consider the induced module $\mathcal{V} := U_\chi(\mathfrak{g}) \otimes_{U_\chi(I^\chi)} V_{I,\chi}$ which can be expressed as a vector space

$$\mathcal{V} = \sum_{(\alpha, \gamma) \in P^s \times E^t} \mathbf{k}e^\alpha f^\gamma \otimes V_{I,\chi},$$

where $\alpha = (a_1, \dots, a_s) \in \mathbb{Z}_{\geq 0}^s$ and $\gamma = (c_1, \dots, c_t) \in \mathbb{Z}_{\geq 0}^t$ are s -tuple and t -tuple of nonnegative integers, respectively, $\mathbf{e}^\alpha := e_1^{a_1} \dots e_s^{a_s}$, and $\mathbf{f}^\gamma = f_1^{c_1} \dots f_t^{c_t}$ with

$$P := \{0, 1, \dots, p-1\}$$

and

$$E := \{0, 1\}.$$

Set

$$\|(\alpha, \gamma)\| := \sum_i a_i + \sum_j c_j$$

and put

$$\mathcal{V}_{(l)} = \sum_{\substack{(\alpha, \gamma) \in P^s \times E^t \\ \|(\alpha, \gamma)\| \leq l}} \mathbf{k} \mathbf{e}^\alpha \mathbf{f}^\gamma \otimes V_{I, \chi}.$$

Consider $U_0 = \sum_{i=1}^s \mathbf{k} e_i$ and $U_1 = \sum_{j=1}^t \mathbf{k} f_j$. Define a linear map φ from U_0 to I_0^* by sending any given $X \in I$ onto the function $B_\chi(X, -)$ on I . By the definition of I^χ , this φ is nondegenerate. Hence, there is a set $\{Z_1, \dots, Z_s\} \subset I_0$ such that $\psi(e_j)(Z_i) = \delta_{ij}$ for $1 \leq i, j \leq s$. Here and further, δ_{ij} denotes the Kronecker function whose value at (i, j) is zero when $i \neq j$ and 1 when $i = j$. Similarly, consider a linear map ψ from U_1 to I_1^* sending Y onto the function $B_\chi(Y, -)$ on I_1 , and this ψ is nondegenerate too. We can choose $\{T_1, \dots, T_t\} \subset I_1$ such that $\psi(f_j)(T_i) = \delta_{ij}$ for $1 \leq i, j \leq t$.

Lemma 4.3 *Keep the notations and assumptions as above. For $\mathbf{e}^\alpha \mathbf{f}^\gamma \otimes v \in \mathcal{V}_{(l)}$ with $v \in V_{I, \chi}$, the following formula holds:*

$$(4.3) \quad \begin{aligned} (Z_i - \chi(Z_i)) \cdot \mathbf{e}^\alpha \mathbf{f}^\gamma \otimes v &\equiv a_i \mathbf{e}^{\alpha - \varepsilon_i} \mathbf{f}^\gamma \otimes v \pmod{\mathcal{V}_{(l-2)}} \text{ if } \alpha \neq 0; \text{ and} \\ T_j \cdot \mathbf{f}^\gamma \otimes v &\equiv c_j \mathbf{f}^{\gamma - \varepsilon_j} \otimes v \pmod{\mathcal{V}_{(l-2)}} \text{ if } \alpha = 0. \end{aligned}$$

Proof We first generally introduce an order relation \leq on $\mathbb{Z}_{\geq 0}^q$ (the set of q -tuples of nonnegative integers) by defining $\kappa \leq \alpha$ for any given $\kappa' = (k'_1, \dots, k'_q)$, $\alpha' = (a'_1, \dots, a'_q) \in \mathbb{Z}_{\geq 0}^q$ if and only if $k'_i \leq a'_i$ for all i .

Now, we turn to the situation for the lemma. Set $\varepsilon_k := (\delta_{1k}, \dots, \delta_{sk})$, $k = 1, \dots, s$. Recall the following formula for $Z \in I_0$:

$$Z \mathbf{e}^\alpha = \sum_{0 \leq \kappa \leq \alpha} (-1)^{\|\kappa\|} \binom{\alpha}{\kappa} \mathbf{e}^{\alpha - \kappa} \text{ad}(e_s)^{k_s} (\dots (\text{ad}(e_1)^{k_1})) Z,$$

here and further, we set

$$\|\kappa\| := \sum_{i=1}^s k_i$$

for $\kappa = (k_1, \dots, k_s) \in \mathbb{Z}_{\geq 0}^s$, and set

$$\binom{\alpha}{\kappa} := \sum_{i=1}^s \binom{a_i}{k_i},$$

and for $T \in I_1$

$$T\mathbf{f}^\gamma = \sum_{0 \leq \kappa \leq \gamma} \mathbf{f}^{\gamma-\kappa} \text{ad}(f_t)^{k_t} (\dots (\text{ad}(f_1)^{k_1})) T.$$

So in the case when $\alpha \neq 0$, we have

$$(Z_i - \chi(Z_i)) \cdot \mathbf{e}^\alpha \mathbf{f}^\gamma \otimes v = \sum_{0 \leq \kappa \leq \alpha} (-1)^{\|\kappa\|} \binom{\alpha}{\kappa} \mathbf{e}^{\alpha-\kappa} \text{ad}(e_s)^{k_s} (\dots (\text{ad}(e_1)^{k_1})) (Z_i - \chi(Z_i)) \mathbf{f}^\gamma \otimes v.$$

Note that any elements from I_1^χ act trivially on v . So we can write

$$\begin{aligned} (Z_i - \chi(Z_i)) \cdot \mathbf{e}^\alpha \mathbf{f}^\gamma \otimes v &\equiv \mathbf{e}^\alpha \mathbf{f}^\gamma (Z_i - \chi(Z_i)) \otimes v - \sum_{k=1}^s a_k \mathbf{e}^{\alpha-\varepsilon_k} \mathbf{f}^\gamma [e_k, Z_i] \otimes v \pmod{\mathcal{V}_{(l-2)}} \\ &\equiv a_i \mathbf{e}^{\alpha-\varepsilon_i} \mathbf{f}^\gamma \otimes v \pmod{\mathcal{V}_{(l-2)}}. \end{aligned}$$

By the same arguments, we can deal with the case when $\alpha = 0$. We finally have

$$\begin{aligned} T_j \mathbf{f}^\gamma \otimes v &\equiv \mathbf{f}^\gamma T_j \otimes v + \sum_{k=1}^t \mathbf{f}^{\gamma-\varepsilon_k} [f_k, T_j] \otimes v \pmod{\mathcal{V}_{(l-2)}} \\ &\equiv \mathbf{f}^{\gamma-\varepsilon_j} \otimes v \pmod{\mathcal{V}_{(l-2)}}. \end{aligned}$$

The proof is completed. ■

Lemma 4.4 *Keep the above notations and assumptions. In particular, let \mathfrak{g} be a solvable restricted Lie superalgebra. Let $\rho : \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ be an irreducible representation of $U_\chi(\mathfrak{g})$ on V . The following statements hold.*

- (1) *The module $V_{l,\chi}$ defined in (4.2) is an irreducible $U_\chi(I^X)$ -module.*
- (2) *Furthermore, if we set $\mathcal{V} := U_\chi(\mathfrak{g}) \otimes_{U_\chi(I^X)} V_{l,\chi}$, then $V \cong \mathcal{V}$.*

Proof (1) It is clear.

(2) Suppose W is a nonzero submodule of $U_\chi(\mathfrak{g})$ in \mathcal{V} . Set $W_0 := \{w \in V_{l,\chi} \mid 1 \otimes w \in W\}$. Obviously, W_0 is a $U_\chi(I^X)$ -submodule of $V_{l,\chi}$. According to part (1), $V_{l,\chi}$ is an irreducible $U_\chi(I^X)$ -module. Hence $W_0 = V_{l,\chi}$ or $W_0 = 0$. If the former case occurs, then $W = \mathcal{V}$. Hence it suffices to show that W_0 is nonzero.

Recall $\mathcal{V} = \sum_{(\alpha,\gamma) \in P^s \times E^t} \mathbf{ke}^\alpha \mathbf{f}^\gamma \otimes V_{l,\chi}$. Obviously, $W = \bigcup_{l \geq 0} (W \cap \mathcal{V}_{(l)})$. On the other hand, we put $W_{(l)} := \sum_{\substack{(\alpha,\gamma) \in P^s \times E^t \\ \|(\alpha,\gamma)\| \leq l}} \mathbf{ke}^\alpha \mathbf{f}^\gamma \otimes W_0$. We claim that

$$W = \sum_{(\alpha,\gamma) \in P^s \times E^t} \mathbf{ke}^\alpha \mathbf{f}^\gamma \otimes W_0.$$

It yields that $W_0 \neq 0$, which is our purpose. In order to verify the claim, it suffices to show that

$$(*) \quad W_{(l)} = \mathcal{V}_{(l)} \cap W \text{ for all } l \geq 0.$$

By definition, $(*)$ is true for $l = 0$. We now prove that $W \cap \mathcal{V}_{(l)} \subset W_{(l)}$ by induction on l . Let $l \geq 1$ and assume that $W \cap \mathcal{V}_{(l-1)} \subset W_{(l-1)}$. Suppose that $v \in W \cap \mathcal{V}_{(l)}$ is arbitrarily given, we intend to show that $w \in W_{(l)}$. Take a cobasis $\{v_1, \dots, v_q\}$ of W_0 in $V_{l,\chi}$. Without loss of generality, we might as well assume

$$v = \sum_{k=1}^q \sum_{\substack{(\alpha, \gamma) \in P^s \times E^t \\ \|(\alpha, \gamma)\| \leq l}} C_{\alpha, \gamma, k} \mathbf{e}^\alpha \mathbf{f}^\gamma \otimes v_k$$

with all $C_{\alpha, \gamma, k} \in \mathbf{k}$. By (4.3), we have for $i = 1, \dots, s$,

$$(Z_i - \chi(Z_i)) \cdot v \equiv \sum_k \sum_{\|(\alpha, \gamma)\| = l} C_{\alpha, \gamma, k} \mathbf{e}^{\alpha - \varepsilon_i} \mathbf{f}^\gamma \otimes v_k \pmod{\mathcal{V}_{(l-2)}}.$$

Thus $(Z_i - \chi(Z_i)) \cdot v \in W \cap \mathcal{V}_{(l-1)}$. By the inductive hypothesis, it lies in $W_{(l-1)}$. Hence, all $C_{\alpha, \gamma, k} = 0$ as long as $\|(\alpha, \gamma)\| = l$ and $\alpha \neq 0$. In the same way, let us deal with the case when $\|(\alpha, \gamma)\| = l$ and $\alpha = 0$. Multiplication by T_j on v , $j = 1, \dots, t$ yields

$$T_j \cdot v \equiv \sum_k \sum_{\|(\alpha, \gamma)\| = l} C_{0, \gamma, k} \mathbf{f}^{\gamma - \varepsilon_j} \otimes v_k \pmod{\mathcal{V}_{(l-2)}}.$$

For the same reason as the previous case, we have that $C_{0, \gamma, k} = 0$ for all γ with $\|(0, \gamma)\| = l$. Hence v must be zero. In summary, $W \cap \mathcal{V}_{(l)} \subset W_{(l)}$. Finally, we have $W \cap \mathcal{V}_{(l)} = W_{(l)}$. This implies $W = \mathcal{V}$. Consequently, \mathcal{V} is an irreducible $U_\chi(\mathfrak{g})$ -module. On the other hand, there is a natural surjective $U_\chi(\mathfrak{g})$ -homomorphism from \mathcal{V} onto V . The irreducibility of both V and \mathcal{V} implies that the natural surjective homomorphism must be an isomorphism. ■

4.3 Irreducible modules of solvable restricted Lie superalgebras

Furthermore, we have the following result.

Proposition 4.5 *Keep the assumptions and notations as above. Suppose $\rho : \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ is a finite-dimensional irreducible representation of $U_\chi(\mathfrak{g})$. Then there is a subalgebra $\mathfrak{h} = \mathfrak{h}_0 + \mathfrak{h}_1$ such that:*

- (i) $\dim V \geq p^{\dim \mathfrak{g}_0 - \dim \mathfrak{h}_0} 2^{\dim \mathfrak{g}_1 - \dim \mathfrak{h}_1}$, and
- (ii) V contains a one-dimensional \mathfrak{h} -submodule.

Proof Consider $\ker \rho$ (the kernel of ρ) which is an ideal of \mathfrak{g} . If $\mathfrak{g}/\ker \rho$ is abelian, then $[\mathfrak{g}, \mathfrak{g}]$ acts trivially on V . Hence, V is certainly one-dimensional with trivial action of $[\mathfrak{g}_0, \mathfrak{g}_0] + \mathfrak{g}_1$. So the proposition is true in this case while we take \mathfrak{h} to be \mathfrak{g} itself.

In the following, we suppose that $\bar{\mathfrak{g}} := \mathfrak{g}/\ker \rho$ is not abelian. We will prove the proposition by induction on $\dim \mathfrak{g}$ by steps.

(1) Keep in mind the assumption that $\bar{\mathfrak{g}} := \mathfrak{g}/\ker \rho$ is not abelian. So the center $C(\bar{\mathfrak{g}})$ is a proper subalgebra of $\bar{\mathfrak{g}}$. So $C(\bar{\mathfrak{g}})$ does not contain all abelian ideal of \mathfrak{g} , due to Lemma 4.1(2). So, there exists an ideal I of \mathfrak{g} such that $I^{(1)} \subset \ker \rho$ and $[\mathfrak{g}, I] \not\subset \ker \rho$. Note that $I^{(1)}$ acts trivially on V . So $\chi(I^{(1)}) = 0$. Hence, there is an irreducible I -submodule in V which is one-dimensional. Still denote $V_{I, \chi} = \{v \in V \mid X \cdot v = \chi(X)v \ \forall X \in I\}$. Then we have $V_{I, \chi}$ is nonzero. Keep the notation $I^\chi = \{X \in \mathfrak{g} \mid \chi([X, I]) = 0\}$. It is already known that I^χ is a solvable restricted subalgebra of \mathfrak{g} . By the arguments in the first paragraph of Section 4.2, I^χ is a proper subalgebra of \mathfrak{g} . By Lemma 4.4, $V_{I, \chi}$ is an irreducible $U_\chi(I^\chi)$ -module, and

$$V \cong U_\chi(\mathfrak{g}) \otimes_{U_\chi(I^\chi)} V_{I, \chi}.$$

Correspondingly,

$$(4.4) \quad \dim V = p^{\dim \mathfrak{g}_0 - \dim I_0^\chi} 2^{\dim \mathfrak{g}_1 - \dim I_1^\chi} \dim V_{I, \chi}.$$

(2) Note that I^χ already turns out to be a proper solvable restricted subalgebra. By the inductive hypothesis, there exists a solvable restricted subalgebra \mathfrak{h} in I^χ such that the requirements (i) and (ii) are satisfied with respect to the irreducible $U_\chi(I^\chi)$ -module $V_{I, \chi}$. Hence, by (4.4), we finally have $\dim V \geq p^{\dim \mathfrak{g}_0 - \dim \mathfrak{h}_0} 2^{\dim \mathfrak{g}_1 - \dim \mathfrak{h}_1}$. ■

Combining Proposition 4.5 and Lemma 4.4(2), we conclude the following result.

Corollary 4.6 *Let \mathfrak{g} be a solvable restricted Lie superalgebra. Then any irreducible module of \mathfrak{g} must be isomorphic to $U_\chi(\mathfrak{g}) \otimes_{U_\chi(\mathfrak{h})} S$ for some restricted subalgebra \mathfrak{h} with $\chi(\mathfrak{h}^{(1)}) = 0$ and for a one-dimensional $U_\chi(\mathfrak{h})$ -module S . Correspondingly, any irreducible module has dimension $p^m 2^n$ for some $m, n \in \mathbb{N}$.*

4.4 Irreducible modules of general finite-dimensional solvable Lie superalgebras which are not necessarily restricted

In this section, we will extend the above results for solvable restricted Lie subalgebras to any solvable ones. Let $\mathfrak{g} = \mathfrak{g}_0 + \mathfrak{g}_1$ be a given solvable Lie superalgebra. Recall that \mathfrak{g} admits a minimal finite-dimensional p -envelope \mathfrak{g}_p , such that $\mathfrak{g}_p = (\mathfrak{g}_0)_p \oplus \mathfrak{g}_1$ and $(\mathfrak{g}_0)_p$ is a p -envelope of \mathfrak{g}_0 (see Lemma A.3 in the appendix). This \mathfrak{g}_p becomes a solvable restricted Lie superalgebra. As mentioned in appendix, \mathfrak{g} is an ideal of \mathfrak{g}_p . More precisely, $\text{ad}(\mathfrak{g}_0)_p(\mathfrak{g}_0) \subset \mathfrak{g}_0^{(1)}$, and $\text{ad}(\mathfrak{g}_0)_p \mathfrak{g}_1 \subset \mathfrak{g}_1^{(1)}$.

For irreducible \mathfrak{g} -module (V, ρ) , ρ can extend to the one over \mathfrak{g}_p (see, for example, [26, 2.5.3]). Hence, V becomes an irreducible \mathfrak{g}_p -module. Hence, there exists a unique $Y \in (\mathfrak{g}_0)_p^*$, with which the irreducible \mathfrak{g}_p -module V is associated. Set $\chi = Y|_{\mathfrak{g}_0}$. Then the irreducible \mathfrak{g} -module V is associated with a unique linear function $\chi \in \mathfrak{g}_0^*$.

Theorem 4.7 *Let \mathfrak{g} be a solvable Lie superalgebra. Any irreducible module of \mathfrak{g} is associated with some $\chi \in \mathfrak{g}_0^*$, which has dimension $p^{\dim \mathfrak{g}_0/\mathfrak{h}_0} 2^{\dim \mathfrak{g}_1/\mathfrak{h}_1}$, where \mathfrak{h} is a subalgebra with $\chi(\mathfrak{h}^{(1)}) = 0$ and V contains a one-dimensional \mathfrak{h} -module.*

Proof For any given irreducible \mathfrak{g} -module (V, ρ) , as arguments above, V becomes an irreducible \mathfrak{g}_p -module associated with $Y \in (\mathfrak{g}_0)_p^*$, and V is associated with $\chi := Y_{\mathfrak{g}_0} \in \mathfrak{g}_0^*$. By Corollary 4.6, there exists a restricted subalgebra \mathfrak{h} of \mathfrak{g}_p with $Y(\mathfrak{h}^{(1)}) = \chi(\mathfrak{h}^{(1)}) = 0$ such that $V \cong U_\chi(\mathfrak{g}_p) \otimes_{U_\chi(\mathfrak{h})} S$, where $S \subset V$ is a one-dimensional \mathfrak{h} -module. Correspondingly, $\dim V = p^{\dim (\mathfrak{g}_0)_p/\mathfrak{h}_0} 2^{\dim \mathfrak{g}_1/\mathfrak{h}_1}$.

Take $\mathfrak{h} = \mathfrak{h} \cap \mathfrak{g}$. By definition, $\mathfrak{h}_0 = \mathfrak{h}_0 \cap \mathfrak{g}_0$, and $\mathfrak{h}_1 = \mathfrak{h}_1 \cap \mathfrak{g}_1 = \mathfrak{h}_1$. Then $\chi(\mathfrak{h}^{(1)}) = 0$, and \mathfrak{h} has one-dimensional module S . Note that $(\mathfrak{g}_0)_p/\mathfrak{h}_0 \cong \mathfrak{g}_0/\mathfrak{h}_0 \cap \mathfrak{g}_0$, as vector space. Hence

$$\dim V = p^{\dim \mathfrak{g}_0/\mathfrak{h}_0} 2^{\dim \mathfrak{g}_1/\mathfrak{h}_1}.$$

The proof is completed. ■

Remark 4.8 (1) The above theorem is an extension of the counterpart result on solvable Lie algebras (see [20] or [26, Section 5.8]).

(2) With the above theorem, we can propose the possibility that super KW property raised by Wang–Zhao in [29] is satisfied with all finite-dimensional solvable Lie superalgebras over \mathbf{k} .

5 Irreducible modules of completely solvable Lie superalgebras

A Lie superalgebra \mathfrak{g} is called completely solvable if $\mathfrak{g}^{(1)}$ is nilpotent. Obviously, a completely solvable \mathfrak{g} is solvable and its even part \mathfrak{g}_0 is a completely solvable Lie algebra.

The following facts are very important to the subsequent arguments.

Lemma 5.1 *Let $\mathfrak{g} = \mathfrak{g}_0 + \mathfrak{g}_1$ be a completely solvable Lie superalgebra. The following statements hold.*

- (1) *Any minimal ideal of \mathfrak{g} is one-dimensional.*
- (2) *There exists a sequence of ideals $\mathfrak{g} = \mathfrak{g}_0 \supset \mathfrak{g}_1 \supset \cdots \supset \mathfrak{g}_{n-1} \supset \mathfrak{g}_n = 0$ such that $\dim_{\mathbf{k}} \mathfrak{g}_i = n - i$.*
- (3) *For $\chi \in \mathfrak{g}_0^*$, if \mathfrak{h} is a subalgebra of codimension one which contains \mathfrak{z}^χ , then $D(\mathfrak{h}, \chi) \subset D(\mathfrak{g}, \chi)$.*
- (4) *Each proper subalgebra of \mathfrak{g} is contained in a subalgebra of codimension one of \mathfrak{g} .*

Proof By definition $\mathfrak{g}^{(1)}$ is nilpotent, it acts nilpotently on \mathfrak{g} under ad-action. By Lemma 4.1(1), the part (1) follows.

As to the part (2), by (1), there must be minimal ideals of \mathfrak{g} which is \mathbb{Z}_2 -homogeneous and one-dimensional. Notice that any subalgebras and quotients of \mathfrak{g} are also completely solvable Lie superalgebras. By induction on dimension, the statement follows.

As to (3), we are given $\mathfrak{m} \in D(\mathfrak{h}, \chi)$, intending to show $\mathfrak{m} \in D(\mathfrak{g}, \chi)$. We first note that $\dim \mathfrak{m} = d(\mathfrak{h}, \chi)$. As \mathfrak{h} has codimension one, either \mathfrak{h}_0 has codimension one in \mathfrak{g}_0 while $\mathfrak{h}_1 = \mathfrak{g}_1$, or \mathfrak{h}_1 has codimension one in \mathfrak{g}_1 while $\mathfrak{h}_0 = \mathfrak{g}_0$. For the first case, $\dim \mathfrak{h}_0 = \dim \mathfrak{g}_0 - 1$. We have that the even entry of the super-datum $d(\mathfrak{h}, \chi)$ is not less than $\frac{\dim \mathfrak{g}_0 - 1 + \dim \mathfrak{z}_0^\chi}{2} = \frac{\dim \mathfrak{g}_0 + \dim \mathfrak{z}_0^\chi}{2} - \frac{1}{2}$, therefore, both are equal because both are integers and $\frac{\dim \mathfrak{g}_0 + \dim \mathfrak{z}_0^\chi}{2}$ is already an integer. The odd entries of $d(\mathfrak{h}, \chi)$ and of $d(\mathfrak{g}, \chi)$ coincide. So in this case, $d(\mathfrak{h}, \chi) = d(\mathfrak{g}, \chi)$.

For the second case, $\dim \mathfrak{h}_1 = \dim \mathfrak{g}_1 - 1$ while $\mathfrak{h}_0 = \mathfrak{g}_0$. We can show by similar arguments as in the first case, that $d(\mathfrak{h}, \chi) = d(\mathfrak{g}, \chi)$ when $\frac{\dim \mathfrak{g}_1 + \dim \mathfrak{z}_1^\chi}{2}$ is an integer.

Suppose $\frac{\dim \mathfrak{g}_1 + \dim \mathfrak{z}_1^\chi}{2}$ is not an integer. Then the odd entries of $d(\mathfrak{h}, \chi)$ is equal to $\frac{\dim \mathfrak{h}_1 + \dim \mathfrak{z}_1^\chi}{2} = \frac{\dim \mathfrak{g}_1 - 1 + \dim \mathfrak{z}_1^\chi}{2}$, which is exactly $\lceil \frac{\dim \mathfrak{g}_1 + \dim \mathfrak{z}_1^\chi}{2} \rceil$, equal to the odd entry of $d(\mathfrak{g}, \chi)$. And the even entries of $d(\mathfrak{h}, \chi)$ and of $d(\mathfrak{g}, \chi)$ coincide already. Hence $d(\mathfrak{h}, \chi) = d(\mathfrak{g}, \chi)$ in this case. Hence, we always have $D(\mathfrak{h}, \chi) \subset D(\mathfrak{g}, \chi)$ in any case. The proof of (3) is completed.

Now, we prove (4) by induction on dimension. Suppose \mathfrak{h} is a proper subalgebra of \mathfrak{g} , and has codimension greater than one. Note that \mathfrak{g}_{n-1} is an ideal of dimension one.

So $\mathfrak{h} + \mathfrak{g}_{n-1}$ must be a proper subalgebra of \mathfrak{g} . If $\mathfrak{h} + \mathfrak{g}_{n-1}$ has codimension one, then we are done. If it has codimension greater than one, we consider $\phi : \mathfrak{g} \rightarrow \bar{\mathfrak{g}} := \mathfrak{g}/\mathfrak{g}_{n-1}$. Then $\phi(\mathfrak{h} + \mathfrak{g}_{n-1})$ has codimension greater than one in $\bar{\mathfrak{g}}$. Now, $\bar{\mathfrak{g}}$ has dimension less than $\dim_{\mathbf{k}} \mathfrak{g}$. By the inductive hypothesis, $\phi(\mathfrak{h} + \mathfrak{g}_{n-1})$ is contained in a subalgebra say $\bar{\mathfrak{p}}$, of codimension one in $\bar{\mathfrak{g}}$. We take $\phi^{-1}(\bar{\mathfrak{p}})$ the preimage of $\bar{\mathfrak{p}}$ in \mathfrak{g} . Then $\phi^{-1}(\bar{\mathfrak{p}})$ is a subalgebra of codimension one in \mathfrak{g} , containing \mathfrak{h} . This subalgebra is desired. ■

Corollary 5.2 *For a completely solvable Lie superalgebra \mathfrak{g} , the set $D(\mathfrak{g}, \chi)$ is not empty.*

Proof If $\mathfrak{z}^{\chi} = \mathfrak{g}$, there is nothing to do because \mathfrak{g} itself belongs to $D(\mathfrak{g}, \chi)$. So we only need to consider the situation where \mathfrak{z}^{χ} is a proper subalgebra of \mathfrak{g} in the following.

By Lemma 5.1, it is easily deduced that there are minimal subalgebras in \mathfrak{g} containing \mathfrak{z}^{χ} and having super-dimension not less than $d(\mathfrak{g}, \chi)$. We take one, say \mathfrak{p} . By Lemma 5.1 again, it is deduced that $D(\mathfrak{p}, \chi) \subset D(\mathfrak{g}, \chi)$. We will show that this \mathfrak{p} exactly belongs to $D(\mathfrak{p}, \chi)$, therefore belongs to $D(\mathfrak{g}, \chi)$.

(1) By applying Lemma 5.1(4) to the completely solvable superalgebra \mathfrak{p} , it is concluded that the proper subalgebra \mathfrak{z}^{χ} is contained in a subalgebra of codimension one in \mathfrak{p} . Hence, the minimality of \mathfrak{p} yields that $\underline{\dim} \mathfrak{p} = d(\mathfrak{g}, \chi)$.

(2) By the above arguments, in \mathfrak{p}_0 (resp. \mathfrak{p}_1) the maximal isotropic space has dimension equal to $\dim \mathfrak{p}_0$ (resp. $\dim \mathfrak{p}_1$). Hence, \mathfrak{p} itself is isotropic, which means $\chi(\mathfrak{p}^{(1)}) = 0$. Hence $\mathfrak{p} \in D(\mathfrak{p}, \chi) \subset D(\mathfrak{g}, \chi)$.

Thus $D(\mathfrak{g}, \chi)$ is not empty. The proof is completed. ■

By Corollary 5.2, there exists $\mathfrak{h} \in D(\mathfrak{g}, \chi)$ for any given $\chi \in \mathfrak{g}_0^*$.

Proposition 5.3 *Let $\mathfrak{g} = \mathfrak{g}_0 + \mathfrak{g}_1$ be a completely solvable Lie superalgebra, and $\chi \in \mathfrak{g}_0^*$ given. Then there is an irreducible module V of \mathfrak{g} such that $\dim V = p^{\frac{b_0^{\chi}}{2}-1} 2^{\lfloor \frac{b_1^{\chi}}{2} \rfloor}$.*

Proof By Corollary 5.2, there exists $\mathfrak{h} \in D(\mathfrak{g}, \chi)$. If \mathfrak{g} coincides with \mathfrak{h} , then all irreducible modules are one-dimensional, we are done.

In the following, we suppose that \mathfrak{h} is a proper subalgebra of \mathfrak{g} . We prove the statement by induction on $\dim \mathfrak{g}$. By Lemma 5.1, \mathfrak{h} is contained in a subalgebra \mathfrak{p} of codimension one in \mathfrak{g} , and $D(\mathfrak{p}, \chi) \subset D(\mathfrak{g}, \chi)$. By the inductive hypothesis along with Remark 2.2(1), there is an irreducible \mathfrak{p} -module W with

$$(5.1) \quad \dim W = \begin{cases} p^{\frac{b_0^{\chi}}{2}-1} 2^{\lfloor \frac{b_1^{\chi}}{2} \rfloor} & \text{if } \underline{\dim} \mathfrak{p} = (\dim \mathfrak{g}_0 - 1 | \dim \mathfrak{g}_1); \\ p^{\frac{b_0^{\chi}}{2}} 2^{\lfloor \frac{b_1^{\chi}}{2} \rfloor - 1} & \text{if } \underline{\dim} \mathfrak{p} = (\dim \mathfrak{g}_0 | \dim \mathfrak{g}_1 - 1). \end{cases}$$

There certainly exists an irreducible module V of \mathfrak{g} such that V contains an irreducible \mathfrak{p} -submodule isomorphic to W . We claim that $\dim V = p^{\frac{b_0^{\chi}}{2}} 2^{\lfloor \frac{b_1^{\chi}}{2} \rfloor}$.

Actually, by Proposition 4.5 and its proof, there is a subalgebra \mathcal{H} such that $\chi(\mathcal{H}^{(1)}) = 0$ and the dimensions of W and of V are respectively formulated as $\dim V = p^{\dim \mathfrak{g}_0 / \mathcal{H}_0} 2^{\dim \mathfrak{g}_1 / \mathcal{H}_1'}$, and $\dim W = p^{\dim \mathfrak{p}_0 / \mathcal{H}_0'} 2^{\dim \mathfrak{p}_1 / \mathcal{H}_1'}$, where $\mathcal{H}_0' = \mathfrak{g}_0 \cap \mathcal{H}_0$, $\mathcal{H}_1' =$

$\mathfrak{g}_{\bar{1}} \cap \mathcal{H}_{\bar{1}}$. Note that $(\mathfrak{p}_i + \mathcal{H}_i/\mathcal{H}_i) \cong \mathfrak{p}_i/\mathfrak{p}_i \cap \mathcal{H}_i$, $\forall i \in \mathbb{Z}_2$, as vector spaces. From (5.1), it yields that $\dim V = p^{\frac{b_0^x}{2}} 2^{\lfloor \frac{b_1^x}{2} \rfloor}$. ■

From now on, we turn to the situation of completely solvable restricted Lie superalgebras.

Proposition 5.4 *Let \mathfrak{g} be a completely solvable restricted Lie superalgebras with p -mapping $[p]$ on \mathfrak{g}_0 , and $\chi \in \mathfrak{g}_0^*$ any given p -character. Then the following statements hold.*

- (1) *Each irreducible $U_\chi(\mathfrak{g})$ -module V is associated with certain subalgebra \mathfrak{h} with $\chi(\mathfrak{h}^{(1)}) = 0$, such that V has dimension*

$$p^{\dim \mathfrak{g}_0/\mathfrak{h}_0} 2^{\dim \mathfrak{g}_0/\mathfrak{h}_0}$$

and there is a one-dimensional \mathfrak{h} -submodule in V .

- (2) *All irreducible modules of $U_\chi(\mathfrak{g})$ have the same dimension.*

Proof (1) follows from Theorem 4.7.

For (2), we first notice that the even center $C(\mathfrak{g})_{\bar{0}}$ is an ideal of \mathfrak{g} , and $C(\mathfrak{g})_{\bar{0}} = C(\mathfrak{g}) \cap \mathfrak{g}_{\bar{0}} \subset C(\mathfrak{g}_{\bar{0}})$. Then, we prove the proposition by different steps.

(2.i) We claim that if $C(\mathfrak{g})_{\bar{0}}^{[p]} = 0$, then every irreducible restricted representation of \mathfrak{g} is one-dimensional. Note that by definition $\mathfrak{g}^{(1)}$ is nilpotent, acting nilpotently on \mathfrak{g} . Hence, there exists a positive integer k such that $X^{[p]^k} \in C(\mathfrak{g})_{\bar{0}}$ for all $X \in \mathfrak{g}_{\bar{0}}$. The assumption that $C(\mathfrak{g})_{\bar{0}}^{[p]} = 0$ entails that $\mathfrak{g}_{\bar{0}}^{(1)} \subset \text{rad}_p(\mathfrak{g}_{\bar{0}})$, where $\text{rad}_p(\mathfrak{g}_{\bar{0}})$ denotes p -radical which is the set of $X \in \mathfrak{g}_{\text{ev}}$ with $X^{[p]^n} = 0$ for some positive integer $n \in \mathbb{N}$ (here n is dependent on X). By the same arguments as in the proof [26, Lemma 5.8.6(1)], one has $\mathfrak{g}_{\bar{0}} = T \oplus \text{rad}_p(\mathfrak{g}_{\bar{0}})$, where T is a maximal torus of $\mathfrak{g}_{\bar{0}}$. In particular, $C(\mathfrak{g})_{\bar{0}} \subset \text{rad}_p(\mathfrak{g}_{\bar{0}})$, and $\mathfrak{g}^{(1)} \subset \text{rad}_p(\mathfrak{g}_{\bar{0}}) + \mathfrak{g}_{\bar{1}}$ while $[\mathfrak{g}_{\bar{1}}, \mathfrak{g}_{\bar{1}}] \subset \text{rad}_p(\mathfrak{g}_{\bar{0}})$. Hence $\mathfrak{g}^{(1)} \subset \text{rad}(\mathfrak{g})$. For any irreducible restricted representation $\rho: \mathfrak{g} \rightarrow \mathfrak{gl}(V)$, $\rho(\text{rad}_p(\mathfrak{g}_{\bar{0}}))$ acts nilpotently on V , therefore acts trivially on V . So $\rho(\mathfrak{g}^{(1)})$ acts trivially on V . By Lemma 4.2, V is one-dimensional, which is of parity 0². This V is completely decided by a function λ with $\lambda(\mathfrak{g}^{(1)}) = 0$ and $\lambda(X^{[p]}) = \lambda(X)^p$ for $X \in \mathfrak{g}_{\bar{0}}$.

(2.ii) Suppose that (W_i, ρ_i) ($i = 1, 2$) are two irreducible modules of $U_\chi(\mathfrak{g})$ with $\rho_1|_{C(\mathfrak{g})_{\bar{0}}} = \rho_2|_{C(\mathfrak{g})_{\bar{0}}}$. Consider $\mathcal{H} := \text{Hom}_{\mathbf{k}}(W_1, W_2)$. Then $\mathcal{H} = \mathcal{H}_{\bar{0}} + \mathcal{H}_{\bar{1}}$ becomes a \mathfrak{g} -module by defining via homogenous elements $X \in \mathfrak{g}_{|X|}$, and $\nu \in \mathcal{H}_{|\nu|}$, where $|X|, |\nu| \in \mathbb{Z}_2$ denote the parities of X and ν , respectively:

$$(5.2) \quad (X.\nu)(v) = \rho_2(X)\nu(v) - (-1)^{|X||\nu|}\nu(\rho_1(X)v).$$

Furthermore, it is readily shown that \mathcal{H} is a restricted \mathfrak{g} -module, on which the representation is denoted by ϑ . By definition, $\vartheta(\mathfrak{g})$ admits a p -mapping satisfying $\vartheta(C(\mathfrak{g})_{\bar{0}})^{[p]} = 0$ because $\rho_1|_{C(\mathfrak{g})_{\bar{0}}} = \rho_2|_{C(\mathfrak{g})_{\bar{0}}}$ and $C(\mathfrak{g})_{\bar{0}} \subset C(\mathfrak{g}_{\bar{0}})$. Hence, there is an irreducible submodule of $U_0(\mathfrak{g})$ in \mathcal{H} , which must be one-dimensional by (2.i),

²For representation categories of $U_\chi(\mathfrak{g})$, one can give parities arising from the given parity of homogeneous generating space to [5, Section 6].

admitting parity 0. We take such one, for example, $\mathbb{S} := \mathbf{k}v$. Then $\mathfrak{g}^{(1)}$ acts trivially on v , and \mathfrak{g} acts on v by a scalar $\lambda \in \mathfrak{g}_0^*$ with $\lambda(\mathfrak{g}^{(1)}) = 0$. Also $\vartheta(X)v = \lambda(X)v$ for all $X \in \mathfrak{g}$, consequently $\lambda(X^{[p]}) = \lambda(X)^p$ for $X \in \mathfrak{g}_0$. By (5.2), $\rho_2(X)(\vartheta(w)) - \vartheta(\rho_1(X)w) = \lambda(X)\vartheta(w)$ for $X \in \mathfrak{g}$ and $w \in W_1$. So the assignment $w \mapsto \vartheta(w) \otimes 1$ for $w \in W_1$ defines a nontrivial \mathfrak{g} -module homomorphism: $W_1 \rightarrow W_2 \otimes_{\mathbf{k}} \mathbf{k}_{-\lambda}$, which is even. By Schur's lemma, this homomorphism must be an isomorphism. Hence $\dim W_1 = \dim W_2$.

(2.iii) Note that $C(\mathfrak{g})_{\bar{0}} \subset C(\mathfrak{g}_{\bar{0}})$. So $\mathfrak{g}_{\bar{0}}$ possesses a p -mapping $[p]'$ with $C(\mathfrak{g})_{\bar{0}}^{[p]'} = 0$ (see [26, Chapter 2]), and for any $y \in \mathfrak{g}_{\bar{0}}$, $\xi(y) := y^{[p]} - y^{[p]'}$ belongs to $C(\mathfrak{g})_{\bar{0}}$. So for any given irreducible representation (ρ, W) of \mathfrak{g} on the space W , by Schur's lemma $\xi(y)$ acts on W by scalar $c_W(y)^p$, where c_W is a linear function on $C(\mathfrak{g})_{\bar{0}}$. Then, we have for any $y \in C(\mathfrak{g})_{\bar{0}}$,

$$(5.3) \quad \rho(y)^p = (\chi + c_W)(y)^p \text{id}_V.$$

Note that c_W can extend to a function on $\mathfrak{g}_{\bar{0}}$ which gives rise to a one-dimensional $U_{c_W}(\mathfrak{g})$ -module, with respect to $(\mathfrak{g}, [p]')$. So for any two irreducible $U_{\chi}(\mathfrak{g})$ -modules W_1 and W_2 , we have two new irreducible representations $\bar{\rho}_i$ on $\bar{W}_i := W_i \otimes_{\mathbf{k}} \mathbf{k}_{-c_{W_i}}$ ($i = 1, 2$), respectively. Then both of them become $U_{\chi}(\mathfrak{g})$ -modules with respect to $(\mathfrak{g}, [p]')$, and $\bar{\rho}_i|_{C(\mathfrak{g})_{\bar{0}}}$ ($i = 1, 2$) coincide. By (2.ii), $\dim \bar{W}_i$ ($i = 1, 2$) have the same dimension. Hence, both W_1 and W_2 have the same dimension.

The proof is completed. ■

Summing up Propositions 5.3 and 5.4 we have the following.

Theorem 5.5 *Let $\mathfrak{g} = \mathfrak{g}_{\bar{0}} + \mathfrak{g}_{\bar{1}}$ be a finite-dimensional completely solvable restricted Lie superalgebra over \mathbf{k} . Then for any given $\chi \in \mathfrak{g}_0^*$, all irreducible \mathfrak{g} -modules associated with χ have dimension*

$$p^{\frac{b_0^{\chi}}{2}} 2^{\lfloor \frac{b_1^{\chi}}{2} \rfloor}.$$

The above theorem is an extension of Kac–Weisfeiler's result on completely solvable restricted Lie algebras (see [31, Theorem 1]). As a corollary to Theorem 5.5, we have the following.

Corollary 5.6 *Conjecture 2.3 holds for completely solvable restricted Lie superalgebras.*

A Appendix: Minimal p -envelopes for a finite-dimensional Lie superalgebra over \mathbf{k}

Let $\mathfrak{g} = \mathfrak{g}_{\bar{0}} \oplus \mathfrak{g}_{\bar{1}}$ be a finite-dimensional Lie superalgebra over \mathbf{k} of characteristic $p > 2$. In this appendix section, we introduce the properties of p -envelopes of \mathfrak{g} . For more details on restricted Lie algebras, the reader may refer to [26, Section 2.5].

A.1 Definition of p -envelopes

Definition A.1 Keep the notations and assumptions as above. A restricted Lie superalgebra $(\mathfrak{g}, [p])$ is said to be a p -envelope of \mathfrak{g} if there exists a homomorphism

of Lie superalgebras $\mathbf{i} : \mathfrak{g} \rightarrow \mathcal{G}$ such that \mathbf{i} is injective and the restricted Lie sub-superalgebra $\mathbf{i}(\mathfrak{g})_p$ of \mathcal{G} generated by \mathfrak{g} coincides with \mathcal{G} .

In the following, we make an example of p -envelopes which is taken from the universal enveloping algebra $U(\mathfrak{g})$. We first recall that an associative superalgebra $\mathfrak{A} = \mathfrak{A}_0 \oplus \mathfrak{A}_1$ can be endowed with structure of a restricted Lie superalgebra, which is denoted by \mathfrak{A}^- , where the underline space of \mathfrak{A}^- is \mathfrak{A} itself, and the Lie bracket is defined via $[u_1, u_2] := u_1 u_2 - (-1)^{|u_1||u_2|} u_2 u_1$ for \mathbb{Z}_2 -homogeneous elements $u_i \in \mathfrak{A}_{|u_i|}$, $|u_i| \in \mathbb{Z}_2$, $i = 1, 2$. And the p -mapping of \mathfrak{A}_0 is just the usual p th power in \mathfrak{A}_0 .

Example A.2 Let $\mathbf{i} : \mathfrak{g} \rightarrow U(\mathfrak{g})$ be the canonical imbedding of \mathfrak{g} into $U(\mathfrak{g})$. Recall that in $U(\mathfrak{g}_0)^- \subset U(\mathfrak{g})^-$, the Lie subalgebra $\mathbf{i}(\mathfrak{g}_0)$ generates a restricted Lie subalgebra $\mathbf{i}(\mathfrak{g}_0)_p = \bigoplus_{i=0}^{\infty} \mathbf{i}(\mathfrak{g})^{p^i}$ of $U(\mathfrak{g}_0)^-$, where $\mathbf{i}(\mathfrak{g})^{p^i} = \{\mathbf{i}(g)^{p^i} \mid g \in \mathfrak{g}_0\}$ (see [38] or [26, Section 5.2]). Take

$$\mathcal{G} = \mathbf{i}(\mathfrak{g}_0)_p \oplus \mathbf{i}(\mathfrak{g}_1) \subset U(\mathfrak{g})^-.$$

Then \mathcal{G} becomes a p -envelope of \mathfrak{g} .

Such a p -envelope as above is usually called the universal p -envelope of \mathfrak{g} , which we denote by $\widehat{\mathfrak{g}}$. By straightforward calculations, it is not hard to see that $\mathbf{i}(\mathfrak{g})$ is an ideal of $\widehat{\mathfrak{g}}$. Consider the superalgebra $\text{SDer}(\mathfrak{g})$ of super derivations on \mathfrak{g} , and the homomorphism $\text{ad} : \widehat{\mathfrak{g}} \rightarrow \text{SDer}(\mathfrak{g})$ defined via sending $x \mapsto \text{ad}x|_{\mathbf{i}(\mathfrak{g})}$. Then $\ker(\text{ad})$ coincides with the center $C(\widehat{\mathfrak{g}})$ of $\widehat{\mathfrak{g}}$.

By the same arguments as in the Lie algebras case (see [26, Section 2.5]), we have the following basic results.

Lemma A.3 *The following statements hold.*

- (1) *There is a finite-dimensional p -envelope \mathcal{G} of \mathfrak{g} such that $\mathcal{G} = (\mathfrak{g}_0)_p \oplus \mathfrak{g}_1$ where $(\mathfrak{g}_0)_p$ is a p -envelope of \mathfrak{g}_0 .*
- (2) *There is a minimal finite-dimensional p -envelope \mathcal{G} of \mathfrak{g} satisfying (1).*
- (3) *Any two minimal dimensional p -envelopes of \mathfrak{g} are isomorphic, as Lie superalgebras.*

Proof For the part (1), we choose a sub-superspace V in the center $C(\widehat{\mathfrak{g}})$ of $\widehat{\mathfrak{g}}$ such that $C(\widehat{\mathfrak{g}}) = V \oplus (C(\widehat{\mathfrak{g}}) \cap \phi(\mathfrak{g}))$. This V naturally becomes an ideal of $\widehat{\mathfrak{g}}$. Consider $\widetilde{\mathfrak{g}} := \widehat{\mathfrak{g}}/V$. It is easily seen that $\widetilde{\mathfrak{g}}$ is endowed with structure of restricted Lie superalgebras arising from the one of $\widehat{\mathfrak{g}}$. Furthermore, $\dim \widetilde{\mathfrak{g}} = \dim \widehat{\mathfrak{g}}/C(\widehat{\mathfrak{g}}) + \dim C(\widehat{\mathfrak{g}} \cap \phi(\mathfrak{g}))$. Note that $\phi(\mathfrak{g})$ is an ideal of $\widehat{\mathfrak{g}}$, and the homomorphism $\text{ad} : \widehat{\mathfrak{g}} \rightarrow \text{SDer}(\mathfrak{g})$ defined via sending $X \mapsto \text{ad}X|_{\phi(\mathfrak{g})}$, admits $\ker(\text{ad}) = C(\widehat{\mathfrak{g}})$. Hence $\dim \widehat{\mathfrak{g}}/C(\widehat{\mathfrak{g}}) \leq \dim \text{SDer}(\mathfrak{g}) < \infty$. Hence $\dim \widetilde{\mathfrak{g}} < \infty$.

By the choice of V , there is an embedding ψ of \mathfrak{g} into $\widetilde{\mathfrak{g}}$, i.e. $\psi = \text{p} \circ \phi$ for the natural surjective homomorphism $\text{p} : \widehat{\mathfrak{g}} \rightarrow \widetilde{\mathfrak{g}}$. Consequently, it is readily known that $\widetilde{\mathfrak{g}}$ is a p -envelope of \mathfrak{g} . Such $\widetilde{\mathfrak{g}}$ satisfies the requirement that $\widetilde{\mathfrak{g}}_0 = \psi(\mathfrak{g}_0)_p$ is a p -envelope of \mathfrak{g}_0 and $\widetilde{\mathfrak{g}} = \psi(\mathfrak{g}_0)_p \oplus \psi(\mathfrak{g}_1)$.

As to (2), note that in the above arguments, the center of $\tilde{\mathfrak{g}}$ lies in $\psi(\mathfrak{g})$. By an analog of the arguments in the proof of [26, Theorem 2.5.8], it can be proved that $\tilde{\mathfrak{g}}$ is a minimal finite-dimensional p -envelope of \mathfrak{g} satisfying (1).

The proof for (3) is also an analog of that of [26, Theorem 2.5.8]. We omit the details. ■

Acknowledgment The author expresses his sincere thanks to the anonymous referee for his/her helpful comments and suggestions, and to Dr. Priyanshu Chakraborty for assistance in English expression.

References

- [1] J. Brundan, *Modular representations of the supergroup $Q(n)$, II*. Pacific J. Math. 224(2006), 65–90.
- [2] J. Brundan and A. Kleshchev, *Modular representations of the supergroup $Q(n)$, I*. J. Algebra 260(2003), 64–98.
- [3] J. Brundan and J. Kujawa, *A new proof of the Mullineux conjecture*. J. Algebraic Combin. 18(2003), 13–39.
- [4] C. Carmeli, L. Caston, and R. Fiorese, *Mathematical foundations of supersymmetry*. Vol. 15, European Mathematical Society, Zürich, 2011.
- [5] S.-J. Cheng, N. Lam, and W. Wang, *The Brundan–Kazhdan–Lusztig conjecture for general linear Lie superalgebras*. Duke Math. J. 164(2015), no. 4, 617–695.
- [6] S.-J. Cheng, B. Shu, and W. Wang, *Modular representations of exceptional supergroups*. Math. Z. 291(2019), 635–659.
- [7] R. Fiorese and F. Gavarini, *Chevalley supergroups*. Mem. Amer. Math. Soc. 215(2012), 1014.
- [8] R. Fiorese and F. Gavarini, *Algebraic supergroups with Lie superalgebras of classical type*. J. Lie Theory 23(2013), 143–158.
- [9] S. M. Goodwin and L. Topley, *Minimal-dimensional representations of reduced enveloping algebras for \mathfrak{gl}_n* . Compos. Math. 155(2019), 1594–1617.
- [10] N. Jacobson, *Lie algebras*, Interscience Tracts in Pure and Applied Mathematics, No. 10, Interscience Publishers (a division of John Wiley & Sons, Inc.), New York, 1962.
- [11] V. G. Kac, *Lie superalgebras*. Adv. Math. 26(1977), 8–96.
- [12] V. G. Kac, Review on [17], MR 1345285 (96g: 17007), MathSciNet, Amer. Math. Soc.
- [13] Y. Li and B. Shu, *Jantzen filtration of Weyl modules for general linear supergroups*. Forum. Math. 35(2023), 1435–1468.
- [14] W. Liu, S. Wang, and J. Yuan, *On restricted simple modules of contact Lie superalgebras of odd type*. Math. Nachr. 290(2017), 2934–2947.
- [15] B. Martin, D. Stewart, and L. Topley, *A proof of the first Kac–Weisfeiler conjecture in large characteristics, with an appendix by T. Akaki*. Represent. Theory 23(2019), 278–293.
- [16] L. Pan and B. Shu, *Jantzen filtration and strong linkage principle for modular Lie superalgebras*. Forum. Math. 30(2018), 1573–1598.
- [17] A. Premet, *Irreducible representations of Lie algebras of reductive groups and the Kac–Weisfeiler conjecture*. Invent. Math. 121(1995), 79–117.
- [18] A. Premet and S. Skryabin, *Representations of restricted Lie algebras and families of associative π -algebras*. J. Reine Angew. Math. 507(1999), 189–218.
- [19] Y. Ren, B. Shu, F. Yang, and A. Zhang, *Modular representations of strange classical Lie superalgebras and the first super Kac–Weisfeiler conjecture*. Preprint, 2023. [arXiv:2307.00483\[math.RT\]](https://arxiv.org/abs/2307.00483)
- [20] J. Schue, *Representations of solvable Lie p -algebras*. J. Algebra 38(1976), 253–267.
- [21] B. Shu and W. Wang, *Modular representations of the ortho-symplectic supergroups*. Proc. Lond. Math. Soc. 96(2008), 251–271.
- [22] B. Shu and Y. Yao, *Character formulas for restricted simple modules of the special superalgebras*. Math. Nachr. 285(2012), 1107–1116.
- [23] B. Shu and C. Zhang, *Restricted representations of the Witt superalgebras*. J. Algebra 324(2010), no. 4, 652–672.
- [24] B. Shu and C. Zhang, *Representations of the restricted Cartan type Lie superalgebra $W(m, n, 1)$* . Algebr. Represent. Theory 14(2011), 463–481.

- [25] B. Shu and L. Zheng, *On Lie superalgebras of algebraic supergroups*. Algebra Colloq. 16(2009), 361–370.
- [26] H. Strade and R. Farnsteiner, *Modular Lie algebras and their representations*, Marcel Dekker, New York, 1988.
- [27] L. Topley, *A non-restricted counterexample to the first Kac–Weisfeiler conjecture*. Proc. Amer. Math. Soc. 45(2016), 1937–1942.
- [28] S. Wang, J. Yuan, and W. Liu, *Restricted Kac modules for special contact Lie superalgebras of odd type*. Front. Math. China 15(2020), 419–434.
- [29] W. Wang and L. Zhao, *Representations of Lie superalgebras in prime characteristic I*. Proc. Lond. Math. Soc. (3) 99(2009), 145–167.
- [30] W. Wang and L. Zhao, *Representations of Lie superalgebras in prime characteristic II: The queer series*. J. Pure Appl. Algebra 215(2011), 2515–2532.
- [31] B. J. Weisfeiler and V. G. Kac, *The irreducible representations of Lie p -algebras*. Funkcial. Anal. i Prilozhen. 5(1971), 28–36.
- [32] Y. Yao, *On restricted representations of the extended special type Lie superalgebra $\bar{S}(m, n, 1)$* . Monatsh. Math. 170(2013), 239–255.
- [33] Y. Yao, *Irreducible representations of the classical Lie superalgebra of type $A(0, n)$ in prime characteristic*. Monatsh. Math. 172(2013), 207–231.
- [34] Y. Yao, *Character formulas for a class of simple restricted modules over the simple Lie superalgebras of Witt type*. Chinese Ann. Math. Ser. B 41(2020), 49–60.
- [35] Y. Yao and B. Shu, *Restricted representations of Lie superalgebras of Hamiltonian type*. Algebr. Represent. Theory 16(2013), 615–632.
- [36] Y. Yao and B. Shu, *A note on restricted representations of the Witt superalgebras*. Chinese Ann. Math. Ser. B 34(2013), 921–926.
- [37] J. Yuan and W. Liu, *Restricted Kac modules of Hamiltonian Lie superalgebras of odd type*. Monatsh. Math. 178(2015), 473–488.
- [38] H. Zassenhaus, *The representations of Lie algebras of prime characteristic*. Proc. Glasgow Math. Assoc. 2(1954), 1–36.
- [39] Y. Zeng and B. Shu, *On Kac–Weisfeiler modules for general and special linear Lie superalgebras*. Israel J. Math. 214(2016), 471–490.
- [40] Y. Zeng and B. Shu, *Finite W -superalgebras and dimensional lower bounds for the representations of basic Lie superalgebras*. Publ. Res. Inst. Math. Sci. 53(2017), 1–63.
- [41] C. Zhang, *On the simple modules for the restricted Lie superalgebra $\mathfrak{sl}(n|1)$* . J. Pure Appl. Algebra 213(2009), 756–765.
- [42] L. Zhao, *Representations of Lie superalgebras in prime characteristic, III*. Pacific J. Math. 248(2010), 493–510.
- [43] L. Zheng and B. Shu, *Representations of $\mathfrak{gl}(m|n)$ and infinitesimal subgroups of $GL(m|n)$* . Chinese Ann. Math. Ser. A 31(2010), 129–142.

School of Mathematical Sciences, Ministry of Education Key Laboratory of Mathematics and Engineering Applications & Shanghai Key Laboratory of PMMP, East China Normal University, No. 500 Dongchuan Road, Shanghai 200241, China

e-mail: bshu@math.ecnu.edu.cn