

ON THE CANONICAL FORM OF TURBULENCE

SEIZÔ ITÔ

§ 0. **Introduction.** In K. Itô's paper [1] on the theory of turbulence, the problem to determine the canonical form of the moment tensor of temporally homogeneous and isotropic turbulence, has not been solved. In the present paper, the author will solve the problem by making use of the result of his preceding paper [2]. We shall treat the turbulence in R^3 , but the similar argument is possible in R^n .

§ 1. **Generalities.** In the theory of *turbulence*, the deviation of the velocity from its mean may be considered as a system of random vectors $u(t, \mathfrak{X}, \omega) = \langle u_p(t, \mathfrak{X}, \omega) / p = 1, 2, 3 \rangle$, where $t \in R^1$ and $\mathfrak{X} \in R^3$ denote the time and the position respectively and $\omega \in (\Omega, P)$ is the probability parameter; we assume naturally that $u_p(t, \mathfrak{X}, \omega)$ is B -measurable in $\langle t, \mathfrak{X}, \omega \rangle$ and belongs to $L^2(\Omega, P)$ for any fixed $\langle p, t, \mathfrak{X} \rangle$.¹⁾ We have clearly

$$(1.1) \quad \mathbf{E}_\omega [u_p(t, \mathfrak{X}, \omega)] = 0 \quad (\mathbf{E}_\omega [.] \text{ denotes the expectation}).$$

Now we define the moment tensor of the turbulence by

$$(1.2) \quad R_{pq}(t, \mathfrak{X}; s, \mathfrak{Y}) = \mathbf{E}_\omega [u_p(t, \mathfrak{X}, \omega) u_q(s, \mathfrak{Y}, \omega)];$$

then

$$(1.3) \quad R_{pq}(t, \mathfrak{X}; s, \mathfrak{Y}) = R_{qp}(s, \mathfrak{Y}; t, \mathfrak{X}), \text{ and}$$

$$(1.4) \quad \sum_{i,j} \alpha_i \bar{\alpha}_j R_{p_i p_j}(t_i, \mathfrak{X}_i; t_j, \mathfrak{X}_j) \cong 0^{2)} \quad (\alpha_i: \text{complex number}).$$

We consider the turbulence satisfying the following three conditions:

$$(1.5) \quad R_{pq}(t + \tau, \mathfrak{X}; s + \tau, \mathfrak{Y}) = R_{pq}(t, \mathfrak{X}; s, \mathfrak{Y}) \quad (\text{temporally homogeneous});$$

$$(1.6) \quad R_{pq}(t, \mathfrak{X} + a; s, \mathfrak{Y} + a) = R_{pq}(t, \mathfrak{X}; s, \mathfrak{Y}) \quad (\text{spatially homogeneous});$$

and

$$(1.7) \quad \sum_{p', q'} k_{p'p} k_{q'q} R_{p'q'}(t, \mathfrak{X}; s, \mathfrak{X} + K(\mathfrak{Y} - \mathfrak{X})) = R_{pq}(t, \mathfrak{X}; s, \mathfrak{Y})$$

for any orthogonal transformation $K \equiv (K_{pq}/p, q = 1, 2, 3)$ (*isotropic*). We can easily prove by (1.3) that the isotropism implies the homogeneity (1.6). Consequently we get

Received June 28, 1950.

¹⁾ See [1].

²⁾ If $R_{pq}(t, \mathfrak{X}; s, \mathfrak{Y})$ is real-valued and satisfies (1.4), then it satisfies (1.3) automatically.

$$(1.8) \quad \sum_{p, q} k_{pp} k_{qq} R_{pq}(t, K\mathfrak{x}; s, K\mathfrak{y}) = R_{pq}(t, \mathfrak{x}; s, \mathfrak{y}).$$

We shall determine a canonical form of $R_{pq}(t, \mathfrak{x}; s, \mathfrak{y})$ which satisfies (1.4), (1.5) and (1.7).

Suppose that $R_{pq}(t, \mathfrak{x}; s, \mathfrak{y})$ satisfies (1.4), (1.5) and (1.7) (and consequently all of (1.3) — (1.8) above); then there exists $u(t, \mathfrak{x}, \omega)$ satisfying (1.1) and (1.2).³⁾ Define

$$(1.9) \quad \begin{cases} u(l; t, \mathfrak{x}, \omega) = (l, u(t, \mathfrak{x}, \omega))_3^{4)}; \text{ and} \\ R(l, t, \mathfrak{x}; m, s, \mathfrak{y}) = \mathbf{E}_\omega [u(l; t, \mathfrak{x}, \omega) u(m; s, \mathfrak{y}, \omega)] \end{cases}$$

for $l \equiv \langle l_1, l_2, l_3 \rangle \in S$ ($=$ the unit sphere whose center is the origin of R^3), and put $l^p = \langle \delta_{p1}, \delta_{p2}, \delta_{p3} \rangle (\in S)$ for $p = 1, 2, 3$ (δ_{pq} is the Kronecker's delta). Then we have, by (1.2) and (1.8),

$$\begin{aligned} \mathbf{E}_\omega [(Kl^p, u(t, K\mathfrak{x}, \omega))_3 \cdot (Kl^q, u(s, K\mathfrak{y}, \omega))_3] \\ = \mathbf{E}_\omega [(l^p, u(t, \mathfrak{x}, \omega))_3 \cdot (l^q, u(s, \mathfrak{y}, \omega))_3]. \end{aligned}$$

From this equality and by simple calculation, we obtain

$$(1.8') \quad R(Kl, t, K\mathfrak{x}; Km, s, K\mathfrak{y}) = R(l, t, \mathfrak{x}; m, s, \mathfrak{y}).$$

And (1.6), (1.5), (1.4) (and (1.9)) imply following relations:

$$(1.6') \quad R(l, t, \mathfrak{x} + a; m, s, \mathfrak{y} + a) = R(l, t, \mathfrak{x}; m, s, \mathfrak{y}),$$

$$(1.5') \quad R(l, t + \tau, \mathfrak{x}; m, s + \tau, \mathfrak{y}) = R(l, t, \mathfrak{x}; m, s, \mathfrak{y}),$$

$$(1.4') \quad \sum_{i, j} \alpha_i \bar{\alpha}_j R(l_i, t_i, \mathfrak{x}_i; l_j, t_j, \mathfrak{x}_j) \cong 0.$$

Now we put $\mathbf{R} = S \times R^1 \times R^3$ and define the transformations T_τ, U_K, V_α on \mathbf{R} by $T_\tau \langle l, t, \mathfrak{x} \rangle = \langle l, t + \tau, \mathfrak{x} \rangle; U_K \langle l, t, \mathfrak{x} \rangle = \langle Kl, t, K\mathfrak{x} \rangle; V_\alpha \langle l, t, \mathfrak{x} \rangle = \langle l, t, \mathfrak{x} + a \rangle$. Let G be the group of transformations on \mathbf{R} generated by T_τ 's, U_K 's and V_α 's defined above. We easily see that $U_K V_\alpha = V_{K\alpha} U_K$ and that T_τ commutes with every element of G . Then \mathbf{R} is a homogeneous space with the locally compact group G of homeomorphisms which is transitive on \mathbf{R} ; for

$P_1 \equiv \langle l^1, 0, 0 \rangle \in \mathbf{R}$, $\{g \in G / gP_1 = P_1\}$ is identical with $\left\{ U_K / K = K_\theta \equiv \begin{pmatrix} 1 & 0 \\ 0 & \cos \theta \\ 0 & \sin \theta \end{pmatrix}; 0 \leq \theta < 2\pi \right\}$, which is compact subgroup of G ; and $R(l, t, \mathfrak{x}; m, s, \mathfrak{y})$ is a positive definite function on \mathbf{R}^2 in the sense of [2] by (1.4'), (1.5'),

³⁾ See [1].

⁴⁾ We denote the inner product and the norm in R^n by $(\dots)_n$ and $\|\cdot\|_n$ respectively.

(1.6') and (1.8'). Hence there exists a cyclic unitary representation⁵⁾ $\{\mathfrak{H}, \mathfrak{U}, f_1\}$ of $\langle \mathbf{R}, G, P_1 \rangle$ such that

$$(1.10) \quad R(t, \mathfrak{x}; m, s, \mathfrak{y}) = (T_t V_{\mathfrak{x}} U_{K(l)} f_1, T_s V_{\mathfrak{y}} U_{K(m)} f_1),^{6)}$$

where $K(l)$ is an orthogonal transformation such as $K(l)l = l$ and we denote the unitary operators ($\in \mathfrak{U}$) corresponding to $T_t, V_{\mathfrak{x}}, U_K (\in G)$ by the same notations respectively.

We shall denote by f_p the element of \mathfrak{H} corresponding to the point $P_p \equiv \langle l^p, 0, 0 \rangle \in \mathbf{R} (p = 1, 2, 3)$.⁷⁾ Then by (1.10) and (1.9) we have

$$\begin{aligned} (T_t V_{\mathfrak{x}} U_{K(l)} f_1, U_K f_p) &= R(t, \mathfrak{x}; K(l^p), 0, 0) \\ &= \sum_{p'} k_{pp'} R(t, \mathfrak{x}; l^{p'}, 0, 0) = (T_t V_{\mathfrak{x}} U_{K(l)} f_1, \sum_{p'} k_{pp'} f_{p'}) \end{aligned}$$

($K = (k_{pq})$). Since $\{T_t V_{\mathfrak{x}} U_{K(l)} f_1 / t \in R^1, \mathfrak{x} \in R^3, l \in S\}$ spans \mathfrak{H} ,⁷⁾ we have $U_K f_p = \sum_{p'} k_{pp'} f_{p'}$; and hence

$$(1.11) \quad U_K f = \sum_{p'} (\sum_p k_{pp'} \alpha_p) f_{p'}, \text{ for } f = \sum_p \alpha_p f_p \quad (\alpha_p: \text{complex number}).$$

Thus we see that, in order to determine a canonical form of $R(t, \mathfrak{x}; m, s, \mathfrak{y})$, it is sufficient to consider cyclic unitary representations $\{\mathfrak{H}, \mathfrak{U}, f_1\}$ satisfying (1.11).

§ 2. Preliminary lemmas.

LEMMA 1. Let $F(\cdot)$ be an additive set function on R^n .⁸⁾ If $\int_{R^n} e^{i(\mathfrak{x}, X)} F(dX) = 0$ for any $\mathfrak{x} \in R^n$, then $F(\cdot) \equiv 0$.

The proof can be achieved by simple calculations and so will be omitted.

Let \mathfrak{G} be a locally compact group, and let \mathfrak{g} be a compact subgroup of \mathfrak{G} ,

⁵⁾ See [2]; — In the paper [2], if we use the notations following after the paper [2], the positive-definiteness of $f(p, q)$ is defined as follows: $f(p, q)$ is measurable in $\langle p, q \rangle$ and essentially bounded and satisfies.

$$(1) \quad f(\sigma p, \sigma q) = f(p, q) \quad (\langle p, q \rangle \in \Omega^2, \sigma \in G), \text{ and}$$

$$(2) \quad \iint_{\Omega^2} f(p, q) x(p) \bar{x}(q) dpdq \cong 0 \quad (x \in L^1(\Omega));$$

and it is noted that if $f(p, q)$ is continuous in $\langle p, q \rangle$ then (2) is equivalent to

$$(3) \quad \sum_{i, j} \alpha_i \bar{\alpha}_j f(p_i, p_j) \cong 0$$

and (3) implies that $f(p, q)$ is bounded ($|f(p, q)| \leq f(p, p) = f(p_0, p_0)$). But it may be proved without any use of the continuity of $f(p, q)$ that (3) implies (2) (cf. [5], pp. 56-57). So we can make use of the results of the paper [2] directly from (1.4') (and (1.5'), (1.6'), (1.8')) in the present paper.

⁶⁾ We denote the inner product and the norm in \mathfrak{H} by (\cdot, \cdot) and $\|\cdot\|$ respectively.

⁷⁾ See [2], § 2.

⁸⁾ Under an additive set function on a topological space \mathbf{S} we understand a (generally complex-valued) countably additive Borel-set function on \mathbf{S} such that the total variation on \mathbf{S} is finite (following after S. Saks [6]), where Borel sets mean such sets as belongs to the minimal countably additive class including all open sets in the space \mathbf{S} .

we introduce a \mathfrak{G} -invariant measure ds on the homogeneous space $\mathfrak{S} \equiv \mathfrak{G}/\mathfrak{g}$ in the natural way so that $ds \cdot dK' = dK$ ³⁾ where dK denotes a left-invariant Haar-measure on the locally compact group \mathfrak{G} and dK' denotes the Haar-measure on the compact group \mathfrak{g} such as $\int_{\mathfrak{g}} dK' = 1$. Let $F(\cdot), F_1(\cdot), \dots, F_n(\cdot)$ be additive set functions on \mathfrak{S} , and $|F|(\cdot), |F_1|(\cdot), \dots, |F_n|(\cdot)$ be their total variations respectively. Then we have the following

LEMMA 2. *Suppose that $F(K\mathcal{A}) (K \in \mathfrak{G})$ is expressible in the form:*

$$F(K\mathcal{A}) = \sum_{\nu=1}^n \int_{\mathcal{A}} \varphi_{\nu}(K; s) F_{\nu}(ds),$$

$\varphi_{\nu}(K; s)$ being summable on \mathfrak{S} with respect to the measure $|F_{\nu}|(\cdot)$, $\nu = 1, 2, \dots, n$. Then $F(\cdot)$ is absolutely continuous with respect to the \mathfrak{G} -invariant measure ds .

Proof. Let $s_0 (\in \mathfrak{S})$ be the image of \mathfrak{g} by the natural mapping $\mathfrak{G} \rightarrow \mathfrak{S} \equiv \mathfrak{G}/\mathfrak{g}$, K_s be an element of \mathfrak{G} such that $K_s s_0 = s$, and $\rho(K)$ be the Neumann's function with respect to the Haar-measure dK on \mathfrak{G} ; it is easy to see that $\rho(K) \equiv 1$ on the compact subgroup \mathfrak{g} of the group \mathfrak{G} , and hence $\rho(K_s)$ depends only on s . Suppose that \mathcal{A} is a Borel set in \mathfrak{S} such that $\int_{\mathcal{A}} ds = 0$, and let $C_{\mathcal{A}}(s)$ be the characteristic function of \mathcal{A} . Then for all $s \in \mathfrak{S}$

$$\begin{aligned} \int_{\mathfrak{G}} C_{\mathcal{A}}(K^{-1}s) dK &= \int_{\mathfrak{G}} C_{\mathcal{A}}(K^{-1}K_s s_0) dK = \int_{\mathfrak{G}} C_{\mathcal{A}}(K^{-1}s_0) dK = \int_{\mathfrak{G}} C_{\mathcal{A}}(Ks_0) \rho(K) dK \\ &= \int_{\mathfrak{G}} \rho(K_s) ds \int_{\mathfrak{g}} \rho(K') C_{\mathcal{A}}(K_s K' s_0) dK' = \int_{\mathfrak{G}} \rho(K_s) C_{\mathcal{A}}(s) ds = 0. \end{aligned}$$

Hence for every ν we have by Fubini's theorem

$$0 = \int_{\mathfrak{G}} |F_{\nu}|(ds) \int_{\mathfrak{G}} C_{\mathcal{A}}(K^{-1}s) dK = \int_{\mathfrak{G}} dK \int_{\mathfrak{G}} C_{\mathcal{A}}(K^{-1}s) |F_{\nu}|(ds) = \int_{\mathfrak{G}} |F_{\nu}|(K\mathcal{A}) dK.$$

Therefore

$$|F_{\nu}|(K\mathcal{A}) = 0, \quad \nu = 1, 2, \dots, n,$$

for almost all K with respect to dK ; and for such a K we obtain by the assumption that

$$F(\mathcal{A}) = F(K^{-1}K\mathcal{A}) = \sum_{\nu=1}^n \int_{K\mathcal{A}} \varphi_{\nu}(K^{-1}; s) F_{\nu}(ds) = 0, \quad q.e.d.$$

§ 3. **Canonical form of isotropic turbulence.** In this paragraph as well as in the next, we represent every point $X \in R^3$ except the origin by the couple

³⁾ See [5], pp. 43-45.

$\langle \rho, \lambda \rangle$ of $\rho = \|X\|_3 (> 0)$ and $\lambda = X/\|X\|_3 (\in \text{the unit sphere } S \text{ with the center } \langle 0, 0, 0 \rangle \in \mathbb{R}^3)$; $X \equiv \rho\lambda = \langle \rho, \lambda \rangle$; and in order to clarify that a set A is a subset of the set $\{X/X \in \mathbb{R}^3\}$, we shall use the notation with the subscript $X: A_X$; similarly Γ_τ and B_ρ denote subsets of $\{\tau / -\infty < \tau < \infty\}$ and $\{\rho / 0 < \rho < \infty\}$ respectively.

The purpose of this paragraph is to prove the following

THEOREM 1. *A necessary and sufficient condition for $(R_{pq}(t, \mathfrak{X}; s, \mathfrak{Y}) / p, q = 1, 2, 3)$ to be a moment tensor of a temporally homogeneous and isotropic turbulence is that $R_{pq}(t, \mathfrak{X}; s, \mathfrak{Y})$ is expressible in the form:*

$$(3.1) \quad R_{pq}(t, \mathfrak{X}; s, \mathfrak{Y}) = \int_{[0, \infty)_\tau \times (0, \infty)_\rho \times S} \cos[(t-s)\tau + (\mathfrak{X} - \mathfrak{Y}, \rho\lambda)_3] \times \left(K(\lambda)^{-1} \mathfrak{P}, \mu(d\tau d\rho) K(\lambda)^{-1} \mathfrak{Q} \right)_3 d\lambda + \delta_{pq} \int_{-9}^{\infty} \cos(t-s)\tau \cdot \nu(d\tau),^{10)}$$

where $\mathfrak{P} = \langle \delta_{p1}, \delta_{p2}, \delta_{p3} \rangle (\in S) (p = 1, 2, 3)$ and $\{K(\lambda) / \lambda \in S\}$ denotes a system of orthogonal transformations such that $K(\lambda)$ is an arbitrarily fixed one such as $K(\lambda) \langle 1, 0, 0 \rangle = \lambda$ for every $\lambda \in S$,¹¹⁾ and $\mu(A)$ is written as

$$(3.2) \quad \mu(A) = \begin{pmatrix} \mu_1(A) & & 0 \\ & \mu_2(A) & \\ 0 & & \mu_3(A) \end{pmatrix}$$

and $\mu_p(A) (p = 1, 2)$ and $\nu(\Gamma_\tau)$ are measures defined on $[0, \infty)_\tau \times (0, \infty)_\rho$ and on $[0, \infty)_\tau$ respectively such that $\mu_p([0, \infty)_\tau \times (0, \infty)_\rho) < \infty (p = 1, 2)$ and $\nu([0, \infty)_\tau) < \infty$.

The sufficiency of this condition is easily proved by simple calculations.

To prove the necessity, we consider the cyclic unitary representation $\{\mathfrak{H}, \Pi, f_j\}$ of $\{R, G, P_i\}$ defined in §1. Since $\{T_t V_{\mathfrak{X}} / t \in R^1, \mathfrak{X} \in R^3\}$ is a unitary representation of the 4-dimensional vector space, we have by Stone's theorem (in the form generalized by W. Ambrose [4])

$$(3.3) \quad T_t V_{\mathfrak{X}} = \int_{R^4} e^{it\tau} e^{i(\mathfrak{X}, X)_3} E(d\tau dX)$$

where $E(\cdot)$ is a resolution of the identity on $R^4 (= R_\tau^1 \times R_X^3)$. The function

¹⁰⁾ The second term of the right-hand side corresponds to the case mentioned in [1] (see [1], Theorem 1).

¹¹⁾ It is true that the above system $\{K(\lambda)\}$ is not uniquely determined, but we may see from (3.2) that $\left(K(\lambda)^{-1} \mathfrak{P}, \mu(A) K(\lambda)^{-1} \mathfrak{Q} \right)_3$ is independent of the special choice of this system and is continuous in λ for any fixed A .

$F_{pq}(\Gamma_\tau \times \Delta_X) = (E(\Gamma_\tau \times \Delta_X) f_p, f_q)$ ($\Gamma_\tau \subseteq R^1, \Delta_X \subseteq R^3$) is an additive set function on R^4 and $F(\Gamma_\tau \times \Delta_X) = (F_{pq}(\Gamma_\tau \times \Delta_X) / p, q = 1, 2, 3)$ is a hermitian 3×3 -matrix. In considering (1.11), we obtain from (1.10) and (3.3)

$$(3.4) \quad R(l, t, \mathfrak{X}; m, s, \mathfrak{Y}) = (T_t V_{\mathfrak{X}} U_{K(l)} f_l, T_s V_{\mathfrak{Y}} U_{K(m)} f_s) \\ = \int_{R^4} e^{i(t-s)\tau} e^{i(\mathfrak{X}-\mathfrak{Y}, X)\mathfrak{s}} (l, F(d\tau dX) m)_{\mathfrak{s}}$$

for $l, m \in S$, and hence, putting $K = (k_{pq})$, we get for any t and \mathfrak{X}

$$\int_{R^4} e^{it\tau} e^{i(\mathfrak{X}, X)\mathfrak{s}} F_{pq}(d\tau d_X(KX)) \\ = \int_{R^4} e^{it\tau} e^{i(\mathfrak{X}, K^{-1}X)\mathfrak{s}} (p, F(d\tau dX) q)_{\mathfrak{s}} = (T_t V_{K\mathfrak{X}} f_p, f_q) \\ = (T_t V_{\mathfrak{X}} U_{K^{-1}} f_p, U_{K^{-1}} f_q) = \int_{R^4} e^{it\tau} e^{i(\mathfrak{X}, X)\mathfrak{s}} \left[\sum_{p', q'} k_{pp'} k_{qq'} F_{p'q'}(d\tau dx) \right].$$

Therefore by Lemma 1

$$(3.5) \quad F_{pq}(\Gamma_\tau \times K\Delta_X) = \sum_{p', q'} k_{pp'} k_{qq'} F_{p'q'}(\Gamma_\tau \times \Delta_X).$$

If we put $R_\tau = (-\infty, \infty)_\tau \times \{0\}_X (\subseteq R^4)$ and define $H(M)$ and $H'(M)$ for every Borel set $M \subseteq R^4$ by

$$(3.6) \quad \begin{cases} H(M) \equiv (H_{pq}(M)) = (F_{pq}(M - R_\tau)) \cdot \\ H'(M) \equiv (H'_{pq}(M)) = (F_{pq}(M \cap R_\tau)), \end{cases}$$

then $H'(M)$ is a hermitian matrix and satisfies $H'(M) = K \cdot H'(M) \cdot K^{-1}$ for any orthogonal transformation $K \equiv (k_{pq})$ by (3.5). Therefore we see that $H'(M)$ is specialized as

$$(3.7) \quad H'(M) = \begin{pmatrix} H'_1(M) & & 0 \\ & H'_1(M) & \\ 0 & & H'_1(M) \end{pmatrix}.$$

Next we consider the matrix $H(M)$. For any Borel set $A \subseteq (-\infty, \infty)_\tau \times (0, \infty)_p$, the function $H_{pq}(A \times A)$ is an additive function of a Borel set $A \subseteq S$, and we have by (3.5) and (3.6)

$$(3.8) \quad H_{pq}(A \times KA) = \sum_{p', q'} k_{pp'} k_{qq'} H_{p'q'}(A \times A).$$

Hence by Lemma 2 (in putting $\mathfrak{S} = S, \varphi_{p'q'}(K; \lambda) \equiv k_{pp'} k_{qq'}$) and by Radon-Nikodym's theorem, we obtain a function $h_{pq}(A; \lambda)$ such that

$$(3.9) \quad H_{pq}(A \times A) = \int_A h_{pq}(A; \lambda) d\lambda,$$

where $d\lambda$ is the K -invariant measure on S such as $\int_S d\lambda = 4\pi$ (i.e., the area in the usual sense). And for any fixed A , we can determine $h_{pq}(A; \lambda)$ for all λ so that

$$(3.10) \quad h_{pq}(A; K\lambda) = \sum_{p', q'} k_{pp'} k_{qq'} h_{p'q'}(A; \lambda),$$

in making use of (3.8) and the fact that the group $\{K\}$ of all orthogonal transformations is transitive on S . Consequently from $H_{pp}(A \times A) = (E(A \times A - R_\tau) f_p, f_p) = \|E(A \times A - R_\tau) f_p\|^2 \cong 0$ (see (3.6)) we get

$$(3.11) \quad h_{pp}(A; \lambda) \cong 0.$$

The matrix $h(A; \lambda) = (h_{pq}(A; \lambda)/p, q = 1, 2, 3)$ is hermitian for every A and λ , and by (3.10) we have

$$(3.10') \quad h(A; K\lambda) = K \cdot h(A; \lambda) \cdot K^{-1}.$$

If we put $\lambda = \lambda^1 \equiv \langle 1, 0, 0 \rangle$ and $K = K(\lambda)$ in (3.10') and define $h(A) \equiv (h_{pq}(A)) = (h_{pq}(A; \lambda^1))$, then we have

$$(3.12) \quad h(A; \lambda) = K(\lambda) h(A) K(\lambda)^{-1}.$$

And if we put $\lambda = \lambda^1$ and $K = K_0 \equiv \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \theta & -\sin \theta \\ 0 & \sin \theta & \cos \theta \end{pmatrix}$ in (3.10'), then we have

$$K_0 h(A) K_0^{-1} = h(A; K_0 \lambda^1) = h(A).$$

Thus we see that $h(A)$ is a hermitian matrix which is invariant under any transformation K_0 of the above form. This fact implies that $h(A)$ is written as

$$(3.13) \quad h(A) = \begin{pmatrix} h_1(A) & & 0 \\ & h_2(A) & \\ 0 & & h_3(A) \end{pmatrix},$$

consequently $K(\lambda) h(A) K(\lambda)^{-1}$ depends only on λ and is independent of the choice of $K(\lambda)$, and hence $h(A; \lambda)$ is continuous in λ by (3.12). From this and (3.11) we get $0 \leq h_p(A) (\equiv h_{pp}(A)) < \infty$ ($p = 1, 2$).

Now we shall show that the set functions $h_p(A) = h_{pp}(A)$ ($p = 1, 2$) have the properties of *measure*. Obviously they are defined for all Borel sets $A \equiv (-\infty, \infty)_\tau \times (0, \infty)_\rho$. We shall prove the countable additivity of $h_p(A)$.

If $A = \sum_{n=1}^\infty A_n$, then we have by (3.11) and Lebesgue's theorem

$$\begin{aligned} \int_A \sum_{n=1}^\infty h_{pp}(A_n; \lambda) d\lambda &= \sum_{n=1}^\infty \int_{A_n} h_{pp}(A_n; \lambda) d\lambda = \sum_{n=1}^\infty H_{pp}(A_n \times A) \\ &= H_{pp}(A \times A) = \int_A h_{pp}(A; \lambda) d\lambda \end{aligned}$$

for any Borel set $A \subseteq S$. Hence

$$(3.14) \quad \sum_{n=1}^\infty h_{pp}(A_n; \lambda) = h_{pp}(A; \lambda)$$

for almost all λ ; since the group $\{K\}$ of all orthogonal transformations is transi-

tive on S , we see from (3.10) that (3.14) is true for *all* λ , especially putting $\lambda = \lambda^i$, we get $\sum_{n=1}^{\infty} h_p(A_n) = h_p(A)$, which proves the countable additivity of $h_p(A)$.

Thus if we put $h'(\Gamma_\tau) = H'_1(\Gamma_\tau \times \langle 0 \rangle_X)$ (see (3.7)) and apply (3.6), (3.7), (3.9) and (3.12) to (3.4), we obtain

$$R(l, t, \mathfrak{X}; m, s, \mathfrak{Y}) = \int_{R^4 - R_\tau} e^{i(t-s)\tau} e^{i(\mathfrak{X}-\mathfrak{Y}, X)_s} \left(l, K(\lambda) h(d\tau d\rho) K(\lambda)^{-1} m \right)_3 d\lambda + \int_{-\infty}^{\infty} e^{i(t-s)\tau} (l, m)_s h'(d\tau), \quad X = \langle \rho, \lambda \rangle.$$

Since $R(l, t, \mathfrak{X}; m, s, \mathfrak{Y})$ is a *real-valued* positive definite function, we can easily prove that

$$(3.15) \quad R(l, t, \mathfrak{X}; m, s, \mathfrak{Y}) = \int_{[0, \infty)_\tau \times (0, \infty)_\rho \times S} \cos[(t-s)\tau + (\mathfrak{X}-\mathfrak{Y}, \rho\lambda)_s] \times \left(l, K(\lambda) \mu(d\tau d\rho) K(\lambda)^{-1} m \right)_3 d\lambda + \int_{-\infty}^{\infty} \cos(t-s)\tau \cdot (l, m)_s \cdot \nu(d\tau),$$

where

$$(3.2) \quad \mu(A) = \begin{pmatrix} \mu_1(A) & & 0 \\ & \mu_2(A) & \\ 0 & & \mu_3(A) \end{pmatrix}$$

and $\mu_p(A)$ ($p = 1, 2$) and $\nu(\Gamma_\tau)$ are measures defined on $[0, \infty)_\tau \times (0, \infty)_\rho$ and on $[0, \infty)_\tau$ respectively such that $\mu_p([0, \infty)_\tau \times (0, \infty)_\rho) < \infty$ ($p = 1, 2$) and $\nu([0, \infty)_\tau) < \infty$; it is true that the transformation $K(\lambda)$ such as $K(\lambda)\lambda^i = \lambda$ is not uniquely determined, but the measure matrix $K(\lambda)\mu(d\tau d\rho)K(\lambda)^{-1}$ depends only on λ and is independent of special choice of $K(\lambda)$, as may be verified from the form of (3.2). Putting $l = l^p$, $m = m^q$ in (3.15) and making use of (1.2) and (1.9), we obtain (3.1), which completes the proof of Theorem 1.

§ 4. Inversion formula. In this paragraph, we shall show the formulas which express the measures $\mu_p(A)$ ($p = 1, 2$) and $\nu(\Gamma_\tau)$ in Theorem 1 by means of corresponding $(R_{pq}(t, \mathfrak{X}; s, \mathfrak{Y}))$, from which we may conclude the uniqueness of the measures $\mu_p(A)$ and $\nu(\Gamma_\tau)$ for the functions $R_{pq}(t, \mathfrak{X}; s, \mathfrak{Y})$; $p, q = 1, 2, 3$. By definition we shall term *continuity points* of μ such ρ and τ as $\mu_p([0, \infty)_\tau \times \{\rho\}) = 0$ and $\mu_p(\{\tau\} \times (0, \infty)_\rho) = 0$ for $p = 1, 2$, and *continuity point* of ν such τ as $\nu(\{\tau\}) = 0$.

THEOREM 2. *The measures $\mu_p(A)$ and $\nu(\Gamma_\tau)$ in Theorem 1 are expressible*

by $R_{pq}(t, \mathfrak{X}; s, \mathfrak{Y})$ as follows: if we put $A_n = \{\lambda \in S / \|\lambda - \lambda^1\|_3 \leq 1/n (\lambda^1 = \langle 1, 0, 0 \rangle)$ and $I_c = \{\langle t, \mathfrak{X} \rangle / |t|, |x_1|, |x_2|, |x_3| \leq c\} (\mathfrak{X} = \langle x_1, x_2, x_3 \rangle)$, then we have for any continuity points $\rho, \rho' (0 < \rho < \rho')$ and τ of μ

$$(4.1) \quad \begin{aligned} \mu_p([0, \tau] \times (\rho, \rho']) &= \lim_{n \rightarrow \infty} \left(\frac{n^2}{\pi^3}\right) \lim_{c \rightarrow \infty} \int_{I_c} R_{pp}(t, \mathfrak{X}; 0, 0) dt d\mathfrak{X} \\ &\quad \times \int_{(-1, \tau] \times (\rho, \rho'] \times \Lambda_n} \cos [t\tau + (\mathfrak{X}, X)_3] d\tau dX,^{12)} \\ &\quad X = \langle \rho, \lambda \rangle, \end{aligned}$$

and for any continuity point τ of ν

$$(4.2) \quad \begin{aligned} \nu([0, \tau]) &= \lim_{\rho \downarrow 0} \left(\frac{1}{\pi^3}\right) \lim_{c \rightarrow \infty} \int_{I_c} R_{11}(t, \mathfrak{X}; 0, 0) dt d\mathfrak{X} \\ &\quad \times \int_{(-1, \tau] \times \{X\|_3 \leq \rho\}} \cos [t\tau + (\mathfrak{X}, X)_3] d\tau dX. \end{aligned}$$

Proof. Define the measures $\sigma_p^1(M)$, $\sigma^2(M)$ and $\sigma_p(M)$ on $R^4 (= R_\tau^1 \times R_X^3)$ by

$$\begin{cases} \sigma_p^1(\Gamma_\tau \times B_\rho \times A) = \int_\Lambda (K(\lambda)^{-1}[\rho, \mu(\Gamma_\tau \times B_\rho) K(\lambda)^{-1}[\rho]) d\lambda, \\ \sigma_p^1(M) = 0 \text{ if } M \cap ([0, \infty)_\tau \times (0, \infty)_\rho \times S) \text{ is empty;} \\ \sigma^2(\Gamma_\tau \times \{0\}_X) = \nu(\Gamma_\tau), \\ \sigma^2(M) = 0 \text{ if } M \cap ([0, \infty)_\tau \times \{0\}_X) \text{ is empty;} \text{ and} \\ \sigma_p(M) = \sigma_p^1(M) + \sigma^2(M). \end{cases}$$

Then it follows from Theorem 1 that

$$R_{pp}(t, \mathfrak{X}; 0, 0) = \int_{R^4} \cos [t\tau + (\mathfrak{X}, X)_3] \sigma_p(d\tau dX).$$

Hence, by Lévy-Haviland's inversion formula [3], for any continuity points $\rho, \rho' (0 < \rho < \rho')$ and τ of μ , we have

$$(4.3) \quad \begin{aligned} \int_{\Lambda_n} (K(\lambda)^{-1}[\rho, \mu([0, \tau] \times (\rho, \rho']) K(\lambda)^{-1}[\rho]) d\lambda \\ = \sigma_p((-1, \tau] \times (\rho, \rho'] \times \Lambda_n) \\ = \frac{1}{\pi^3} \lim_{c \rightarrow \infty} \int_{I_c} R_{pp}(t, \mathfrak{X}; 0, 0) dt d\mathfrak{X} \\ \quad \times \int_{(-1, \tau] \times (\rho, \rho'] \times \Lambda_n} \cos [t\tau + (\mathfrak{X}, X)_3] d\tau dX, \\ X = \langle \rho, \lambda \rangle. \end{aligned}$$

¹²⁾ “-1” in the domain $(-1, \tau]$ of the integral in the right-hand side has no particular sense, and may be replaced by an arbitrary negative number, as will easily be seen from the proof of this theorem.

Since $\left(K(\lambda)^{-1}l^p, \mu([0, \tau] \times (\rho, \rho']) K(\lambda)^{-1}l^p \right)_3$ is continuous in λ , we get

$$(4.4) \quad \begin{aligned} \mu_p([0, \tau] \times (\rho, \rho']) &= \left(l^p, \mu([0, \tau] \times (\rho, \rho']) l^p \right)_3 \\ &= \left(K(\lambda)^{-1}l^p, \mu([0, \tau] \times (\rho, \rho']) K(\lambda)^{-1}l^p \right)_3 \\ &= \lim_{n \rightarrow \infty} \frac{1}{|A_n|} \int_{A_n} \left(K(\lambda)^{-1}l^p, \mu([0, \tau] \times (\rho, \rho']) K(\lambda)^{-1}l^p \right)_3 d\lambda \end{aligned}$$

where $|A_n| = \int_{A_n} d\lambda$. From (4.4), (4.3) and the fact that $\lim_{n \rightarrow \infty} (|A_n| n^2/\pi) = 1$, we obtain (4.1).

Next we put

$$(4.5) \quad \begin{aligned} \psi(\tau, \rho) &= \lim_{\sigma \rightarrow \infty} \frac{1}{\pi^\sigma} \int_{I_\sigma} R_{11}(t, \mathfrak{X}; 0, 0) dt d\mathfrak{X} \\ &\quad \times \int_{(-1, \tau] \times \{ \|X\|_3 \leq \rho \}} \cos [t\tau + (\mathfrak{X}, X)_3] d\tau dX; \end{aligned}$$

then, by Lévy-Haviland's inversion formula, we have

$$(4.6) \quad \begin{aligned} \sigma_1^1 \left((-1, \tau) \times (0, \rho) \times S \right) + \sigma^2 \left((-1, \tau) \times \{0\}_X \right) &\leq \psi(\tau, \rho) \\ &\leq \sigma_1^1 \left((-1, \tau] \times (0, \rho] \times S \right) + \sigma^2 \left((-1, \tau] \times \{0\}_X \right) \end{aligned}$$

for any τ and ρ . If τ is a continuity point of ν , then $\sigma^2 \left((-1, \tau) \times \{0\}_X \right) = \sigma^2 \left((-1, \tau] \times \{0\}_X \right) = \nu([0, \tau])$; and hence, if ρ tend to 0 in (4.6), we obtain

$$(4.7) \quad \nu([0, \tau]) = \lim_{\rho \downarrow 0} \psi(\tau, \rho).$$

Thus (4.2) follows from (4.5) and (4.7), *q.e.d.*

LITERATURE

- [1] K. Itô: A kinematic theory of turbulence, Proc. Imp. Acad. Tokyo, **20**, No.3 (1944), pp. 120-122.
- [2] S. Itô: Positive definite functions on homogeneous spaces, Proc. Japan Acad. **26**, No.1 (1950), pp.17-28.
- [3] E. K. Haviland: On the inversion formula for Fourier-Stieltjes transforms in more one dimension, Amer. Journ. Math. **57** (1935), pp.94-100 & 382-388.
- [4] W. Ambrose: Spectral resolution of groups of unitary operators, Duke Math. Journ. **11**, No.3 (1944).
- [5] A. Weil: L'integration dans les groupes topologiques et ses applications, Act. Sci. Ind. Paris, **869** (1940).
- [6] S. Saks: Theory of the integral, Warsaw (1937).

*Mathematical Institute,
Nagoya University*