

HOLD IT!



For the treatment of Parkinson Syndrome –
“levodopa, combined with a decarboxylase
inhibitor, remains the best treatment for
most patients.”¹

In most Parkinsonian patients ‘Prolopa’:

- improves motor movement rapidly²
- achieves high serum levels quickly²
- minimizes common side effects like nausea
and vomiting
- all three ‘Prolopa’ dose forms contain
the established 4:1 ratio

The use of the 4:1 levodopa/decarboxylase
inhibitor combination has been shown to
reduce significantly the incidence of side
effects attributed to the 10:1 ratio^{3,4,5}

Prolopa[®]
(levodopa/benserazide) 4:1

Helps return the simple pleasures of living.



For brief prescribing information see page xiii

<https://doi.org/10.1017/S0317167100035356> Published online by Cambridge University Press

New **Tegretol**[®] **Chewtabs**[™] (carbamazepine)

100mg and 200mg*

- the only chewable carbamazepine
- easier titration for both children and adults
- increased convenience provided by an easily administered chewable formulation
- improved compliance arising from a pleasant tasting cherry-mint flavour

Now indicated in children aged 6 years and over



For brief prescribing information see page ix

Geigy

Mississauga, Ontario

<https://doi.org/10.1111/j.1469-7529.1100035356> Published online by Cambridge University Press

PAAB
CCPP

G-5088