

Title page

Children's adherence to a healthy and environmentally sustainable dietary pattern based on the EAT-Lancet recommendations: the role of the family environment

Healthy and Sustainable Diet in childhood

Beatriz Teixeira (Master)^{1,2,3}, Cláudia Afonso (PhD)^{1,2,3}, Andreia Oliveira (PhD)^{2,3,4}

¹Faculty of Nutrition and Food Sciences, University of Porto, Rua do Campo Alegre, 823/4150-180 Porto, Portugal;

²EPIUnit ITR, Instituto de Saúde Pública da Universidade do Porto, Universidade do Porto, [*EPIUnit ITR, Institute of Public Health of the University of Porto*] Rua das Taipas, n° 135, 4050-600 Porto, Portugal

³Laboratório para a Investigação Integrativa e Translacional em Saúde Populacional (ITR), [*Laboratory for Integrative and Translational Research in Population Health (ITR)*], Rua das Taipas 135, 4050-600 Porto, Portugal;

⁴Faculty of Medicine, University of Porto, Alameda Prof. Hernâni Monteiro, 4200-319 Porto, Portugal

Keywords: maternal characteristics; food sustainability; healthy diets; cohort studies; childhood.

Corresponding author: Beatriz Teixeira, beatrizteixeira.nutricao@gmail.com

Abbreviations: BMI: Body Mass Index; CFQ Child-Feeding Questionnaire; EFSA: European Food Safety Authority; FAO: Food and Agriculture Organization; FFQ Food Frequency Questionnaire; WISH: World Index for Sustainability and Health



This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI

10.1017/S0007114525000157

The British Journal of Nutrition is published by Cambridge University Press on behalf of The Nutrition Society

Abstract

This study aimed to investigate the association between family characteristics and adherence to the EAT-Lancet dietary recommendations in 7-year-old children. This is a prospective birth cohort study with 2062 children from Generation XXI (Porto, Portugal), who provided 3-day food diaries at age 7, used to assess habitual food consumption. At the age of 4, maternal diet was assessed using a Food Frequency Questionnaire, and a diet quality score was calculated (higher scores indicating a better maternal diet), and parental-child feeding practices were assessed with the Child-Feeding Questionnaire. Adherence to the EAT-Lancet recommendations was evaluated using the World Index for Sustainability and Health (WISH) at 7, previously adapted to pediatric age. Hierarchical linear regression models (consecutive addition of blocks of variables based on a theoretical framework) were employed to evaluate the associations between family characteristics and adherence to the WISH at age 7 [β regression coefficients and the respective 95% confidence intervals (95%CI)]. Higher maternal age and education at child's birth were associated with increased adherence to the WISH at age 7 ($\beta=0.018$, 95%CI: 0.005, 0.031; $\beta=0.038$, 95%CI: 0.024, 0.053, respectively). A better maternal diet quality, and using more restrictive practices on child's diet, at 4 years old, were both associated with higher scoring in the WISH at 7 years old ($\beta=0.033$, 95%CI: 0.018, 0.049; $\beta=0.067$, 95%CI: 0.009, 0.125, respectively). Early maternal sociodemographic and diet quality play a significant role in influencing the adherence to a healthy and environmentally sustainable dietary pattern at school-age.

Introduction

Childhood is a critical period for the development of healthy eating habits that potentially track into adulthood ⁽¹⁾. Currently, children's diets have been characterized by a high intake of energy-dense and nutritionally poor foods (as pastry products and soft drinks), and a low consumption of fruit and vegetables ⁽²⁾. Moreover, the excessive consumption of meat and dairy products, which have a high environmental footprint, coupled with an inadequate intake of plant-based foods, suggests that children's dietary patterns may not be environmentally sustainable ⁽³⁾.

The Food and Agriculture Organization (FAO) defines sustainable diets as diets that minimize environmental impacts, contribute to food and nutrition security, and promote a healthy life for present and future generations. These diets should be nutritionally adequate, safe, and health-promoting while optimizing natural and human resources ⁽⁴⁾. Various organizations have recognized the growing importance of sustainability in food policy-making ^(5; 6). Although there is a general understanding of the environmental impact of dietary patterns, based on a systematic review, there is no established tool for assessing the healthiness and sustainability of diets in the pediatric age group ⁽⁷⁾.

The EAT-Lancet recommendations on healthy diets from sustainable food systems are designed to promote a balanced and nutritious diet while considering the environmental impact of food production, aiming to achieve a sustainable food system for the future ⁽⁸⁾. Recently, the World Index for Sustainability and Health (WISH) was developed for adults, building upon the EAT-Lancet dietary recommendations ⁽⁹⁾.

Family background, and mothers in particular, play an important role in the early development of children's eating habits ⁽¹⁰⁾. This influence is recognized to begin even before birth ⁽¹¹⁾, as the diet of pregnant women, along with their body mass index (BMI), seem to influence the quality of their children's diet later in life ⁽¹²⁾. In addition, some parental feeding practices, defined as specific strategies used by parents to regulate the types and amounts of foods consumed by their children, may interfere with the development of healthy eating habits ⁽¹³⁾.

Although the association between these family characteristics and healthy dietary patterns has been previously studied ^(14; 15), to the best of our knowledge, this association has not been explored in the context of a healthy and, at the same time, environmentally sustainable dietary pattern in pediatric age. Given the importance of

prioritizing adherence to these specific diets, and understanding the associated health benefits⁽⁸⁾, it is crucial to understand the hierarchy of associated factors that may play a role in the acquisition of children's dietary patterns. This insight is of particular importance for the effective planning of public health initiatives, which aim to enhance the quality of children's diets⁽¹⁶⁾.

The objective of this study was to investigate the association between family characteristics (organized into a hierarchical theoretical framework) and adherence to the EAT-Lancet dietary recommendations in 7-year-old children from a population-based birth cohort.

Material & Methods

Study design and population

The present study included participants, who are part of the ongoing population-based birth cohort Generation XXI, which has been described in detail elsewhere^(17; 18). Mothers' resident in the Porto Metropolitan area (northern Portugal) who delivered a live-born child, with a gestational age ≥ 24 weeks, in one of the five public maternity units, between April 2005 and August 2006, were eligible for enrolment in this study. Mothers were invited to participate within 24 to 72 hours of delivery and, among those invited, 91.4% accepted. A total of 8647 infants and 8495 mothers were enrolled in the study. At the ages of 4 and 7, the families of these children were invited to participate in evaluations, with participation rates of 86% and 80%, respectively. Data were collected through face-to-face interviews or via telephone using an abbreviated version of the questionnaire when the family was unable to be physically present (20% and 15% at 4 and 7 years old, respectively).

The present study included children at the age of 7, who completed 3-day food diaries ($n = 3587$). From the initial sample, children with incomplete information on variables of interest (such as maternal feeding practices and dietary intake) were excluded from the study ($n = 1462$), resulting in a final sample size of 2125 children. The flowchart of the participant's selection is available in Figure 1. A comparison of baseline characteristics between the included participants and the remaining cohort ($n = 6522$) showed that participating mothers were slightly older (mean = 30.4 years; SD = 4.8 vs. mean=28.5, SD=5.8) and more educated (mean = 11.8 years of complete schooling years; SD = 4.3 vs. mean=10.0, SD=4.2). However, the magnitude of the differences was not substantial, as indicated by Cohen's effect size values (0.36 and

0.42, respectively)⁽¹⁹⁾. This means that the reported significant differences were likely a result of the large sample size rather than variations in the participants' characteristics.

All phases of the study complied with the Ethical Principles for Medical Research Involving Human Subjects expressed in the Declaration of Helsinki. The baseline and follow-up evaluations at 4 and 7 years of age were approved by the University of Porto Medical School/ S. João Hospital Centre Ethics Committee. Each participant's legal representative received a comprehensive explanation of the study's objectives and methodology, and written informed consent was obtained at the baseline and subsequent follow-up assessments. The baseline evaluation was approved by the Data Protection National Commission and the study follows the present EU General Data Protection Regulation under close supervision of the Data Protection Office of ISPUP.

Data Collection

A wide range of characteristics were assessed, including socioeconomic status, healthcare utilization, and lifestyle behaviors, at the child's birth, 4 and 7 years of age. These data were retrieved from clinical records or collected through in-person interviews and physical examinations, performed by trained professionals. A detailed description of the data included in this study can be found below.

At the child's birth

Birth data such as the child's sex, gestational age (in weeks), and birth weight (in grams) were retrieved from clinical records at baseline. Maternal age and completed years of schooling were asked to mothers, as well as marital status, household income, and employment status. Maternal marital status was assessed as married or cohabiting vs. single, divorced, or widowed. Monthly household income was categorized as ≤ 1000 €, 1001-2000 €, and > 2000 €. Maternal current employment status was defined as working (part or full-time, student, and working student) vs. not working (unemployed, retired, and housewife) at the time the questionnaire was completed.

At the child's birth, data on smoking habits during pregnancy were collected, categorizing mothers as smokers or non-smokers. Information on self-reported gestational diabetes during the current pregnancy was also gathered, along with

maternal height and pre-pregnancy weight to calculate BMI, as weight divided by the squared height (kg/m^2).

At the child's 4 years of age

Maternal weight was measured and recorded to the nearest 0.1 kg and height was measured without shoes to the nearest 0.1 cm to calculate maternal BMI when the child was 4 years of age.

Maternal dietary intake was evaluated using a qualitative Food Frequency Questionnaire (FFQ), adapted from a previously validated questionnaire for the Portuguese adult population⁽²⁰⁾. This FFQ assesses consumption over the past 12 months and comprises 18 items with response options on a 9-point frequency scale, ranging from “never” to “ ≥ 4 times per day”. The consumption frequencies were converted into daily frequencies (e.g., one instance per week was converted to 1/7 of a day or 0.14 instances per day).

Definition of a maternal diet quality score

Based on maternal dietary intake at the child's 4 years of age, a diet quality score previously described⁽²¹⁾ was calculated in this study. The score was defined based on 8 food components: milk, fish, red and processed meat, bread, fruit, vegetables, cakes and salty pastries, and sugar-sweetened beverages. Each component was divided into quartiles, and a scoring system was applied. Scores were assigned on a scale of 1 to 4, according to increasing quartiles of consumption (milk, fish, bread, fruits, and vegetables) or decreasing quartiles of consumption (red meat, cakes and salty pastries, and sugar-sweetened beverages). The total points assigned were summed up to derive a continuous score of diet quality, which could range from 8 to 32 points. A higher score indicates a better diet quality.

Parental child-feeding practices

Parental child-feeding practices were assessed using a combined version of the Child-Feeding Questionnaire (CFQ)⁽²²⁾ and the scales measuring overt and covert control⁽²³⁾. These were completed by the caregivers themselves (95.3% of mothers). The maternal feeding practices subscales included the following: restriction (the extent to which parents control the child's access to foods or opportunities to consume those foods) (3 items), pressure to eat (parents' insistence or demands that their children eat

more food) (4 items), monitoring (the extent to which parents track what and how much the child is eating) (3 items), overt control (represented by a firm attitude from parents about what, how much, where, and when the child eats, which can be perceived by the child) (5 items), and covert control (in which the child is unable to detect the control; for example, avoiding buying energy-dense foods) (4 items). Participants were assigned mean scores for each dimension if they provided at least 50% of the answers. The scores ranged from 1 to 5, with higher scores indicating a greater degree of practice. CFQ has previously been adapted and validated in Portuguese children from the Generation XXI cohort ⁽²⁴⁾. Furthermore, the dimensions of this questionnaire have demonstrated satisfactory reliability in a previous study ⁽²⁵⁾.

Definition of patterns of maternal child-feeding practices

Previously, principal component analysis identified three patterns of maternal child-feeding practices ⁽²⁶⁾, considering the subscales dimensions. The three patterns of maternal child-feeding practices identified by principal component analysis were labeled: "perceived monitoring" (identifying mothers with higher levels of monitoring, perceived responsibility, and overt control), "restriction" (describing mothers with higher covert control, concern about child's weight and restriction) and "pressure to eat" (defining mothers with higher levels of pressuring the child to eat more and overt control). The scores for these factors were on a standard deviation scale, ranging from -3 to 3. A score of 0 represented mothers' ratings that were close to the sample's average, while a score of -2 corresponded to maternal ratings that were two standard deviations below the sample mean ⁽²⁶⁾. These scores were subsequently calculated to each participant in the study.

At the child's 7 years of age

The family structure was classified into two types of families: "family with both parents" and "single-parent family" (meaning living only with the mother or the father) at the age 7. Information about whether they lived or not with siblings was also obtained. Monthly household income was updated and categorized as ≤ 1000 €, 1001-2000 €, and > 2000 €. In addition, information on children's lifestyles, such as their engagement in regular sports activities, were collected.

A physical examination of the children was performed, including their height and weight, measured by trained professionals following standard procedures ⁽²⁷⁾.

Weight was measured to the nearest tenth of a kilogram (Tanita® digital scale, Arlington Heights, IL, USA) in light clothing and without shoes, and height was measured to the nearest tenth of a centimeter (Seca® wall stadiometer, Hamburg, Germany). Body mass index (BMI) was calculated and children were classified according to age- and sex-specific BMI z-scores (BMIZ) developed by the World Health Organization (WHO). Overweight (pre-obesity plus obesity) was defined as a BMI for age z-score $>+1$ SD ⁽²⁸⁾.

At age 7, dietary intake of children were collected by a 3-day food diary (2 weekdays and 1 weekend day), completed by parents or other main caregivers, prior to the face-to-face interview, as previously described ⁽²⁹⁾. Data were collected throughout all four seasons (winter, autumn, spring, and summer) from 2012 to 2014. Oral (by telephone) and written instructions were provided for the correct completion of the food diaries and for the quantification of food portions. Parents were instructed on the use of household measures and standard units for the quantification of food portions, and photographic examples were provided for more accurate assessment. They were also asked to provide detailed descriptions of each food and drink consumed by the child, including the method of preparation, recipes and place of consumption, whenever possible. It was advised to let children follow their usual diet and to ask for the help of other caregivers in case the child was out of the home during the day. During the face-to-face interview, the fieldwork team was responsible for receiving and thoroughly inspecting the food diaries to ensure their completeness. In cases where any information was missing, the team would request a revised report to be sent by mail at a later time.

After, the information was handled by a team of trained nutritionists. Energy and nutrient intake were calculated using the methodology of the software eAT24 (Electronic Assessment Tool for 24-hours recall) ⁽³⁰⁾. This software was previously validated ⁽³⁰⁾ and includes the harmonized food classification and description system proposed by the European Food Safety Authority (EFSA) - the FoodEx2 classification system ⁽³¹⁾ and food composition tables from Portugal and other sources to allow the conversion into nutrients ⁽³²⁾.

Definition of the WISH dietary pattern

To measure the adherence to the EAT-Lancet recommendations, an adapted version for pediatric age of the WISH score ⁽³³⁾ was applied to the food consumption data. The WISH, originally developed for adults ⁽⁹⁾, consists of 13 food groups: grains,

vegetables, fruits, dairy, red meat, fish, eggs, chicken and other poultry, legume grains, nuts, unsaturated fats, saturated fats, soft drinks, and added sugars. Scores for each component can range from 0 to 10, reflecting adherence to the EAT-Lancet recommendations. The total score, ranging from 0 to 130, indicates the overall health and environmental sustainability of the diet, with higher scores indicating a healthier and more environmentally sustainable diet ⁽⁹⁾. WISH was estimated for each participant based on the individual consumption of each food group and the recommended thresholds of the EAT-Lancet Commission. The details of each component, scoring criteria, and their respective recommendations are presented in Table A of supplementary material.

Statistical analysis

Continuous variables were summarized by means and standard deviations (SD) and categorical variables by counts and proportions.

To address the aim of this study more effectively, a theoretical framework (Figure 2) based on previous models ^(34; 35; 36) and ensuring time precedence between blocks of variables was established. Firstly, we conducted a comprehensive review of the literature to identify relevant variables related to adherence to a healthy diet in childhood ^(12; 13; 35; 37; 38). Then, we checked which variables were available within the Generation XXI cohort with potential to be included in this framework, based on the theoretical and empirical relevance found. This predefined theoretical framework was based on (a) socio-economic characteristics at the child's birth, (b) maternal characteristics before and during pregnancy, (c) maternal diet and feeding practices at the child's 4 years of age, and (d) family characteristics at the child's 7 years of age. Variables were organized to reflect temporal precedence, ensuring that socio-economic characteristics at the child's birth influence maternal characteristics during pregnancy, which in turn impact feeding practices and, finally, family characteristics at 7 years of age.

Associations between exposures (mother and other family characteristics) and adherence to the adapted WISH at 7 years of age were measured through hierarchical linear regressions models, obtaining crude and adjusted β -regression coefficients and the respective 95% confidence intervals (95% CI). A step-by-step approach was used. Firstly, univariate models were applied for each variable considered in the framework. Only variables that reached statistical significance were then conceptually grouped into

blocks (blocks a, b, c and d previously described). Multivariable models were then fitted for these blocks, with variables within each block being mutually adjusted. Considering a high correlation between maternal education years and the household income both at baseline and when the child reached 7 years of age, the inclusion of maternal education was chosen for the adjusted models as it yielded the highest improvement in Nagelkerke's R^2 . Models were further adjusted for child's sex and sports practice at the age of 7. No significant child's sex interaction was observed in the associations under study; hence, stratification was not applied in the models.

The hierarchical linear regression analysis used the coefficient of determination, Nagelkerke's R^2 , to evaluate the factors (i.e. characteristics) with the greatest relevance for the children's dietary pattern, meaning a higher score (greater adherence) to the WISH⁽³⁹⁾.

A sensitivity analysis was conducted to examine the associations between mother and other family characteristics and the WISH score into quartiles (instead of a continuous variable), using a multinomial logistic regression, where all four blocks were mutually adjusted.

A significance level of 5% was assumed, as well as independence between observations. Analyses were performed in the IBM® SPSS® Statistics version 28.

Results

At baseline, mothers had a mean age of 30.4 years (SD=4.8) and 11.8 years of schooling (SD=4.3). The majority of mothers were married or cohabiting (96.4%) and employed (84.3%). Before pregnancy, mothers had a mean BMI of 23.9 kg/m² (SD=4.2), and 15.9% smoked during pregnancy. At the child's 4 years of age, mothers had a mean diet quality score of 21.2 points (SD=4.0, ranging from 8 to 31; possible range 8-32). Approximately half of the children were males (51.8%), and at 7 years of age the majority of them lived with both parents (85.7%) and had siblings (60.8%). The average WISH score was 59.6 points (SD=13.8, ranging from 10.6 to 100; possible range 0-130) (Table 1).

The food consumption of this sample, in g/day, according to the 13 food groups evaluated in the WISH score, is presented in Table B of supplementary material. The high children's consumption (in grams) of red and processed meats (Median = 65.2 percentile 25-75=41.6-89.6) and dairy (Median = 504.4 percentile 25-75=385.2-626.7)

is highlighted, as well as the low consumption (in grams) of legume grains (Median = 6.5 percentile 25-75=0.0- 14.4) and nuts (Median = 0.0 percentile 25-75=0.0-0.0).

The consumption of these 13 food groups according to the distribution of the WISH score by quartiles is also described in Table C of the supplementary material. In addition, the sample characteristics according to the WISH quartiles are described in Supplementary Table D.

Table 2 presents the associations between maternal and other family characteristics and adherence to the adapted WISH at the age of 7. In statistical analysis, a predefined theoretical framework (Figure 2) was followed by the sequential addition of blocks of variables into the model. Firstly, the socioeconomic characteristics at the child's birth were analyzed (Model 1). It was found that maternal age and education had a significant overall effect, with increasing age and education, resulting in a significantly higher score of the adapted WISH at age 7. These characteristics (maternal age, education, marital status, and work status) collectively accounted for 31% of the variance of the outcome under study (WISH score), as determined by the Nagelkerke's R^2 . The inclusion of pre-pregnancy BMI did not yield significant effects on the associations under study, as the previous estimates remained unchanged (Model 2). The addition of maternal diet and feeding practices at the age of 4 (Model 3) showed positive associations between the maternal diet quality score and the use of restrictive feeding practices and adherence to the adapted WISH at age 7. The inclusion of this set of variables led to an 11% increase in the model's explanatory power (Nagelkerke's $R^2=42\%$). Lastly, family characteristics at 7 years old were added (Model 4). However, the inclusion of these variables did not result in any significant changes in the model estimates.

In the final model, maternal age and education at baseline (i.e. child's birth) were associated with a higher adherence to the adapted WISH at age 7 ($\beta=0.018$, 95%CI: 0.005, 0.031; $\beta=0.038$, 95%CI: 0.024, 0.053, respectively). Also, when the child was 4, the maternal diet quality score and the use of more restrictive feeding practices were associated with increased scoring in the adapted WISH at age 7 ($\beta=0.033$, 95%CI: 0.018, 0.049; $\beta=0.067$, 95%CI: 0.009, 0.125, respectively). Despite the absence of a sex interaction in the associations under study, particularly in this analysis, distinct results were exhibited for girls and boys, with more restrictive feeding practices being significantly associated with a healthier and more environmentally

sustainable dietary pattern only in girls ($\beta=0.093$, 95% CI: 0.014, 0.172) (results only shown in text).

In a sensitivity analysis, associations were tested considering the WISH score distributed into quartiles (instead of a continuous variable) and results are summarized in Supplementary Table E D. Similar results were obtained for the upper quartile of the WISH score, giving robustness to our findings.

To identify the key factors influencing the children's dietary pattern, variables significantly associated with the WISH in Model 4 were systematically removed one by one from the final model. Maternal education (assessed at baseline, in completed years of schooling) and the maternal diet quality (at the child's 4 years old) were the key factors associated with the adherence to the adapted WISH at age 7. The removal of these variables from the final model produced the biggest drop in Nagelkerke's R^2 (full vs. reduced model: $R^2=42\%$ vs. $R^2=33\%$ and $R^2=42\%$ vs. $R^2=37\%$, respectively) (results presented only in text).

Discussion

The main findings from this study suggest that maternal characteristics are associated with greater adherence to a healthy and environmentally sustainable diet at age 7. Maternal age and education at the child's birth were positively associated with the adapted WISH score. At age 4, the maternal diet quality score and the use of more restrictive feeding practices (including dimensions of covert control, i.e. a type of control not perceived by the child, concerns about the child's weight and food restriction) were associated with a higher adherence to the WISH when they were 7 years old. A methodology of consecutive addition of blocks of variables into the models suggests that maternal education at the child's birth and maternal diet quality at the child's 4 years of age were the key factors associated with children's adherence to the adapted WISH.

According to the WHO, socioeconomic characteristics significantly influence lifestyles and overall well-being⁽³⁵⁾. Individuals' socioeconomic status determines their exposure and vulnerability to health-compromising conditions⁽³⁵⁾. Similarly, it is well documented that maternal sociodemographic characteristics, including age, education level, and income, are positively correlated with the promotion of healthier dietary habits during childhood⁽¹⁶⁾. Indeed, in this study, these characteristics account for

approximately one-quarter of the variance of the WISH score. Specifically, mothers who are older and more educated have children with healthier and more environmentally sustainable diets at school age. Additionally, maternal education at the child's birth was identified as one of the key factors influencing children's diet. These findings are consistent with those reported in the literature regarding healthy dietary patterns^(21; 37).

A recent systematic review⁽⁴⁰⁾ has showed that a higher pre-pregnancy BMI in mothers has been associated with an increased risk of their children developing unhealthy eating habits. These include consuming a diet high in ultra-processed foods, added sugars, and unhealthy fats, which are typically considered foods with a high environmental footprint. In this study, a higher pre-pregnancy BMI was associated with a lower adherence to the EAT-Lancet dietary recommendations. However, in the adjusted model, this result lost both magnitude and significance, suggesting that the initial association might be explained by maternal age and education at baseline (covariates in the adjusted model).

In the present study, there was an inverse association between smoking during pregnancy and adherence to the adapted WISH at age 7, although this result was not statistically significant. According to the literature, smoking during pregnancy has been associated with childhood obesity⁽³⁸⁾ which, in turn, is known to be closely linked to inadequate eating habits⁽⁴¹⁾. Despite this, this effect appears to be greater in children up to 5 years old⁽⁴²⁾. Considering pregnancy-related events, children born from mothers with gestational diabetes mellitus are at a high risk of developing various health problems, including obesity⁽⁴³⁾. In this study, no significant association was found between maternal smoking during pregnancy and the dietary patterns of children at 7 years of age. Further studies are necessary to elucidate these associations.

Maternal diet quality after pregnancy can have a significant impact on the dietary pattern of the child as they grow. Mothers who maintain a healthy and balanced diet and good eating behaviors are more likely to have children who adopt similar dietary patterns⁽⁴⁴⁾. Indeed, in this study, the maternal diet quality at the child's 4 years of age was identified as one of the key factors influencing a greater adherence to the WISH at age 7. Despite not assessing children or mothers' dietary patterns, a previous study⁽⁴⁵⁾ has showed similar associations considering only some food groups.

It is important to highlight that not only the maternal diet during pregnancy plays a pivotal role in the optimal growth and development of the child⁽¹²⁾. This importance persists over time. The preschool period is a period of heightened vulnerability and dependence for parents, particularly mothers, who serve as important role models⁽¹⁰⁾. During this period of vulnerability and importance, children not only learn social and behavioral cues from their parents, particularly their mothers, but they also consolidate their relationship with food⁽¹⁶⁾. The dietary choices and eating behaviors observed and adopted by children from their caregivers can have a long-lasting impact on their future eating habits and overall approach to nutrition⁽¹⁶⁾. Consequently, it is recommended that intervention strategies be implemented to improve the dietary habits of mothers, with the objective of promoting better dietary habits over time.

It is well-established that parental child-feeding practices are closely related to eating habits in childhood⁽¹³⁾. The present study observed that children whose mothers employed more restrictive feeding practices at 4 years of age exhibited a higher score on the adapted WISH at 7 years of age. In a previous study conducted in Generation XXI, it was observed that restricting the child's access to unhealthy foods, using covert control, a kind of control not perceived by the child, and expressing concerns about the child's weight at the age of 4 were inversely associated with the 'Energy-dense foods' dietary pattern (a dietary pattern with foods typically characterized by a high environmental footprint) after 3 years⁽⁴⁶⁾. These findings are consistent with those of a Dutch birth cohort study that observed that following a restrictive feeding practice was associated with a lower intake of sweets, cookies, and chocolates, and a higher intake of fruits and vegetables⁽⁴⁷⁾. This dietary pattern aligns with a more environmentally sustainable pattern. In light of the potential for a sex interaction in the associations under study, previous research has documented that parents tend to employ more restrictive practices with girls than with boys⁽⁴⁸⁾. In this study, when this analysis was stratified by sex, this association remained as statistically significant only among girls, suggesting a sex modification effect on this association in particular, as previously reported in the literature.

Nevertheless, research on the relationship between parental restriction and children's diet has yielded conflicting results, suggesting a complex nature. When critically analyzing this association, it is important to mention that some studies have found an association between restriction and higher childhood BMI⁽⁴⁹⁾. In other words,

it has been proposed that mothers are more likely to adopt these feeding practices in response to their child's BMI, rather than the other way around ⁽⁴⁹⁾. The practice of restriction may originate from parental concerns about unhealthy weight development in children ⁽⁵⁰⁾. However, its potential beneficial effects remain uncertain, and it has been linked to the emergence of disordered eating behaviors in adolescence, particularly among girls ⁽⁵¹⁾. Consequently, while these practices may have a positive impact on children's diets in the short term, they may also have adverse effects in the long term.

When interpreting the results of this study, it should be taken into account several limitations. The exclusion of participants due to missing data could introduce a selection bias. However, we conducted a sensitive analysis comparing participants and non-participants and the Cohen's effect size values were not particularly high ⁽¹⁹⁾. This leads us to conclude that the significance of differences between this sample and the remaining cohort are likely due to the large sample size rather than substantial differences between participants. Indeed, when compared to the national data from the National Institute of Statistics, the mean age of women at the time of birth is close to the average age of mothers in this study (29.2 vs. 30.4 years) ⁽⁵²⁾. Conversely, this sample exhibits a slightly higher level of education than the national data for women aged 25 to 34 years (35.3% vs. 33.1% with higher education) ⁽⁵³⁾. It should be noted that this sample is not representative of Portuguese children, thus caution should be taken when extrapolating the results. Also, it is important to note that the cohort was recruited in 2005/2006, and the assessment at 7 years took place over 10 years ago. Therefore, the socioeconomic characteristics of postpartum women in 2024 may differ, and it is also possible that infant feeding practices have changed. As such, the results of this study should be interpreted carefully.

In addition, it is important to acknowledge the limitation of using a short-list FFQ to assess maternal diet, which relies heavily on participants' memory and trustworthiness, potentially leading to recall bias and inaccurate reporting. FFQs typically offer a limited range of food items, potentially omitting fewer common foods and underestimating true consumption. However, it is important to note that the FFQ used in this study was adapted from a previously validated questionnaire ^(20; 54), and its food groups were combined, with less desegregation than the original FFQ, but covered the same food groups, which makes its usage acceptable. Notably, the CFQ consists of parent-report measures, which could also introduce measurement error owing to social

desirability effects and inherent subjectivity⁽⁵⁵⁾. However, the questionnaire has shown good psychometric properties in children from this cohort⁽²⁴⁾.

Finally, despite several variables have been tested in the multivariable analysis, we cannot fail to exclude any residual confounding. For instance, the influence of the school environment on children's diet was not assessed, and we know that schools play a crucial role in developing healthy eating habits during childhood⁽⁵⁶⁾.

This study is notable for its robust methodology, including a relatively large sample size and comprehensive assessment of potential confounding variables. Furthermore, the study is prospective, which ensures temporal sequence between variables. A significant benefit of this study is the use of a predefined theoretical framework for analysis and the assessment of children's diets through a reference method, such as the 3-day food records. This approach has been described as one of the most accurate methods for estimating food consumption in children⁽⁵⁷⁾. Moreover, to the best of our knowledge, this study is the first to assess maternal and other family determinants influencing a healthy and environmentally sustainable dietary pattern in childhood, specifically aligned with the EAT-Lancet recommendations on healthy diets from sustainable food systems. Given the growing importance of sustainability in shaping food policies⁽⁶⁾, this study contributes valuable insights for policymakers and researchers seeking to promote sustainable and healthy dietary habits among children. However, further research with larger and more diverse samples is necessary to confirm and extend our findings.

In conclusion, this study suggests that a higher maternal education (at the child's birth) and the maternal diet quality (at the child's 4 years of age) were the key factors associated with increased adherence to the EAT-Lancet recommendations on healthy diets from sustainable food systems at age 7. Given the longitudinal nature of these data, it would be interesting in future studies to analyze the main determinants of children's diets as they grow, as it is expected that the contextual determinants will change throughout the life course.

Our results seem to emphasize the importance of promoting better maternal dietary habits, particularly among less educated mothers, to foster healthier dietary patterns in school-age children.

Acknowledgement: We gratefully acknowledge the families enrolled in Generation XXI for their kindness, the participating hospitals and their staff for their help and

support, and all previous and current members of the research and field team for their enthusiasm and perseverance.

Funding statement: Generation XXI was funded by Programa Operacional de Saúde–Saúde XXI, Quadro Comunitário de Apoio III and Administração Regional de Saúde Norte (Regional Department of Ministry of Health). It has support from the Portuguese Foundation for Science and Technology (FCT) through the projects with references UIDB/04750/2020 and LA/P/0064/2020 and DOI identifiers <https://doi.org/10.54499/UIDB/04750/2020> and <https://doi.org/10.54499/LA/P/0064/2020>. The first author (BT) had a PhD grant 2021.05133.BD (<https://doi.org/10.54499/2021.05133.BD>), funded by the Foundation for Science and Technology (FCT) and FSE (Fundo Social Europeu) Program. The authors also acknowledge the SYSTEMIC project (n°696295), a knowledge hub on Nutrition and Food Security, in collaboration with JPI-HDHL (Healthy Diet for a Healthy Life), FACCE-JPI (Joint Programming Initiative for Agriculture, Climate Change, and Food Security), JPI-OCEANS (Joint Programming Initiative Healthy and Productive Seas and Oceans).

Declaration of interests: The authors declare none.

Author Contribution: **BT:** Conceptualization, formal analysis, investigation, methodology, writing - original draft; **CA:** investigation, methodology, supervision, validation, visualization, writing - review & editing; **AO:** Conceptualization, formal analysis, investigation, methodology, project administration, supervision, validation, visualization, writing - review & editing. All authors read and approved the final version of the manuscript.

References

1. Craigie AM, Lake AA, Kelly SA *et al.* (2011) Tracking of obesity-related behaviours from childhood to adulthood: A systematic review. *Maturitas* **70**, 266-284. doi: 10.1016/j.maturitas.2011.08.005
2. Williams J, Buoncristiano M, Nardone P *et al.* (2020) A Snapshot of European Children's Eating Habits: Results from the Fourth Round of the WHO European Childhood Obesity Surveillance Initiative (COSI). *Nutrients* **12**, 2481. doi: 10.3390/nu12082481.
3. Stehfest E, Bouwman L, van Vuuren DP *et al.* (2009) Climate benefits of changing diet. *Climatic Change* **95**, 83-102. doi: 10.1007/s10584-008-9534-6.
4. Food Agriculture Organization (2012) *Sustainable Diets and Biodiversity Directions and Solutions for Policy, Research and Action*. FAO Bioversity International. E-ISBN 978-92-5-107288-2.
5. IOM (Institute of Medicine). (2014) *Sustainable diets: Food for healthy people and a healthy planet: Workshop summary*. Washington, DC: The National Academies Press. ISBN-13: 978-0-309-29667-0
6. Joseph H, Clancy K (2015) Dietary guidelines and sustainable diets: pathways to progress. In: *Advancing health and well-being in food systems: strategic opportunities for funders*. *Global Alliance for the Future of Food*, 88-107.
7. Teixeira B, Afonso C, Rodrigues S *et al.* (2021) Healthy and Sustainable Dietary Patterns in Children and Adolescents: A Systematic Review. *Advances in Nutrition*. **13**, 1144-1185. doi: 10.1093/advances/nmab148.
8. Willett W, Rockström J, Loken B *et al.* (2019) Food in the Anthropocene: the EAT Lancet Commission on healthy diets from sustainable food systems. *The Lancet* **393**, 447-492. doi: 10.1016/S0140-6736(18)31788-4.
9. Trijsburg L, Talsma EF, Crispim SP *et al.* (2020) Method for the Development of WISH, a Globally Applicable Index for Healthy Diets from Sustainable Food Systems. *Nutrients* **13**, 93. doi: 10.3390/nu13010093.
10. Savage JS, Fisher JO, Birch LL (2007) Parental influence on eating behavior: conception to adolescence. *J Law Med Ethics* **35**, 22-34. doi: 10.1111/j.1748-720X.2007.00111.x.
11. Ventura AK, Worobey J (2013) Early influences on the development of food preferences. *Curr Biol* **23**, R401-408. doi: 10.1016/j.cub.2013.02.037.

12. Biagi C, Nunzio MD, Bordoni A *et al.* (2019) Effect of Adherence to Mediterranean Diet during Pregnancy on Children's Health: A Systematic Review. *Nutrients* **11**, 997. doi: 10.3390/nu11050997.
13. Yee AZ, Lwin MO, Ho SS (2017) The influence of parental practices on child promotive and preventive food consumption behaviors: a systematic review and meta-analysis. *Int J Behav Nutr Phys Act* **14**, 47. doi: 10.1186/s12966-017-0501-3.
14. Vidhyashree MD, Raveendran SR, Priya RL *et al.* (2015) A Review of Family and Social Determinants of Children's Eating Patterns and Diet Quality. *Research Journal of Pharmaceutical Biological and Chemical Sciences* **6**, 1196-1201.
15. Favara G, Maugeri A, San Lio RM *et al.* (2024) Exploring Gene-Diet Interactions for Mother-Child Health: A Systematic Review of Epidemiological Studies. *Nutrients* **16**, 994. doi: 10.3390/nu16070994.
16. Scaglioni S, De Cosmi V, Ciappolino V *et al.* (2018) Factors Influencing Children's Eating Behaviours. *Nutrients* **10**, 706. doi: 10.3390/nu10060706.
17. Larsen PS, Kamper-Jørgensen M, Adamson A *et al.* (2013) Pregnancy and birth cohort resources in europe: a large opportunity for aetiological child health research. *Paediatr Perinat Epidemiol* **27**, 393-414. doi: 10.1111/ppe.12060.
18. Alves E, Correia S, Barros H *et al.* (2012) Prevalence of self-reported cardiovascular risk factors in Portuguese women: a survey after delivery. *Int J Public Health* **57**, 837-847. doi: 10.1007/s00038-012-0340-6.
19. Husted JA, Cook RJ, Farewell VT *et al.* (2000) Methods for assessing responsiveness: a critical review and recommendations. *J Clin Epidemiol* **53**, 459-468. doi: 10.1016/s0895-4356(99)00206-1.
20. Lopes C, Aro A, Azevedo A *et al.* (2007) Intake and adipose tissue composition of fatty acids and risk of myocardial infarction in a male Portuguese community sample. *J Am Diet Assoc* **107**, 276-286. doi: 10.1016/j.jada.2006.11.008.
21. Durão C, Severo M, Oliveira A *et al.* (2017) Association of maternal characteristics and behaviours with 4-year-old children's dietary patterns. *Matern Child Nutr* **13**. doi: 10.1111/mcn.12278.
22. Birch LL, Fisher JO, Grimm-Thomas K *et al.* (2001) Confirmatory factor analysis of the Child Feeding Questionnaire: a measure of parental attitudes, beliefs and practices about child feeding and obesity proneness. *Appetite* **36**, 201-210. doi: 10.1006/appe.2001.0398.

23. Ogden J, Reynolds R, Smith A (2006) Expanding the concept of parental control: a role for overt and covert control in children's snacking behaviour? *Appetite* **47**, 100-106. doi: 10.1016/j.appet.2006.03.330.
24. Real H, Oliveira A, Severo M *et al.* (2014) Combination and adaptation of two tools to assess parental feeding practices in pre-school children. *Eating Behaviors* **15**, 383-387. doi: 10.1016/j.eatbeh.2014.04.009.
25. Durão C, Andreozzi V, Oliveira A *et al.* (2015) Maternal child-feeding practices and dietary inadequacy of 4-year-old children. *Appetite* **92**, 15-23. doi: 10.1016/j.appet.2015.04.067.
26. Moreira I, Severo M, Oliveira A *et al.* (2016) Social and health behavioural determinants of maternal child-feeding patterns in preschool-aged children. *Matern Child Nutr* **12**, 314-325. doi: 10.1111/mcn.12132.
27. Gibson R (2005) *Principles of Nutritional Assessment: Oxford University Press*. Oxford, New York: Oxford University Press 2nd ed. ISBN: 9780195171693.
28. de Onis M OA, Borghi E, Siyam A *et al.* (2007) Development of a WHO growth reference for school-aged children and adolescents. *Bull World Health Organ* **85**, 660-7. doi: 10.2471/blt.07.043497.
29. Moreira T, Severo M, Oliveira A *et al.* (2015) Eating out of home and dietary adequacy in preschool children. *Br J Nutr* **114**, 297-305. doi: 10.1017/S0007114515001713.
30. Goios AC, Severo M, Lloyd AJ *et al.* (2020) Validation of a new software eAT24 used to assess dietary intake in the adult Portuguese population. *Public Health Nutr* **23**, 3093-3103. doi: 10.1017/S1368980020001044.
31. European Food Safety Authority (2011) *Report on the development of a Food Classification and Description System for exposure assessment and guidance on its implementation and use*. European Food Safety Authority doi: 10.2903/j.efsa.2011.2489.
32. Lopes C, Torres D, Oliveira A *et al.* (2018) National Food, Nutrition, and Physical Activity Survey of the Portuguese General Population (2015-2016): Protocol for Design and Development. *JMIR Research Protocols* **7**, e42. doi: 10.2196/resprot.8990.
33. Teixeira B, Afonso C, Severo M *et al.* (2024) Exploring dietary patterns and their association with environmental sustainability and body mass index in children and adolescents: Insights from the National Food, nutrition and physical activity survey 2015-2016. *Sci Total Environ*, 174051. doi: 10.1016/j.scitotenv.2024.174051

34. Victora CG, Huttly SR, Fuchs SC *et al.* (1997) The role of conceptual frameworks in epidemiological analysis: a hierarchical approach. *Int J Epidemiol* **26**, 224-227. doi: 10.1093/ije/26.1.224.
35. World Health Organization (2010) A conceptual framework for action on the social determinants of health. *World Health Organization*
<https://apps.who.int/iris/handle/10665/44489>.
36. UNICEF (1990) Strategy for improved nutrition of children and women in developing countries. *Indian J Pediatr.* **58**, 13–24. doi: 10.1007/BF02810402.
37. Wen X, Kong KL, Eiden RD *et al.* (2014) Sociodemographic differences and infant dietary patterns. *Pediatrics* **134**, e1387-1398. doi: 10.1542/peds.2014-1045.
38. Suzuki K, Ando D, Sato M *et al.* (2009) The association between maternal smoking during pregnancy and childhood obesity persists to the age of 9-10 years. *J Epidemiol* **19**, 136-142. doi: 10.2188/jea.JE20081012.
39. NAGELKERKE NJD (1991) A note on a general definition of the coefficient of determination. *Biometrika* **78**, 691-692. doi: 10.1093/biomet/78.3.691.
40. Mannino A, Sarapis K, Moschonis G (2022) The Effect of Maternal Overweight and Obesity Pre-Pregnancy and During Childhood in the Development of Obesity in Children and Adolescents: A Systematic Literature Review. *Nutrients* **14**. doi: 10.3390/nu14235125.
41. Liberali R, Kupek E, Assis MAA (2020) Dietary Patterns and Childhood Obesity Risk: A Systematic Review. *Child Obes* **16**, 70-85. doi: 10.1089/chi.2019.0059.
42. Suzuki K, Sato M, Ando D *et al.* (2013) Differences in the effect of maternal smoking during pregnancy for childhood overweight before and after 5 years of age. *J Obstet Gynaecol Res* **39**, 914-921. doi: 10.1111/jog.12025.
43. Dugas C, Perron J, Kearney M *et al.* (2017) Postnatal Prevention of Childhood Obesity in Offspring Prenatally Exposed to Gestational Diabetes mellitus: Where Are We Now. *Obesity Facts* **10**, 396-406. doi: 10.1159/000477407.
44. Juton C, Lerin C, Homs C *et al.* (2021) Prospective Associations between Maternal and Child Diet Quality and Sedentary Behaviors. *Nutrients* **13**. doi: 10.3390/nu13051713.
45. Johnson SL (2016) Developmental and Environmental Influences on Young Children's Vegetable Preferences and Consumption. *Adv Nutr* **7**, 220s-231s. doi: 10.3945/an.115.008706.

46. Barbosa C, Lopes C, Costa A *et al.* (2023) Parental child-feeding practices at 4 years of age are associated with dietary patterns of 7-year-olds. *Journal of Human Nutrition and Dietetics* **36**, 1339-1348. doi: 10.1111/jhn.13151.
47. Gubbels JS, Kremers SP, Stafleu A *et al.* (2009) Diet-related restrictive parenting practices. Impact on dietary intake of 2-year-old children and interactions with child characteristics. *Appetite* **52**, 423-429. doi: 10.1016/j.appet.2008.12.002.
48. Gubbels JS, Kremers SP, Stafleu A *et al.* (2011) Association between parenting practices and children's dietary intake, activity behavior and development of body mass index: the KOALA Birth Cohort Study. *Int J Behav Nutr Phys Act* **8**, 18. doi: 10.1186/1479-5868-8-18.
49. Quah PL, Ng JC, Fries LR *et al.* (2019) Longitudinal Analysis Between Maternal Feeding Practices and Body Mass Index (BMI): A Study in Asian Singaporean Preschoolers. *Front Nutr* **6**, 32. doi: 10.3389/fnut.2019.00032.
50. Beckers D, Karssen LT, Vink JM *et al.* (2021) Food parenting practices and children's weight outcomes: A systematic review of prospective studies. *Appetite* **158**, 105010. doi: 10.1016/j.appet.2020.105010.
51. Loth KA, MacLehose RF, Fulkerson JA *et al.* (2014) Are food restriction and pressure-to-eat parenting practices associated with adolescent disordered eating behaviors? *Int J Eat Disord* **47**, 310-314. doi: 10.1002/eat.22189.
52. Instituto Nacional de Estatística (2011) Maternal age at birth. *Instituto Nacional de Estatística*
https://www.inept.pt/xportal/xmain?xpid=INE&xpgid=ine_indicadores&indOcorrCod=0008219&contexto=bd&selTab=tab2&xlang=pt (accessed Jul 2023).
53. Instituto Nacional de Estatística (2011) Rate of higher education enrollment among women aged between 25 and 34 years.
https://www.inept.pt/xportal/xmain?xpid=INE&xpgid=ine_indicadores&contexto=pi&indOcorrCod=0006400&selTab=tab0 (accessed Jul 2023).
54. Pinto E, Severo M, Correia S *et al.* (2010) Validity and reproducibility of a semi-quantitative food frequency questionnaire for use among Portuguese pregnant women. *Matern Child Nutr* **6**, 105-119. doi: 10.1111/j.1740-8709.2009.00199.x.
55. Kimberlin CL, Winterstein AG (2008) Validity and reliability of measurement instruments used in research. *Am J Health Syst Pharm* **65**, 2276-2284. doi: 10.2146/ajhp070364.

56. Cohen JFW, Hecht AA, McLoughlin GM *et al.* (2021) Universal School Meals and Associations with Student Participation, Attendance, Academic Performance, Diet Quality, Food Security, and Body Mass Index: A Systematic Review. *Nutrients* **13**. doi: 10.3390/nu13030911.

57. Burrows TL, Martin RJ, Collins CE (2010) A systematic review of the validity of dietary assessment methods in children when compared with the method of doubly labeled water. *J Am Diet Assoc* **110**, 1501-1510. doi: 10.1016/j.jada.2010.07.008.

Figure Captions





Figure 1. Flowchart of participant's selection (Generation XXI).

**Caption:**

This theoretical framework was based on previous models (34, 35, 36).

BMI: Body Mass Index; WISH: World Index for Sustainability and Health

Figure 2. Theoretical framework of maternal and other family determinants of children's dietary pattern.

Table 1. Characteristics of families and their 7-year-olds from the Generation XXI birth cohort with 3-day food diaries and complete data in variables of interest (n=2125).

<i>Family socioeconomic characteristics at child's birth</i>	
Maternal age*, years, mean (SD)	30.4 (4.8)
Maternal education*, school years, mean (SD)	11.8 (4.3)
Household monthly income, n (%) [n=2059]	
≤ 1000 €	590 (28.7)
1001 – 2000 €	949 (46.1)
>2000 €	385 (18.7)
Does not know	109 (5.3)
Prefer not to say	26 (1.3)
Marital status*, n (%)	
Married or cohabiting	2049 (96.4)
Single, divorced or widowed	76 (3.6)
Work status*, n (%)	
Working	1791 (84.3)
Not working	334 (15.7)
<i>Maternal characteristics before/during pregnancy</i>	
BMI before pregnancy*, kg/m ² , mean (SD)	23.9 (4.2)
Smoking during pregnancy*, n (%) [n=2047]	335 (15.9)
Gestational diabetes*, n (%)	154 (7.2)
<i>Maternal characteristics at child's 4 years</i>	
Diet quality score* ^a , mean (SD)	21.2 (4.0)
Feeding practices*, z-scores ^b	
Perceiving monitoring, median (p25, p75)	0.08 (-0.6, 0.8)
Restriction, median (p25, p75)	0.08 (-0.6, 0.7)
Pressure to eat, median (p25, p75)	0.06 (-0.7, 0.6)
Maternal BMI*, kg/m ² , mean (SD)	25.0 (4.6)
<i>Family characteristics at 7 years</i>	
Family structure*, n (%)	
Single-parent family	304 (14.3)
Family with both parents	1821 (85.7)
Living with siblings*, n (%)	1292 (60.8)

Household monthly income*, n (%)	
≤ 1000 €	452 (21.3)
1001 – 2000 €	1058 (49.8)
>2000 €	585 (27.5)
Does not know	25 (1.2)
Prefer not to say	5 (0.2)
<i>Children's characteristics</i>	
Sex*, n (%)	
Female	1024 (48.2)
Male	1101 (51.8)
Birth weight*, grams, mean (SD)	3159.4 (522.1)
Gestational age*, weeks, mean (SD)	38.5 (1.9)
BMI at 7 years ^c , n (%) [n=2120]	
Overweight (z-score >+1 SD)	736 (34.7)
Normal weight and thinness (z-score <+1 SD)	1384 (65.3)
Sports practice at 7 years*, n (%)	1832 (86.2)
WISH score at 7 years ^{*,d} , mean (SD)	59.6 (13.8)

* Data on these variables are available for all the participants in this analysis (n=2125).

^a possible range from 8 to 32 (21); ^b possible range from -3 to 3 (26); ^c defined according to the World Health Organization (28); ^d possible range from 0 to 130 (33).

BMI, Body Mass Index; p: percentile; WISH, World Index for Sustainability and Health; SD: standard deviations; %: percentage; €: euro

Table 2. Hierarchical multivariable linear regression associations between maternal and other family characteristics and adherence to the World Index for Sustainability and Health (WISH) in 7-year-olds from the Generation XXI birth cohort (n=2125).

		Unadjusted Model	Model 1*	Model 2*	Model 3*	Model 4*
		β (95% CI)				
BLOCK 1	<i>Socioeconomic characteristics at the child's birth</i>					
	Maternal age, years	0.032 (0.019, 0.044)	0.024 (0.012, 0.037)	0.024 (0.012, 0.037)	0.018 (0.006, 0.031)	0.018 (0.005, 0.031)
	Maternal education, years	0.048 (0.034, 0.062)	0.048 (0.035, 0.062)	0.048 (0.034, 0.062)	0.041 (0.027, 0.056)	0.038 (0.024, 0.053)
	Marital status					
	Single, divorced or widowed	Ref.	Ref.	Ref.	Ref.	Ref.
	Married or cohabiting	0.347 (0.011, 0.684)	0.008 (-0.314, 0.330)	0.007 (-0.316, 0.330)	0.060 (-0.261, 0.382)	0.067 (-0.268, 0.402)
	Maternal work status					
	Not working	Ref.	Ref.	Ref.	Ref.	Ref.
Working	0.188 (0.022, 0.354)	-0.026 (-0.191, 0.138)	-0.026 (-0.191, 0.138)	-0.020 (-0.183, 0.144)	-0.019 (-0.183, 0.144)	
BLOCK 2	<i>Maternal characteristics before/during pregnancy</i>					
	Pre-pregnancy BMI, kg/m ²	-0.013 (-0.024, -	----	-0.001 (-0.015,	-0.004 (-0.018,	-0.004 (-0.018,

		0.002)		0.014)	0.010)	0.011)
	Smoking status during pregnancy					
	Non-smokers	Ref.	----	----	----	----
	Smokers	-0.113 (-0.279, 0.053)	----	----	----	----
	Gestational diabetes					
	No	Ref.	----	----	----	----
	Yes	-0.303 (-1.536, 0.930)	----	----	----	----
	<i>Maternal characteristics at the child's 4 years of age</i>					
BLOCK 3	Diet quality score	0.055 (0.040, 0.070)	----	----	0.034 (0.019, 0.049)	0.033 (0.018, 0.049)
	Feeding practices patterns					
	Restriction	0.083 (0.023, 0.143)	----	----	0.069 (0.011, 0.127)	0.067 (0.009, 0.125)
	Perceived monitoring	0.059 (-0.003, 0.121)	----	----	----	----
	Pressure to eat	0.022 (-0.039, 0.084)	----	----	----	----
	Maternal BMI, kg/m ²	-0.011 (-0.024, 0.002)	----	----	----	----

			0.002)			
BLOCK 4	<i>Maternal and other family characteristics at the child's 7 years of age</i>					
	Family structure					
	Single-parent family	Ref.	----	----	----	Ref.
	Family with both parents	0.200 (0.025, 0.374)	----	----	----	-0.051 (-0.224, 0.122)
	Living with siblings					
	No	Ref.	----	----	----	Ref.
Yes	-0.107 (-0.200, -0.014)	----	----	----	-0.039 (-0.161, 0.184)	
	Nagelkerke's R ²		0.31	0.31	0.42	0.44

CI: Confidence Interval; BMI: Body Mass Index. Statistically significant associations are highlighted in bold.

*Blocks of variables (1. socioeconomic characteristics of mothers before the child's birth; 2. maternal characteristics before/during pregnancy; 3. maternal characteristics at the child's 4 years of age; 4. family characteristics at the child's 7 years of age) were sequentially added into the analysis, if significantly associated with the WISH score in the unadjusted model. Maternal age, education, marital and work status (Block 1), were added do Model 1. Pre-pregnancy BMI (Block 2) was added to Model 2. Mothers' diet quality score and restrictive feeding practices (Block 3) were added to model 3. Family structure and living with siblings (Block 4) at the child's 7 years of age were added to Model 4. Child's sex and practice of sports at 7 years old were added as additional co-variates in all models.

Nagelkerke's R²: explained variance of each model in the step-by-step approach (block of variables sequentially added).