

THE JOURNAL OF AGRICULTURAL SCIENCE

EDITED BY

G. D. H. BELL, C.B.E., PH.D., F.R.S., Plant Breeding Institute, Cambridge

K. L. BLAXTER, PH.D., N.D.A. (HONS.), D.SC., F.R.S.E.,
Rowett Research Institute, Bucksburn, Aberdeen

G. W. COOKE, PH.D., F.R.I.C., Rothamsted Experimental Station, Harpenden

JOHN HAMMOND, Jr., M.A., School of Agriculture, Cambridge

PROF. SIR J. B. HUTCHINSON, C.M.G., SC.D., F.R.S., School of Agriculture,
Cambridge

H. L. PENMAN, O.B.E., PH.D., F.R.S., Rothamsted Experimental Station, Harpenden

H. H. ROGERS, B.SC., DIP.AG.SCI., Plant Breeding Institute, Cambridge

PROF. E. W. RUSSELL, C.M.G., PH.D., F.INST.P., Department of Soil Science,
University of Reading

F. YATES, C.B.E., SC.D., F.R.S., Rothamsted Experimental Station, Harpenden



CAMBRIDGE UNIVERSITY PRESS

BENTLEY HOUSE, 200 EUSTON ROAD, LONDON, N.W. 1

AMERICAN BRANCH: 32 EAST 57TH STREET, NEW YORK, N.Y. 10022

Price 35s. net
(U.S.A. \$5.50)

INSTRUCTIONS TO AUTHORS

Papers intended for publication should be submitted to Dr G. D. H. Bell, C.B.E., F.R.S., Plant Breeding Institute, Cambridge, or to one of the Associate Editors. It must be understood that, if accepted by the Editorial Board, the paper will not be published elsewhere in the same form.

MANUSCRIPTS. Papers, written in English, are accepted from any country and should be typed in double-spacing on one side of the paper with a margin at least 4 cm wide on the left-hand side. Authors should instruct typists on the style of manuscript required. A top copy and one carbon copy should be submitted.

TABLES must be self-explanatory. They should be typed on separate sheets, numbered consecutively and carry an appropriate title. Wherever possible, tables should be arranged so that they can be printed in the normal orientation of the text and without the need for rules.

LINE DRAWINGS (with photocopies) should be 25 cm. wide and drawn in black waterproof ink on Bristol board, graph paper with blue lines or tracing paper. Legends should be typed on a separate copy and numbering inserted lightly and clearly in soft pencil on the drawing. Tables and figures should not reproduce the same data.

The approximate position of tables and figures should be noted in the text.

PLATES should make a definite contribution to the value of the paper and the number submitted should be kept to a minimum. They should be good quality, unmounted, glossy prints and be lightly numbered in pencil on the reverse side. If several, or coloured, plates are submitted the author may be asked to contribute to the cost of reproducing them.

TITLE. The title must be specific and suitable for indexing by the mechanical methods now being employed. The full name and address of the institution in which the research has been carried out should be stated. Change of address may be given as a footnote. A short title, not exceeding 50 characters, should be provided for the running headlines.

STYLE. Experimental details and results should be recorded in the past tense and there should be no unnecessary repetition or loose phrases. Manuscripts are likely to be returned for modification if the presentation is not clear and precise.

LAYOUT. Authors are recommended to study '*General Notes on the Preparation of Scientific Papers*'

(Royal Society, London, 2nd edn., 1965). The Editorial Board do not insist upon a rigid format but it is usually convenient to divide the paper into sections, e.g. Introduction, Materials and Methods, Results and Discussion. An excess of headings and sub-headings should be avoided. A Summary should always be included.

Authors are advised to note the following points: a detailed review of literature is not necessary; relevant details should be given of the plant or animal material, the experimental design and chemical or other techniques employed; mean results with their relevant standard errors should be presented rather than detailed data; the statistical methods used should be clearly stated; the discussion should relate the author's experiments to other work on the subject and give the author's conclusions; the summary should be factual and suitable for use in abstracting journals. Footnotes should be avoided.

REFERENCES. The bibliography must be given in the form—Surname of authors, initials, year of publication (in parentheses), title of paper, name of journal (abbreviated according to the *World List of Scientific Periodicals*, 4th edn, Butterworths, London), volume and pages of reference. References should be in alphabetical order. In the text a reference should be quoted by the author's name and date (in parentheses). Where there are more than two authors, the initial reference in the text should include the names of all authors but subsequent citations, should be in the form—first author followed by *et al.* Authors should check that all references in the text appear at the end of the paper and vice versa, and that names and dates correspond in the two places.

PROOF CORRECTION. Standard proof correction marks (British Standard 1219) should be made as legibly as possible in ink, not pencil. Directions to the printer which are not to be set up in type should be encircled. Captions to illustrations and all references should be checked. Queries marked by the printer should be answered. Proofs are provided in order that authors can check the correctness of the type-setting—excessive alterations may be charged to the author.

OFFPRINTS. Contributors will receive 25 copies of their papers free and can order others when they receive the proofs.

Failure to comply with the 'Instructions to Authors' may delay publication.

ANIMAL PRODUCTION

JOURNAL OF THE BRITISH SOCIETY OF ANIMAL PRODUCTION

Contents of Volume 9, Part 1, February 1967 include the following

- | | |
|---|---|
| LEVI, SOLLER and SHILO. Dressing percentage and fat content in bulls. | OAKES. The assessment of a variety of chicken. |
| BROADBENT and WATSON. Sire and progeny performance. | DALTON, PEARSON and SHEARD. Behaviour of dairy bulls. |
| VANSCHOU BROEK, DE WILDE and LAMPO. Feed restriction in fattening pigs. | PURSER and KARAM. Lamb performance and birthcoat type. |
| STOBO, ROY and GASTON. Protein requirement of ruminant calf. | ROLLINS and CASADY. Pre-weaning deaths in rabbits. |
| TAYLOR and CRAIG. Growth of twin cattle. | BROADBENT, DODSWORTH, BALL and SAMPFORD. Semi-intensively reared Charolais crosses. |
| OWEN and RIDGMAN. Dietary energy and voluntary intake. | RUSSEL. Goitre in lambs grazing rape. |
| | SMITH. Selecting on components of a trait. |

The Journal is published quarterly in February, May, August and November. The annual subscription rate is 90s. (U.S.A. \$14.00) single copy 25s. (U.S.A. \$4.00). Orders and subscriptions may be sent to any bookseller or direct to:

OLIVER & BOYD LTD.

Tweeddale Court, 14 High Street, Edinburgh, 1

FULL PARTICULARS OF THE

JOURNALS

PUBLISHED BY THE

CAMBRIDGE
UNIVERSITY PRESS

MAY BE HAD FROM

THE MANAGER
CAMBRIDGE UNIVERSITY PRESS
BENTLEY HOUSE
200 EUSTON ROAD
LONDON, N.W. 1

CROP SCIENCE

Crop breeders, plant geneticists and physiologists, and workers in related areas will find *Crop Science* a source of valuable articles in their branches of science. This bimonthly journal carries reports of research in the genetics, physiology, ecology, breeding and management of field crops, turf-grasses, pastures and ranges, and in seed technology. It is published by the Crop Science Society of America.

\$16.00 per year in U.S. and Canada.
\$17.00 per year elsewhere.

Crop Science Society of America
677 S. Segoe Rd,
Madison, Wisconsin, U.S.A., 53711

(i)



Safe from insects for 10 to 20 days – with dieldrin

Once dieldrin has been applied, foliage is safe from destructive pests for up to 20 days. Dieldrin is especially useful in both temperate and Mediterranean regions for control of such insects as Colorado beetle, flea beetle, blossom beetle, leafminer and fly pests of crops. In tropical and sub-tropical regions, dieldrin is highly effective for protection against the stem borers in tea and coffee. It is also most effective against locusts, grasshoppers and termites. Dieldrin is complementary to endrin and aldrin. Together they control a large variety of major destructive pests, ranging from soil insects to foliage pests. They have excellent residual action, are stable in the presence of other chemicals and do not affect germination or cause taint. Furthermore their use, according to recommendation, is completely safe to the applicator and to the consumer of treated crops.

Aldrin, dieldrin and endrin are available as technical materials and high concentrate dusts suitable for further formulation, and also in forms suitable for the grower: emulsifiable concentrates, wettable powders, field-strength dusts and granules. For details of approved uses and advice on application consult the Shell company in your country.

