

Congruences on distributive pseudocomplemented lattices

William H. Cornish

Let $B_0 \subset B_1 \subset B_2 \subset \dots \subset B_n \subset \dots \subset B_\omega$ be all the non-trivial varieties of distributive pseudocomplemented lattices $(L; \vee, \wedge, *, 0, 1)$ considered as algebras of type

$(2, 2, 1, 0, 0)$. A subset J of such an algebra L is a congruence-kernel if and only if it is a lattice-ideal and $x^{**} \in J$ for each $x \in J$. The smallest congruence having J as its kernel is $\Theta(J)$, where $a \equiv b (\Theta(J))$, $(a, b \in L)$ if and only if $a \wedge c^* = b \wedge c^*$ for some $c \in J$. For given $0 \leq n \leq \omega$, let $\Sigma_n(J)$ be the smallest congruence having J as its kernel and such that the associated quotient algebra is in B_n . Of course, $\Sigma_\omega(J) = \Theta(J)$ and the main result of this paper shows that for $1 \leq n < \omega$,

$$\Sigma_n(J) = \Pi \{ \Theta(P_1 \cap P_2 \cap \dots \cap P_n) : J \subseteq P_1, P_2, \dots, P_n \subset L \\ \text{are minimal prime ideals} \}.$$

It is also shown that $\Sigma_\omega(J)$ is the smallest congruence on the lattice $(L; \vee, \wedge, 0, 1)$ having J as its kernel for each congruence kernel J if and only if $L \in B_1$. On the other hand, for any congruence kernel J , $\Sigma_0(J)$ is always the largest congruence on the lattice $(L; \vee, \wedge, 0, 1)$ having J as its kernel and so $\Sigma_0(J)$ is the unique congruence with J as its kernel and such that quotient algebra is boolean.

Received 10 October 1972.

1. Congruence-kernels

A congruence θ on a distributive pseudocomplemented lattice $(L; \vee, \wedge, *, 0, 1)$ is a congruence of the lattice $(L; \vee, \wedge, 0, 1)$ which also has the substitution property for the $*$ -operation. At times it will be convenient to refer to such a congruence as a $*$ -congruence. For such a congruence θ , $\ker(\theta) = \{x \in L : x \equiv 0(\theta)\}$ and $\text{coker}(\theta) = \{x \in L : x \equiv 1(\theta)\}$ are called, respectively, the *kernel* and *cokernel* of θ . A subset J of L is called a *congruence-kernel* (respectively *congruence-cokernel*) if $J = \ker(\theta)$ (respectively $J = \text{coker}(\theta)$) for some $*$ -congruence θ on L .

The purpose of this section is to characterize the congruence-kernels of a given algebra $(L; \vee, \wedge, *, 0, 1)$ and also to describe the smallest and largest congruences on L which have a given congruence-kernel as their common kernel. The necessary prerequisite information on the variety of all distributive pseudocomplemented lattices can be obtained from Chapter 3 of Grätzer's recent book [4]. We begin with some preliminary results on distributive lattices which will be useful both in this section and in the later sections.

Let J be an ideal in a distributive lattice L . Then a prime ideal P in L is called a *minimal prime ideal belonging to J* if P is a minimal member of the set of all prime ideals in L which contain J . If L has a smallest member, 0 , then a *minimal prime ideal* of L is simply a minimal prime ideal belonging to the zero-ideal (0) . The following lemma summarizes some useful information which appears in the literature.

LEMMA 1.1. *The following results hold in a distributive lattice L .*

- (i) *A prime ideal P containing an ideal J is a minimal prime ideal belonging to J if and only if, for each $x \in P$, there exists $y \in L \setminus P$ such that $x \wedge y \in J$.*
- (ii) *Suppose that $0 \in L$ and that Q is a prime ideal in L . Then, $0(Q) = \{x \in L : x \wedge y = 0 \text{ for some } y \in L \setminus Q\}$ is the intersection of all the minimal prime ideals of L which are contained in Q .*
- (iii) *If L is pseudocomplemented then a prime ideal P is a minimal prime ideal if and only if $x^{**} \in P$ for each $x \in P$; and the set of all minimal prime ideals of L endowed with the*

hull-kernel topology, is a compact Hausdorff space.

Proof. (i) follows from [7, Lemma 3.1] and the distributivity of L .

(ii) is established in [1, Proposition 2.2].

The first part of (iii) follows from (i), see also [4, Lemma 5, p. 169], while the second part is a consequence of [9, Proposition 3.2]. \square

If J is an ideal in a distributive lattice L then $\Theta_{\text{Lat}}(J)$ and $R(J)$ will be used to respectively denote the smallest congruence and the largest congruence on L possessing J as a congruence class. It is well-known that $x \equiv y (\Theta_{\text{Lat}}(J))$, $(x, y \in L)$, if and only if $x \vee a = y \vee a$ for some $a \in J$, while $x \equiv y (R(J))$, $(x, y \in L)$, if and only if, for any $b \in L$, $x \wedge b \in J$ if and only if $y \wedge b \in J$. For detailed information on $\Theta_{\text{Lat}}(J)$ and $R(J)$ we refer respectively to [4] and [10]. In [10, Proposition 4.1] it is shown that if $1 \in L$ then $\{x \in L : x \equiv 1 (R(J))\}$ is the filter $D(J)$ where $D(J) = \{a \in L : J:a = J\}$ and $J:a$ is the ideal $\{b \in L : a \wedge b \in J\}$. When $0 \in L$, R will be used to denote $R(\{0\})$ and D will be used to denote $D(\{0\})$. If L is pseudocomplemented then $x \equiv y (R)$, $(x, y \in L)$, if and only if $x^{**} = y^{**}$ and $D = \{x \in L : x^{**} = 1\}$, the filter of dense elements.

LEMMA 1.2. *Let $h(J)$ denote the set of all minimal prime ideals belonging to a given ideal J in a distributive lattice L . Then*

(i) $R(J) = \cap \{R(P) : P \in h(J)\}$, and

(ii) $\Theta_{\text{Lat}}(J) = \cap \{\Theta_{\text{Lat}}(P) : P \in h(J)\}$.

Proof. (i) Suppose that $x, y \in L$ and $x \equiv y (R(J))$. If $x \in P \in h(J)$ then, according to Lemma 1.1 (i), $x \wedge z \in J$ for some $z \in L \setminus P$. Then $y \wedge z \in J$ and so $y \in P$. By symmetry, we conclude that $x \in P$ if and only if $y \in P$, that is, $x \equiv y (R(P))$. Thus, $R(J) \subseteq \cap \{R(P) : P \in h(J)\}$. But $J = \cap \{P : P \in h(J)\}$ and so J is a congruence class of L modulo $\cap \{R(P) : P \in h(J)\}$. Hence $\cap \{R(P) : P \in h(J)\} \subseteq R(J)$. Part (i) is now proved.

(ii) For any $P \in h(J)$, it is clear that $\Theta_{\text{Lat}}(J) \subseteq \Theta_{\text{Lat}}(P)$.

Suppose that $x, y \in L$ and $x \not\equiv y (\Theta_{\text{Lat}}(J))$ and consider $F = \{a \in L : x \vee a = y \vee a\}$. Then F is non-empty since $x \vee y \in F$, and F is a filter. By the hypothesis on x and y , it follows that J and F are disjoint. Using the Birkhoff-Stone Theorem and the dual form of Zorn's Lemma, it follows that there exists $Q \in \mathcal{h}(J)$ such that Q and F are disjoint. Hence, $x \not\equiv y (\Theta(Q))$. The equality asserted in (ii) is now readily established. \square

If F is a filter in an arbitrary lattice L then the relation $\Psi(F)$ defined by: $x \equiv y (\Psi(F))$ ($x, y \in L$) if and only if $x \wedge f = y \wedge f$ for some $f \in F$, is always an equivalence relation. When L is a distributive lattice, $\Psi(F)$ is the smallest congruence having F as a congruence class. Special cases of the following lemma are well-known, for example, [5, Lemma 7, p. 36] and [6, Theorem 2]; we include a simple direct proof.

LEMMA 1.3. *For any two filters F_1 and F_2 in a lattice L , the equivalence relations $\Psi(F_1)$ and $\Psi(F_2)$ are permutable.*

Proof. Suppose that $x, y \in L$ and that $x \equiv y (\Psi(F_1) \circ \Psi(F_2))$. Then, $x \wedge f_1 = z \wedge f_1$ and $z \wedge f_2 = y \wedge f_2$ for some $z \in L$, $f_1 \in F_1$ and $f_2 \in F_2$. Thus, $x \wedge f_1 \wedge f_2 = y \wedge f_1 \wedge f_2$. Put $w = (x \wedge f_2) \vee (y \wedge f_1)$. Then

$$x = x \vee (x \wedge f_1 \wedge f_2) = x \vee (y \wedge f_1 \wedge f_2) \\ \equiv (x \wedge f_2) \vee (y \wedge f_1) = w (\Psi(F_2)),$$

and $w = (x \wedge f_2) \vee (y \wedge f_1) \equiv (x \wedge f_1 \wedge f_2) \vee y = y (\Psi(F_1))$. That is, $x \equiv y (\Psi(F_2) \circ \Psi(F_1))$. The lemma now follows from symmetry. \square

The next result describes congruence-cokernels in a distributive pseudocomplemented lattice; it will be useful throughout the entire paper.

PROPOSITION 1.4. *Let $(L; \vee, \wedge, *, 0, 1)$ be a distributive pseudocomplemented lattice. A subset F of L is a congruence-cokernel if and only if F is a filter in L . Moreover, the smallest $*$ -congruence having a given filter F as its cokernel is $\Psi(F)$.*

Proof. It is sufficient to prove that, for a given filter F , $\Psi(F)$ has the substitution property for the $*$ -operation. Thus suppose that $x, y \in L$ and $x \equiv y(\Psi(F))$, that is, $x \wedge f = y \wedge f$ for some $f \in F$. Then $x^{**} \wedge f^{**} = (x \wedge f)^{**} = (y \wedge f)^{**} = y^{**} \wedge f^{**}$ so that $x^* \vee f^* = y^* \vee f^*$ in the skeleton $(S(L); \vee, \wedge, *, 0, 1)$ of L . Then $x^* \wedge f^{**} = y^* \wedge f^{**}$, since $S(L)$ is a boolean algebra, and hence $x^* \wedge f = y^* \wedge f$. \square

If J is an ideal in a distributive pseudocomplemented lattice L , let $J_* = \{x \in L : x \geq a^* \text{ for some } a \in J\}$. It is easy to see that J_* is a filter. We now come to our first characterization of congruence-kernels.

THEOREM 1.5. *Let J be a non-empty subset of a distributive pseudo-complemented lattice $(L; \vee, \wedge, *, 0, 1)$. Then the following conditions are equivalent:*

- (i) J is a congruence-kernel,
- (ii) J is an ideal of the lattice L and $x^{**} \in J$ for each $x \in J$,
- (iii) J is an ideal of the lattice L and each minimal prime ideal belonging to J is a minimal prime ideal,
- (iv) J is an intersection of minimal prime ideals of the lattice L .

Moreover, if J satisfies any of the equivalent conditions (i)-(iv) then the smallest congruence on $(L; \vee, \wedge, *, 0, 1)$ having J as its kernel is $\Theta(J)$, where $x \equiv y(\Theta(J))$ ($x, y \in L$) if and only if $x \wedge a^* = y \wedge a^*$ for some $a \in J$. Thus, $\Theta(J) = \Psi(J_*)$.

Proof. (i) \Rightarrow (ii). Suppose that $J = \ker(\Theta)$ for some $*$ -congruence Θ . Then J is certainly a lattice-ideal and, for $x \in J$, $x \equiv 0(\Theta)$, so that $x^{**} \equiv 0^{**} = 0(\Theta)$, and hence (ii) holds for J .

(ii) \Rightarrow (i). From (ii) and Proposition 1.4, $\Psi(J_*)$ is a $*$ -congruence. Because $x^{**} \in J$ for $x \in J$, $\ker(\Psi(J_*)) = J$. If Φ is a $*$ -congruence with $\ker(\Phi) = J$ then $x \wedge a^* = y \wedge a^*$ for $a \in J$ implies $x \equiv y(\Phi)$ since $a^* \equiv 0^* = 1(\Phi)$. Thus, (ii) \Rightarrow (i), and we have also described the smallest $*$ -congruence having J as its kernel.

The implication $(iv) \Rightarrow (ii)$ follows from Lemma 1.1 (iii) so that it remains to establish that $(ii) \Rightarrow (iii)$ and $(iii) \Rightarrow (iv)$.

$(ii) \Rightarrow (iii)$. Let J satisfy (ii) and suppose that P is a minimal prime ideal belonging to J . Let $x \in P$. By Lemma 1.1 (i) , $x \wedge z \in J$ for some $z \in L \setminus P$. Then $x^{**} \wedge z^{**} = (x \wedge z)^{**} \in J$ and so $x^{**} \in P$, since P is prime and $z \leq z^{**}$. Then, by Lemma 1.1 (iii) , P is a minimal prime ideal.

$(iii) \Rightarrow (iv)$. Each ideal in a distributive lattice is the intersection of all the minimal prime ideals belonging to it. Hence (iv) follows from (iii) . \square

THEOREM 1.6. *Let J be a given congruence-kernel in a distributive pseudocomplemented lattice $(L; \vee, \wedge, *, 0, 1)$. The following conditions relating to an equivalence relation Φ on L are equivalent:*

- (i) Φ is the largest $*$ -congruence such that $J = \ker \Phi$,
- (ii) $\Phi = R(J)$,
- (iii) for any $x, y \in L$, $x \equiv y (\Phi)$ if and only if $x^{**} \wedge a^* = y^{**} \wedge a^*$ for some $a \in J$,
- (iv) $\Phi = \Theta(J) \vee R$ in the lattice of $(*)$ -congruences,
- (v) for any $x, y \in L$, $x \equiv y (\Phi)$ if and only if $x \wedge (b \vee b^*) \wedge a^* = y \wedge (b \vee b^*) \wedge a^*$ for some $b \in L$ and $a \in J$,
- (vi) Φ is the smallest $*$ -congruence with $J = \ker \Phi$ and such that L/Φ is a boolean algebra,
- (vii) Φ is the unique $*$ -congruence with $J = \ker \Phi$ and such that L/Φ is a boolean algebra.

Proof. $(i) \Leftrightarrow (ii)$. By Theorem 1.5, each minimal prime ideal belonging to J is a minimal prime ideal. Hence, by Lemma 1.1, $R(P)$ is a $*$ -congruence for each $P \in h(J)$ and so Lemma 1.2 implies that $R(J)$ is a $*$ -congruence. It is now evident that (i) and (ii) are equivalent.

$(iii) \Leftrightarrow (v)$. Because $(b \vee b^*)^{**} = 1$ for each $b \in L$, (iii) follows from (v) . On the other hand suppose that $x^{**} \wedge a^* = y^{**} \wedge a^*$ for some $a \in J$. Since $x \vee x^*$ and $y \vee y^*$ are dense,

$(x \vee x^*) \wedge (y \vee y^*) = b \vee b^*$ for some $b \in L$ and then

$$\begin{aligned} x \wedge (b \vee b^*) \wedge a^* &= (x^{**} \wedge (x \vee x^*)) \wedge (b \vee b^*) \wedge a^* \\ &= x^{**} \wedge (b \vee b^*) \wedge a^* = y^{**} \wedge (b \vee b^*) \wedge a^* = y \wedge (b \vee b^*) \wedge a^* . \end{aligned}$$

Thus, (iii) and (v) are equivalent.

(iv) \Leftrightarrow (v). Because L is pseudocomplemented, $R = \Psi(D)$, (cf. [10, Proposition 5.4]). Thus, from Theorem 1.5,

$$\Theta(J) \vee R = \Psi(J_*) \vee \Psi(D) = \Psi(J_* \vee D) ,$$

where

$$J_* \vee D = \{x \in L : x = b \wedge d, b \in J_*, d \in D\}$$

is the join of J_* and D in the lattice of filters on L . Hence (iv) and (v) are equivalent.

(ii) \Leftrightarrow (iii). Because of the equivalence of (iii), (iv), and (v) it is clear that the relation Φ of (iii) is a $*$ -congruence with $\ker \Phi = J$ and hence $\Phi \subseteq R(J)$. It remains to establish the reverse inequality.

Let P be a minimal prime ideal and suppose that $x \equiv y (R(P))$ for given $x, y \in L$. Then $x, y \in P$ or $x, y \in L \setminus P$. In the first case $x \vee y \in P$ and $x^{**} \wedge (x \vee y)^* = x^{**} \wedge x^* \wedge y^* = 0 = y^{**} \wedge (x \vee y)^*$, while in the second case $x \wedge y \in L \setminus P$ so that $(x \wedge y)^* \in P$ and $x^{**} \wedge ((x \wedge y)^*)^* = x^{**} \wedge (x \wedge y)^{**} = x^{**} \wedge y^{**} = y^{**} \wedge ((x \wedge y)^*)^*$. Thus, for any minimal prime ideal P , $x \equiv y (R(P))$ ($x, y \in L$) if and only if $x^{**} \wedge p^* = y^{**} \wedge p^*$ for some $p \in P$.

Now suppose $x, y \in L$ and $x \equiv y (R(J))$. Let $h(J) = \{P_\lambda : \lambda \in \Lambda\}$ be the set of all minimal prime ideals containing J . By Lemma 1.2 and Theorem 1.5, $x \equiv y (R(P_\lambda))$ for each $\lambda \in \Lambda$. It follows from the preceding paragraph that, for each $\lambda \in \Lambda$, there exists $p_\lambda \in P_\lambda$ such that $x^{**} \wedge p_\lambda^* = y^{**} \wedge p_\lambda^*$. Hence, for each $P_\lambda \in h(J)$, $P_\lambda \in g(p_\lambda^*) = \{Q : Q \text{ is minimal prime ideal and } p_\lambda^* \notin Q\}$. That is, $\{g(p_\lambda^*) : \lambda \in \Lambda\}$ is an open cover of the closed subset $h(J)$ of the space of minimal prime ideals. Because of Lemma 1.1 (iii), $h(J)$ is compact and so there exist $\lambda_1, \lambda_2, \dots, \lambda_n \in \Lambda$ such that

$$h(J) \subseteq g(p_{\lambda_1}^*) \cup g(p_{\lambda_2}^*) \cup \dots \cup g(p_{\lambda_n}^*) = g(p_{\lambda_1}^* \vee p_{\lambda_2}^* \vee \dots \vee p_{\lambda_n}^*) .$$

Hence

$$p_{\lambda_1}^{**} \wedge \dots \wedge p_{\lambda_n}^{**} = (p_{\lambda_1}^* \vee \dots \vee p_{\lambda_n}^*)^* \in \cap \{Q : Q \in g(p_{\lambda_1}^* \vee \dots \vee p_{\lambda_n}^*)\} \\ \subseteq \cap \{Q : Q \in h(J)\} = J ,$$

by Theorem 1.5. Now $x^{**} \wedge p_{\lambda_i}^* = y^{**} \wedge p_{\lambda_i}^*$ for $i = 1, \dots, n$ and so

$$x^{**} \wedge (p_{\lambda_1}^* \underline{\vee} \dots \underline{\vee} p_{\lambda_n}^*) = y^{**} \wedge (p_{\lambda_1}^* \underline{\vee} \dots \underline{\vee} p_{\lambda_n}^*)$$

in $S(L)$. Hence

$$x^{**} \wedge (p_{\lambda_1}^{**} \wedge \dots \wedge p_{\lambda_n}^{**})^* = y^{**} \wedge (p_{\lambda_1}^{**} \wedge \dots \wedge p_{\lambda_n}^{**})^*$$

and $p_{\lambda_1}^{**} \wedge \dots \wedge p_{\lambda_n}^{**} \in J$. Thus $x \equiv y (\Phi)$, where Φ is the congruence of (iii). And (ii) and (iii) are equivalent.

Thus conditions (i) to (v) are equivalent.

(v) \Leftrightarrow (vi). If Φ is the congruence of (v) then $\ker(\Phi) = J$ and $b \vee b^* \in \text{coker}(\Phi)$ for each $b \in L$ and so L/Φ is boolean. It is clear that Φ is the smallest $*$ -congruence with these properties and so (v) and (vi) are equivalent.

(vi) \Leftrightarrow (vii). Since (i) and (vi) are equivalent, (vii) is equivalent to (vi).

The theorem has now been established. \square

The preceding theorem shows that the congruence $R(J)$ has a very simple description when J is a congruence-kernel. In fact many of the properties of the congruence $R(J)$ which are implied by that theorem are characteristic of congruence-kernels.

THEOREM 1.7. *The following conditions on an ideal J of a distributive pseudocomplemented lattice L are equivalent:*

- (i) J is a congruence-kernel,
- (ii) $R(J)$ is a $*$ -congruence,

- (iii) for any $x, y \in L$, $x \equiv y (R(J))$ if and only if $x^{**} \wedge a^* = y^{**} \wedge a^*$ for some $a \in J$,
- (iv) for any $x, y \in L$, $x \equiv y (R(J))$ if and only if $x \wedge (b \vee b^*) \wedge a^* = y \wedge (b \vee b^*) \wedge a^*$ for some $b \in L$ and $a \in J$,
- (v) $R(J) = \Psi(D(J))$,
- (vi) $D(J) = D \vee J_*$ in the lattice of filters,
- (vii) $D \subseteq D(J)$,
- (viii) for any filter F , the congruences $R(J)$ and $\Psi(F)$ are permutable,
- (ix) the congruences $R(J)$ and R are permutable,
- (x) $R \subseteq R(J)$.

Proof. The equivalence of conditions (i), (ii), (iii) and (iv) is an easy consequence of Theorems 1.6 and 1.5.

(iv) \Rightarrow (v). Condition (iv) implies that $R(J) = \Psi(\text{coker}R(J))$, and, as has been previously mentioned, $\text{coker}(R(J)) = D(J)$.

(v) \Rightarrow (ii). This follows from Proposition 1.4.

(v) \Rightarrow (vi) and (vi) \Rightarrow (vii) are obvious.

(vii) \Rightarrow (i). Suppose that $D \subseteq D(J)$ and let $x \in J$. Then $x \vee x^* \in D(J)$ and so $x^{**} \wedge (x \vee x^*) = x$ implies $x^{**} \in J : x = J$. Thus (i) holds.

(v) \Rightarrow (viii) follows from Lemma 1.3.

(viii) \Rightarrow (ix) since $R = \Psi(D)$ holds in any pseudocomplemented lattice.

(ix) \Rightarrow (i). Let $x \in J$. Then $0 \equiv x (R(J))$ and $x \equiv x^{**} (R)$ so that $0 \equiv x^{**} (R(J) \circ R)$. By (ix), $0 \equiv z (R)$ and $z \equiv x^{**} (R(J))$. Hence $z = 0$ and so $x^{**} \in J = \ker R(J)$.

(ii) \Rightarrow (x) is clear and (x) \Rightarrow (i) since $x \equiv x^{**} (R)$ for any $x \in L$. The theorem is proved. \square

Some other consequence of Theorem 1.5 will be given in the next sections.

2. The congruences $\Sigma_n(J)$

Because of Theorem 1.5, the congruence-kernels of a distributive pseudocomplemented lattice are precisely its α -ideals, in the sense of [2]. The following lemma is thus an easy consequence of [2, Propositions 3.2, 3.3]; it is easy to provide a direct proof and so details are omitted.

LEMMA 2.1. *The set $K(L)$ of all congruence-kernels of a distributive pseudocomplemented lattice $(L; \vee, \wedge, *, 0, 1)$ is a complete distributive lattice, where the meet is set-theoretic intersection and the join of a family $\{J_\lambda : \lambda \in \Lambda\}$ of congruence-kernels is*

$$\bigvee \{J_\lambda : \lambda \in \Lambda\} = \left\{ x \in L : x \leq \left(a_{\lambda_1} \vee \dots \vee a_{\lambda_n} \right)^{**} \right. \\ \left. \text{for some } a_{\lambda_1} \in J_{\lambda_1}, \dots, a_{\lambda_n} \in J_{\lambda_n} \text{ and } \lambda_1, \dots, \lambda_n \in \Lambda \right\}.$$

Moreover, the maps $J \rightarrow \{b \in S(L) : b = x^{**} \text{ for some } x \in J\}$ and $I \rightarrow \{x \in L : x^{**} \in I\}$ are mutually inverse lattice-isomorphisms between $K(L)$ and the lattice of ideals of the skeleton $(S(L); \underline{\vee}, \wedge, *, 0, 1)$. \square

LEMMA 2.2. *Let θ be a $*$ -congruence on a distributive pseudocomplemented lattice $(L; \vee, \wedge, *, 0, 1)$ and let $\theta : L \rightarrow L/\theta$ be the canonical epimorphism. Then the set-function induced by θ and the inverse image map θ^+ are mutually inverse lattice isomorphisms between the interval $[\ker\theta, L] = \{J \in K(L) : \ker\theta \leq J\}$ of the lattice $K(L)$ of congruence-kernels of L and the lattice $K(L/\theta)$ of congruence-kernels of L/θ . Hence the map $P \rightarrow \theta(P)$ is a bijection of the set of minimal prime ideals of L which contain $\ker\theta$ onto the set of minimal prime ideals of L/θ .*

Proof. Let $J \in K(L)$ be such that $\ker\theta \subseteq J$ and suppose that $x \in \theta^+ \theta(J)$. Hence $\theta(x) = \theta(a)$ for some $a \in J$, and so $\theta(x \wedge a^*) = \theta(x) \wedge \theta(a)^* = 0$ implies that $x \wedge a^* \in \ker\theta \subseteq J$. If P is a minimal prime ideal containing J it follows from Lemma 1.1 that $x \in P$. Because of Theorem 1.5, $x \in J$ and so $\theta^+ \theta(J) = J$.

It is now clear that the first statement of the lemma holds and from this we immediately obtain the second statement, because Theorem 1.5 and

Lemma 2.1 imply that the minimal prime ideals of L are precisely the meet-irreducible elements of $K(L)$. \square

REMARK. Lemma 2.2 gives an estimate of the way in which the lattice of congruence-kernels fails to determine the lattice of $*$ -congruences. Of course, the map $\theta \rightarrow \ker(\theta)$ is a bijection (lattice-isomorphism) of the lattice of $*$ -congruences onto the lattice of congruence-kernels of L if and only if $(L; \vee, \wedge, *, 0, 1)$ is precisely the skeleton $(S(L); \underline{\vee}, \wedge, *, 0, 1)$ that is, if and only if L is a boolean algebra. This is because $\ker(R) = \{0\}$ if and only if $x = x^{**}$ for each $x \in L$.

Let $\{B_n : 0 \leq n \leq \omega\}$ be all the non-trivial varieties of distributive pseudocomplemented lattices (cf. [4, Chapter 3] and [8]). Thus B_0 is the class of boolean algebras, B_1 is the class of Stone lattices and B_ω is the class of all distributive pseudocomplemented lattices. We will need the following characterization of B_n ($1 \leq n < \omega$) due to Lee [8, Theorems 4, 5].

LEMMA 2.3. *Let $(L; \vee, \wedge, *, 0, 1)$ be a distributive pseudocomplemented lattice and let $1 \leq n < \omega$ be a positive integer. Then the following conditions are equivalent:*

- (i) $L \in B_n$,
- (ii) any $x_1, x_2, \dots, x_n \in L$ satisfy the identity

$$\left(\bigwedge_{i=1}^n x_i \right)^* \vee \bigvee_{i=1}^n (x_1 \wedge \dots \wedge x_{i-1} \wedge x_i^* \wedge x_{i+1} \wedge \dots \wedge x_n)^* = 1,$$
- (iii) each prime ideal in L contains at most n distinct minimal prime ideals. \square

LEMMA 2.4. *Let $(L; \vee, \wedge, *, 0, 1)$ be a distributive pseudocomplemented lattice. Then*

- (i) for any n (not necessarily distinct) minimal prime ideals P_1, P_2, \dots, P_n in L , $L/\theta(P_1 \cap P_2 \cap \dots \cap P_n) \in B_n$,
- (ii) for any $1 \leq n < \omega$, $L \in B_n$ implies that

$$\bigcap \{ \theta(P_1 \cap P_2 \cap \dots \cap P_n) : P_1, P_2, \dots, P_n \text{ are } n \text{ minimal prime ideals} \} = \omega_L$$

(the smallest congruence on L).

Proof. (i) $L/\theta(P_1 \cap P_2 \cap \dots \cap P_n)$ has, at most, n distinct minimal prime ideals. This follows from Lemma 2.2 since P_1, P_2, \dots, P_n are the only minimal prime ideals of L containing $P_1 \cap P_2 \cap \dots \cap P_n$. Thus part (i) follows from Lemma 2.3.

(ii) The dual of Lemma 1.2 (ii) implies that

$$\bigcap \{ \Psi(F) : F \text{ is a prime filter in } L \} = \omega_L.$$

Recall that ideal P is prime if and only if $L \setminus P$ is a prime filter. By Lemma 1.4, $\Psi(L \setminus P)$ is a $*$ -congruence and it is clear that $\ker(\Psi(L \setminus P)) = O(P)$ and so $\theta(O(P)) \subseteq \Psi(L \setminus P)$. Combining this information we obtain: $\bigcap \{ \theta(O(P)) : P \text{ is a prime ideal} \} = \omega_L$. Then, part (ii) is an easy consequence of Lemma 2.3 (iii) and Lemma 1.1 (ii). \square

Let J be a congruence-kernel in a distributive pseudocomplemented lattice L . For any $0 \leq n \leq \omega$, let $\Sigma_n(J)$ denote the smallest of all $*$ -congruences θ on L such that $\ker(\theta) = J$ and $L/\theta \in B_n$, so that $L/\Sigma_n(J)$ is the maximal homomorphic image of L which lies in B_n . Even though $\Sigma_\omega(J) = \theta(J)$ and $\Sigma_0(J) = R(J)$ (Theorem 1.6), it will at times be convenient to use that notation $\Sigma_\omega(J)$ and $\Sigma_0(J)$. We will also use Σ_n in place of $\Sigma_n(\{0\})$. The existence and description of $\Sigma_n(J)$ is the aim of the next theorem. Granting the existence it is clear that

$$\theta(J) = \Sigma_\omega(J) \subseteq \dots \subseteq \Sigma_n(J) \subseteq \Sigma_{n-1}(J) \subseteq \dots \subseteq \Sigma_1(J) \subseteq \Sigma_0(J) = R(J).$$

For $1 \leq n < \omega$, let D_n be the filter of L generated by all elements of the form

$$\left(\bigwedge_{i=1}^n x_i \right)^* \vee \bigvee_{i=1}^n (x_1 \wedge \dots \wedge x_{i-1} \wedge x_i^* \wedge x_{i+1} \wedge \dots \wedge x_n)^*$$

where $x_1, \dots, x_n \in L$. Thus

$$D_n = \left\{ x \in L : x \geq \bigwedge_{j=1}^m \left(\left(\bigwedge_{i=1}^n x_{ij} \right)^* \vee \bigvee_{i=1}^n (x_{1j} \wedge \dots \wedge x_{i-1j} \wedge x_{i+1j}^* \wedge \dots \wedge x_{nj})^* \right) \right. \\ \left. \text{for some } x_{ij}, (i = 1, \dots, n, j = 1, \dots, m), \text{ in } L \right\}.$$

For any $x_1, \dots, x_n \in L$, it is easy to see that

$$(x_1 \wedge \dots \wedge x_n)^* \vee \bigvee_{i=1}^n (x_1 \wedge \dots \wedge x_{i-1} \wedge x_{i+1}^* \wedge \dots \wedge x_n)^* \\ \geq (x_1 \wedge \dots \wedge x_n)^* \vee (x_1 \wedge \dots \wedge x_n)^{**}$$

and so $D_n \supseteq D_1 \supseteq D$ for any $1 \leq n \leq \omega$.

THEOREM 2.5. *Let $1 \leq n < \omega$ be a given positive integer and let J be a congruence-kernel in a distributive pseudocomplemented lattice $(L; \vee, \wedge, *, 0, 1)$. The following conditions relating to a $*$ -congruence Φ are equivalent:*

(i) $\Phi = \Sigma_n(J)$,

(ii) for any $x, y \in L$, $x \equiv y(\Phi)$ if and only if $x \wedge b \wedge a^* = y \wedge b \wedge a^*$, where

$$b = \bigwedge_{j=1}^m \left(\left(\bigwedge_{i=1}^n b_{ij} \right)^* \vee \bigvee_{i=1}^n (b_{1j} \wedge \dots \wedge b_{i-1j} \wedge b_{i+1j}^* \wedge \dots \wedge b_{nj})^* \right) \\ \text{for some } b_{ij} (i = 1, \dots, n, j = 1, \dots, m) \text{ in } L, \text{ and } \\ a \in J,$$

(iii) $\Phi = \Psi(D_n \vee J_*)$,

(iv) $\Phi = \Sigma_n \vee \Theta(J)$ in the lattice of $(*)$ -congruences,

(v)

$\Phi = \cap \{ \Theta(P_1 \cap P_2 \cap \dots \cap P_n) : P_1, P_2, \dots, P_n \text{ are (not necessarily distinct) minimal prime ideals in } L \text{ which contain } J \}$.

Proof. It is clear that (ii) and (iii) are equivalent. The congruence Φ given in both is, indeed, a \ast -congruence (Proposition 1.4) and has J as its kernel since $D_n \subseteq D$. As $D_n \subseteq \text{coker}(\Phi)$, $L/\Phi \in B_n$ in view of Lemma 2.3 (ii). It follows that $\Phi = \Sigma_n(J)$ and so (i), (ii) and (iii) are equivalent.

The equivalence of (iii) and (iv) follows from Theorem 1.5 and the equivalence of (i) and (iii) for the case $J = \{0\}$.

Now let Φ be the congruence given in (v). The theorem will be established if we can show that $\Phi = \Sigma_n(J)$.

For any n minimal prime ideals P_1, \dots, P_n , $L/\theta(P_1 \cap \dots \cap P_n) \in B_n$, by Lemma 2.4 (i), and since B_n is a variety, $L/\Phi \in B_n$, as L/Φ is a subdirect product of algebras in B_n . But $\ker \Phi = J$ (Theorem 1.5) and so $\Sigma_n(J) \subseteq \Phi$ by the definition of $\Sigma_n(J)$. Hence it remains to show that $\Phi \subseteq \Sigma_n(J)$.

Let $\sigma : L \rightarrow L/\Sigma_n(J)$ denote the canonical epimorphism and, for congruence $\theta \supseteq \Sigma_n(J)$, let θ' denote the congruence on $L/\Sigma_n(J)$ induced by θ . For $\sigma(x), \sigma(y)$ in $L/\Sigma_n(J)$, $(x, y \in L)$, $x \equiv y (\theta')$ if and only if $x \equiv y (\theta)$; for details of this construction see [3, §11, Chapter 3].

Since $\Sigma_n(J) \subseteq \Phi$, Φ' is a well-defined congruence on $L/\Sigma_n(J)$. Using Theorem 1.5 it follows that

$$\Phi' \subseteq \cap \{ \theta(\sigma(P_1 \cap P_2 \cap \dots \cap P_n)) : P_1, P_2, \dots, P_n \text{ are minimal primes in } L \text{ containing } J \}.$$

As $\ker(\Sigma_n(J)) = J$, Lemma 2.2 implies that

$$\Phi' \subseteq \cap \{ \theta(Q_1 \cap Q_2 \cap \dots \cap Q_n) : Q_1, Q_2, \dots, Q_n \text{ are minimal prime ideals in } L/\Sigma_n(J) \}.$$

Hence, by Lemma 2.4 (ii), $\Phi' = \omega_{L/\Sigma_n(J)}$ and so $\Phi = \Sigma_n(J)$. \square

Some easy consequences of the above theorem are:

COROLLARY 2.6. For any $0 \leq n < \omega$, $D_{n+1} \subseteq D_n \subseteq D$. \square

COROLLARY 2.7. For any $0 \leq n \leq \omega$, $\Sigma_0(J) = \Sigma_n(J) \vee \Sigma_0$. \square

Proof. Combine Theorems 1.6 and 2.5. \square

COROLLARY 2.8. For any $0 \leq n, m \leq \omega$ and congruence-kernels I and J , the congruences $\Sigma_n(I)$ and $\Sigma_m(J)$ are permutable.

Proof. Combine Lemma 1.3, Theorem 1.5, Theorem 1.6 and Theorem 2.5. \square

COROLLARY 2.9. For any $1 \leq n < \omega$ and congruence-kernel J , $L/\Theta(J) \in B_n$ if and only if

$$\Theta(J) = \bigcap \{ \Theta(P_1 \cap P_2 \cap \dots \cap P_n) : P_1, P_2, \dots, P_n \text{ are minimal prime ideals containing } J \} .$$

We end this section with the following result.

THEOREM 2.10. For any fixed $0 \leq n \leq \omega$, the map $J \mapsto \Sigma_n(J)$ is an injection of the lattice of congruence-kernels into the lattice of $*$ -congruences which preserves arbitrary joins. When $n = 0$, the map preserves finite meets.

Proof. It easily follows from Theorem 1.5 and Lemma 2.1 that

$$\Theta\left(\bigvee \{J_\lambda : \lambda \in \Lambda\}\right) = \bigvee \{\Theta(J_\lambda) : \lambda \in \Lambda\}$$

for any set $\{J_\lambda : \lambda \in \Lambda\}$ of congruence-kernels. As a consequence, Theorem 2.5 (iv) and Theorem 1.6 (iv) imply that $J \mapsto \Sigma_n(J)$ preserves arbitrary joins for any $0 \leq n \leq \omega$.

The last statement of the theorem follows from Theorem 1.6 (iii). \square

In the next section we will consider conditions on L which ensure that the map $J \mapsto \Sigma_n(J)$ preserves finite meets even when $1 \leq n \leq \omega$.

3. Stone lattices

LEMMA 3.1. Let J be a congruence-kernel in the distributive pseudocomplemented lattice L . Then $\Theta(J) = \Theta_{\text{Lat}}(J)$ if and only if $J = \{x \in J : J \vee (x^*) = L\}$.

Proof. Suppose that $\Theta(J) = \Theta_{\text{Lat}}(J)$. Thus, $x \in J$ implies

$x^* \equiv 1 \ (\Theta_{\text{Lat}}(J))$ and so $a \vee x^* = a \vee 1 = 1$ for some $a \in J$.

Consequently, $\{x \in J : J \vee (x^*) = L\} = J$.

Conversely, suppose that $J = \{x \in J : J \vee (x^*) = L\}$. Suppose that $a, b \in L$ are such that $a \equiv b \ (\Theta(J))$. Then, by Theorem 1.5, $a \wedge x^* = b \wedge x^*$ for some $x \in J$. Using the hypothesis imposed on J and working in the lattice of ideals of L , we obtain

$$\begin{aligned} (a] \vee J &= ((a] \vee J) \cap L = ((a] \vee J) \cap ((x^*) \vee J) = ((a] \cap (x^*)) \vee J \\ &= ([b] \cap (x^*)) \vee J = [b] \vee J. \end{aligned}$$

Hence $a \equiv b \ (\Theta_{\text{Lat}}(J))$ and so $\Theta(J) = \Theta_{\text{Lat}}(J)$. \square

It should be noted that any ideal J of the distributive pseudo-complemented lattice L is necessarily a congruence-kernel if $J = \{x \in J : J \vee (x^*) = L\}$, and we will refer to such ideals as *special congruence-kernels*. It is easy to see that L is a Stone lattice, that is, $L \in B_1$, if and only if each congruence-kernel is a special congruence-kernel. Using this observation we now come to our final result.

THEOREM 3.2. *The following conditions on a distributive pseudo-complemented lattice $(L; \vee, \wedge, *, 0, 1)$ are equivalent:*

- (i) L is a Stone lattice,
- (ii) each congruence-kernel in L is a special congruence-kernel,
- (iii) $\Theta(J) = \Theta_{\text{Lat}}(J)$ for each congruence-kernel J ,
- (iv) $\Theta_{\text{Lat}}(J)$ is a $*$ -congruence for each congruence-kernel J ,
- (v) $\Theta(P) = \Theta_{\text{Lat}}(P)$ for each minimal prime ideal P ,
- (vi) $\Theta_{\text{Lat}}(P)$ is a $*$ -congruence for each minimal prime ideal P ,
- (vii) each minimal prime ideal is a special congruence-kernel,
- (viii) $\cap\{\Theta(P) : P \text{ is a minimal prime ideal}\} = \omega_L$,
- (ix) for any congruence-kernel J and any filter F , $\Theta_{\text{Lat}}(J)$ and $\Psi(F)$ are permutable congruences,
- (x) for any congruence-kernels I and J , $\Theta(I \cap J) = \Theta(I) \cap \Theta(J)$,

(xi) $L \in B_n$ for some $1 < n < \omega$ and, given any congruence-kernels I and J , $\Sigma_n(I) \cap \Sigma_n(J) = \Sigma_n(I \cap J)$,

(xii) for any $0 \leq n \leq \omega$, the map $J \rightarrow \Sigma_n(J)$ is a lattice monomorphism of the lattice of congruence-kernels into the lattice of $*$ -congruences.

Proof. The equivalence of conditions (i), (ii), (iii) and (iv) follows from Lemma 3.1 as does the equivalence of conditions (v), (vi) and (vii).

Condition (iii) implies (v) by specialization and (v) \Rightarrow (viii) follows from Lemma 1.2 (ii).

Condition (viii) implies (i) by Theorem 2.5, and so the first eight conditions are equivalent.

Condition (ix) follows from (iv), by virtue of Lemma 1.3 and Theorem 1.5.

Suppose that (ix) holds. Let $x \in L$ and set $J = (x^{**}]$ and $F = [x)$. Then J is a congruence-kernel, $0 \equiv x \ (\Theta_{\text{Lat}}(J))$, and $x \equiv 1 \ (\Psi(F))$ so that $0 \equiv 1 \ (\Theta_{\text{Lat}}(J) \circ \Psi(F))$. By (ix), there exists $a \in L$ such that $0 \equiv a \ (\Psi(F))$ and $a \equiv 1 \ (\Theta_{\text{Lat}}(J))$. Hence $0 = a \wedge x$, that is, $a \leq x^*$, and $a \vee x^{**} = 1$. Thus (ix) \Rightarrow (i) holds and the first nine conditions are equivalent.

(iii) \Rightarrow (x) is clear and (x) \Rightarrow (vii) follows from Theorem 2.5 (iv), Theorem 1.6 (iv) and the distributivity of the lattice of ($*$ -)congruences. Of course, (xii) \Rightarrow (x) and (x) \Rightarrow (i), because setting $I = (x^*]$ and $J = (x^{**}]$ for given $x \in L$ implies that

$$x^* \vee x^{**} \equiv 1 \ (\Theta(I) \cap \Theta(J)) = \Theta(I \cap J) = \Theta(\{0\}) = \omega_L.$$

Since (i) and (xii) are equivalent, (xi) follows from (i) and the theorem will have been proved if we show that (xi) implies (i).

Suppose that (xi) holds. Put $I = (x^*]$ and $J = (x^{**}]$ for given $x \in L$. Then for $1 < n < \omega$ (fixed),

$$\Sigma_n(I) \cap \Sigma_n(J) = \Sigma_n(I \cap J) = \Sigma_n(\{0\}) = \omega_L,$$

as $L \in B_n$ (Theorem 2.5). By Theorem 2.10,

$$\Sigma_n(I) \vee \Sigma_n(J) = \Sigma_n(I \underline{\vee} J) = \Sigma_n(L) ,$$

as $x^* \underline{\vee} x^{**} = 1$, and so $\Sigma_n(I) \vee \Sigma_n(J) = 1_L$, the largest $*$ -congruence on L . Corollary 2.8 says that $\Sigma_n(I)$ and $\Sigma_n(J)$ are permutable. Thus, from the general theory of universal algebra ([3, §19, p. 120]), the map $a \mapsto ([a]_{\Sigma_n(I)}, [a]_{\Sigma_n(J)})$ is an isomorphism of L onto the direct product $L/\Sigma_n(I) \times L/\Sigma_n(J)$. Under this map $x^* \vee x^{**}$ and 1 both correspond to $(1, 1)$ and so $x^* \vee x^{**} = 1$. That is, $L \in B_1$ and so (xi) implies (i). \square

COROLLARY 3.3. *Let J be a congruence-kernel in a Stone lattice L . Then, $x \equiv y (R(J))$, $(x, y \in L)$, if and only if $(x \vee a)^{**} = (y \vee a)^{**}$ for some $a \in J$.*

Proof. By the previous theorem, R permutes with $\Theta_{\text{Lat}}(J) = \Theta(J)$ and so $R(J) = R \circ \Theta_{\text{Lat}}(J)$ (Theorem 1.6 (iv)). Thus $x \equiv y (R(J))$ implies that $x^{**} = z^{**}$ and $z \vee a = y \vee a$ for some $a \in J$. In any Stone lattice, $(s \vee t)^{**} = s^{**} \vee t^{**}$ is an identity, and so we obtain $(x \vee a)^{**} = (y \vee a)^{**}$ for some $a \in J$. The converse follows from the same identity and Theorem 1.6 (iii). \square

References

- [1] William H. Cornish, "Normal lattices", *J. Austral. Math. Soc.* 14 (1972), 200-215.
- [2] William H. Cornish, "Annulets and α -ideals in a distributive lattice", *J. Austral. Math. Soc.* (to appear).
- [3] George Grätzer, *Universal algebra* (Van Nostrand, Princeton, New Jersey; Toronto; London; Melbourne; 1968).
- [4] George Grätzer, *Lattice theory; first concepts and distributive lattices* (Freeman, San Francisco, 1971).

- [5] G. Grätzer and E.T. Schmidt, "Standard ideals in lattices", *Acta Math. Acad. Sci. Hungar.* 12 (1961), 17-86.
- [6] Iqbalunnisa, "Permutable congruences in a lattice", *Illinois J. Math.* 10 (1966), 235-239.
- [7] Joseph Kist, "Minimal prime ideals in commutative semigroups", *Proc. London Math. Soc.* (3) 13 (1963), 31-50.
- [8] K.B. Lee, "Equational classes of distributive pseudo-complemented lattices", *Canad. J. Math.* 22 (1970), 881-891.
- [9] T.P. Speed, "Some remarks on a class of distributive lattices", *J. Austral. Math. Soc.* 9 (1969), 289-296.
- [10] T.P. Speed, "Two congruences on distributive lattices", *Bull. Soc. Roy. Sci. Liège* 38 (1969), 86-95.

School of Mathematical Sciences,
Flinders University of South Australia,
Bedford Park,
South Australia.