

- 1 Zigmond T. Pressures to adhere to treatment: observations on 'leverage' in English mental healthcare. *Br J Psychiatry* 2011; **199**: 90–1.
- 2 Burns T, Yeeles K, Molodynski A, Nightingale H, Vazquez-Montes M, Sheehan K, et al. Pressure to adhere to treatment ('leverage') in English mental healthcare. *Br J Psychiatry* 2011; **199**: 145–50.
- 3 Buchy L, Ad-Dab'bagh Y, Malla A, Lepage C, Bodnar M, Joober R, et al. Cortical thickness is associated with poor insight in first-episode psychosis. *J Psychiatr Res* 2011; **45**: 781–7.
- 4 Buchy L, Czechowska Y, Chochol C, Malla A, Joober R, Pruessner J, et al. Toward a model of cognitive insight in first-episode psychosis: verbal memory and hippocampal structure. *Schizophr Bull* 2010; **36**: 1040–9.
- 5 Buchner F, Firestone M. Where the public peril begins: 25 years after Tarasoff. *J Leg Med* 2000; **21**: 25.
- 6 Becker T, Kilian R. Psychiatric services for people with severe mental illness across western Europe: what can be generalized from current knowledge about differences in provision, costs and outcomes of mental health care? *Acta Psychiatr Scand Suppl* 2006; **429**: 9–16.
- 7 Glancy DR, Glancy GD. The case that has psychiatrists running scared: Ahmed v. Stefaniu. *J Am Acad Psychiatry Law* 2009; **37**: 250–6.
- 8 Glancy GD, Chaimowitz G. Tarasoff warnings result in criminal charges. *J Am Acad Psychiatry Law* 2003; **31**: 524–5.
- 9 Gutheil TG. In search of true freedom: drug refusal, involuntary medication, and 'rotting with your rights on'. *Am J Psychiatry* 1980; **137**: 327–8.
- 10 Kress K. Rotting with their rights on: why the criteria for ending commitment or restraint of liberty need not be the same as the criteria for initiating commitment or restraint of liberty, and how the restraint may sometimes justifiably continue after its prerequisites are no longer satisfied. *Behav Sci Law* 2006; **24**: 573–98.

Mansfield Mela, Department of Psychiatry, College of Medicine, University of Saskatchewan, Canada. Email: mansfieldmela@gmail.com

doi: 10.1192/bjp.200.1.81a

Author's reply: Thank you to Drs Scott-Orr and Mela for their interest. It seems to me that there are two issues here. First, should the law be discriminatory between patients with a physical illness and those with a mental illness? I think not and I'm pleased to say the United Nations Convention on the Rights of Persons with Disabilities (2006), to which the UK is a signatory, supports this view. The convention obligates States to (among many other things) 'take all appropriate measures, including legislation, to modify or abolish existing laws, regulations, customs and practices that constitute discrimination against persons with disabilities'. To explain this further, the UN High Commissioner for Human Rights said,¹

'Legislation authorizing the institutionalization of persons with disabilities on the grounds of their disability without their free and informed consent must be abolished. . . This should not be interpreted to say that persons with disabilities cannot be lawfully subject to detention for care and treatment or to preventive detention, but that the legal grounds upon which restriction of liberty is determined must be de-linked from the disability and neutrally defined so as to apply to all persons on an equal basis.'

Second, should the law (for everyone) favour patient autonomy, medically determined best interest or a mixture?

In other words, either everyone, with the capacity to make the decision, should be permitted to 'die (or rot) with their rights on' or nobody should. Or the authority to overrule capacious refusal could be based on a neutral factor such as risk to other people. It should not be dependent on the stigma associated with certain terminology (a mental illness diagnosis).

- 1 United Nations. *Annual Report of the United Nations High Commissioner for Human Rights and Reports of the Office of the High Commissioner and the Secretary-General: A/HRC/10/48, 26 January 2009*. United Nations, 2009.

Tony Zigmond, Newsam Centre, Leeds LS14 6WB, UK. email: azigmond@doctors.net.uk

doi: 10.1192/bjp.200.1.82

Family psychoeducation for major depression: randomised controlled trial

The paper by Shimazu *et al*¹ adds robustness to already existing evidence for the role of family psychoeducation in psychiatric disorders. The study has sound methodology (i.e. randomised controlled trial) with adequate masking, in addition to being the first ever study to examine the effect of family psychoeducation for major depressive disorder. The authors describe the possible limitations of the study honestly. A source of funding (Grant-in-Aid for Scientific Research, Ministry of Health, Labour and Welfare, 2004) is also mentioned. However, there are some issues which should be further looked into. The aim was to examine family psychoeducation in the maintenance treatment of depression.

However, the patients included were either on continuation or maintenance treatment. Patients who are in partial remission cannot be considered as being in a continuation/maintenance phase.² Also it was not mentioned how many patients had single or recurrent episodes (patients with single episodes need not receive maintenance phase treatment). Any other psychiatric comorbidity (substance misuse or personality disorder) in the participants was not mentioned, even though it has treatment implications. The health status and intellectual functioning of the primary family member included in the study was not mentioned, although these might compromise their active participation in psychoeducation sessions. The authors are silent on the ethical clearance of the study. For four caregivers psychoeducation sessions were done in the individual's home and not in group sessions, which were not included in the final analysis and not part of the methodology mentioned – this could also have had an effect on the efficacy of the study. Remission was defined by the authors as a Hamilton Rating Scale for Depression (HRSD) score <6, but the normal score is mentioned as <7.³ The authors have not mentioned the reason for keeping a low score of HRSD in the study. Last, it could be a printing mistake, in the last line of Table 1 it is mentioned that high expressed emotion (as per FMSS) was seen in seven patients in the intervention group and none in the control group, but the results mention that it is seen in six patients in the intervention group and ten in the control group.

- 1 Shimazu K, Shimodera S, Mino Y, Nishida A, Kamimura N, Sawada K, et al. Family psychoeducation for major depression: randomised controlled trial. *Br J Psychiatry* 2011; **198**: 385–90.
- 2 American Psychiatric Association. *Practice Guideline for the Treatment of Patients with Major Depressive Disorder, Third Edition*. APA, 2010.
- 3 Blacker D. Psychiatric rating scales. In *Kaplan and Sadock's Comprehensive Textbook of Psychiatry, Ninth Edition, Volume 1* (eds BJ Sadock, VA Sadock, P Ruiz): 1047. Lippincott Williams and Wilkins, 2009.

Bichitra N. Patra, Department of Psychiatry, Postgraduate Institute of Medical Education and Research, Chandigarh – 160012, India. Email: patrab.aalms@gmail.com; **B. N. Subodh**, Department of Psychiatry, Postgraduate Institute of Medical Education and Research, India

doi: 10.1192/bjp.200.1.82a

Author's reply: We thank Patra & Subodh for their interest in and their very thorough reading of our study.¹ Most of their questions are factual ones and we are grateful that we have been given an opportunity to clarify them. First, whether to call further treatment of patients in partial or full remission after the fully syndromatic episode, as in our study, continuation/maintenance treatment is a terminological issue and not a medically substantive one. And we think our usage of the terms is in consonance with the majority of psychiatrists of the world, as for example done by Paykel *et al* in their famous study of cognitive therapy to prevent relapse after acute episode of major depression.²