

VOL. VII.—1893.

THE JOURNAL
OF
LARYNGOLOGY,
RHINOLOGY, AND OTOTOLOGY;

*AN ANALYTICAL RECORD OF CURRENT LITERATURE RELATING TO
THE THROAT, NOSE, AND EAR.*

EDITED BY

R. NORRIS WOLFENDEN, M.D. Cantab.,
JOHN MACINTYRE, M.B. Glasgow,

AND

DUNDAS GRANT, M.A., M.D., F.R.C.S. Eng.,

WITH THE ASSISTANCE OF

Dr. BARCLAY J. BARON (Bristol). Dr. SANDFORD (Cork).

And with the co-operation of

DR. RICARDO BOTEY (BARCELONA)
DR. BRYSON DELAVAN (NEW YORK)
DR. DRAISPUL (ST. PETERSBURG)
PROF. GUYE (AMSTERDAM)
DR. MIDDLEMASS HUNT (LIVERPOOL)
DR. VALERIUS IDELSON (BERNE)
DR. JOAL (PARIS)
DR. KARWOWSKI (WARSAW)
DR. JOHN N. MACKENZIE (BALTIMORE)
DR. GRAZZI (FLORENCE)
DR. SANDFORD (CORK)
DR. LUBET-BARBON (PARIS)

DR. G. W. MAJOR (MONTREAL)
PROF. MASSEI (NAPLES)
DR. W. ROBERTSON (NEWCASTLE-ON-TYNE)
DR. J. MICHAEL (HAMBURG)
DR. HOLGER MYGIND (COPENHAGEN)
DR. PORTER (ST. LOUIS)
DR. SAJOUS (PHILADELPHIA)
DR. SENDZIAK (WARSAW)
PROF. RAMON DE LA SOTA (SEVILLE)
DR. HICGUET (BRUSSELS)
DR. MILLIGAN (MANCHESTER)
DR. LICHTWITZ (BORDEAUX)

Published Monthly.

LONDON:
F. J. REBMAN,
Albion Chambers, 11, Adam Street, Strand.

PHILADELPHIA, PA.:
THE F. A. DAVIS COMPANY,
1914-16, Cherry Street.

ENTERED AT STATIONERS' HALL.

CONTENTS.

Original Article:—

Simplest and Most Practical Electric Apparatus, especially for Galvano-Cautery Operations. By
Dr. A. G. Hobbs (Atlanta). Page 519.

Therapeutics and Diphtheria.—Electric Lamp, *Winckler*—Therapeutic Value of Hydro-Carbons, *Bayer*—Cocaine for Local Anesthesia, *Rhodes*—Diphtheria and Infected Milk Supply, *Applegate*
—Muscular Anomalies in Diphtheria, *Rauberg*. Page 521.

Mouth, Pharynx, &c.—Treatment of Severe Hare Lip, *Thomas*—Cleft Palate and Hare Lip, *Forbes*—Lymphoid Growths in the Vault of the Pharynx, *French*—Organic Stricture of the Esophagus, *Brown*. Page 522.

Nose and Naso-Pharynx.—Empyema of Frontal Sinus, *Lichtwitz*—Diagnosis and Treatment of Suppuration of the Accessory Cavities of the Nose, *Oaks*—Draining the Antrum of Highmore, *Major*—Operative Treatment of Deviations, *Hamburger*—Electrolysis for Deviations, *Schmidt*—Unilateral Membranous Rhinitis, *Dunn*—Hypertrophic Rhinitis and Asthenopia, *Bernstein*—Treatment of Atrophic Rhinitis, *Cox*—Ozone in Atrophic Catarrh, *Rice*—Diameters of the Septum and Naso-Pharynx, *Hopmann*—The Galvano-Cautery in Diseases of the Nose, etc., *Heymann*—Lateral Naso-Pharyngeal Diverticula, *Schmidt*—Adenoid Vegetations, *Sympson, Hardie*. Pages 523 to 527.

Larynx.—Onodi's Researches on Laryngeal Paralysis, *Neumann*—Paralysis of Recurrent Laryngeal Nerve by Scrofulous Glands, *Finlayson*—Glottic Spasm, with Unconsciousness, in the Adult, *Porcher*—Tuberculous Laryngeal Tumours, *Hajek*—Sarcoma of the Larynx, *Bruck*—Foreign Body in the Lung, *Schild*. Pages 527 to 529.

Neck.—Goitre, *Sulzer, Kronlein*—Basedow's Disease, *Marcus, Freiberg*—Struma Maligna, *Speyer*
—Metastases of Bones in Thyroid Cancer, *Eiselberg*—Angina Ludovici, *Trumpp*. Page 529.

Ear.—Foreign Body for Thirty-nine Years, *Conners*—Aural Syphilis, *Toeplitz*—Clinical Lecture on Middle-Ear Suppuration, *Politzer*—Treatment of Chronic Tympanic Vertigo, *Burnett*—Mastoid Measurements, *Coues*. Pages 530 to 533.

Society Meetings:—

American Laryngological Association—Report of Meeting.	Pages 534 to 543.
Pan-American Medical Congress—Report of Meeting.	Pages 543 to 556.
Australasian Medical Congress—Report of Meeting.	Pages 557 to 575.
Sixty-fifth <i>Versammlung Deutscher Naturforscher und Aerzte</i> .	Page 575.
Swedish Medical Society.	Page 577.
Finnish Medical Society.	Page 578.

Notes. Page 578.

GUIDE TO ADVERTISEMENTS.

	PAGE		PAGE
Allen & Hanburys	vi.	Ingram & Royle	v.
Balcomb	vii.	Ivins	iii.
Bullock & Co.	iii.	Lambert Pharmaceutical Co.	ii.
Burroughs, Wellcome & Co.	viii., xii.	Parke, Davis & Co.	iv.
Central London Throat Hospital	xiv.	Rebman	iv., ix., x.
Godfrey & Cooke	iv.		

The Kepler Solution of Cod Liver Oil in Malt Extract.

The Kepler Solution of Cod Liver Oil in Extract of Malt is free from the many objectionable constituents and characteristics of ordinary emulsions, and since its introduction the use of Cod Liver Oil in Emulsion has steadily decreased. This change in the method of administering this invaluable remedial agent was evidently because medical men found that emulsions, however well prepared, created such a disturbance of the digestive functions that the very idea of taking another dose would cause nausea in the patient; consequently the physician in many cases was seriously hampered in the successful control of cases by the disinclination and frequent positive refusal on the part of the sufferer to take the preparation prescribed. All this is changed in the case of the Kepler Solution of Cod Liver Oil. It has but little if any odour or taste of the oil; consequently patients—even fastidious invalids and children—will take it readily, and even with avidity, after having once tasted it. In the case of emulsions made on the saponification method—for the process can be nothing else where alkalies are used—the first result of its ingestion must naturally be that a large percentage of the Oil becomes free, and the fact is too well known that the digestion of uncombined oil is a task hopelessly beyond the power of the general run of those for whom Cod Liver Oil is necessary. Drs. Leaffenen and Beam recently investigated the various Cod Liver Oil Emulsions in the market, and if their work shows anything it is that emulsions are not to be relied upon as to the asserted percentage of Oil. It would have been equally instructive to know what quality Oil had been used and what were the emulsifying or so-called emulsifying media. The latter point is a highly important one to medical men, as it bears directly upon the value of the preparation to the patient, the index of its therapeutic worth being exactly the percentage of Oil taken up by the organism. Again, there is nothing in simple emulsions to assist the digestion and assimilation as in the case of Kepler Solution—a question of immense importance. And if it be a question of combinations such as are frequently necessary in the treatment of the consumptive, the scrofulous, the invalid, and the convalescent, we are assured that many indispensable adjuncts are inadmissible with emulsions, since they either disturb the balance in the emulsion itself, or else are themselves being acted upon, become decomposed, and consequently the scheme of the physician is at once broken up. The association of a first-class Malt Extract such as the Kepler with the very best Cod Liver Oil obtainable (without regard to price) is in accordance with the natural requirements of the stomach, for, as Dr. Lauder Brunton, one of the highest authorities one can quote, very pithily says, when discussing the administration of fat, “If anyone were to swallow a lump of butter by itself, it would probably make him sick; but if he spread the same butter upon pieces of bread he could take it, not only without discomfort, but with actual enjoyment.” Here is at once the reason why the Kepler Solution is so easily borne, so readily taken, so quickly digested, and so highly nutritious.

Kepler Solution and Kepler Extract of Malt supplied to the Medical Profession by all Chemists in $\frac{3}{4}$ lb. and $\frac{1}{2}$ lb. bottles at 1s. 8d. and 3s. each.

BURROUGHS, WELLCOME & CO., Manufacturing Chemists,
SNOW HILL BUILDINGS, LONDON, E.C.