

Results:

Features		Total	26	%
Clinical	Diagnosis	Dementia	19	73.1
		Schizophrenia	2	
		Depression	4	
		Obsessive Compulsive Disorder	1	
	Behavioral difficulties	18	69.2	
	Activities of Daily Living (ADLs) assistance	22	84.6	
Social	Caregiver burnout	18	69.2	
	Poor relationships	3	11.5	
	Inadequate additional caregiver support	3	11.5	

Demographic profile: Out of 57 inpatients (28 male patients and 29 female patients), 26 patients (46%) were over-stayers. Out of these 26 over-stayers, 18 patients (69%) were female and 14 patients (54%) were above age 70.

Clinical profile (n=26): 19 patients (73.1%) were diagnosed with dementia. 18 patients (69.2%) had severe Behavioral and Psychological Symptoms of Dementia (BPSD). 22 patients (84.6%) required assistance in their basic activities of daily living.

Social profile (n=26): Caregivers of 18 patients (69%) were burnt out by patient's behavior problems, which is commonly seen in caregivers for patients with dementia[ii]. Family members of 3 patients (11.5%) were estranged from them. Caregivers of 3 patients (11.5%) had difficulty engaging additional caregiver support for ADL assistance.

[ii] Reuben DB, Romero T, Evertson LC, Jennings LA. Overwhelmed: a Dementia Caregiver Vital Sign. *J Gen Intern Med.* 2022 Aug;37(10):2469-2474. doi: 10.1007/s11606-021-07054-3. Epub 2021 Aug 13. PMID: 34389938; PMCID: PMC9360256.

Conclusions: The above profiles enabled the department of Geriatric Psychiatry in IMH to identify elderly patients at risk of prolonged hospital stay at the beginning of their admission and improve the care of these patients to reduce their length of stay.

Disclosure of Interest: None Declared

Others 01**EPP0219****The effect of chronobiology on frontal functions and social functionality in remitted patients with schizophrenia**

B. Çınar^{1*}, S. Gıca² and N. Karamustafaloğlu³

¹psychiatry, Ardahan Public Hospital, Ardahan; ²psychiatry, Necmettin Erbakan Üniversitesi Meram Tıp Fakültesi, Konya and ³psychiatry, Istanbul Bakirkoy Prof. Dr. Mazhar Osman Research

and Training Hospital for Psychiatry, Neurology and Neurosurgery, İstanbul, Türkiye

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.549

Introduction: Due to the effects of sleep on the central nervous system, it is thought that sleep problems have a special importance in the onset, course and treatment of psychiatric diseases. Although the negative effects of sleep problems on the occurrence, recurrence and clinical course of psychiatric disorders are well known, it is reported that clinicians do not spend enough time for sleep problems in practice. To our knowledge there is no study in the literature which examining its effect on frontal lobe functions or social functionality in schizophrenia.

Objectives: In current study, it is aimed i- to examine the chronobiological characteristics of remitted patients with schizophrenia, ii- to determine the effect of chronobiology on sleep quality, frontal lobe functions, depressive symptoms, interpersonal relationships and social functionality in patients with schizophrenia, iii- thus to improve the quality of life and the treatment outcome of patients with schizophrenia.

Methods: 185 patients with schizophrenia who met the Andreasen remission criteria were included in the study. The patients were evaluated with Positive and Negative Syndrome Scale (PANSS), Morningness & Eveningness Scale (MEQ), Calgary Depression Index (CDI), Pittsburgh Sleep Quality Index (PSQI), Barratt Impulsivity Scale Short Form (BIS-11-SF), Frontal Assessment Battery (FAB), Personal Social Performance Scale (PSPS). Each patient was classified either as morning type (MT) or evening type (ET) or intermediate type (IT) according to MEQ scores.

Results: The obtained MEQ responses indicated that 29 (15.7%) of the patients were "ET", 124 (67.0%) were "IT", and 32 (17.3%) were "MT". In the "ET" group, the mean of CDI total score, the mean of PSQI total score, the mean of BIS-11-SF total and the mean score of BIS-11-SF planlessness and inattention subscale were higher than in both the "IT" and "MT". The mean total scores PSPS of "ET" patients were found to be lower than the mean scores of both the "IT" and "MT" patients. There was no significant difference between the groups in terms of FAB total scores. According to Multiple Linear Regression Analysis, total MEQ score and PANSS-negative symptoms subscale score were found to have an effect on PSPS[1] Personal Social Relationship subscale scores.

Conclusions: Although all patients included in current study were in remission, it was found that chronotype characteristics were effective on many clinical manifestations and comorbid conditions. The findings obtained from our study emphasize how important it is to question chronotypic features in daily psychiatric practice. In this context, being aware of the importance of chronotypic features in the treatment of schizophrenia patients, questioning the patients in this respect and taking necessary interventions may have the potential to improve functionality, which is one of the main treatment goals in patients with schizophrenia.

Disclosure of Interest: None Declared