

NOTES FOR AUTHORS

Papers submitted for publication in the *Bulletin of Entomological Research* should be sent, in duplicate, to the Director, C·A·B International Institute of Entomology, 56 Queen's Gate, London SW7 5JR. They must deal with original research concerning insects, mites or ticks of economic importance in the agricultural, medical or veterinary fields; they must be of more than purely local interest, and taxonomic papers will not be accepted unless the organisms dealt with are of definite importance in the field of applied entomology. Review articles may also be submitted, but will normally be by invitation. Papers must not have been published or accepted for publication elsewhere. They will normally be published in order of their receipt in definitive form.

For general guidance on presentation and style, including the preparation of illustrations, contributors are urged to consult *Writing Scientific Papers in English. An ELSE-Ciba Foundation Guide for Authors* (by Maeve O'Connor & E. Peter Woodford). 116 pp., Amsterdam, Associated Scientific Publishers, 1975, ISBN 90-219-4035-3.

Text

Papers should be typewritten (double spaced, with ample margins and on one side of the paper only) and a current number of the *Bulletin* should be consulted for details of style and lay-out. Titles should be as brief as is consistent with clarity. Abbreviations should conform to *British Standard 1991, Part 1, 1967*, and are alike in singular and plural; stops will be used only where ambiguity might occur, e.g. 'in.' not 'in' for inch, but 'in²' for square inch. Metric units are preferred; if non-metric units are used, metric equivalents should be given at least once in each article.

Tables

Tables should be reduced to the simplest form, and should not be used where text or illustrations give the same information. They should be submitted on separate sheets at the end of the article and must on no account be of a size or form that will not conveniently fit into the page of the *Bulletin*.

Illustrations

Line drawings and graphs must be in jet-black indian ink, preferably on Bristol-board or tracing cloth. Graphs may be drawn on paper ruled in pale blue or light grey; paper having green, red, yellow or brown lines should not be used. Any stippling or hatching, as well as any numerals or lettering, on the text-figures must be completed, by the author, in a form suitable for reproduction; this work is *not* undertaken by the printer. Text-figures, when reproduced, cannot exceed 12·7 cm in width; if, therefore, the original drawings are larger, the lettering on them must be in proportion, so that it is still legible when reduced; special attention should be paid to the relative scales of lettering and line thicknesses in relation to the figure as a whole. Photographs for reproduction as figures are admissible when they are a real contribution to the text. Captions to figures should be typed on a separate sheet.

Abstract

Each paper must contain a carefully prepared and accurate informative abstract that shall be complete in itself and intelligible without reference to the text or figures. It should not exceed 250 words, and should be included at the beginning immediately after the title, author and author's affiliation.

References

References must be based on the name and year system and conform to the present usages of the *Bulletin*. Titles of journals should be abbreviated as in the 4th edition of *The World List of Scientific Periodicals*. References to unpublished information should normally be included only in the text—(A. B. Jones, pers. comm.) or (A. B. Jones, Control of sandflies by insecticides. Unpublished report. Ross Institute, London, 1950).

Offprints

Free copies of offprints are allowed as follows: One author—20 copies, two or more authors—30 copies. Further copies may be obtained on payment, and the number required should be specified when the paper is submitted.

BULLETIN OF ENTOMOLOGICAL RESEARCH

(World List Abbreviation: *Bull. ent. Res.*)

ISSN 0007-4853

	PAGE
VORLEY, V. T. & WRATTEN, S. D. Migration of parasitoids (Hymenoptera: Braconidae) of cereal aphids (Hemiptera: Aphididae) between grassland, early-sown cereals and late-sown cereals in southern England	555
LUGURU, S. M., CHIZYUKA, H. G. B. & MUSISI, F. L. A survey for resistance to acaricides in cattle ticks (Acari: Ixodidae) in three major traditional cattle areas in Zambia	569
ERZINÇLIOĞLU, Y. Z. The larval instars of the African blowfly, <i>Calliphora croceipalpis</i> Jaenicke, with a key to the genera of the third instars of African carrion-breeding Calliphoridae (Diptera)	575
FORRESTER, N. W. Diapause development in the sorghum midge, <i>Contarinia sorghicola</i> (Coquillett) (Diptera: Cecidomyiidae)	581
SAUL, A. Estimation of survival rates and population size from mark-recapture experiments of bait-caught haematophagous insects	589
MANSOUR, F., KARCHI, Z. & OMARI, N. Resistance of melon to the carmine spider mite, <i>Tetranychus cinnabarinus</i> (Boisduval) (Acari: Tetranychidae)	603
BROWNE, L. B., VAN GERWEN, A. C. M. & SMITH, P. H. Relationship between mated status of females and their stage of ovarian development in field populations of the Australian sheep blowfly, <i>Lucilia cuprina</i> (Wiedemann) (Diptera: Calliphoridae)	609
THORNHILL, W. A. & HEATHCOTE, G. D. Aphid (Hemiptera: Aphididae) populations and the incidence of virus yellows on insecticide-free sugarbeet in south-eastern England in 1978-81	617
AALBERSBERG, Y. K., DU TOIT, F., VAN DER WESTHUIZEN, M. C. & HEWITT, P. H. Development rate, fecundity and lifespan of apterae of the Russian wheat aphid, <i>Diuraphis noxia</i> (Mordvilko) (Hemiptera: Aphididae), under controlled conditions	629
AALBERSBERG, Y. K., VAN DER WESTHUIZEN, M. C. & HEWITT, P. H. A simple key for the diagnosis of the instars of the Russian wheat aphid, <i>Diuraphis noxia</i> (Mordvilko) (Hemiptera: Aphididae)	637
MOORE, D. & ALEXANDER, L. Aspects of migration and colonization of the coconut palm by the coconut mite, <i>Eriophyes guerreronis</i> (Keifer) (Acari: Eriophyidae)	641
GUNN, A. & GATEHOUSE, A. G. The influence of larval phase on metabolic reserves, fecundity and lifespan of the African armyworm moth, <i>Spodoptera exempta</i> (Walker) (Lepidoptera: Noctuidae)	651
CRANSTON, P. S. A non-biting midge (Diptera: Chironomidae) of horticultural significance	661
RUBIA, E. G. & SHEPARD, B. M. Biology of <i>Metioche vittaticollis</i> (Stål) (Orthoptera: Gryllidae), a predator of rice pests	669
CARTER, N. Management of cereal aphid (Hemiptera: Aphididae) populations and their natural enemies in winter wheat by alternate strip spraying with a selective insecticide	677
WEBB, M. D. Species recognition in <i>Cicadulina</i> leafhoppers (Hemiptera: Cicadellidae), vectors of pathogens of Gramineae	683
BLACKMAN, R. L. Morphological discrimination of a tobacco-feeding form from <i>Myzus persicae</i> (Sulzer) (Hemiptera: Aphididae), and a key to New World <i>Myzus</i> (<i>Nectarosiphon</i>) species	713
RIDS DILL-SMITH, T. J., HAYLES, L. & PALMER, M. J. Mortality of eggs and larvae of the bush fly, <i>Musca vetustissima</i> Walker (Diptera: Muscidae), caused by scarabaeine dung beetles (Coleoptera: Scarabaeidae) in favourable cattle dung	731
INDEX OF AUTHORS	737
INDEX OF GENERA AND SPECIES	743

© C·A·B International, 1987

All rights reserved. No part of this publication may be reproduced, in any form or by any means, electronically, mechanically, by photocopying, recording or otherwise, without prior permission of the copyright owner.