




# Impact of crown architecture on light availability, gas exchange, flowering and fruiting in jamun (*Syzygium cumini* [L.] Skeels)

Ajaya Kumar Trivedi , Anand Kumar Singh, Krishna Kumar Mishra and Gaurav Singh Vishen

ICAR – Central Institute for Subtropical Horticulture, Lucknow, Uttar Pradesh, India

## Crops and Soils Research Paper

**Cite this article:** Trivedi AK, Singh AK, Mishra KK, Vishen GS (2024). Impact of crown architecture on light availability, gas exchange, flowering and fruiting in jamun (*Syzygium cumini* [L.] Skeels). *The Journal of Agricultural Science* **162**, 458–467. <https://doi.org/10.1017/S0021859624000443>

Received: 23 January 2024  
Revised: 24 April 2024  
Accepted: 12 May 2024  
First published online: 27 November 2024

### Keywords:

canopy; fruit yield; leaf area index; light interception; photosynthesis

### Corresponding author:

Ajaya Kumar Trivedi;  
Email: [ak.trivedi@icar.gov.in](mailto:ak.trivedi@icar.gov.in)

### Abstract

Experiments were conducted to assess the impact of crown architecture on light availability beneath the trees, flowering, fruiting, yield and quality of jamun (*Syzygium cumini* [L.] Skeels). Trees were maintained as control, palmette and open centre crown. Impact was evaluated for three consecutive years, i.e. 2017–2019. Diffuse light beneath the trees ranged from  $69.7 \pm 2.22$  to  $45.9 \pm 1.45\%$ , whereas direct light varied from  $30.4 \pm 0.97$  to  $54.1 \pm 1.78\%$ . At flowering and fruit development stage (June), photosynthesis rate (A) in control trees was  $12.5 \pm 0.43 \mu\text{mol CO}_2/\text{m}^2/\text{s}$ ; however, at fruit maturity and dormancy (August), it was only  $9.5 \pm 0.35 \mu\text{mol CO}_2/\text{m}^2/\text{s}$ . Similarly, in palmette and open centre trees, photosynthesis rate at flowering and fruit development stage was  $13.5 \pm 0.46$  and  $15.7 \pm 0.54 \mu\text{mol CO}_2/\text{m}^2/\text{s}$ , respectively; whereas at fruit maturity and dormancy, photosynthesis rate dropped to  $10.5 \pm 0.39$  and  $11.7 \pm 0.43 \mu\text{mol CO}_2/\text{m}^2/\text{s}$ , respectively. Substantial variation in stomatal conductance (gs), vapour pressure deficit (VPD) and transpiration rate (E) was also found. Days to start flowering ranged from  $92 \pm 0.33$  to  $98 \pm 0.33$ . Similarly, days to end flowering varied from  $99 \pm 0.07$  to  $107 \pm 0.36$ , days to fruit set  $132 \pm 0.33$  to  $139 \pm 0.33$  and days to fruit maturity  $176 \pm 0.48$  to  $184 \pm 0.63$ . Significant variation in fruit length, fruit width and fruit weight was also found. Total soluble solids in fruit pulp varied from  $9.0 \pm 0.15$  to  $12.2 \pm 0.149^\circ\text{Brix}$  and fruit yield  $62.3 \pm 1.5$  to  $86.7 \pm 1.33$  kg per tree. Noteworthy variation in fruit quality traits was also recorded. This study illustrates that crown architecture has considerable impact on gas exchange parameters, flowering, fruiting, yield and quality of jamun.

### Introduction

The jamun (*Syzygium cumini* [L.] Skeels), a member of family Myrtaceae is one of the important underutilized evergreen fruits widely distributed throughout tropical and sub-tropical region in India as well as in certain pockets of the lower Himalayan ranges up to an elevation of 1600 m (Mishra *et al.*, 2014). It is an important indigenous fruit tree of India, originated from Indonesia and India, now growing abundantly in Southern Asia (Periyathambi, 2007) and the Pacific Islands. It is widely cultivated in the Indo-Gangetic plains. Jamun is also known by other common names viz., Java plum, black plum, jambul and Indian blackberry, etc. Jamun has been attributed in the Indian folklore medicine system to possess several medicinal properties (Warrier *et al.*, 1996). The fruit is a good source of anthocyanins, iron, pectin, phenols and protein (Ghosh *et al.*, 2017). Leaves and small twigs are used for feeding cattle, particularly during summer and dry spell.

Jamun tree canopy is characterized by profuse foliage which substantially affects light availability in different inner canopy layers. Owing to cross-pollination and seed propagation, enormous variability is available in jamun with respect to vegetative growth, aspects of spread, leaf shape, tree shape, canopy architecture, fruiting habit, maturity (June–August), fruit yield and quality. New vegetative shoots emerge as terminal growth on the previous season twigs in two distinct flushes, i.e. from February to May and August to October. The flush, which appears in the month of February, provides maximum growth and flowering. Flowering occurs in terminal as well as auxiliary inflorescences on 5-month to 1-year-old branches.

It is one of the most hardy fruit crops that can easily be grown even in areas where other fruits fail to establish (Singh *et al.*, 2007). Due to wider adaptability, jamun may be a suitable fruit crop in semiarid, arid, saline, sodic, ravine, degraded and wasteland areas where it is difficult to grow other fruit crops. In climate change scenario, it may be a potential fruit crop for resource-poor areas of tropical and subtropical region.

Jamun is a vigorous tree; hence, canopy management is important for ensuring maximum utilization of light, ease of cultural operations and maximizing the productivity and quality. Crowded canopy in jamun trees causes decreased light availability in different inner canopy layers. This in turn affects flowering, fruiting and productivity. Positive effects of diffuse

light on plant growth have been well recognized and have been applied to natural communities (Norman and Miller, 1971; Norman and Arkebauer, 1991). However, so far no information is available about the relative ability of individual leaves to utilize direct vs. diffuse light for photosynthesis (Brodersen *et al.*, 2008). Effects of direct vs. diffuse light may be different at leaf level as compared to canopy level. Within canopy, light distribution is influenced by direction of light, fraction of diffuse or direct light incident on the canopy as well as canopy architecture (Li *et al.*, 2014). Canopy architecture has substantial impact on light distribution and photosynthesis (Sarlikioti *et al.*, 2011).

In tree crops having multilayered canopy, light availability in different layers is affected by foliage structure as well as canopy architecture (Maddonna *et al.*, 2001; Acreche *et al.*, 2009). To enhance gross primary production, direct diffuse light ration is important rather than absolute direct or diffuse light (Bodin and Franklin, 2012). However, so far no systematic work has been carried out to find out the impact of crown architecture on light availability, flowering, fruiting, yield and quality of jamun. Hence, the present study was conducted with the objective to study the impact of the different tree crown architectures on flowering, fruiting, yield and fruit quality of jamun in subtropical region.

## Materials and methods

The present study was conducted for three consecutive years (2017–2019) at the experimental farm of ICAR – Central Institute for Subtropical Horticulture, Lucknow (India), located at 26° 45' to 27° 10' N latitude, 80° 30' to 80° 55' E longitude and 123 m above mean sea level. The study area falls under humid subtropical region. Fully grown, 10-year-old jamun trees of the genotype CISH J-37 (Jamwant) planted at 5 m × 5 m spacing (400 plants/ha) were selected for the study.

Experimental plants were planted in pits (90 × 90 × 90 cm) dug during the summer months. Young plants were allowed 3–5 well-spaced scaffold branches 60 cm above from the ground level to develop the main framework. It was followed by pruning to regulate tree size and shape to achieve the desired architecture of the canopy with a network of primary, secondary and tertiary branches. Treatments (i.e. tree architectures – control, open centre and palmette) were imposed from the beginning of the experiment in young plants by removing the undesired branches to properly maintain the desired crown architecture.

### Light interception study

Light availability (direct vs. diffuse light) study was conducted with the help of hemispherical photography done by HemiView canopy analyser Version 2.1 (Delta-T Devices Limited, Cambridge, UK). Canon SLR camera with fisheye lens for DCM9 camera was used in the device with linear 180 lens selection. The basic model for the estimation of solar radiation was used for the estimation of direct and diffuse light. The instrument was placed beneath the tree canopy on a tripod 1.5 m above ground level. Light intensity was measured with lux meter (PCE instruments, PCE-LM 4).

### Photosynthesis and gas exchange parameters

For the study of gas exchange parameters viz., photosynthesis rate, stomatal conductance, vapour pressure deficit and transpiration

rate, CIRAS-3 portable photosynthesis system (PP Systems International, Inc. Amesbury, MA, USA), i.e. infrared gas analyser was used. In CIRAS-3 portable photosynthesis system, ambient conditions were selected for CO<sub>2</sub> and H<sub>2</sub>O reference (chemicals were removed – both CO<sub>2</sub> and H<sub>2</sub>O absorber columns were empty). After approximately 10 min, warm up time system performed several periodic zero and differential balance cycles and gradually stabilized and frequency reduced to every 30 min. After approximately 5 min, CO<sub>2</sub> reference, CO<sub>2</sub> analysis, H<sub>2</sub>O reference and H<sub>2</sub>O analysis were stable. CO<sub>2</sub> and H<sub>2</sub>O differential was 0.0 (±1). Ambient and leaf temperature was same (±0.2°C). Light availability and gas exchange parameters were recorded weekly between 10.00 h and 11.00 h from April to August. Forth fully expanded mature leaf was selected for recording observations. Observations were recorded in 20 leaves (five leaves in each direction viz., east, west, north and south) and five readings were recorded in each leaf. Average of weekly data of each month was calculated and expressed as mean of the particular month. Year-to-year variability in the averaged monthly data was non-significant. Thus, mean monthly data of both the year were pooled and mean values are given in the figures and tables. Performance of trees was consistent during both the years; hence, average of monthly data of both the years has been presented in the tables and figures.

### Flowering and fruiting characters

For flowering data, 100 inflorescences in each tree (25 inflorescences in each direction, i.e. east, west, north and south) were tagged in each plant and observations were recorded from tagged inflorescences. Finally, 100 fruits from each tree, i.e. 25 fruits from each direction viz., east, west, north and south were taken for observations and analyses.

Number of days from 1 January to 5% flower appearance has been expressed as days to start flowering and number of days from 1 January to 95% flower appearance has been expressed as days to end flowering. Number of days from 1 January to fruit at pin head stage has been denoted as days to fruit set and number of days from 1 January to fruit at harvesting stage has been mentioned as days to maturity. Fruit length and fruit width were measured with the help digital Vernier calliper. Weight of ten representative fruits of different sizes at ripe stage was recorded with the help of electronic pan balance and average is expressed as fruit weight (g).

### Biochemical parameters

Total soluble solids at full ripe stage (ready to eat) was measured with the help of the refractometer and expressed as °Brix. The edible portion of fruits was crushed and homogenized by the pestle and mortar method without peeling off the skin. The homogenized sample was transferred into a 100 ml volumetric flask and 50% ethanol was added to maintain the volume up to the mark. The mixture was shaken manually and then filtered. Filtrate was centrifuged to obtain a clear supernatant liquid, which was subsequently used for the various assays. The extracts were stored at –20°C. All tests were performed within a week. Total anthocyanin contents in fresh (ready to eat) fruits were measured by pH differential method using cyanidin hydrochloride as standard (Cheng and Breen, 1991). Total phenol contents (TPC) were estimated by spectrophotometric method (Shimadzu UV 2550 spectrophotometer) using Folin–Ciocalteu reagent method (Slinkard and Singleton, 1977). Ascorbic acid

estimation has been done by spectrophotometric method using dichlorophenol indophenol dye solution (Davis and Masten, 1991). Scavenging DPPH free radical activity was determined by the method described by Yue and Xu (2008). An aliquot of 0.2 ml of the test solution was mixed with 1.8 ml of DPPH solution (0.1 mmol/l) in a spectrophotometer cuvette for 30 min at 25°C in the dark. The absorbance was measured at 0 and 30 min under wavelength 517 nm, respectively. Then, the difference of the absorbance was calculated and then converted to  $\mu\text{mol}$  of Trolox equivalent/litre based on the standard curve of Trolox. Total dietary fibre content was determined using the AOAC method (2005).

The fruit yield of each individual tree was recorded at harvest and average is expressed as fruit yield per tree.

### Statistical analysis

The experiment was conducted for three consecutive years (2017–2019) in randomized block design with three treatments (viz., control, open centre and palmette) and seven replications. Two plants were maintained in each replication. Observations were recorded from the same plant for three consecutive years. Data presented are mean of three consecutive year data. Data for each parameter were evaluated for statistical significance using two-way analysis of variance (ANOVA) to compare means, considering training system and parameter as independent variables. The individual parameter among three crown architectures (viz., control, palmette and open centre) has been assessed by computation of least significant difference taking 't' values for error D.F. at the 5% level of significance. Letters given as superscripts in the data indicate statistically significant parameter at 5% ( $P \leq 0.05$ ) level of significance. WASP (Web Agri Stat Package) of ICAR – Central Coastal Agricultural Research Institute, India (<https://ccari.res.in/waspnew.html>) was used for statistical analyses. Group comparison of diffuse light, direct light, leaf area index (LAI), photosynthesis rate (A), stomatal conductance (gs), vapour pressure deficit (VPD), transpiration rate (E) as seven groups, tested for significance over months (April to August) and clustering of these groups, was done. Membership probability of these groups for significance was tested and confirmed by canonical discriminate function.

### Results

Significant variation in diffuse light availability beneath the trees of different crown architecture was found. In control trees, diffuse light availability beneath the tree varied from  $69.7 \pm 2.22\%$  in April to  $65.6 \pm 2.07\%$  in August. In the palmette canopy, diffuse light availability ranged from  $59.3 \pm 1.90\%$  in April to  $50.4 \pm$

$1.59\%$  in August; whereas in trees with open centre canopy architecture, it varied from  $55.0 \pm 1.76\%$  in April to  $46.0 \pm 1.45\%$  in August (Table 1). Quite the reverse, direct light availability in control trees was found to vary from  $30.4 \pm 0.97\%$  in April to  $34.4 \pm 1.13\%$  in August. For the palmette trees, it ranged from  $40.7 \pm 1.30\%$  in April to  $49.6 \pm 1.64\%$  in August, and in open centre trees, it ranged from  $45.0 \pm 1.44\%$  in April to  $54.1 \pm 1.78\%$  in August (Table 2). A non-significant difference was found in the intensity of direct light measured 1.5 m above ground level beside treated and control trees. Intensity of direct light ranged from  $139.1 \pm 6.05$  to  $139.7 \pm 6.07$  watt/m<sup>2</sup>/s in April to  $147.9 \pm 6.43$  to  $148.3 \pm 6.45$  watt/m<sup>2</sup>/s in August (Table 3). Miniscule difference in light intensity between treatments might be due to slight temporal variation, wind velocity as well as other environmental factors. In contrast, significant variation in the intensity of diffuse light beneath the trees was observed. During the month of April, it ranged from  $2.9 \pm 0.12$  watt/m<sup>2</sup>/s in control to  $3.5 \pm 0.15$  watt/m<sup>2</sup>/s in open centre. Similarly, in August diffuse light ranged from  $5.4 \pm 0.24$  watt/m<sup>2</sup>/s in control to  $5.9 \pm 0.26$  watt/m<sup>2</sup>/s in open centre (Table 4). Moreover, radical difference in LAI was also found in control, palmette and open centre trees. In control trees, LAI ranged from  $4.1 \pm 0.13$  in June to  $4.6 \pm 0.16$  in April, in palmette canopy architecture LAI ranged from  $2.5 \pm 0.08$  in June to  $3.6 \pm 0.12$  in April, whereas in open centre trees it was  $2.02 \pm 0.06$  in June and  $2.9 \pm 0.10$  in April (Table 5).

Ample variation in photosynthesis rate of leaves in control, palmette and open centre trees was found. Instantaneous photosynthesis rate in control trees varied from  $9.5 \pm 0.35$   $\mu\text{mol CO}_2/\text{m}^2/\text{s}$  in August to  $12.5 \pm 0.43$   $\mu\text{mol CO}_2/\text{m}^2/\text{s}$  in June. In palmette canopy architecture, it was  $10.4 \pm 0.39$   $\mu\text{mol CO}_2/\text{m}^2/\text{s}$  in August and  $13.5 \pm 0.46$   $\mu\text{mol CO}_2/\text{m}^2/\text{s}$  in June. Similarly, in open centre trees, it ranged from  $11.68 \pm 0.43$   $\mu\text{mol CO}_2/\text{m}^2/\text{s}$  in August to  $15.7 \pm 0.54$   $\mu\text{mol CO}_2/\text{m}^2/\text{s}$  in June (Table 6). Correspondingly, maximum stomatal conductance was found in open centre trees. Stomatal conductance increased from April to June and then a gradual decline was recorded during the month of July and August. In control trees stomatal conductance ranged from  $35.8 \pm 1.30$  mmol H<sub>2</sub>O/m<sup>2</sup>/s in August to  $53.7 \pm 1.90$  mmol H<sub>2</sub>O/m<sup>2</sup>/s in June. Similarly, in palmette canopy architecture, it ranged from  $47.0 \pm 1.69$  mmol H<sub>2</sub>O/m<sup>2</sup>/s in August to  $62.7 \pm 2.21$  mmol H<sub>2</sub>O/m<sup>2</sup>/s in June, and in open centre trees, it varied from  $61.3 \pm 2.21$  mmol H<sub>2</sub>O/m<sup>2</sup>/s in August to  $88.90 \pm 3.14$  mmol H<sub>2</sub>O/m<sup>2</sup>/s in June (Table 7). Similarly, leaf-to-air vapour pressure deficit (VPD) was maximum in June and minimum in August. In control trees, VPD ranged from  $1.1 \pm 0.04$  kPa in August to  $2.3 \pm 0.08$  kPa in June. In palmette trees, VPD was found to vary from  $1.3 \pm 0.04$  kPa in August to  $2.4 \pm 0.08$  kPa in June. Maximum VPD was found in open centre trees which varied from  $1.9 \pm 0.06$  kPa in August to  $3.9 \pm 0.13$  kPa in June (Table 8). In accordance with

**Table 1.** Diffuse light availability at ground level beneath the tree in jamun (*Syzygium cumini* [L.] Skeels)

Crown architecture	Availability of diffuse light (%) beneath the tree				
	April	May	June	July	August
Control	$69.7 \pm 2.22^a$	$69.4 \pm 2.35a$	$68.3 \pm 2.24^a$	$68.1 \pm 2.08^a$	$65.6 \pm 2.07^a$
Palmette	$59.3 \pm 1.90^{ab}$	$57.0 \pm 1.93^{ab}$	$55.5 \pm 1.81^{ab}$	$52.7 \pm 1.62^{ab}$	$50.4 \pm 1.6^{ab}$
Open centre	$55.0 \pm 1.76^b$	$52.9 \pm 1.79^b$	$49.1 \pm 1.60^b$	$47.2 \pm 1.45^b$	$45.9 \pm 1.5^b$

Values represent means  $\pm$  s.e.

Different superscripts denote statistically significant difference ( $P < 0.05$ ).

Data presented are mean of three consecutive year data (2017–2019).

**Table 2.** Direct light availability at ground level beneath the tree in jamun (*Syzygium cumini* [L.] Skeels)

Crown architecture	Availability of direct light (%) beneath the tree				
	April	May	June	July	August
Control	30.5 ± 0.97 <sup>b</sup>	30.7 ± 0.94 <sup>b</sup>	31.7 ± 1.08 <sup>b</sup>	31.9 ± 1.07 <sup>b</sup>	34.4 ± 1.13 <sup>b</sup>
Palmette	40.7 ± 1.30 <sup>ab</sup>	43.0 ± 1.32 <sup>ab</sup>	44.5 ± 1.51 <sup>ab</sup>	47.3 ± 1.58 <sup>ab</sup>	49.6 ± 1.64 <sup>ab</sup>
Open centre	45.0 ± 1.44 <sup>a</sup>	47.1 ± 1.45 <sup>a</sup>	51.0 ± 1.73 <sup>a</sup>	52.8 ± 1.77 <sup>a</sup>	54.1 ± 1.78 <sup>a</sup>

Values represent means ± s.e.

Different superscripts denote statistically significant difference ( $P < 0.05$ ).

Data presented are mean of three consecutive year data (2017–2019).

**Table 3.** Intensity of direct light in jamun (*Syzygium cumini* [L.] Skeels) orchard

Crown architecture	Intensity of direct light (watt/m <sup>2</sup> /s)				
	April	May	June	July	August
Control	139.1 ± 6.05	156.2 ± 6.79	161.0 ± 7.00	153.4 ± 6.67	147.9 ± 6.43
Palmette	139.6 ± 6.07	156.6 ± 6.81	161.3 ± 7.01	153.4 ± 6.67	148.2 ± 6.44
Open centre	139.7 ± 6.07	156.6 ± 6.82	161.5 ± 7.02	153.6 ± 6.68	148.3 ± 6.45

Values represent means ± s.e.

Data presented are mean of three consecutive year data (2017–2019).

**Table 4.** Intensity of diffuse light in jamun (*Syzygium cumini* [L.] Skeels) orchard

Crown architecture	Intensity of diffuse light (watt/m <sup>2</sup> /s)				
	April	May	June	July	August
Control	2.9 ± 0.12 <sup>a</sup>	4.0 ± 0.18 <sup>a</sup>	6.1 ± 0.27 <sup>a</sup>	5.6 ± 0.25 <sup>a</sup>	5.4 ± 0.24 <sup>a</sup>
Palmette	3.0 ± 0.13 <sup>ab</sup>	4.2 ± 0.18 <sup>ab</sup>	6.7 ± 0.29 <sup>ab</sup>	6.2 ± 0.27 <sup>ab</sup>	5.6 ± 0.24 <sup>ab</sup>
Open centre	3.5 ± 0.15 <sup>b</sup>	4.6 ± 0.20 <sup>b</sup>	7.4 ± 0.32 <sup>b</sup>	6.8 ± 0.29 <sup>b</sup>	5.9 ± 0.26 <sup>b</sup>

Values represent means ± s.e.

Different superscripts denote statistically significant difference ( $P < 0.05$ ).

Data presented are mean of three consecutive year data (2017–2019).

this, minimum transpiration rate was found in August and maximum in June. In control trees, transpiration rate varied from 1.2 ± 0.05 mmol H<sub>2</sub>O/m<sup>2</sup>/s in August to 1.9 ± 0.06 mmol H<sub>2</sub>O/m<sup>2</sup>/s in June. In palmette trees, it was found to vary from 1.6 ± 0.069 mmol H<sub>2</sub>O/m<sup>2</sup>/s in August to 2.1 ± 0.07 mmol H<sub>2</sub>O/m<sup>2</sup>/s in June and in open centre trees 1.8 ± 0.07 mmol H<sub>2</sub>O/m<sup>2</sup>/s in August to 2.6 ± 0.08 mmol H<sub>2</sub>O/m<sup>2</sup>/s in June (Table 9).

Modulation of direct–diffuse light ratio was found to induce early flowering. In open centre trees, flowering was recorded at day 92 ± 1.33 from 1 January, whereas in palmette at day 95 ± 1.33 and in control at day 98 ± 1.67. Similarly, early completion of flowering was also found in open centre trees, i.e. at day 99 ± 1.67 as compared with palmette (at day 103 ± 1.67) and control trees (at day 107 ± 1.67). In open centre trees, fruit set was observed at day 132 ± 1.33, whereas in palmette and control trees at day 135 ± 1.33 and at day 139 ± 1.33, respectively. Days to fruit maturity is important for harvesting and post-harvest management of fruits. In open centre canopy architecture, early fruit maturity was found, i.e. fruits matured at day 176 ± 1.67; whereas in palmette and control trees, fruits matured at day 180 ± 1.67 and 186 ± 1.67, respectively (Table 10).

In open centre, palmette and control trees, average fruit length was 3.4 ± 0.08, 2.9 ± 0.07 and 2.3 ± 0.07 cm, respectively. Similarly, average fruit width in open centre, palmette and control trees was 2.6 ± 0.07, 2.4 ± 0.07, 2.2 ± 0.07 cm, respectively. Significant variation in fruit weight was also found among all the three canopy architectures. Average fruit weight was 8.3 ± 0.15, 9.6 ± 0.15 and 10.4 ± 0.15 g in control, palmette and open centre trees, respectively. Furthermore, fruit yield per tree was found to vary from 62.3 ± 1.53 kg in control, 73.5 ± 1.65 kg in palmette to 86.7 ± 1.33 kg in open centre trees. Increase in fruit length, width, weight and total yield of open centre canopy architecture might be due to congenial microclimate leading to proper hormonal regulation. In the fruits of different crown architectures, substantial variation in fruit quality traits was also found. Noteworthy, variation was recorded in total soluble solid content of fruit of control, palmette and open centre trees which was 9.0 ± 0.15, 9.9 ± 0.14 and 12.2 ± 0.14°Brix, respectively (Table 11).

Anthocyanin and TPC in control, palmette and open centre crown architecture varied from 207.7 ± 6.63, 208.3 ± 6.65, 212.5 ± 6.8 mg/100 g and 209.7 ± 7.20, 212.3 ± 7.29, 219.6 ± 7.54 mg gallic acid equivalent (GAE)/g, respectively. In addition, ascorbic acid content was found to vary from 51.3 ± 1.65 mg/100 g in

**Table 5.** Leaf area index (LAI) of jamun (*Syzygium cumini* [L.] Skeels) tree canopy

Crown architecture	Leaf area index (LAI)				
	April	May	June	July	August
Control	4.6 ± 0.16 <sup>a</sup>	4.6 ± 0.15 <sup>a</sup>	4.1 ± 0.13 <sup>a</sup>	4.3 ± 0.134 <sup>a</sup>	4.5 ± 0.15 <sup>a</sup>
Palmette	3.6 ± 0.12 <sup>ab</sup>	3.2 ± 0.11 <sup>ab</sup>	2.5 ± 0.08 <sup>ab</sup>	2.6 ± 0.08 <sup>ab</sup>	2.9 ± 0.10 <sup>ab</sup>
Open centre	2.9 ± 0.10 <sup>b</sup>	2.6 ± 0.09 <sup>b</sup>	2.0 ± 0.06 <sup>b</sup>	2.2 ± 0.07 <sup>b</sup>	2.3 ± 0.08 <sup>b</sup>

Values represent means ± s.e.  
Different superscripts denote statistically significant difference ( $P < 0.05$ ).  
Data presented are mean of three consecutive year data (2017–2019).

**Table 6.** Net photosynthesis rate ( $\mu\text{mol CO}_2/\text{m}^2/\text{s}$ ) of jamun (*Syzygium cumini* [L.] Skeels) tree leaves

Crown architecture	Net photosynthesis rate ( $\mu\text{mol CO}_2/\text{m}^2/\text{s}$ )				
	April	May	June	July	August
Control	9.6 ± 0.32 <sup>b</sup>	11.8 ± 0.42 <sup>b</sup>	12.5 ± 0.43 <sup>b</sup>	12.3 ± 0.40 <sup>b</sup>	9.5 ± 0.35 <sup>b</sup>
Palmette	10.5 ± 0.35 <sup>ab</sup>	12.6 ± 0.45 <sup>ab</sup>	13.5 ± 0.46 <sup>ab</sup>	13.4 ± 0.43 <sup>ab</sup>	10.4 ± 0.39 <sup>ab</sup>
Open centre	12.1 ± 0.41 <sup>a</sup>	14.9 ± 0.54 <sup>a</sup>	15.7 ± 0.54 <sup>a</sup>	15.2 ± 0.49 <sup>a</sup>	11.7 ± 0.43 <sup>a</sup>

Values represent means ± s.e.  
Different superscripts denote statistically significant difference ( $P < 0.05$ ).  
Data presented are mean of three consecutive year data (2017–2019).

**Table 7.** Stomatal conductance ( $\text{mmol H}_2\text{O}/\text{m}^2/\text{s}$ ) of jamun (*Syzygium cumini* [L.] Skeels) tree leaves

Crown architecture	Stomatal conductance ( $\text{mmol H}_2\text{O}/\text{m}^2/\text{s}$ )				
	April	May	June	July	August
Control	45.3 ± 1.45 <sup>b</sup>	49.2 ± 1.53 <sup>b</sup>	53.7 ± 1.90 <sup>b</sup>	43.0 ± 1.47 <sup>b</sup>	35.8 ± 1.29 <sup>b</sup>
Palmette	55.9 ± 1.79 <sup>ab</sup>	61.3 ± 1.90 <sup>ab</sup>	62.7 ± 2.21 <sup>ab</sup>	54.0 ± 1.859 <sup>ab</sup>	47.0 ± 1.69 <sup>ab</sup>
Open centre	64.4 ± 2.06 <sup>a</sup>	78.8 ± 2.45 <sup>a</sup>	88.9 ± 3.14 <sup>a</sup>	76.0 ± 2.60 <sup>a</sup>	61.3 ± 2.21 <sup>a</sup>

Values represent means ± s.e.  
Different superscripts denote statistically significant difference ( $P < 0.05$ ).  
Data presented are mean of three consecutive year data (2017–2019).

control,  $52.9 \pm 1.70$  mg/100 g in palmette to  $54.2 \pm 1.74$  mg/100 g in open centre. Total antioxidant capacity (DPPH value) ranged from  $33.5 \pm 1.169$  in control,  $35.7 \pm 1.24$  in palmette to  $41.3 \pm 1.43$  in open centre crown architecture. Considerable variation in the fibre content of fruit pulp was also found which ranged from  $0.51 \pm 0.02$ ,  $0.53 \pm 0.02$  and  $0.59 \pm 0.02\%$  in control, palmette and open centre crown architecture, respectively (Table 12).

Pictorial illustration of different training systems depicts that open centre canopy architecture harvests maximum available solar radiation (Fig. 1). Group comparison of diffuse light, direct light, leaf area index (LAI), net assimilation rate (A), stomatal conductance (gs), vapour pressure deficit (VPD) and transpiration rate (E) as seven groups tested for significance over months, i.e. April to August, inferred that diffuse light, direct light, LAI, A, gs, VPD and E have significant variations over months (Table 13). ANOVA-based clustering showed that crown architecture had prominent impact on groups 3 and 4 (leaf area index and net assimilation rate) and groups 6 and 7 (vapour pressure deficit and transpiration rate), which were found to aggregate together (Fig. 2). Membership probability again shows high probability of groups 3 and 4 (leaf area index and net assimilation rate) as

well as groups 6 and 7 (vapour pressure deficit and transpiration rate) (Fig. 3). In canonical discriminant function, groups 3 and 4 (leaf area index and net assimilation rate) and groups 6 and 7 (vapour pressure deficit and transpiration rate) group together separately which proves the impact of crown architecture on these parameters (Fig. 4).

## Discussion

Solar radiation reaching tree top is composed of light (direct and diffuse light) scattered by the atmosphere (Bird and Riordan, 1986). Intensity of light coming to tree top remains same in all the trees. Leaves in different layers of canopy enhance diffuse light component. Diffuse light increases photosynthesis rate at community level, because of more even distribution of light (Geider *et al.*, 2001; Urban *et al.*, 2007). However, positive effect of diffuse light may not be able to compensate for the reduction in the light transmission, as found at leaf level in the present study. Effects of direct and diffuse light on photosynthetic process are different at the leaf and canopy level (Brodersen and Vogelmann, 2007). Due to reduction in light level, plant growth

**Table 8.** Vapour pressure deficit (kPa) of jamun (*Syzygium cumini* [L.] Skeels) tree leaves

Crown architecture	Vapour pressure deficit (kPa)				
	April	May	June	July	August
Control	1.2 ± 0.04 <sup>b</sup>	1.7 ± 0.05 <sup>b</sup>	2.3 ± 0.08 <sup>b</sup>	1.4 ± 0.053 <sup>b</sup>	1.1 ± 0.04 <sup>b</sup>
Palmette	1.8 ± 0.07 <sup>ab</sup>	2.0 ± 0.07 <sup>ab</sup>	2.4 ± 0.08 <sup>ab</sup>	2.0 ± 0.07 <sup>ab</sup>	1.3 ± 0.04 <sup>ab</sup>
Open centre	3.2 ± 0.12 <sup>a</sup>	3.8 ± 0.12 <sup>a</sup>	3.9 ± 0.13 <sup>a</sup>	2.3 ± 0.09 <sup>a</sup>	1.9 ± 0.06 <sup>a</sup>

Values represent means ± s.e.

Different superscripts denote statistically significant difference ( $P < 0.05$ ).

Data presented are mean of three consecutive year data (2017–2019).

**Table 9.** Transpiration rate (mmol H<sub>2</sub>O/m<sup>2</sup>/s) of jamun (*Syzygium cumini* [L.] Skeels) tree leaves

Crown architecture	Transpiration rate (mmol H <sub>2</sub> O/m <sup>2</sup> /s)				
	April	May	June	July	August
Control	1.3 ± 0.04 <sup>b</sup>	1.8 ± 0.06 <sup>a</sup>	1.9 ± 0.06 <sup>a</sup>	1.5 ± 0.06 <sup>a</sup>	1.2 ± 0.05 <sup>a</sup>
Palmette	1.8 ± 0.06 <sup>ab</sup>	1.9 ± 0.07 <sup>ab</sup>	2.1 ± 0.07 <sup>ab</sup>	1.9 ± 0.07 <sup>ab</sup>	1.6 ± 0.06 <sup>ab</sup>
Open centre	2.3 ± 0.07 <sup>a</sup>	2.4 ± 0.09 <sup>a</sup>	2.6 ± 0.08 <sup>a</sup>	2.2 ± 0.08 <sup>a</sup>	1.8 ± 0.07 <sup>a</sup>

Values represent means ± s.e.

Different superscripts denote statistically significant difference ( $P < 0.05$ ).

Data presented are mean of three consecutive year data (2017–2019).

**Table 10.** Impact of crown architecture on days to flowering, fruit set and maturity of jamun (*Syzygium cumini* [L.] Skeels) pulp

Crown architecture	Days to start flowering	Days to end flowering	Days to fruit set	Days to fruit maturity
Control	98 ± 1.19	107 ± 2.38	139 ± 2.14	184 ± 1.63
Palmette	95 ± 1.26	103 ± 2.34	135 ± 2.26	180 ± 1.48
Open centre	92 ± 1.22	99 ± 2.41	132 ± 2.32	176 ± 1.36

Values represent means ± s.e.

Different superscripts denote statistically significant difference ( $P < 0.05$ ).

Data presented are mean of three consecutive year data (2017–2019).

and fruit development do not accelerate. Light scattered by canopy has different spectral composition as compared to that diffused by environmental factors (clouds, aerosols and air pollutants, etc.) (Brodersen *et al.*, 2008), hence this differently affects fruit growth and yield. Under intense direct light, chloroplasts move to periclinal walls, shading other chloroplasts and thus reduce photoinhibition (Gorton *et al.*, 1999); this helps to maintain photosynthesis as found in open centre crown architecture. Under diffuse light, chloroplast movement to the periclinal walls is not complete (Williams *et al.*, 2003); this causes lower absorptance (Brodersen and Vogelmann, 2007) as evident in control trees. Variation in the availability of direct and diffuse light as well as light intensity in control, palmette and open centre trees affects leaf pigments, metabolites, primary productivity and fruit quality. More leaf area index (LAI) in control trees affects light availability and tree growth in the inner canopy layers (Calvo-Rodriguez and Sanchez-Azofeifa, 2016).

Although diffused light increases productivity at the community level, leaf-level photosynthesis rates may be lower as found in the present study (Brodersen *et al.*, 2008). Studies of canopy light environments (light intensity and diffuseness) and photosynthesis at leaf and canopy scales are challenging (Williams *et al.*, 2014). In fact, no studies have been performed in crops

grown at different levels of diffuseness with similar incident light intensity on the top of the crop (Li *et al.*, 2014). Different processes are functioning at different levels within the plant community (Brodersen *et al.*, 2008). At leaf level, diffuse light leads shallow penetration than direct light in sun-grown leaves like jamun. A greater proportion of diffuse light reflects from the surface of leaves, and less light enters the leaf for photosynthesis. The unequal absorptance of direct and diffuse light and the extent to which a change in the directional quality and intensity of light affects photosynthesis at the leaf level is yet not known clearly (Brodersen and Vogelmann, 2007). Hence, such studies in fruit tree orchards will help to manage tree crown architecture for enhancing production. Weaker penetration of diffuse light into the mesophyll of sun-grown leaves leads to a more heterogeneous saturation of electron transport capacity and lowers its CO<sub>2</sub> concentration drawdown capacity in the intercellular airspace and chloroplast stroma. This decoupling of light availability from photosynthetic capacity under diffuse light generates a decline in photosynthesis (Earles *et al.*, 2017) as found in the present study.

Morpho-physiological characteristics of plant organs such as stomata are affected by their prevailing microclimate (Sultan, 2000; Niinemets, 2007). Sensitivity of stomata to different light

**Table 11.** Impact of crown architecture on fruit characters of jamun (*Syzygium cumini* [L.] Skeels) pulp

Crown architecture	Fruit length (cm)	Fruit width (cm)	Fruit weight (g)	Fruit yield (kg/tree)	Total soluble solids ( <sup>o</sup> Brix)
Control	2.3 ± 0.07	2.2 ± 0.07	8.3 ± 0.25	62.3 ± 1.50	9.0 ± 0.15
Palmette	2.9 ± 0.07	2.4 ± 0.07	9.6 ± 0.25	73.5 ± 1.65	9.9 ± 0.15
Open centre	2.4 ± 0.08	2.6 ± 0.07	10.4 ± 0.34	86.7 ± 2.33	12.2 ± 0.14

Values represent means ± s.e.

Different superscripts denote statistically significant difference ( $P < 0.05$ ).

Data presented are mean of three consecutive year data (2017–2019).


**Table 12.** Variation in chemical characteristics of jamun (*Syzygium cumini* [L.] Skeels) pulp

Crown architecture	Anthocyanin (mg/100 g)	Total phenol content (mg GAE/g)	Ascorbic acid (mg/100 g)	Antioxidant capacity (DPPH value)	Fibre (%)
Control	207.7 ± 6.64 <sup>b</sup>	209.7 ± 7.20 <sup>b</sup>	51.3 ± 1.65 <sup>b</sup>	33.5 ± 1.16 <sup>b</sup>	0.51 ± 0.02 <sup>b</sup>
Palmette	208.3 ± 6.65 <sup>ab</sup>	212.3 ± 7.29 <sup>ab</sup>	52.9 ± 1.70 <sup>ab</sup>	35.7 ± 1.24 <sup>ab</sup>	0.53 ± 0.02 <sup>ab</sup>
Open centre	212.5 ± 6.78 <sup>a</sup>	219.6 ± 7.54 <sup>a</sup>	54.2 ± 1.74 <sup>a</sup>	41.3 ± 1.43 <sup>a</sup>	0.59 ± 0.02 <sup>a</sup>

Values represent means ± s.e.

Different superscripts denote statistically significant difference ( $P < 0.05$ ).

Data presented are mean of three consecutive year data (2017–2019).

**Figure 1.** Pictorial presentation of training systems.

levels varies in different species (Li *et al.*, 2016); however, within-species variation in stomatal response to direct and diffused light might be due to variation in microclimate, as found in the present study.

Diffuse and direct light condition differently affects growth and development of reproductive organs. Development of flowers and fruits is adversely affected under shaded condition in inner canopy layers. Delayed flowering and fruiting and poor fruit quality traits in the diffuse light condition (control) found in the present study are in conformity with earlier findings viz., reduced flowering (Jackson, 1980), increased fruit abscission (Byers *et al.*, 1991), reduction in fruit size (Warrington *et al.*, 1996) and reduction in internal fruit quality (Warrington *et al.*, 1996) under shaded/low light condition.

Increase in anthocyanin content of fruits due to crown architecture management might be vital to improve fruit quality. Anthocyanins provide pigmentation to fruits and serve as natural antioxidants (Bagchi *et al.*, 2004). Edible anthocyanins possess a broad spectrum of therapeutic and anti-carcinogenic properties.

In addition, an increase in phenolic compounds was also found which have beneficial effects on health including anti-inflammatory, antiviral, antimicrobial and antioxidant activity. Jamun fruits are good source of ascorbic acid and other natural antioxidants (Benvenuti *et al.*, 2006); an increase in these compounds indicates production of better quality fruits. Slight increase in fibre content in fruits was found, such fruits may help to lower blood cholesterol and reduce the risk of heart disease (Jones *et al.*, 2006) and diabetes (Abdul-Hamid and Luan, 2000).


The present study revealed that there is considerable variation in light distribution within the canopy due to crown architecture management which might be contributing to fruit size, yield and quality. Understanding about how diffuse vs. direct light affects photosynthesis and other plant processes in a particular tree crop ultimately helps to determine the proportion of diffuse or direct light a plant receives and needed for enhancing production (Brodersen and Vogelmann, 2007). Therefore, it becomes increasingly important to understand how much direct and diffuse light

**Table 13.** ANOVA for group (diffuse light, direct light, LAI, A, gs, VPD, E) over 5 months

Month	Group	Sum of squares	D.F.	Mean square	F	Sig.
April	Between groups	12 410	6	2068.4	69.24	0.00
	Within groups	418	14	29.9		
	Total	12 828	20			
May	Between groups	13 562	6	2260.3	42.36	0.00
	Within groups	747	14	53.4		
	Total	14 309	20			
June	Between groups	14 528	6	2421.4	31.97	0.00
	Within groups	1060	14	75.7		
	Total	15 588	20			
July	Between groups	12 206	6	2034.4	27.43	0.00
	Within groups	1038	14	74.2		
	Total	13 245	20			
August	Between groups	10 760	6	1793.3	33.17	0.00
	Within groups	757	14	54.1		
	Total	11 517	20			

Significance (value < 0.05).


Data presented for each month are the mean of data for that month from three consecutive years (2017–2019).



**Figure 2.** ANOVA-based clustering of seven groups (diffuse light, direct light, LAI, A, gs, VPD, E). \*Groups on the 'X' axis: (1) diffuse light, (2) direct light, (3) leaf area index (LAI), (4) photosynthesis rate (A), (5) stomatal conductance (gs), (6) vapour pressure deficit (VPD), (7) transpiration (E).

penetrates leaves and how the directional quality of light affects photosynthesis at the leaf level as well as during flowering, fruiting and fruit quality traits.


ANOVA-based clustering approach for similarity aggregation was applied and similar traits were found to cluster together. Moreover, membership probability analysis showed that there was a high probability of the impact of crown architecture on these groups (groups 3 and 4 [leaf area index and net assimilation



**Figure 3.** Membership probability of seven groups (diffuse light, direct light, LAI, A, gs, VPD, E). \*Vertical columns show probability of seven member components (groups). Each group is represented by three vertical columns. Groups 1, 2 and 5 have 100% probability, whereas groups 3, 4, 6 and 7 have partial probability. Group 1: diffuse light, group 2: direct light, group 3: leaf area index (LAI), group 4: photosynthesis rate (A), group 5: stomatal conductance (gs), group 6: vapour pressure deficit (VPD), group 7: transpiration (E).

rate] and groups 6 and 7 [vapour pressure deficit and transpiration rate]). Canonical discriminant analysis, a multivariate technique was used to determine the correlation between these variables and showed that crown architecture significantly affects leaf area index, net assimilation rate, vapour pressure deficit and transpiration rate.





**Figure 4.** Canonical discriminate function of seven groups. Group 1: diffuse light, group 2: direct light, group 3: leaf area index (LAI), group 4: photosynthesis rate (A), group 5: stomatal conductance (gs), group 6: vapour pressure deficit (VPD), group 7: transpiration (E). Groups 1, 2 and 5 are depicted separately, these groups and their centroid are scattered. Groups 3, 4, 6 and 7 have highest possible multiple correlation and have overlapping centroid.

## Conclusion

Crown architecture of the tree affects direct vs. diffuse light available to leaves in the inner canopy layers. The effects of direct and diffuse light on plant processes are different at leaf and community level. Understanding the difference in plant processes under direct vs. diffuse light condition may help to predict the response of fruit crops to changing light/environmental conditions and standardize canopy architecture accordingly. Crown architecture management may be a useful strategy to enhance fruit quality and production of underutilized fruit crops like jamun.

**Data availability statement.** This manuscript is based on basic study; no specific software/data have been used. Data will be made available on reasonable request.

**Acknowledgements.** The authors are thankful to the Director, ICAR – Central Institute for Subtropical Horticulture, Lucknow (India) for providing necessary facility and keen interest in the study.

**Author contributions.** A. K. Trivedi: conceptualization, investigation, collection and processing of data, writing original draft. A. K. Singh: curation, writing, review and editing. K. K. Mishra: data collection, methodology, analysis. G. S. Vishen: data collection, methodology, analysis.

**Funding statement.** The present study was conducted under institute project ‘Conservation and utilization of genetic resources for improvement of Jamun fruits for higher productivity and quality’. Hence, no external financial support was involved.

**Competing interests.** On behalf of all authors, the corresponding author states that there is no conflict of interest.

**Ethical standards.** Not applicable.

## References

Abdul-Hamid A and Luan YS (2000) Functional properties of dietary fibre prepared from defatted rice bran. *Food Chemistry* **68**, 15–19.

Acreche MM, Briceno-Felix G, Sanchez JAM and Slafer GA (2009) Radiation interception and use efficiency as affected by breeding in Mediterranean wheat. *Field Crops Research* **110**, 91–97.

Association of Official Analytical Chemists (2005) Methods 2001.03 Dietary fiber in foods containing resistant maltodextrin. In Horwithz W (ed.), *Official Methods of Analysis of AOAC International*, 18th Edn. Gaithersburg, MD: AOAC International, p 44–54.

Bagchi DC, Sen K, Bagchi M and Atalay M (2004) Anti-angiogenic, antioxidant and anti-carcinogenic properties of a novel anthocyanin-rich berry extract formula. *Biochemistry* **69**, 75–80.

Benvenuti S, Pellati F, Melegari M and Bertelli D (2006) Polyphenols, anthocyanins, ascorbic acid, and radical scavenging activity of Rubus, Ribes, and Aronia. *Journal of Food Science* **69**, 164–169.

Bird R and Riordan C (1986) Simple solar spectral model for direct and diffuse irradiance on horizontal and tilted planes at the earth’s surface for cloudless atmospheres. *Journal of Climate and Applied Meteorology* **25**, 87–97.

Bodin P and Franklin O (2012) Efficient modeling of sun/shade canopy radiation dynamics explicitly accounting for scattering. *Geoscientific Model Development* **5**, 535–541.

Brodersen C and Vogelmann T (2007) Do epidermal lens cells facilitate the absorbance of diffuse light? *American Journal of Botany* **94**, 1061–1066.

Brodersen CR, Voglmann TC, Williams WE and Gorton HL (2008) A new paradigm in leaf-level photosynthesis: direct and diffuse lights are not equal. *Plant, Cell and Environment* **31**, 159–164.

Byers RE, Carbaugh DH, Presley CN and Wolf TK (1991) The influence of low light on apple fruit abscission. *Journal of Horticultural Science* **66**, 7–17.

Calvo-Rodriguez S and Sanchez-Azofeifa GA (2016) Light diffusion in the tropical dry forest of Costa Rica. In *The International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences, Volume XLI-B8, 2016 XXIII ISPRS Congress*, 12–19 July 2016, Prague: Czech Republic.

Cheng GW and Breen PJ (1991) Activity of phenylalanine ammonia-lyase (PAL) and concentrations of anthocyanins and phenolics in developing strawberry fruit. *Journal of the American Society for Horticultural Science* **116**, 865–869.

Davis SHR and Masten SJ (1991) Spectrophotometric method for ascorbic acid using dichlorophenolindophenol: elimination of the interference due to iron. *Analytica Chimica Acta* **248**, 225–221.

Earles JM, Théroux-Rancourt G, Gilbert ME, McElrone AJ and Brodersen CR (2017) Excess diffuse light absorption in upper mesophyll limits CO<sub>2</sub> drawdown and depresses photosynthesis. *Plant Physiology* **174**, 1082–1096.

- Geider RJ, Delucia EH, Falkowski PG, Finzi AC, Grime JP, Grace J, Kana TM, Roche JL, Long SP, Osborne BA, Platt T, Prentice IC, Raven JA, Schlesinger WH, Smetacek V, Stuart V, Sathyendranath S, Thomas RB, Vogelmann TC, Williams P and Woodward FI (2001) Primary productivity of planet earth: biological determinants and physical constraints in terrestrial and aquatic habitats. *Global Change Biology* 7, 849–882.
- Ghosh P, Pradhan RC, Mishra S, Patel AS and Kar A (2017) Physicochemical and nutritional characterization of jamun (*Syzygium cumini*). *Current Research in Nutrition and Food Science* 5, 25–35.
- Gorton H, Williams W and Vogelmann T (1999) Chloroplast movement in *Alocasia macrorrhiza*. *Physiologia Plantarum* 106, 421–428.
- Jackson JE (1980) Light interception and utilization by orchard systems. *Horticultural Reviews* 2, 208–267.
- Jones JR, Lineback DM and Levine MJ (2006) Dietary reference intakes: implications for fibre labelling and consumption: a summary of the international life sciences institute. North America fibre workshop, June 1–2, 2004, Washington, DC. *Nutrition Reviews* 64, 31–38.
- Li T, Heuvelink E, Dueck TA, Janse J, Gort G and Marcelis LFM (2014) Enhancement of crop photosynthesis by diffuse light: quantifying the contributing factors. *Annals of Botany* 114, 145–156.
- Li T, Kromdijk J, Heuvelink E, van Noort FR, Kaiser E and Marcelis LFM (2016) Effects of diffuse light on radiation use efficiency of two anthurium cultivars depend on the response of stomatal conductance to dynamic light intensity. *Frontiers in Plant Science* 7, 56. doi: 10.3389/fpls.2016.00056
- Maddonni GA, Otegui ME and Cirilo AG (2001) Plant population density, row spacing and hybrid effects on maize canopy architecture and light attenuation. *Field Crops Research* 71, 183–193.
- Mishra DS, Singh A, Kumar R, Singh S, Singh AK and Swamy GSK (2014) Jamun. In Ghosh SN (ed.), *Tropical and Sub Tropical Fruit Crops: Crop Improvement and Varietal Wealth Part-2*. New Delhi: Jaya Publishing House, pp. 375–390.
- Niinemets U (2007) Photosynthesis and resource distribution through plant canopies. *Plant Cell and Environment* 30, 1052–1071. doi: 10.1111/j.1365-3040.2007.01683.x.
- Norman J and Arkebauer T (1991) Predicting canopy light use efficiency from leaf characteristics. *Agronomy Monographs* 31, 125–143.
- Norman J and Miller E (1971) Light intensity and sunfleck size distributions in plant canopies. *Agronomy Journal* 63, 743–748.
- Periyathambi R (2007) *Jamun - The Potential Untapped*. In Kerala Calling, February 2007 pp 30–32. Public Relations Department, Govt. of Kerala, Thiruvananthapuram (India).
- Sarlikoti V, de Visser PH, Buck-Sorlin GH and Marcelis LFM (2011) How plant architecture affects light absorption and photosynthesis in tomato: towards an ideotype for plant architecture using a functional-structural plant model. *Annals of Botany* 108, 1065–1073.
- Singh AK, Bajpai A, Singh A and Reddy BMC (2007) Evaluation of variability in jamun using morphological and physico-chemical characterization. *Indian Journal of Agricultural Sciences* 77, 845–848.
- Slinkard K and Singleton VL (1977) Total Phenol Analysis: Automation and Comparison with Manual Methods. *American Journal of Enology and Viticulture* 28, 49–55.
- Sultan SE (2000) Phenotypic plasticity for plant development, function and life history. *Trends in Plant Science* 5, 537–542. doi: 10.1016/S1360-1385(00)01797-0.
- Urban O, Janous D, Acosta M, Czerny R, Markova I, Navrátil M, Pavelka M, Pokorný R, Šprtová M, Zhang R, Špunda V, Grace J and Marek M (2007) Ecophysiological controls over the net ecosystem exchange of mountain spruce stand, comparison of the response in direct vs. diffuse solar radiation. *Global Change Biology* 13, 157–168.
- Warrier PK, Nambiar VPK and Ramankutty C (eds) (1996) *Indian Medicinal Plants: A Compendium of 500 Species*, Vol. 5. Hyderabad: Orient Longman Ltd., pp. 225–228.
- Warrington IJ, Stanley CJ, Tustin DS, Hirst PM and Cashmore WM (1996) Light transmission, yield distribution, and fruit quality in six tree canopy forms of 'Granny Smith' apple. *Journal of Tree Fruit Production* 1, 27–54. doi: 10.1300/J072v01n0103
- Williams W, Gorton H and Witiak S (2003) Chloroplast movements in the field. *Plant, Cell and Environment* 26, 2005–2014.
- Williams M, Rastetter EB, Pol LV and Shaver GR (2014) Arctic canopy photosynthetic efficiency enhanced under diffuse light, linked to a reduction in the fraction of the canopy in deep shade. *New Phytologist* 202, 1266–1276.
- Yue X and Xu Z (2008) Changes of anthocyanins, anthocyanidins, and antioxidant activity in bilberry extract during dry heating. *Journal of Food Science* 73, C494–C499.