

## THE RING OF INVARIANTS OF MATRICES

YASUO TERANISHI

### § 1. Introduction

We denote by  $M(n)$  the space of all  $n \times n$ -matrices with their coefficients in the complex number field  $\mathcal{C}$  and by  $G$  the group of invertible matrices  $GL(n, \mathcal{C})$ . Let  $W = M(n)^l$  be the vector space of  $l$ -tuples of  $n \times n$ -matrices. We denote by  $\rho: G \rightarrow GL(W)$  a rational representation of  $G$  defined as follows:

$$\rho(S)(A(1), A(2), \dots, A(l)) = (SA(1)S^{-1}, SA(2)S^{-1}, \dots, SA(l)S^{-1})$$

if  $S \in G$ ,  $A(i) \in M(n)$  ( $i = 1, 2, \dots, l$ ).

This action of  $G$  defines an action of  $G$  on an algebra  $\mathcal{C}[W] = \mathcal{C}[x_{ij}(1), \dots, x_{ij}(l)]$  of all polynomial functions on  $W$ . We denote by  $\mathcal{C}[W]^G$  the subalgebra of  $G$  invariant polynomials. This is a finitely generated subalgebra of  $\mathcal{C}[W]$ .

If  $l = 1$  it is a classical result that this ring of invariants is a polynomial ring in  $n$  variables. In fact the coefficients of characteristic polynomial of the matrix  $X(1) = (x_{ij}(1))$  are algebraically independent invariants and the ring of invariants is generated by them. By the Newton's formula all coefficients of characteristic polynomial of  $X(1)$  are expressed by  $n$  traces

$$\text{Tr}(X(1)), \text{Tr}(X^2(1)), \dots, \text{Tr}(X(1)^n),$$

and hence  $\mathcal{C}[x_{ij}(1)]^G$  is the polynomial ring generated by these traces.

Procesi [5] has shown the following important

**THEOREM 1.1.** *The ring of invariants  $\mathcal{C}[W]^G$  is generated by all traces  $\text{Tr}(X(i_1) \cdots X(i_j))$  ( $j = 1, 2, \dots$ ), where  $X(i_1) \cdots X(i_j)$  runs all possible non-commutative monomials.*

The object of this paper is to determine the Poincaré series of  $\mathcal{C}[W]^G$  and to determine generators of  $\mathcal{C}[W]^G$  for some cases.

---

Received June 13, 1985.

The following notations are fixed throughout:

- $\mathbf{C}$  the field of complex numbers
- $\mathbf{N}$  additive semigroup of nonnegative integers
- $\mathbf{Q}$  the field of rational numbers

For a complex number  $z$ , we denote by  $\bar{z}$  its complex conjugate and set  $e(z) = \exp 2\pi\sqrt{-1}z$ .

The author expresses his hearty thanks to Dr. Daniele Montanari who pointed out a mistake in the earlier version of this paper.

### §2. Poincaré series

We give  $\mathbf{C}[W]$  the structure of  $\mathbf{N}^l$ -graded algebra by defining  $\deg x_{ij}(k)$  to be the  $k$ -th unit coordinate vector  $\varepsilon_k$  in  $\mathbf{N}^l$ . Let

$$\mathbf{C}[W] = \bigoplus_{d \in \mathbf{N}^l} \mathbf{C}[W]_d,$$

where  $\mathbf{C}[W]_d$  is a vector space spanned over  $\mathbf{C}$  by the monomials in  $\mathbf{C}[W]$  of degree  $d \in \mathbf{N}^l$ . Then  $\mathbf{C}[W]^G$  has the structure

$$\mathbf{C}[W]^G = \bigoplus_{d \in \mathbf{N}^l} \mathbf{C}[W]_d^G,$$

of an  $\mathbf{N}^l$ -graded algebra given by

$$\mathbf{C}[W]_d^G = \mathbf{C}[W]^G \cap \mathbf{C}[W]_d.$$

The Poincaré series of  $\mathbf{C}[W]^G$  is the formal power series  $P(z_1, \dots, z_l)$  in  $l$ -variables  $z_1, \dots, z_l$  defined by

$$P(z_1, \dots, z_l) = \sum_{d \in \mathbf{N}^l} \dim_{\mathbf{C}} \mathbf{C}[W]_d^G z^d$$

where  $z^d = z_1^{d_1} \dots z_l^{d_l}$  with  $d = (d_1, \dots, d_l)$ .

A theorem of Hilbert-Serre implies that  $P(z_1, \dots, z_l)$  is a rational function in  $l$  variables  $z_1, \dots, z_l$ . By using a classical method of Molien-Weyl, we shall calculate this rational function.

For each diagonal unitary matrix  $\varepsilon$  with diagonal entries

$$\varepsilon_1, \varepsilon_2, \dots, \varepsilon_n,$$

since  $|\varepsilon_i| = 1$  ( $i = 1, 2, \dots, n$ ), we can put  $\varepsilon_i = e(\varphi_i)$  ( $0 \leq \varphi_i \leq 1$ ). We set

$$\Delta = \prod_{i < j} (e(\varphi_i) - e(\varphi_j)).$$

Then the normalized volume element on the group consisting of diagonal unitary matrices is given by

$$\frac{1}{n!} \Delta \bar{\Delta} d\varphi_1 \cdots d\varphi_n, [8].$$

We define polynomials in one variable  $z$  by

$$\Delta(z) = \prod_{i < j} (e(\varphi_i) - ze(\varphi_j))$$

and

$$\bar{\Delta}(z) = \prod_{i < j} (\bar{e}(\bar{\varphi}_i) - z\bar{e}(\bar{\varphi}_j)).$$

**THEOREM 2.1.** *The Poincaré series  $P(z_1, \dots, z_n)$  is*

$$\frac{1}{n! \prod_{i=1}^n (1 - z_i)^n} \int_0^1 \cdots \int_0^1 \frac{\Delta \bar{\Delta}}{\prod_{i=1}^n \Delta(z_i) \bar{\Delta}(z_i)} d\varphi_1 \cdots d\varphi_n, \\ |z_i| < 1, \dots, |z_i| < 1.$$

*Proof.* Let  $f(z)$  be a polynomial in one variable  $z$  defined as

$$f(z) = \det(I_n - \rho(\varepsilon)z), \quad I_n = \text{the } n \times n\text{-identity matrix,} \\ = \prod_{1 \leq i < j \leq n} (1 - z\varepsilon_i\varepsilon_j^{-1}) \\ = (1 - z)^n \Delta(z) \bar{\Delta}(z).$$

Then by the Molien-Weyl formula [8], the Poincaré series  $P(z_1, \dots, z_n)$  equals

$$\frac{1}{n!} \int_0^1 \cdots \int_0^1 \frac{\Delta \bar{\Delta}}{f(z_1) \cdots f(z_n)} d\varphi_1 \cdots d\varphi_n, \quad |z_i| < 1.$$

By changing variables from  $\varphi_1, \dots, \varphi_n$  to  $\varepsilon_1, \dots, \varepsilon_n$ , we have

$$P(z_1, \dots, z_n) \\ = \left( \frac{1}{2\pi\sqrt{-1}} \right)^n \frac{1}{n! \prod_{i=1}^n (1 - z_i)^n} \int_{C_1} \cdots \int_{C_n} \frac{\Delta \bar{\Delta}}{\prod_{i=1}^n \Delta(z_i) \bar{\Delta}(z_i)} d\varepsilon_1 \cdots d\varepsilon_n,$$

where  $C_k$  denotes the unit circle  $|\varepsilon_k| = 1$  in the complex  $\varepsilon_k$ -plane. Thus the Poincaré series  $P(z_1, \dots, z_n)$  can be calculated in principle by means of residues. Since

$$\Delta(z) \bar{\Delta}(z) = (-z)^{(n(n-1))/2} (\varepsilon_1 \cdots \varepsilon_n)^{1-n} \prod_{i < j} (\varepsilon_i - z\varepsilon_j) \left( \varepsilon_i - \frac{1}{z}\varepsilon_j \right),$$

we have

$$\frac{\Delta \bar{\Delta}}{\prod_{i=1}^n \Delta(z_i) \bar{\Delta}(z_i)} = (-1)^{n(n-1)(n-2)/2} (z_1 \cdots z_n)^{(n(n-1))/2} (\varepsilon_1 \cdots \varepsilon_n)^{(n-1)(n-2)} \\ \times \frac{D(\varepsilon_1, \dots, \varepsilon_n)}{\prod_{p=1}^n \prod_{i < j} (\varepsilon_i - z_p\varepsilon_j)(\varepsilon_i - (1/z_p)\varepsilon_j)},$$

where  $D(\varepsilon_1, \dots, \varepsilon_n) = \prod_{i < j} (\varepsilon_i - \varepsilon_j)^2$ . And so we can rewrite Theorem 2.1 as

$$(2.2) \quad P(z_1, \dots, z_l) = (-1)^{n(n-1)(l-1)/2} \frac{1}{n! \prod_{i=1}^l (1-z_i)^n (z_1 \dots z_l)^{n(n-1)/2}} \left( \frac{1}{2\pi\sqrt{-1}} \right)^n \times \int \dots \int \frac{(\varepsilon_1 \dots \varepsilon_n)^{(n-1)(l-1)-1} D(\varepsilon_1, \dots, \varepsilon_n)}{\prod_{p=1}^l \prod_{i < j} (\varepsilon_i - z_p \varepsilon_j)(\varepsilon_i - (1/z_p) \varepsilon_j)} d\varepsilon_1 \dots d\varepsilon_n.$$

**PROPOSITION 2.3.** *The Poincaré series  $P(z_1, \dots, z_l)$  ( $l \geq 2$ ) satisfies the following functional equation*

$$P(z_1^{-1}, \dots, z_l^{-1}) = (-1)^{n(l-1)+1} (z_1, \dots, z_l)^{n^2} P(z_1, \dots, z_l).$$

*Proof.* Consider a rational function  $I(z_1, \dots, z_l)$  defined in  $|z_1| < 1, \dots, |z_l| < 1$  as follows

$$I(z_1, \dots, z_l) = \int_{c_1} \dots \int_{c_n} F_{z_1, \dots, z_l}(\varepsilon_1, \dots, \varepsilon_n) d\varepsilon_1 \dots d\varepsilon_n,$$

where

$$F_{z_1, \dots, z_l}(\varepsilon_1, \dots, \varepsilon_n) = \frac{(\varepsilon_1 \dots \varepsilon_n)^{(n-1)(l-1)-1} D(\varepsilon_1, \dots, \varepsilon_n)}{\prod_{p=1}^l \prod_{i < j} (\varepsilon_i - z_p \varepsilon_j)(\varepsilon_i - (1/z_p) \varepsilon_j)}.$$

Set inductively

$$\begin{aligned} I_1(\varepsilon_1, \dots, \varepsilon_n) &= F_{z_1, \dots, z_l}(\varepsilon_1, \dots, \varepsilon_n), \\ I_{i+1}(\varepsilon_{i+1}, \dots, \varepsilon_n) &= \int_{|\varepsilon_i|=1} I_i(\varepsilon_i, \varepsilon_{i+1}, \dots, \varepsilon_n) d\varepsilon_i, \\ &\quad (i = 1, \dots, n-1). \end{aligned}$$

Then we find that  $I_i(\varepsilon_i, \dots, \varepsilon_n)$  is, as a function of  $\varepsilon_i$ , holomorphic at  $\varepsilon_i = \infty$ . If  $|z_1| > 1, \dots, |z_l| > 1$ , we have

$$\begin{aligned} I(z_1^{-1}, \dots, z_l^{-1}) &= \int_{c_1} \dots \int_{c_n} F_{z_1, \dots, z_l}(\varepsilon_1, \dots, \varepsilon_n) d\varepsilon_1 \dots d\varepsilon_n \\ &= (-1)^{n-1} \int_{c_1}^{-1} \dots \int_{c_{n-1}}^{-1} \int_{c_n} F_{z_1, \dots, z_l}(\varepsilon_1, \dots, \varepsilon_n) d\varepsilon_1 \dots d\varepsilon_n. \end{aligned}$$

By the Cauchy integral formula we have

$$I(z_1^{-1}, \dots, z_l^{-1}) = (-1)^{n-1} I(z_1, \dots, z_l),$$

and hence we obtain the result by 2.2.

We consider  $C[W]$  as a  $N$ -graded algebra

$$C[W] = \bigoplus_{d \in N} C[W]_d$$

by defining  $\deg x_{i,j}(k) = 1$  and define the Poincaré series  $P(z)$  in one variable  $z$  by

$$P(z) = P(z, \dots, z) = \sum_{d \in \mathbb{N}} \dim_{\mathbb{C}} C[W]_d^g z^d.$$

Then it follows from (2.2) that the Poincaré series  $P(z)$  equals

$$(2.4) \quad (-1)^{(n(n-1)(l-1))/2} \frac{1}{n!(1-z)^{nl} z^{(n(n-1)l)/2}} \left( \frac{1}{2\pi\sqrt{-1}} \right)^n \\ \times \int \dots \int \frac{(\varepsilon_1 \dots \varepsilon_n)^{(n-1)(l-1)-1} D(\varepsilon_1, \dots, \varepsilon_n)}{(\prod_{i < j} (\varepsilon_i - z\varepsilon_j)(\varepsilon_i - (1/z)\varepsilon_j))^l} d\varepsilon_1 \dots d\varepsilon_n.$$

Let  $f_1, \dots, f_m$  be a homogeneous system of parameters of the  $N$ -graded algebra  $C[W]^g$ . By a theorem of Hochster and Roberts [4],  $C[W]^g$  is a free module over the polynomial ring  $C[f_1, \dots, f_m]$ . Let  $\varphi_1, \dots, \varphi_r$  be a homogeneous system of generators of this module,

$$C[W]^g = \bigoplus_{i=1}^r \varphi_i C[f_1, \dots, f_m].$$

We claim that  $m = (l - 1)n^2 + 1$ . For  $w \in W$ , we denote by  $G_w$  the isotropy subgroup of  $GL(n, \mathbb{C})$  at  $w$ . If  $l \geq 2$ , there exists a dense open subset  $U$  of  $w$  such that  $G_w = \{e\}$ . Then it follows from a theorem of Rosenlicht [6] that the transcendence degree of  $C[W]^g$  is equals  $\dim W - \dim G + 1$ . This shows that  $m = (l - 1)n^2 + 1$ . Formanek [1] has shown that the field of rational invariants  $C(W)^g$  is unirational of transcendence degree  $(l - 1)n^2 + 1$ .

We set

$$\deg f_i = d_i, \quad d_1 \leq \dots \leq d_m \\ \deg \varphi_j = e_j, \quad 0 = e_1 \leq \dots \leq e_r.$$

By Proposition 2.3,  $P(z)$  satisfies the following functional equation

$$P(z^{-1}) = (-1)^{(l-1)n^2+1} z^{n^2l} P(z).$$

This equation is equivalent to

$$d_1 + \dots + d_m - e_{r-i+1} = n^2l + e_i, \quad i = 1, \dots, r.$$

In particular we have

$$e_i + e_{r-i+1} = e_r, \quad i = 1, \dots, l, \\ e_r = d_1 + \dots + d_m - n^2l$$

and

$$(2.5) \quad n^2 l = \sum_{j=1}^m d_j - \frac{2}{r} \sum_{i=1}^r e_i .$$

Let  $\alpha$  and  $\beta$  be the first and second Laurant coefficients of  $P(z)$  respectively. Then the Laurant expansion of the Poincaré series  $P(z)$  begins with

$$P(z) = \frac{\alpha}{(1-z)^m} + \frac{\beta}{(1-z)^{m-1}} + \dots .$$

By 2.5.9 Lemma (7), it follows that

$$(2.6) \quad \alpha = \frac{r}{d_1 \cdots d_m}$$

and

$$\beta = \frac{r \sum_{j=1}^m (d_j - 1) - 2 \sum_{i=1}^r e_i}{2d_1 \cdots d_m} .$$

Then it follows from (2.5) that

$$(2.7) \quad \frac{\beta}{\alpha} = \frac{n^2 - 1}{2} .$$

We shall need the following important theorem due to Hilbert [3].

**THEOREM 2.8.** *Assume that some invariants  $I_1, \dots, I_\mu$  have a property that their vanishing implies the vanishing of all invariants. Then the ring of invariants is integral over the subring generated by  $I_1, \dots, I_\mu$ .*

**§ 3. The ring of invariants of  $2 \times 2$  matrices**

In this section we shall be concerned with the ring of invariants of  $2 \times 2$  matrices. Throughout this section we assume that  $l \geq 2$ .

**PROPOSITION 3.1.** (1) *The Poincaré series  $P_2(z)$  is given by*

$$P_2(z) = (-1)^{l-1} \frac{1}{2(l-1)!(1-z)^{2l}} \left( \frac{d}{d\varepsilon} \right)^{l-1} \frac{\varepsilon^{l-2}(\varepsilon-1)^2}{(z\varepsilon-1)^l} \Big|_{\varepsilon=z} .$$

(2) *The Laurant expansion of  $P_2(z)$  at  $a = 1$  begins with*

$$P_2(z) = \frac{[l-1]_{l-2}}{(l-1)! 2^{2l-1} (1-z)^{4l-3}} + \frac{3[l-1]_{l-2}}{(l-1)! 2^{2l} (1-z)^{4l-4}} + \dots ,$$

where  $[l-1]_{l-2} = (l-1)l(l+1) \cdots (2l-4)$ .

(3) If  $C[X(1), \dots, X(l)]^{GL(2)} = \bigoplus_{i=1}^r \varphi_i C[f_1, \dots, f_{4l-3}]$ , where  $f_1, \dots, f_{4l-3}$  is a system of parameters of  $C[X(1), \dots, X(l)]^{GL(2)}$ , we have

$$r = \frac{[l-1]_{l-2}}{(l-1)!} \prod_{i=1}^{4l-3} \frac{\deg(f_i)}{2^{2i-1}}.$$

*Proof.* (1) follows from (2.4). By a direct computation, we see that the first Laurant coefficient at  $z = 1$  equals

$$\frac{[l-1]_{l-2}}{(l-1)! 2^{2l-1}}.$$

Then (2) follows from (2.7). (3) is an immediate consequence from (2) and (2.6).

We denote by  $C_i$  a subring of  $C[X(1), \dots, X(l)]^{GL(2)}$  generated by traces  $\text{Tr}(X(i)X(j))$ ,  $1 \leq i, j \leq l$ ,  $\text{Tr}(X(i))$ ,  $1 \leq i \leq l$ .

**PROPOSITION 3.2.** *The ring of invariants  $C[X(1), \dots, X(l)]^{GL(2)}$  is integral over  $C_i$ .*

*Proof.* By Theorem 1.1, it is enough to show

$$(*) \quad \begin{aligned} &\text{if } \text{Tr}(A_i A_j) = \text{Tr}(A_i) = 0 \quad (A_i, A_j \in M(2, C), 1 \leq i, j \leq l), \\ &\text{Tr}(A_{i_1} A_{i_2} \cdots A_{i_k}) = 0 \quad \text{for any } k, 1 \leq i_1, \dots, i_k \leq l. \end{aligned}$$

We shall prove (\*) by induction on  $l$ . By making the substitution  $A_i \rightarrow BA_i B^{-1}$  ( $B \in GL(2, C)$ ), we can assume  $A_1 = 0$  or  $A_1 = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ .

If  $A_1 = 0$ , by the inductive hypothesis (\*) is true. If  $A_1 = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ , we have  $A_i = \begin{pmatrix} 0 & \alpha_i \\ 0 & 0 \end{pmatrix}$ ,  $\alpha_i \in C$  ( $1 \leq i \leq l$ ). Because  $\text{Tr}(A_1 A_i) = 0$  and  $A_i^2 = 0$ ,  $1 \leq i \leq l$ . This shows that  $\text{Tr}(A_{i_1} A_{i_2} \cdots A_{i_k}) = 0$ . This completes the proof.

If  $l = 2$  or  $3$ ,  $\text{Tr}(X(i)X(j))$  ( $1 \leq i, j \leq l$ ),  $\text{Tr}(X(i))$  ( $1 \leq i \leq l$ ) is a homogeneous system of parameters of  $C[X(1), \dots, X(l)]^{GL(2)}$ .

**PROPOSITION 3.3.** (E. Formanek, P. Halpin and W.C.W. Li [2])

$$\begin{aligned} &C[X(1), X(2)]^{GL(2)} \\ &= C[\text{Tr}(X(1)), \text{Tr}(X(2)), \text{Tr}(X(1)^2), \text{Tr}(X(2)^2), \text{Tr}(X(1)X(2))] \end{aligned}$$

*Proof.* By (3) Proposition 3.1, we have  $r = 1$  and we obtain the result.

#### §4. The ring of invariants $C[X(1), X(2)]^{GL(3)}$

In this section we treat the case:  $n = 3$  and  $l = 2$ . Set

$$\begin{aligned} f_1 &= \text{Tr}(X(1)), & f_2 &= \text{Tr}(X(1)^2), & f_3 &= \text{Tr}(X(1)^3), \\ f_4 &= \text{Tr}(X(2)), & f_5 &= \text{Tr}(X(2)^2), & f_6 &= \text{Tr}(X(2)^3), \\ f_7 &= \text{Tr}(X(1)X(2)), & f_8 &= \text{Tr}(X(1)X(2)^2), & f_9 &= \text{Tr}(X(1)^2X(2)), \\ f_{10} &= \text{Tr}(X(1)^2X(2)^2), & f_{11} &= \text{Tr}(X(1)X(2)X(1)^2X(2)^2). \end{aligned}$$

We denote by  $C$  the subring of  $C[X(1), X(2)]^{GL(3)}$  generated by ten invariants  $f_1, \dots, f_{10}$  which are algebraically independent.

**THEOREM 4.1.**  $f_1, \dots, f_{10}$  is a system of parameters of the ring  $C[X(1), X(2)]^{GL(3)}$  and

$$C[X(1), X(2)]^{GL(3)} = C \oplus f_{11}C.$$

*Proof.* Let  $A_1$  and  $A_2$  be  $3 \times 3$ -matrices which satisfy the following condition:  $f_1(A_1, A_2) = \dots = f_{10}(A_1, A_2) = 0$ .

Since  $\text{Tr}(A_i) = \text{Tr}(A_i^2) = \text{Tr}(A_i^3) = 0$ ,  $i = 1, 2$ , we have  $A_1^3 = A_2^3 = 0$ . If  $A_1^2 = A_2^2 = 0$ , it follows from the Cayley-Hamilton theorem that  $A_1A_2A_1 = A_2A_1A_2 = 0$  and hence we have, for any  $k$ ,  $\text{Tr}(A_{i_1}A_{i_2} \dots A_{i_k}) = 0$ ,  $1 \leq i_1, \dots, i_k \leq 2$ . Assume now that  $A_1^2 \neq 0$ . Then, by making the substitution

$$A_i \longrightarrow BA_iB^{-1}, \quad i = 1, 2,$$

we can assume that  $A_1$  and  $A_2$  are of the form

$$A_1 = \begin{pmatrix} 0 & 1 & \\ & 0 & 1 \\ & & 0 \end{pmatrix}, \quad A_2 = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}.$$

The equations  $\text{Tr}(A_1A_2) = \text{Tr}(A_1^2A_2) = \text{Tr}(A_2) = 0$  imply  $a_{11} + a_{22} + a_{33} = a_{21} + a_{32} = a_{31} = 0$  and  $\text{Tr}(A_1^2A_2^2) = 0$  implies  $a_{21}a_{32} = 0$ . Hence we have  $a_{31} = a_{21} = a_{32} = 0$ . This shows that  $A_2$  is an upper triangular matrix with zero diagonal entries. Consequently  $\text{Tr}(A_{i_1}A_{i_2} \dots A_{i_k}) = 0$ ,  $i_1, i_2, \dots, i_k = 1, 2$  for any  $k$ .

If  $A_1$  or  $A_2$  is the zero matrix, all traces are zero by our assumption. Therefore  $C[X(1), X(2)]^{GL(3)}$  is integral over  $C$ . Since the transcendence degree of the ring  $C[X(1), X(2)]^{GL(3)}$  is ten,  $f_1, \dots, f_{10}$  is a homogeneous system of parameters.

Consider the Poincaré series  $P(z_1, z_2)$ . By the theorem of Hochster and Roberts  $C[X(1), X(2)]^{GL(3)}$  is a free module over the subring  $C$ . Therefore



there is a polynomial  $F(z_1, z_2)$  in two variables such that

$$P(z_1, z_2) = \frac{F(z_1, z_2)}{(1-z_1)(1-z_1^2)(1-z_1^3)(1-z_2)(1-z_2^2)(1-z_2^3)(1-z_1z_2)(1-z_1^2z_2)(1-z_1^3z_2^2)(1-z_1^2z_2^2)}.$$

It follows from the functional equation of  $P(z_1, z_2)$  that  $F(z_1, z_2)$  satisfies the following relation

$$F(z_1, z_2) = (z_1z_2)^3F(z_1^{-1}, z_2^{-1}).$$

And it is easily shown that  $F(z_1, z_2) = 1 + z_1^3z_2^3$ . Therefore  $C[X(1), X(2)]^{GL(3)}$  is generated by  $f_1, \dots, f_{10}$  and an invariant  $\varphi$  of degree  $(3, 3)$ .

Invariants  $\text{Tr}(X(1)X(2)X(1)^2X(2)^2)$ ,  $\text{Tr}(X(2)X(1)X(2)^2X(1)^2)$  and  $\text{Tr}(X(1) \cdot X(2)X(1)X(2)X(1)X(2))$  span the vector space  $C[X(1), X(2)]_{(3,3)}^{GL(3)}$  consisting of invariants of degree  $(3, 3)$ . By the Cayley-Hamilton theorem, we find that  $\text{Tr}(X(1)X(2)X(1)X(2)X(1)X(2)) \in C$  and  $\text{Tr}(X(1)X(2)X(1)X(2)^2) + \text{Tr}(X(2) \cdot X(1)X(2)^2X(1)^2) \in C$ . Therefore the ring of invariants  $C[X(1), X(2)]^{GL(3)}$  is generated by  $f_1, \dots, f_{11}$  and  $C[X(1), X(2)]^{GL(3)} = C \oplus f_{11}C$ . This completes the proof.

**§ 5. The ring of invariants  $C[X(1), X(2)]^{GL(4)}$**

We denote by  $\text{Sym}(n)$  the symmetric group of  $n$  letters and recall the multi-linearized Cayley-Hamilton theorem for  $n \times n$ -matrices  $Y_1, \dots, Y_n$ :

$$\sum_{\pi \in \text{Sym}(n)} Y_{\pi(1)} \cdots Y_{\pi(n)} + \sum_{k=1}^n \sum_u \sum_{\pi \in \text{Sym}(n)} q_u \text{Tr}(Y_{\pi(1)} \cdots Y_{\pi(u_1)}) \cdots Y_{\pi(n-k+1)} Y_{\pi(n-k+2)} \cdots Y_{\pi(n)} = 0,$$

for suitable  $q_u \in \mathbf{Q}$  and suitable  $j$ -tuples  $u = (u_1, \dots, u_j)$  such that  $1 \leq u_1 \leq u_2 \leq \dots \leq u_j$  and  $u_1 + \dots + u_i = k$ .

**PROPOSITION 5.1.** *The ring of invariants  $C[X(1), X(2)]^{GL(4)}$  is generated by invariants of the form*

$$\begin{aligned} &\text{Tr}(X(1)^{\alpha_1}X(2)^{\alpha_2}X(1)^{\alpha_3}X(2)^{\alpha_4}), \quad 0 \leq \alpha_1, \alpha_2, \alpha_3 \leq 3, \\ &\text{Tr}(X(1)X(2)X(1)^2X(2)^2X(1)^3X(2)^3), \quad \text{Tr}(X(1)X(2)X(1)X(2)^2X(1)X(2)^3), \\ &\text{Tr}(X(2)X(1)X(2)X(1)^2X(2)X(1)^3). \end{aligned}$$

*Proof.* We claim that any invariant  $\text{Tr}(X(1)^{\alpha_1}X(2)^{\alpha_2} \cdots X(1)^{\alpha_{2r-1}}X(2)^{\alpha_{2r}})$ ,  $0 \leq \alpha_1, \dots, \alpha_{2r} \leq 3$  ( $r > 6$ ), can be written as a polynomial in  $T(X(1)^{\beta_1}X(2)^{\beta_2} \cdots X(1)^{\beta_5}X(2)^{\beta_6})$ ,  $0 \leq \beta_1, \dots, \beta_6 \leq 3$ . We work by induction on  $r$ . We assume

that, for any  $r' < r$ , this assertion is true. Apply the multi-linearized Cayley-Hamilton theorem for  $4 \times 4$ -matrices  $X_1, X_2, X_3, X_4$  to the case  $X_1 = X(1)^{\alpha_1}, X_2 = X(2)^{\alpha_2}, X(1)^{\alpha_3}, X_3 = X(2)^{\alpha_4}, X_4 = X(1)^{\alpha_5}X(2)^{\alpha_6}$ . Then by the inductive hypothesis we conclude the assertion. A similar argument shows that any invariant of the form

$$\text{Tr}(X(1)^{\alpha_1}X(2)^{\alpha_2}X(1)^{\alpha_3}X(2)^{\alpha_4}X(1)^{\alpha_5}X(2)^{\alpha_6}), \quad 1 \leq \alpha_1, \alpha_2, \dots, \alpha_6 \leq 3,$$

is written as a polynomial in  $\text{Tr}(X(1)^{\alpha_1}X(2)^{\alpha_2}X(1)^{\alpha_3}X(2)^{\alpha_4}), 0 \leq \alpha_1, \dots, \alpha_4 \leq 3, \text{Tr}(X(1)X(2)X(1)^2X(2)^2X(1)^3X(2)^3), \text{Tr}(X(1)X(2)X(1)X(2)^2X(1)X(2)^3), \text{Tr}(X(2) \cdot X(1)X(2)X(1)^2X(2)X(1)^3)$ . The proposition is proved.

Set

$$\begin{aligned} f_1 &= \text{Tr}(X(1)), & f_2 &= \text{Tr}(X(1)^2), & f_3 &= \text{Tr}(X(1)^3), & f_4 &= \text{Tr}(X(1)^4), \\ f_5 &= \text{Tr}(X(2)), & f_6 &= \text{Tr}(X(2)^2), & f_7 &= \text{Tr}(X(2)^3), & f_8 &= \text{Tr}(X(2)^4), \\ f_9 &= \text{Tr}(X(1)X(2)), & f_{10} &= \text{Tr}(X(1)^2X(2)^2), & f_{11} &= \text{Tr}(X(1)X(2)^2), \\ f_{12} &= \text{Tr}(X(1)^2X(2)), & f_{13} &= \text{Tr}(X(1)X(2)^3), & f_{14} &= \text{Tr}(X(1)^3X(2)), \\ f_{15} &= \text{Tr}(X(1)X(2)X(1)X(2)), & f_{16} &= \text{Tr}(X(1)X(2)^2X(1)X(2)^2), \\ f_{17} &= \text{Tr}(X(2)X(1)^2X(2)X(1)^2). \end{aligned}$$

We denote by  $C$  a subring of  $C[X(1), X(2)]^{GL(4)}$  generated by  $f_1, \dots, f_{17}$ .

**PROPOSITION 5.2.**  $f_1, \dots, f_{17}$  is a homogeneous system of parameters of the ring of invariants  $C[X(1), X(2)]^{GL(4)}$ .

*Proof.* Since the transcendence degree of the ring  $C[X(1), X(2)]^{GL(4)}$  is 17, it is enough to show that, for  $4 \times 4$ -matrices  $A_1$  and  $A_2, f_1(A_1, A_2) = \dots = f_{17}(A_1, A_2) = 0$  imply  $\text{Tr}(A_{i_1}, A_{i_2} \dots A_{i_k}) = 0, i_1, \dots, i_k = 1, 2$  for any  $k$ .

Notice that  $A_1^4 = A_2^4 = 0$ , since  $f_4(A_1, A_2) = \dots = f_8(A_1, A_2) = 0$ . Assume that  $A_1^3 \neq 0$ . Then, by the substitution  $A_i \rightarrow BA_iB^{-1}, B \in GL(4)$  and  $i = 1, 2$ , we can assume that

$$A_1 = \begin{pmatrix} 0 & 1 & & \\ & 0 & 1 & \\ & & 0 & 1 \\ & & & 0 \end{pmatrix} \quad \text{and} \quad A_2 = \begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{pmatrix}$$

It follows from the equations  $\text{Tr}(A_1^2A_2) = \text{Tr}(A_1^3A_2) = 0$  that  $a_{41} = a_{31} + a_{42} = 0$  and the Cayley-Hamilton theorem shows that the equation  $\text{Tr}(A_1^2A_2A_1^2A_2) = 0$  implies  $\text{Tr}(A_1^2A_1A_2A_1A_2) = 0$ .

Since

$$A_1 A_2 = \begin{pmatrix} a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ 0 & a_{42} & a_{43} & a_{44} \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

it follows from the equation  $\text{Tr}(A_1^2 A_1 A_2 A_1 A_2) = 0$  that  $a_{31} a_{42} = 0$  and hence we have  $a_{31} = a_{42} = 0$ . Then it follows from the relation  $\text{Tr}(A_1 A_2) = a_{21} + a_{32} + a_{43} = 0$  that  $\text{Tr}(A_1^2 A_2^2) = a_{21} a_{32} + a_{32} a_{43} = -a_{32}^2$  and we obtain  $a_{32} = 0$ .

Since

$$\begin{aligned} \text{Tr}(A_1 A_2 A_1 A_2) &= \text{Tr} \left( \begin{pmatrix} a_{21} & a_{22} & a_{23} & a_{24} \\ 0 & 0 & a_{33} & a_{34} \\ 0 & 0 & a_{43} & a_{44} \\ 0 & 0 & 0 & 0 \end{pmatrix}^2 \right) \\ &= a_{21}^2 + a_{43}^2, \end{aligned}$$

$a_{21} = a_{43} = a_{32} = 0$  and hence  $A_2$  is a  $4 \times 4$  upper triangular matrix with zero diagonal entries. Consequently we can conclude that  $\text{Tr}(A_{i_1}, A_{i_2}, \dots, A_{i_k}) = 0, 1 \leq i_1, i_2, \dots, i_k \leq 2$  for any  $k$ . By the same argument, we obtain the same conclusion if  $A_2^3 \neq 0$ .

We next assume that  $A_1^3 = A_2^3 = 0$  and either  $A_1^2$  or  $A_2^2$  is not zero. Then we can take  $A_1$  as

$$A_1 = \begin{pmatrix} 0 & 1 & & \\ & 0 & 1 & \\ & & 0 & \\ & & & 0 \end{pmatrix} \quad \text{or} \quad \begin{pmatrix} 0 & & & \\ & 0 & 1 & \\ & & 0 & 1 \\ & & & 0 \end{pmatrix},$$

and divide into two cases:

Case 1.

$$A_1 = \begin{pmatrix} 0 & 1 & & \\ & 0 & 1 & \\ & & 0 & \\ & & & 0 \end{pmatrix}, \quad A_2 = \begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{pmatrix}.$$

In this case, it follows from the equations  $\text{Tr}(A_1^2 A_2) = 0, \text{Tr}(A_1 A_2 A_1 A_2) = 0$  and  $\text{Tr}(A_1 A_2) = 0$  that  $a_{21} = a_{31} = a_{32} = 0$ .

Therefore  $A_1A_2$  and  $A_1^2A_2$  are upper triangular matrices with zero diagonal entries. Similarly, replacing  $A_2$  by  $A_2^2$ , we see that  $A_1A_2^2$  and  $A_1^2A_2^2$  are also upper triangular matrices with zero diagonal entries. This shows that  $\text{Tr}(A_{i_1}A_{i_2} \cdots A_{i_k}) = 0$ ,  $1 \leq i_1, i_2, \dots, i_k \leq 2$  for any  $k$ .

Case 2.

$$A_1 = \begin{pmatrix} 0 & & & \\ & 0 & 1 & \\ & & 0 & 1 \\ & & & 0 \end{pmatrix} \quad A_2 = \begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{pmatrix}.$$

In this case, by the equation  $\text{Tr}(A_1^2A_2) = 0$ , we have  $a_{42} = 0$ .

Since

$$A_1A_2 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

and  $\text{Tr}(A_1A_2A_1A_2) = 0$ , we have  $a_{32} = a_{43} = 0$ . Then we find that  $A_1A_2A_1 = a_{33}A_1^2$  and, replacing  $A_2$  by  $A_2^2$ ,  $A_1A_2^2A_1 = bA_1^2$ . Here  $b$  denotes the  $(3, 3)$ -entry of the matrix  $A_2^2$ .

Notice that, for any  $4 \times 4$ -matrix  $X = (x_{ij})$ ,

$$A_1^2X = \begin{pmatrix} 0 & x_{32} & x_{33} & x_{34} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

Therefore we can conclude that  $\text{Tr}(A_{i_1}A_{i_2} \cdots A_{i_k}) = 0$  for any  $k$ .

If  $A_1^2 = A_2^2 = 0$ , we have evidently  $\text{Tr}(A_{i_1}A_{i_2} \cdots A_{i_k}) = 0$ . This completes the proof.

Proposition 5.2 shows that  $C$  is a polynomial ring in 17 variables and  $C[X(1), X(2)]^{GL(4)}$  is a free module over  $C$ .

REFERENCES

[ 1 ] E. Formanek, The center of the ring  $3 \times 3$  generic matrices, *Linear and Multilinear Algebra*, **7** (1979), 203–212.  
 [ 2 ] —, P. Halpin and W.-C. W. Li, The Poincaré series of the ring of  $2 \times 2$  generic matrices, *J. Algebra*, **69** (1981), 105–112.

- [ 3 ] D. Hilbert, Über die vollen Invariantensysteme, Ges. Abh., II, 287–344, Springer-Verlag, 1970.
- [ 4 ] M. Hochster and J. Roberts, Rings of invariants of reductive groups acting on regular rings are Cohen-Macaulay, Adv. in Math., **13** (1974), 115–175.
- [ 5 ] C. Procesi, The invariant theory of  $n \times n$  matrices, Adv. in Math., **19** (1976), 306–381.
- [ 6 ] M. Rosenlicht, A remark on quotient spaces, An. Acad. Brasil Ciênc., **35**, (1963), 487–489.
- [ 7 ] T. A. Springer, Invariant theory Springer Lecture note 585, 1977.
- [ 8 ] H. Weyl, The classical groups, Princeton Univ. Press, Princeton, N. J., 1946.
- [ 9 ] ———, Zur Darstellungstheorie und Invariantenabzählung der projection, der Komplex und Drehungsgrope, Ges. Abh., III, 1–25, Springer-Verlag, 1968.

*Department of Mathematics*  
*Faculty of Science*  
*Nagoya University*  
*Chikusa-ku, Nagoya, 464*  
*Japan*