

Erratum

Cite this article: Goenjian AK, Steinberg AM, Walling D, Bishop S, Karayan I, Pynoos R (2021). 25-year follow-up of treated and not-treated adolescents after the Spitak earthquake: course and predictors of PTSD and depression—ERRATUM. *Psychological Medicine* **51**, 989–990. <https://doi.org/10.1017/S0033291720001348>

Received: 28 April 2019

Revised: 29 July 2019

Accepted: 6 December 2019

First published online: 8 May 2020

Key words:


Depression; disaster; earthquake; longitudinal study; PTSD; risk and protective factors

Author for correspondence:

Armen K. Goenjian,

E-mail: agoenja@aol.com

25-year follow-up of treated and not-treated adolescents after the Spitak earthquake: course and predictors of PTSD and depression—ERRATUM

Armen K. Goenjian , Alan M. Steinberg, David Walling, Sheryl Bishop, Ida Karayan and Robert Pynoos

doi.org/10.1017/S0033291719003891, Published online by Cambridge University Press, 14 January 2020

The above mentioned article was published in *Psychological Medicine* with some errors in Table 3. The correct version of [Table 3](#) is shown below:

Cambridge University Press apologises for this error.

Reference

Goenjian, A., Steinberg, A., Walling, D., Bishop, S., Karayan, I., & Pynoos, R. (2020). 25-year follow-up of treated and not-treated adolescents after the Spitak earthquake: Course and predictors of PTSD and depression. *Psychological Medicine*, 1–13. [doi:10.1017/S0033291719003891](https://doi.org/10.1017/S0033291719003891)

© The Author(s), 2020. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.

Table 3. The full model and the final stepwise model for PTSD-RI, PCL, and CES-D scores at 25-year follow-up among adults who were early-adolescents in 1988

Outcome variables and predictors	Final model (best predictors)			Full model		
	<i>B</i>	β	<i>p</i>	<i>B</i>	β	<i>p</i>
PTSD-RI score at 25 years	$F_{7, 127} = 9.98, p < \mathbf{0.001}, R^2 = 0.36$			$F_{13, 121} = 5.71, p < 0.001, R^2 = 0.38$		
1. Home destruction	-3.826	-0.146	0.055	-3.867	-0.148	0.063
2. PTSD-RI at 1½ years	0.263	0.212	0.005	0.199	0.160	0.099
3. Treatment	4.898	0.250	0.032	5.175	0.174	0.027
4. Post-EQ adversities	-8.128	-0.250	0.001	-7.838	-0.241	0.001
5. Post-EQ chronic illnesses	1.877	0.224	0.003	2.229	0.266	0.001
6. Social support	-0.415	-0.265	<0.001	-0.382	-0.244	0.002
7. Sex	3.606	0.138	0.059	2.593	0.099	0.195
8. Nuclear family deaths				-1.535	-0.085	0.259
9. Severe injuries due EQ				-3.543	-0.096	0.225
10. Post-EQ trauma				0.411	0.052	0.498
11. DSRS at 1½ years				0.025	0.009	0.915
12. SAD. at 1½ years				0.611	0.111	0.172
13. Education				-1.689	-0.064	0.403
PCL score at 25 years	$F_{4, 130} = 15.78, p < \mathbf{0.001}, R^2 = 0.33$			$F_{13, 121} = 5.56, p < 0.001, R^2 = 0.37$		
1. Home destruction				-2.245	-0.075	0.345
2. PTSD-RI at 1½ years				0.096	0.068	0.486
3. Treatment	6.542	0.192	0.009	5.194	0.153	0.053
4. Post-EQ adversities	-5.186	-0.140	0.053	-5.120	-0.138	0.064
5. Post-EQ chronic illnesses	3.396	0.354	<0.001	3.482	0.364	<0.001
6. Social support	-0.526	-0.294	<0.001	-0.501	-0.280	<0.001
7. Sex				1.294	0.043	0.572
8. Nuclear family deaths				-1.230	-0.059	0.430
9. Severe injuries due EQ				-4.119	-0.098	0.220
10. Post-EQ trauma				1.004	0.111	0.150
11. DSRS at 1½ years				-0.115	-0.039	0.664
12. SAD at 1½ years				0.652	0.104	0.205
13. Education				-1.047	-0.035	0.651
CES-D score at 25 years	$F_{6, 128} = 14.08, p < \mathbf{0.001}, R^2 = 0.40$			$F_{13, 121} = 6.94, p < 0.001, R^2 = 0.43$		
1. Home destruction	-3.160	-0.131	0.065	-2.537	-0.105	0.169
2. PTSD-RI at baseline				0.131	0.114	0.222
3. Treatment				3.272	0.119	0.114
4. Post-EQ adversities	-6.785	-0.226	0.002	-7.068	-0.235	0.001
5. Post-EQ chronic illnesses	2.798	0.362	<0.001	2.798	0.362	<0.001
6. Social support	-0.420	-0.290	<0.001	-0.428	-0.296	<0.001
7. Sex				0.467	0.019	0.792
8. Nuclear family deaths				1.316	0.079	0.276
9. Severe injuries due EQ				-0.808	-0.024	0.755
10. Post-EQ trauma	0.880	0.120	0.093	0.754	0.103	0.162
11. DSRS at 1½ years				0.075	0.031	0.715
12. SAD at 1½ years	0.791	0.155	0.026	0.931	0.183	0.020
13. Education				-1.85	-0.076	0.302

PTSD-RI, Posttraumatic Stress Disorder-Reaction Index; PCL, PTSD Checklist; CES-D, Center for Epidemiologic Studies-Depression; EQ, earthquake; SAD, separation anxiety disorder. Reference groups used for the analyses: no home destruction; no treatment; no adversities; no death; and no injury.