

Correspondence

Letters for publication in the Correspondence columns should be addressed to:

The Editor, British Journal of Psychiatry, 17 Belgrave Square, London SW1X 8PG

THE SCOPE OF PSYCHIATRY

DEAR SIR,

During a recent meeting of the North Western division of the College there was considerable discussion of the possible dangers of colleagues in associated disciplines, such as psychology, acting independently of psychiatrists in the treatment of patients within the Health Service. Disquiet may well be appropriate, but should perhaps be mitigated by consideration of the sort of problem that these non-medical professionals seek to treat. It is too easy to consider that an erosion of the boundaries of the proper medical sphere of responsibility is taking place, when in fact it could be equally argued that those boundaries have already become inflated beyond the limits of medical competence.

Demands are being made on the psychiatrist for help with problems for which his training equips him ill, and for which he can find little time in an overcrowded time-table. One movement towards mitigation of this position is expressed in a call for more and better trained psychiatrists, but the evidence is that, even were this wholly desirable, the call is unlikely to be answered in the foreseeable future. Moreover recent criticisms of the workings of our mental hospitals suggest that some retrenchment in the core areas of our responsibility should take precedence over demarcation disputes more peripherally.

This said, perhaps you would allow me to advance a rather more abstract speculation regarding the roots of some of our current dilemmas in respect of our responsibility to those people who differ from the accepted social norms in a way which causes them trouble and unhappiness. These people may define themselves as 'patients' by the simple expedient of consulting their doctors. I am thinking here of those with addictions, sexual abnormalities, poor control of aggression, and parasuicidal tendencies, etc, the vast majority of whom manifest no major psychiatric syndrome.

In relation to these people I would hold that medicine, and psychiatry in particular, has been a social tool (not necessarily in a pejorative sense) in

bringing about a major change of public attitude. Many actions considered to merit punishment, and states of mind considered morally reprehensible or despicable, have, by inclusion within the framework of respectable medical practice, become transformed into conditions meriting sympathy and treatment. In historical perspective I think it yet remains to be seen whether this change has brought wholly desirable results or increased the sum of happiness of the age in which we live.

More parochially, it seems to me that we have reached a position which holds considerable dangers for our profession. Having, by our tacit acceptance of so much common human disability and unhappiness as illness, facilitated a change in social attitude, we need to take care that we do not unthinkingly follow through the medical model in which we were trained and too easily accept it as our responsibility to provide 'treatment' for a very significant proportion of the population.

Two serious risks to the psychiatric profession seem to lie in the areas I have mentioned. Firstly, our credibility with our medical colleagues and the public at large may be put into such danger that our advocacy on behalf of the seriously mentally sick and handicapped will lose force. Secondly, our own self-esteem and job satisfaction must be at hazard when we spread our efforts so thinly that we can perceive only a relatively small part of our work as well done.

M. W. FORTH

*West Cheshire Hospital,
Liverpool Road,
Chester CH1 3ST*

LIMBIC LEUCOTOMY

DEAR SIR,

We welcome the useful paper by Mitchell-Heggs *et al* (*Journal*, March 1976, **128**, pp 226-40) on limbic leucotomy, but we would like to question some of the conclusions. It is stated (p 237) that 'In the present series, however, where lower medial quadrant lesions were combined with cingulate lesions, the results are superior', i.e. superior to the results after lesions in the lower