

STARKSTEIN, S. E., FEDOROFF, P., BERTHIER, M. L., *et al* (1991) Manic-depressive and pure manic states after brain lesions. *Biological Psychiatry*, **29**, 149–158.

MARCELO L. BERTHIER

*Virgen de la Victoria University Hospital
Apartment 3091, 29010 Málaga, Spain*

JAIME KULISEVSKY

Sant Pau Hospital

Autonomous University of Barcelona, Spain

Relapse following withdrawal of drug addiction

SIR: There are few published follow-up studies from developing countries on relapse following withdrawal of opiate addiction. I wish to report part of a follow-up study which was carried out on detoxified patients discharged from the Post Graduate Hospital, Dhaka, Bangladesh. The pattern of relapse, effect of group therapy, and probable variables that predict relapse were examined.

Case report. We studied 50 subjects consecutively admitted to the psychiatric unit from December 1987 to July 1988 for detoxification. All were men with a mean age of 28.6 years. In 77% of cases heroin was smoked. Treatment included a two-week withdrawal period with symptomatic medication (chlorpromazine and benzodiazepines) followed by out-patient group therapy for six months. Data were obtained through a questionnaire interview schedule. We measured variables such as peer pressure, causes of relapse stated by patient, family and work situation, type and amount of drugs used in relapse, and physical and sexual difficulties after discharge on a five-point scale.

Subjects were interviewed on admission to the unit before detoxification had begun. The interview covered socio-demographic features including details of drug history. Further interviews were made on the first, third and sixth month following discharge. An operational definition of relapse and abstinence was formulated for the project. We considered relapse as a single use of addiction-forming drugs during follow-up (Hall *et al*, 1991). Abstinence was defined as complete absence of any addictive drug during the period. Only 12 patients voluntarily attended follow-up interviews; information about the rest were obtained by home visits.

By the end of the sixth month, seven subjects remained abstinent. Twenty-eight subjects remained drug free until the one-month follow-up point. At the end of the third month this figure was reduced to less than half (11).

A recent report of outcome by Gossop *et al* (1987) is rather optimistic. Our high relapse rate has to be weighed against the absence of any rehabilitation package. Most patients used less amounts of the addictive drug at relapse than before treatment.

We were interested in the causes of relapse as stated by the subjects. Our results show evidence of peer pressure in 15 (34.88%) of 43 relapses in the form of constant exposure to drug-related cues and offering of

drugs (Marlatt & Gordon, 1985). Craving as a cause of relapse was mentioned by another 11 subjects.

Our results, both in terms of the high relapse rate and the psychological factors that may contribute to this, are strikingly similar to those reported from the West and illustrate the truly international nature of the future of opiate addiction.

GOSSOP, M., GREEN, L., PHILLIPS, G., *et al* (1987) What happens to opiate addicts immediately after treatment: a prospective study. *British Medical Journal*, **294**, 1377–1380.

HALL, S. M., WASSERMAN, D. A. & HAVASSY, B. E. (1991) Relapse prevention. *NIDA Research Monograph*, **106**, 279–292.

MARLATT, G. A. & GORDON, J. R. (1985) *Relapse Prevention*. New York: Guilford Press.

REZAUL ISLAM

*Institute of Psychiatry
De Crespigny Park
Denmark Hill
London SE5 8AF*

Percutaneous endoscopic gastrostomy and severe endogenous depression

SIR: Percutaneous endoscopic gastrostomy (PEG) is a method for obtaining access to the stomach in patients requiring long-term tube-feeding (Bussone *et al*, 1992). The tube is inserted using a gastroscope while the patient is sedated. Since its first description, PEG has become one of the favoured techniques of feeding patients with persistent swallowing difficulty due to neurological or oropharyngeal disorders (Gauderer *et al*, 1980). We recently used this method of feeding in a patient who presented with severe endogenous depression with secondary anorexia and weight loss who improved markedly after electroconvulsive therapy (ECT) after stabilisation of her nutritional status with enteric feeding via PEG.

Case report. A 56-year-old woman was admitted from her home to the psychiatric hospital under Section 2 of the Mental Health Act 1983 with a diagnosis of severe endogenous depression. There was a recent concern by the community psychiatric nurse (CPN) over her poor and intermittent intake of foods and fluids. She had irrational ideas about her digestive processes and her bowels, and nihilistic delusions of what happened to food once she swallowed it. She had lost 19 kg in weight (her body weight was now 32 kg) and her physical state on admission was of marked cachexia and dehydration with a urea concentration of 13.6 mmols/l.

The intensity of her delusion made it impossible to obtain informed consent. A second opinion was obtained from an independent psychiatrist who authorised a course of ECT. Rehydration was started intravenously but her nutritional status continued to decline and she was referred for consideration of parenteral nutrition. It was decided that PEG would be a safer and more effective method of feeding. The