

Original Article

Cite this article: Gillan PC, Johnston S (2024). Nursing students satisfaction and self-confidence with standardized patient palliative care simulation focusing on difficult conversations. *Palliative and Supportive Care*. <https://doi.org/10.1017/S1478951523001918>

Received: 17 April 2023
Revised: 21 September 2023
Accepted: 4 December 2023


Keywords:


Difficult conversations; end-of-life care; palliative care; standardized patient simulation; undergraduate nursing students

Corresponding author:

Pauline Catherine Gillan;
Email: pgillan@csu.edu.au

Nursing students satisfaction and self-confidence with standardized patient palliative care simulation focusing on difficult conversations

Pauline Catherine Gillan, PH.D., M.N. (PALLIATIVE CARE)¹  and

Sandra Johnston, PH.D., M.B.A., B.H.S. (NURSING)² 

¹School of Nursing, Paramedicine and Healthcare Sciences, Charles Sturt University Faculty of Science and Health, Port Macquarie, NSW, Australia and ²School of Nursing, Queensland University of Technology, Brisbane, QLD, Australia

Abstract

Objectives. Lack of experience communicating with patients and families at the end of life are key concerns for nursing students. Palliative care simulation using standardized patients (SPs) focusing on difficult conversations may lead to increased self-confidence in providing palliative and end-of-life care in clinical practice. There is currently a paucity of research on SP palliative care simulations in undergraduate nursing education. The objective of this research was to assess 3rd year undergraduate nursing students' levels of satisfaction and self-confidence with palliative and end-of-life care simulations focusing on difficult conversations, as measured by the Student Satisfaction and Self-Confidence in Learning Scale (SSSCLS) and the Simulation Design Scale (SDS).

Methods. A descriptive post-intervention study. Four palliative care simulation days, consisting of 2 clinical scenarios, were conducted over 4 weeks. The first simulation was an outpatient palliative care clinic scenario, and the second was an inpatient hospital scenario. Nursing students enrolled in a 3rd year nursing palliative care elective ($n = 51$) at an Australian university were invited to participate. Students who attended simulation days were eligible to participate ($n = 31$). Immediately post-simulation, students were invited to complete the SSSCLS and the SDS. Fifty-seven surveys were completed (simulation 1, $n = 28$; simulation 2, $n = 29$). The data were analyzed using descriptive statistics.

Results. Results showed that students had high levels of self-confidence in developing palliative care and communication skills after both simulation experiences and high levels of satisfaction with the SP simulations.

Significance of results. The lack of published literature on palliative care and end-of-life SP simulation highlights the need to collect further evidence to support this as an innovative approach to teaching palliative care. SP palliative care simulation focusing on difficult conversations assists in developing students' communication skills and improves satisfaction and self-confidence with palliative and end-of-life care.

Introduction

Nurses are the primary providers of end-of-life care (Ristevski et al. 2022), and being with someone who is dying is not only a privilege but an essential part of nursing practice (Cheong et al. 2019). However, research indicates that undergraduate nursing students are largely unprepared to provide competent palliative and end-of-life care in clinical practice (Gillan et al. 2021a). For many undergraduate nursing students, providing end-of-life care is a challenging and anxiety-provoking prospect (Fabro et al. 2014). Students' key concerns relate to their lack of knowledge and experience in communicating with patients and families and are reported to have difficulty initiating and engaging in difficult conversations at the end of life (Bloomfield et al. 2015; Kantor and Stadelman 2020; Kirkpatrick et al. 2017). Students also report not knowing what to say and having difficulties dealing with emotional responses (Gillett et al. 2016). It is imperative that nursing students are given opportunities to develop effective communication skills, have confidence to initiate difficult conversations, and are able to provide emotional support for patients in their care (Kantor and Stadelman 2020).

It is widely acknowledged that the most valuable learning experiences are obtained through experiential learning within clinical practice settings (Garrino et al. 2017). However, challenges in nursing education continue due to the limited availability of clinical placements (Sarabia-Cobo et al. 2016), especially in speciality areas of practice such as palliative care, where there

© The Author(s), 2024. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

are limited opportunities to develop communication skills focusing on difficult conversations (Bloomfield et al. 2015).

In response to the limited access to clinical placements internationally, nursing educators are increasingly utilizing simulation in areas such as critical care (Bautista and Bartos 2021), resuscitation (Demirtas et al. 2021), and teaching and assessment of psychomotor skills (Hoang et al. 2022). Introducing simulated palliative and end-of-life care into undergraduate nursing curricula has the potential to provide students with end-of-life care experiences that may be absent in clinical placements. Palliative and end-of-life care simulation has the potential to enhance nursing students' experiences with end-of-life care (Gillan et al. 2014a, 2016; Kirkpatrick et al. 2017; Kunkel et al. 2016) and can facilitate nursing students' self-confidence and competence in providing palliative and end-of-life care in clinical practice (Alconero-Camarero et al. 2018).

In Australian nursing education, nursing students are widely exposed to simulation in a wide range of contexts; however, palliative care and end-of-life care simulations are still developing (Gillan et al. 2014b). A recent review by Kirkpatrick et al. (2017) investigating the use of palliative care simulation in undergraduate nursing education found only 2 ($n = 2$) of the 19 articles originating from Australia (Gillan et al. 2013, 2016); and Hoang et al.'s (2022) scoping review of end-of-life care simulation in undergraduate nursing education identified 42 articles, with only 3 originating from Australia from the same authors (Gillan et al. 2013, 2016, 2021b).

A method of simulation that is underutilized in nursing education is standardized patient (SP) simulation. SP simulation involves trained actors who act out a particular case scenario and take on many roles in the simulation, including patients, family members, or health professionals (Coates 2021). In learning situations focusing on psychosocial settings where a human voice and emotion are needed, the use of SPs can be particularly valuable (Fink et al. 2014). The realism of a situation created when using SPs has the potential to consolidate the linking of theory to practice more effectively than when interacting with a manikin (Tamaki et al. 2019). Scenarios involving patient actors allow students the opportunity to practice and improve on therapeutic communication skills (Halliday et al. 2022), psychomotor skills (Sarmasoglu et al. 2016), and students self-efficacy (Escribano et al. 2021) and are widely known as a better choice when compared to manikins when learning involves patients' psychosocial or emotional responses (Fink et al. 2014).

Due to the increased realism of SPs, this is an exciting mode of simulation for the application of palliative care and end-of-life care education. However, there is limited research on the use of SPs in palliative and end-of-life care simulation for undergraduate nursing students, with only 3 research articles available in the literature, none of these studies arising from Australia. These studies used either a randomized controlled method (Tamaki et al. 2019) or a quasi-experimental approach (Escribano et al. 2021; Fink et al. 2014). The results of these studies showed that the SP simulation improved communication skills (including empathy, respect, and assertiveness) (Escribano et al. 2021), increased self-efficacy and self-confidence with communication (Escribano et al. 2021; Fink et al. 2014), and physical assessment and psychological care (Tamaki et al. 2019). Knowledge and skill performance scores also increased significantly (Tamaki et al. 2019). While these studies provided evidence to support SP as an effective method of teaching practical skills and communication skills, there was no focus on the psychosocial aspects of difficult conversations that occur as part of end-of-life care.

While there is a growing repertoire of palliative care simulation, the use of SPs is sparsely utilized in undergraduate nursing education in Australia. A recent scoping review on the use of simulation in undergraduate nursing education in Australia identified 9 out of 44 journal articles using SP simulation (Kunst et al. 2018), with only one focusing on palliative or end-of-life care (Gillan et al. 2013) and none with a focus on communication at the end of life. Although the focus of this review was not specifically relating to palliative care or communication, this recent review does provide a snapshot of the types of simulation approaches used in Australian undergraduate nursing programs.

These studies indicate an increased interest in palliative care or end-of-life care simulation using SPs; however, there is a gap in simulation, specifically focusing on difficult conversations at the end of life. Only 3 research articles, one qualitative (Jeffers et al. 2022), one quantitative (Cannity et al. 2021), and one mixed-method study (Bloomfield et al. 2015), were found to directly address end-of-life care conversations. Major findings from these studies were increased self-confidence with communication at the end of life following SP simulation (Cannity et al. 2021), increased confidence and competence with communicating with dying patients and their families (Bloomfield et al. 2015), and "delivering bad news is difficult" (Jeffers et al. 2022: 202).

Self-confidence is an important construct to develop for nursing students during their undergraduate degree. Nursing students who are self-confident in their knowledge and clinical skills are able to cope with complex clinical situations (Guerrero et al. 2022), including managing difficult conversations at the end of life. Measuring self-confidence was an important consideration for some of the identified studies; however, all but one of these studies used researcher-developed self-confidence scales (Escribano et al. 2021). Escribano et al. (2021) used the "Self-efficacy in communication skills SE12 Scale," which has shown internal consistency (Cronbach's α 0.95) and test-retest reliability (interclass correlation coefficient 0.71).

Student satisfaction is important to evaluate as it is associated with greater engagement and motivation, which subsequently facilitates the learning process (Alconero-Camarero et al. 2018). Student satisfaction with the method of simulation is also important, as this has not been fully explored in previous studies in the context of palliative care. Only one study reported on satisfaction with SPs simulation (Fink et al. 2014). Satisfaction with the SP method in this study was measured by only 2 questions on the 10-item satisfaction survey, providing minimal evaluation data (Fink et al. 2014). While the results of the other studies demonstrated students' acquisition of skills, there was no evaluation of students' satisfaction with an SP being used as the method of simulation, which is an area lacking in nursing education (Escribano et al. 2021; Tamaki et al. 2019).

While the findings from these studies are positive, there are some limitations to be considered. One study was a pilot study with small numbers ($n = 15$ nursing students pre-test/post-test responses) (Bloomfield et al. 2015). The SP experience in Cannity et al.'s (2021) study was only one small component of a day-long training session, therefore findings may be attributed to the whole course and not solely to the SP simulation. The primary purpose of the study by Jeffers et al. (2022) was to use simulation to reflect on communication competence, therefore findings excluded any details on the student experience of interacting with the SP. Due to the limited number of studies on this topic, further studies involving the use of SP in end-of-life discussions are still needed (Bloomfield et al. 2015).

The results of these studies show a significant gap in the literature on the use of SP simulation in palliative and end-of-life care, with a focus on difficult conversations at the end of life from an Australian perspective.

Thus the objective of our study was to assess nursing students' satisfaction and self-confidence with palliative care and end-of-life care simulation using the Student Satisfaction and Self-Confidence in Learning Scale (SSSCLS) (National League of Nursing [NLN], 2005a) and the Simulation Design Scale (SDS) (NLN, 2005b), scales not previously used in SP palliative care or end-of-life care simulation, and address the gap in knowledge on palliative care and end-of-life care simulation using SPs focusing on difficult conversations.

Hypothesis

The hypothesis proposed for this study was: palliative care simulation using SP patients leads to high levels of satisfaction with the simulation methods used and high levels of self-confidence in mastering palliative care skills and managing difficult conversations relating to end-of-life care.

Methods

Study design

A descriptive post-intervention quantitative design.

Ethical considerations

The study received ethical approval from the university's Human Research and Ethics Committee (approval no. deidentified). Students were advised on the participant information sheet, provided through email, that completing surveys was completely voluntary and anonymous, and that participation or non-participation had no impact on their studies within the degree. Once surveys were submitted, students were unable to withdraw as no identifying information was collected and individual responses were not identifiable.

Participation in the study involved post-simulation completion of the SSSCLS and SDS, which were distributed and collected by a staff member who was not involved in the research. Surveys were completed immediately following the simulation debriefing. Printed surveys reinforced anonymous survey completion/non-completion. The completion of surveys implied consent to participate.

Participants and setting

All nursing students enrolled in the 3rd year Palliative Care Nursing elective unit in the first semester of their final year of study in 2021 ($n = 51$) were invited to participate. Only students enrolled in this unit of study and who participated in the non-compulsory simulation days ($n = 31$) were eligible to participate in the study.

Four palliative care simulation days were conducted over 4 weeks at a large metropolitan university in Australia. Simulations consisted of 2 different clinical scenarios, each delivered twice. The first simulation was set in an outpatient palliative care clinic (delivered during the first 2 simulation weeks), and the second simulation was an inpatient hospital scenario (delivered during the last 2 simulation weeks). Students self-enrolled into simulation sessions according to their own availability. Repeating simulation sessions gave students some flexibility in attending simulation days.

Intervention

Palliative care simulation was introduced into the palliative care elective unit in 2020 with the aim of improving student experiences and preparation for providing palliative and end-of-life care in clinical practice. The simulation intervention was underpinned by the Palliative Care Curriculum for Undergraduates (PCC4U) teaching resources and clinical case scenarios. Two clinical scenarios were delivered as SP simulation during 4 face-to-face workshops. The first scenario was female in her thirties with advanced breast cancer attending an outpatient palliative care clinic for advice on symptom management and psychosocial support. The second case was a male in his sixties with advanced metastatic bowel cancer in the hospital for pain and symptom management (more information can be found at <https://pcc4u.org.au>).

The NLN Jeffries Simulation Theoretical Framework (Jeffries and Rogers 2012) was used to guide the development and delivery of the simulation experiences. Simulation design characteristics included the simulation fidelity using trained SPs, scenario development using the evidence-based resources from PCC4U, pilot testing of simulation scenarios prior to the study, clearly identified learning objectives, and student support during the simulation experience (Jeffries et al. 2015).

Pre-simulation, students had access to the simulation scenarios, pre-simulation readings, and the relevant PCC4U modules and videos, this is consistent with the International Nursing Association for Clinical Simulation and Learning (INACSL) best practice standards for simulation (INACSL Standards Committee, McDermott et al. 2021).

On the simulation days and prior to the simulation, students attended a lecture and a tutorial on the assessment and management of palliative care symptoms and communication in preparation for the application of knowledge during the simulation experience.

The simulation experience consisted of a 10-min pre-brief to orientate students to the clinical simulation space, reassure psychological safety by establishing a safe and non-judgmental environment within the simulation experience, and reaffirm the simulation learning objectives.

Students completed simulations in groups of up to 8 that consisted of 4 active participants and up to 4 observers. The simulations began with a verbal handover of the patient, followed by the clinical scenario, which lasted approximately 20 min.

Simulations were followed by a 20–25 min simulation debriefing led by the simulation facilitator, an experienced nursing educator with training in simulation and extensive experience with palliative care in clinical practice, using a standardized approach to simulation debriefing that included an adaptation of the Plus-Delta model, as aligned with the INACSL best practice standards for debriefing (INACSL Standards Committee, Decker et al. 2021). All participants participated in simulation debriefing.

Instruments

The instrument consisted of 2 surveys: the SSSCLS and the SDS. The SSSCLS was used to measure student satisfaction and self-confidence in learning through simulation. This 13-item instrument using a 5-point Likert scale (1 = strongly disagree to 5 = strongly agree) is designed to measure student satisfaction with the simulation activity (5 items) and self-confidence in learning (8 items). The original scale had a Cronbach's α of 0.94 for student satisfaction and 0.87 for student self-confidence (NLN 2019). In this study, Cronbach's α was 0.92 for student satisfaction and

0.86 for self-confidence. The SDS, a 20-item instrument using a 5-point scale (1 = strongly disagree to 5 = strongly agree), was designed to evaluate 5 design features of the simulations. The features rated are objectives/information (5 items), student support (4 items), problem-solving (5 items), guided reflection or feedback (4 items), and fidelity (2 items). The original scale had a Cronbach's α of 0.92 for the presence of design features and 0.96 for the importance of features (Jeffries 2012). In this study, Cronbach's α was 0.98 for the design feature element and 0.97 for importance.

Data analysis

Post-intervention SSSCLS and SDS scores were loaded into Excel and Statistical Package for the Social Sciences (SPSS) Version 26 to generate survey datasets. Post-test SSSCLS and SDS data were analyzed using descriptive statistics (mean; standard deviation [SD]). Participant characteristics are not included in the analysis, as this was perceived to potentially impact participant confidentiality (Rosenzweig et al. 2008).

Results

A total of $n = 57$ post-simulation surveys were completed (simulation 1, $n = 28/31$; simulation 2, $n = 29/31$), with a response rate of 90.32% and 93.54% (simulation 1 and simulation 2, respectively). The descriptive statistics for the SSSCLS are presented in Table 1. Due to the nature of this study being a descriptive post-intervention study and lacking a control group, it is difficult to assess if the high evaluative scores were directly related to the simulation experience, and thus the results should be interpreted with caution.

We hypothesized that using SPs in palliative care simulation would lead to high levels of satisfaction with the simulation method and high levels of self-confidence in mastering palliative care skills and managing difficult end-of-life conversations, and our results support this hypothesis. The results from the SSSCLS and SDS show that overall, students had a high level of satisfaction and self-confidence in caring for a simulated palliative patient. Results also indicate that both the simulation and the teaching methods used during the simulated learning experience were also very positive.

In simulation 1, of note are the high mean scores (greater than 4 in the Likert scale 1–5) for all but one question, “it is the instructor's responsibility to tell me what I need to learn of the simulation activity” (3.88 ± 1.15). The highest score (5.00 ± 0.00) was for the question “I enjoyed how my instructor taught the simulation.”

In simulation 2, there was a similar pattern of high mean scores. The highest score in this group was for the item “it is my responsibility as the student to learn what I need to know from this simulation activity” (4.64 ± 0.49). The item “it is the instructor's responsibility to tell me what I need to learn of the simulation activity” in this group also received the lowest score (3.79 ± 1.25).

The students in this study reported the simulation design elements (SDS) positively. Three items had equally high scores for simulation 1. These were “independent problem solving was facilitated” (4.93 ± 0.47), “there was an opportunity after the simulation to obtain guidance/feedback in order to build knowledge to another level” (4.93 ± 0.47), and “the scenario resembled a real-life situation” (4.93 ± 0.27). The lowest score for simulation 1 was for the item “the simulation provided me an opportunity to goal set for my patient” (4.50 ± 0.65).

Table 1. Student Satisfaction and Self-Confidence in Learning Scale

	Simulation 1 ($n = 28$)	Simulation 2 ($n = 29$)
Satisfaction with current learning		
1. The teaching methods used in this simulation were helpful and effective.	4.88 (0.33)	4.43 (0.51)
2. The simulation provided me with a variety of learning materials and activities to promote my learning.	4.59 (0.62)	4.07 (0.73)
3. I enjoyed how my instructor taught the simulation.	5.00 (0.00)	4.50 (0.52)
4. The teaching materials used in this simulation were motivating and helped me to learn.	4.82 (0.39)	4.36 (0.49)
5. The way my instructor/s taught the simulation was suitable to the way I learn.	4.76 (0.56)	4.55 (0.52)
Self-confidence in learning		
6. I am confident that I am mastering the content of the simulation activity that my instructors presented to me.	4.18 (0.53)	4.07 (0.62)
7. I am confident that this simulation covered critical content necessary for the mastery of the curriculum.	4.64 (0.49)	4.29 (0.47)
8. I am confident that I am developing the skills and obtaining the required knowledge from this simulation to perform the necessary tasks in a clinical setting.	4.41 (0.62)	4.21 (0.43)
9. My instructors used helpful resources to teach the simulation.	4.59 (0.62)	4.36 (0.63)
10. It is my responsibility as the student to learn what I need to know from this simulation activity.	4.65 (0.49)	4.64 (0.49)
11. I know how to get help when I do not understand the concepts covered in the simulation.	4.71 (0.47)	4.29 (0.47)
12. I know how to use simulation activities to learn critical aspects of these skills.	4.53 (0.51)	4.07 (0.47)
13. It is the instructor's responsibility to tell me what I need to learn of the simulation activity content during class time.	3.88 (1.15)	3.79 (1.25)

For simulation 2, 2 items scored equally highly, “the scenario resembled a real-life situation” (4.50 ± 1.09) and “real life factors, situations and variables were built into the simulation scenario” (4.50 ± 1.09). The lowest score was for item “there was enough information provided at the beginning of the simulation to provide direction and encouragement” (3.75 ± 1.18).

In terms of items of importance, “feedback provided was constructive” was rated the highest in both simulations. Simulation 1 had a slightly higher mean (4.86 , $SD \pm 0.36$) compared with simulation 2 (4.71 ± 0.61). Of lowest importance was “my need for help was recognised” (4.33 ± 0.82) for simulation 1 and “independent problem solving was facilitated” (4.14 ± 0.66) for simulation 2. The findings of the SDS portion of the survey are presented in Table 2.

Table 2. Simulation Design Scale

	Simulation 1 (n = 28)		Simulation 2 (n = 29)	
	Design elements Mean (SD)	Importance Mean (SD)	Design elements Mean (SD)	Importance Mean (SD)
Objectives and information				
1. There was enough information provided at the beginning of the simulation to provide direction and encouragement.	4.59 (0.62)	4.59 (0.62)	3.75 (1.18)	4.38 (0.81)
2. I clearly understood the purpose and objectives of the simulation.	4.76 (0.56)	4.59 (0.62)	4.13 (1.02)	4.50 (0.73)
3. The simulation provided enough information in a clear matter for me to problem-solve the situation.	4.65 (0.5)	4.83 (0.40)	4.31 (1.01)	4.56 (0.51)
4. There was enough information provided to me during the simulation.	4.70 (0.47)	4.76 (0.44)	4.27 (1.03)	4.44 (0.73)
5. The cues were appropriate and geared to promote my understanding	4.88 (0.33)	4.71 (0.47)	4.19 (0.75)	4.50 (0.52)
Support				
6. Support was offered in a timely manner.	4.76 (0.56)	4.50 (0.73)	4.31 (0.87)	4.38 (0.81)
7. My need for help was recognized.	4.69 (1.01)	4.33 (0.82)	4.25 (1.24)	4.27 (0.70)
8. I felt supported by the teacher's assistance during the simulation.	4.75 (0.68)	4.50 (0.90)	4.19 (0.98)	4.44 (0.63)
9. I was supported in the learning process.	4.75 (0.45)	4.63 (0.62)	4.19 (0.98)	4.50 (0.63)
Problem-solving				
10. Independent problem-solving was facilitated.	4.93 (0.47)	4.50 (0.65)	3.93 (0.73)	4.14 (0.66)
11. I was encouraged to explore all possibilities of the simulation	4.71 (0.47)	4.36 (0.63)	4.21 (0.90)	4.43 (0.65)
12. The simulation was designed for my specific level of knowledge and skills.	4.71 (0.47)	4.58 (0.65)	4.00 (0.68)	4.43 (0.51)
13. The simulation allowed me the opportunity to prioritize nursing assessments and care.	4.64 (0.74)	4.58 (0.65)	3.93 (0.83)	4.43 (0.51)
14. The simulation provided me an opportunity to goal set for my patient.	4.50 (0.65)	4.43 (0.76)	4.07 (1.21)	4.36 (0.50)
Feedback/Guided Reflection				
15. Feedback provided was constructive.	4.78 (0.42)	4.86 (0.36)	4.43 (1.09)	4.71 (0.61)
16. Feedback was provided in a timely manner.	4.86 (0.36)	4.64 (0.63)	4.43 (1.09)	4.64 (0.50)
17. The simulation allowed me to analyse my own behaviour and actions.	4.64 (0.63)	4.72 (0.47)	4.21 (1.19)	4.50 (0.65)
18. There was an opportunity after the simulation to obtain guidance/feedback from the teacher in order to build knowledge to another level.	4.93 (0.47)	4.77 (0.60)	4.21 (1.05)	4.50 (0.65)
Fidelity (Realism)				
19. The scenario resembled a real-life situation.	4.93 (0.27)	4.57 (0.65)	4.50 (1.09)	4.71 (0.61)
20. Real life factors, situations, and variables were built into the simulation scenario.	4.86 (0.36)	4.43 (0.76)	4.50 (1.09)	4.71 (0.61)

Discussion

The simulation scenarios used in this research were based on the PCC4U resources. The PCC4U project, funded by the Australian Government Department of Health, aims to improve the confidence and skills of the Australian health-care workforce to meet the population's palliative care needs (PCC4U 2022). The resources developed by PCC4U are evidence-based and provide a range of clinical scenarios to support learning (PCC4U 2022). During this research, we used 2 clinical scenarios to support student's learning, with a focus on communication when approaching difficult

conversations at the end of life. The simulation used SPs playing the roles of Michelle, who was based in an outpatient clinic setting, and William, who was an inpatient. The use of SPs gave students the opportunity to undertake difficult conversations in the context of both settings and provided students the chance to practice vital communication skills required by all health-care professionals (Paal et al. 2019). These findings support other research using SPs that have shown improvement in communication skills in the context of palliative care or end-of-life care communication (Bloomfield et al. 2015; Jeffers et al. 2022).

Both groups were particularly satisfied with the instructor's teaching method. Facilitation of simulation requires a skilled instructor who has the education, skill, and ability to guide, support, and seek out ways to assist participants in achieving expected outcomes (INACSL Standards Committee, Persico et al. 2021). In this study, the instructor had content knowledge of palliative care in addition to simulation pedagogy and therefore could support students to achieve the required learning outcomes. Both groups also indicated overall high levels of self-confidence in relation to mastering palliative care skills and the palliative care content of the simulation activities, and that they were developing the skills and obtaining the knowledge required for clinical practice. This is a positive finding, while simulation has been found to increase nursing students' satisfaction and self-confidence (Cura et al. 2020; de Oliveira Costa et al. 2020; Dincer and Ataman 2020; Olaussen et al. 2020; Zapko et al. 2018; Zulkosky et al. 2021), to date there is limited research on student satisfaction or improved self-confidence with palliative care or end-of-life care simulation.

The overall trend of higher scores in student's satisfaction and self-confidence was evident for simulation 1 where students contributed to addressing both psychological and physical needs of the patient and were challenged and supported to address difficult conversations as the scenario unfolded. This scenario, based in an outpatient setting, focused on communication and symptom management compared with a hospital-based pain management scenario in simulation 2. A possible explanation for this is that simulations based in primary care-type contexts have been found to be well liked by students, in part due to the scenarios not being as time-critical and dependent on checklists as more acute scenarios (Lunde et al. 2021).

In this study, the results of the SDS indicated that perceptions of simulation design elements for both simulations were high overall. Both simulations used an SP, and consistent with other research, the use of a simulation methodology, which is the most realistic, enables students to feel as though they are learning in a real clinical context and offers an active learning experience (Escribano et al. 2021; Rosenzweig et al. 2008; Tamaki et al. 2019). Although SPs were used for both simulations, the scenarios were different, and this may be reflected in the slightly higher fidelity and realism score for the outpatient scenario, which is possibly easier to recreate than a hospital scene. These positive findings are of value in confirming that SPs are useful in creating a realistic clinical situation to prepare students for real-world practice in both care settings. It also provides justification for the expense in an educational era, which may have limited funding for activities such as simulation (So et al. 2019).

In terms of the importance of items on the SDS, receiving feedback that was constructive rated the highest for both simulations. Students may not have had exposure to palliative care during their clinical placements, therefore, they value being guided through purposeful discussion of their actions during the simulation through debriefing (Thidemann and Söderhamn 2013). By increasing confidence and competence in end-of-life care among students, this may, in turn, be translated into clinical practice and has the potential to improve end-of-life care for patients and their families.

Both clinical scenarios focused on patients with different cancers and of different age groups. Michelle, a young woman with a young family undergoing treatment for breast cancer, and William, aged in his early 60s with metastatic bowel cancer, are clinical scenarios rarely seen in the sparse palliative care simulation literature. Recent palliative care simulation literature using SPs and

approaching difficult conversations have used non-cancer diagnoses such as heart failure and Chronic Obstructive Pulmonary Disease (COPD) (Jeffers et al. 2022). The only research of this kind that used cancer diagnosis also used metastatic bowel cancer diagnosis of a middle-aged male (Bloomfield et al. 2015). To date, this study is the first of its kind to use a scenario with a younger female patient with advanced breast cancer. This study gave participants the opportunity to apply clinical skills to patient symptoms beyond non-cancer diagnoses and of varying age groups.

Of the available studies assessing self-confidence and satisfaction with palliative or end-of-life care simulation, there is very limited research using validated instruments such as SSSCLS. Alconero-Camarero et al. (2018), using low-fidelity simulation, in their descriptive, observational, and correlational study used 3 scales (SSSCLS Spanish version; Questionnaire for dealing with stress; Trait Meta-Mood Scale-24) with 2nd year nursing students ($n = 74$). Fabro et al. (2014) used the SSSCLS in conjunction with the Educational Practices Questionnaire to evaluate student learning ($n = 21$) from palliative care simulation using a high-fidelity manikin and the student playing the role of the patient's daughter.

While these studies add to the limited available research on palliative care simulation, the simulation modalities used are either low- or high-fidelity simulation. Our study is the first that we are aware of that has used SPs and been evaluated using the SSSCLS. There were no studies found on palliative care or end-of-life care simulation using the SDS. To our knowledge, this is the first study to use both the NLN-developed SSSCLS and SDS for assessing palliative care or end-of-life care simulation.

Recommendations for research and practice

The INACSL 2020–2023 simulation research priorities provide targeted areas of focus to address known gaps in the literature on simulation and to address the INACSL vision that simulation and innovation transform lives (Franklin and Luctkar-Flude 2020). Among these research priorities are interprofessional simulation, psychomotor skill retention, and transition to practice (Franklin and Luctkar-Flude 2020).

To address these key research priorities and gaps in the literature, from this research, it is recommended that palliative care and end-of-life care interprofessional simulation, which include undergraduate health students from a range of professions, focusing on patient outcomes would be of benefit to further develop skills surrounding communication at the end of life. In addition, longitudinal studies that evaluate specific time points post-simulation along a time continuum may give more insight into the immediate and long-term effects that palliative care and end-of-life care simulation have on nursing students' skill retention and on new graduates transitioning to clinical practice.

While palliative care and end-of-life care simulation are developing, there are few studies that explore the impact that palliative care and end-of-life care simulation have on clinical practice for both undergraduate nursing students and new graduates. Furthermore, there are few studies on SP simulation and even fewer with a focus on communication at the end of life. This study on SP simulation focusing on difficult conversations at the end of life is the first of its kind, originating from Australia. The results of our study showed that nursing students reported high levels of self-confidence and satisfaction with this novel and innovative mode of simulation. A recommendation from our research is a change in simulation practice with the adoption of SPs in the roles of patients, family members, and caregivers.

Limitations

The small convenience sample of nursing students enrolled in an elective unit of study from a single site restricts the generalizability of the results. The post-intervention only study design limits the interpretation of results. While this was a restriction imposed by factors outside the authors' control, future studies will aim to include pre- and post-intervention surveys response to allow for comparison of results. In addition, a mixed-method approach that provides the opportunity for qualitative responses would also have strengthened this research and will be considered for future research on this topic. While this study included students who actively participated in the simulation scenarios as well as observers, there was no opportunity to rotate roles so each student could actively participate in the simulations, nor did this research involve comparison between active participants and observers. Future research is planned to include a comparison of roles between participant and observer to determine differences in participants' experiences and whether there is a difference in knowledge retention between the 2 groups.

Conclusion

Nursing students have limited exposure to palliative care and end-of-life care experiences during their clinical placements, and students report feeling unprepared to provide quality end-of-life care in clinical practice. Little is known of student's self-confidence and satisfaction with palliative and end-of-life care simulation. This study is contributing to the limited available research on palliative care simulation and shows promising results in the development of self-confidence and satisfaction in learning about palliative and end-of-life care. SP simulation focusing on difficult conversations has been found to assist in developing students' communication skills with patients and families at the end of life.

Acknowledgments. The authors wish to thank the nursing students who participated in this research project and would like to acknowledge David Klug and Kylie Ash from PCC4U.

Funding. This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

Competing interests. The authors report no competing interests.

References

- Alconero-Camarero A, Sarabia-Cobo C, Gonzalez-Gomez S, *et al.* (2018) Nursing students' emotional intelligence, coping styles and learning satisfaction in clinically simulated palliative care scenarios: An observational study. *Nurse Education Today* **61**, 94–100. doi:10.1016/j.nedt.2017.11.013
- Bautista C and Bartos S (2021) Innovative classroom upgrade: Simulating an intensive care unit environment in an undergraduate academic classroom. *Nursing Education Perspectives* **42**(4), 250–251. doi:10.1097/01.NEP.0000000000000609
- Bloomfield J, O'Neill B and Gillett K (2015) Enhancing student communication during end-of-life care: A pilot study. *Palliative and Supportive Care* **13**, 1651–1661. doi:10.1017/S147895151500022X
- Cannity K, Banerjee S, Hichenberg S, *et al.* (2021) Acceptability and efficacy of a communication skills training for nursing students: Building empathy and discussing complex situations. *Nurse Education in Practice* **50**, 102928. doi:10.1016/j.nepr.2020.102928
- Cheong C, Ha N, Tan L, *et al.* (2019) Attitudes towards the dying and death anxiety in acute care nurses – can a workshop make any difference? A mixed-methods evaluation. *Palliative and Supportive Care* **18**, 164–169. doi:10.1017/S1478951519000531
- Coates J (2021) The effectiveness of a simulation program to enhance readiness to engage in difficult conversations in clinical practice. *Dimensions of Critical Care Nursing* **40**(5), 275–279. doi:10.1097/DCC.0000000000000489
- Cura U, Kocatepe V, Yildirim D, *et al.* (2020) Examining knowledge, skill, stress, satisfaction, and self-confidence levels of nursing students in three different simulation modalities. *Asian Nursing Research* **14**, 158–164. doi:10.1016/j.anr.2020.07.001
- International Nursing Association for Clinical Simulation and Learning Standards Committee, Decker S, Alinier G, Crawford S, *et al.* (2021) Healthcare Simulation Standards of Best Practice™ the debriefing process. *Clinical Simulation in Nursing* **58**, 27–32. doi:10.1016/j.ecns.2021.08.011
- Demirtas A, Guvenc G, Aslan O, *et al.* (2021) Effectiveness of simulation-based cardiopulmonary resuscitation training programs on fourth-year nursing students. *Australasian Emergency Care* **24**, 4–10. doi:10.1016/j.auec.2020.08.005
- de Oliveira Costa R, de Medeiros S, Dias Coutinho V, *et al.* (2020) Satisfaction and self-confidence in the learning of nursing students: Randomised clinical trial. *Escola Anna Nery* **24**(1), e20190094. doi:10.1590/2177-9465-EAN-2019-0094
- Dincer B and Ataman H (2020) The effect of high reality simulation on nursing students' knowledge, satisfaction, and self-confidence levels in learning. *International Journal of Caring Sciences* **13**(2), 894–900.
- Escribano S, Cabanero-Martinez M, Fernandez-Alcantara M, *et al.* (2021) Efficacy of a standardised patient simulation programme for chronicity and end-of-life care training in undergraduate nursing students. *International Journal of Environmental Research & Public Health* **18**, 11673. doi:10.3390/ijerph182111673
- Fabro K, Schaffer M and Scharton J (2014) The development, implementation and evaluation of an end-of-life simulation experience for baccalaureate nursing students. *Nursing Education Perspectives* **35**(1), 19–26. doi:10.5480/11-593.1
- Fink M, Linnard-Palmer L, Ganley B, *et al.* (2014) Evaluating the use of standardised patients in teaching spiritual care at the end-of-life. *Clinical Simulation in Nursing* **10**, 559–566. doi:10.1016/j.ecns.2014.09.003
- Franklin A and Luctkar-Flude M (2020) 2020 to 2023 Research priorities advance INACSL core values (editorial). *Clinical Simulation in Nursing* **47**, 82–83. doi:10.1016/j.ecns.2020.09.001
- Garrino L, Contratto C, Massariello P, *et al.* (2017) Caring for dying patient and their families: The lived experiences of nursing students in Italy. *Journal of Palliative Care* **32**(3–4), 127–133. doi:10.1177/0825859717745169
- Gillan P, Jeong S and van der Riet P (2014b) End of life care simulation: A review of the literature. *Nurse Education Today* **34**(5), 766–774. doi:10.1016/j.nedt.2013.10.005
- Gillan P, Jeong S and van der Riet P (2021a) Undergraduate nursing students' transformative learning through disorientating dilemmas associated with end-of-life care simulation: A narrative inquiry study. *Nurse Education in Practice* **55**, 103174. doi:10.1016/j.nepr.2021.103174
- Gillan P, Jeong S and van der Riet P (2021b) Embodied good deaths and disembodied bad deaths: Undergraduate nursing students' narratives of experience. *Nurse Education Today* **97**, 1–5. doi:10.1016/j.nedt.2020.104674
- Gillan P, van der Riet P and Jeong S (2013) The experience of end of life care simulation at a rural Australian university. *Nurse Education Today* **33**, 1435–1439. doi:10.1016/j.nedt.2012.11.015
- Gillan P, van der Riet P and Jeong S (2014a) End of life care education, past and present: A review of the literature. *Nurse Education Today* **34**, 331–342. doi:10.1016/j.nedt.2013.06.009
- Gillan P, van der Riet P and Jeong S (2016) Australian nursing students' stories of end-of-life care simulation. *Nursing & Health Sciences* **18**(1), 64–69. doi:10.1111/nhs.12233
- Gillett K, O'Neill B and Bloomfield J (2016) Factors influencing the development of end-of-life communications skills: A focus group study of nursing and medical students. *Nurse Education Today* **36**, 395–400. doi:10.1016/j.nedt.2015.10.015
- Guerrero JG, Ahmed SA and Attallah DM (2022) The acquired critical thinking skills, satisfaction and self confidence of nursing students and staff nurses

- through high-fidelity simulation experience. *Clinical Simulation in Nursing* **64**, 24–30. doi:10.1016/j.ecns.2021.11.008
- Halliday D, Dorney P and Landis J** (2022) Difficult conversations in palliative care: Innovations in learning. *Nursing Education Perspectives* **43**(3), 201–202. doi:10.1097/01.NEP.0000000000000831
- Hoang C, Copnell B, Lawrence K, et al.** (2022) Undergraduate nursing education and end-of-life simulation: A scoping review. *Clinical Simulation in Nursing* **70**, 56–73. doi:10.1016/j.ecns.2022.06.009
- Jeffers S, Lippe M, Justice A, et al.** (2022) Nursing student perceptions of end-of-life communication competence. *Journal of Hospice & Palliative Nursing* **24**(3), 199–205. doi:10.1097/NJH.0000000000000849
- Jeffries P, Rodgers B and Adamson K** (2015) NLN Jeffries simulation theory: Brief narrative description. *Nursing Education Perspectives* **36**(5), 292–293.
- Jeffries P and Rogers K** (2012) Theoretical framework for simulation design. In Jeffries P (ed), *Simulation in Nursing Education: from Conceptualization to Evaluation*, 2nd edn. New York: National League for Nursing, 266.
- Kantor D and Stadelman A** (2020) Preparing nursing graduates to engage in difficult conversations with patients. *American Journal of Nursing* **120**(8), 11. doi:10.1097/01.NAJ.0000694496.25141.a1
- Kirkpatrick A, Cantrell M and Smeltzer S** (2017) Palliative care simulations in undergraduate nursing education: An integrative review. *Clinical Simulation in Nursing* **13**(9), 414–431. doi:10.1016/j.ecns.2017.04.009
- Kunkel C, Kopp W and Hanson M** (2016) A matter of life and death: End-of-life simulation to develop confidence in nursing students. *Nursing Education Perspectives* **37**(5), 285–286. doi:10.1097/01.NEP.0000000000000029
- Kunst E, Henderson A and Johnston A** (2018) A scoping review of the use and contribution of simulation in Australian undergraduate nurse education. *Clinical Simulation in Nursing* **19**, 17–29. doi:10.1016/j.ecns.2018.03.003
- Lunde L, Moen A, Jakobsen R, et al.** (2021) Exploring healthcare students' interprofessional teamwork in primary care simulation scenarios: Collaboration to create a shared treatment plan. *BMC Medical Education* **21**(1), 1–14. doi:10.1186/s12909-021-02852-z
- International Nursing Association for Clinical Simulation and Learning Standards Committee, McDermott D, Ludlow J, Horsley E, et al.** (2021) Healthcare Simulation Standards of Best Practice™ prebriefing: Preparation and briefing. *Clinical Simulation in Nursing* **58**, 9–13. doi:10.1016/j.ecns.2021.08.008
- National League for Nursing** (2005a) Student Satisfaction and Self-Confidence in Learning Scale®. <https://www.nln.org/education/teaching-resources/tools-and-instruments> (accessed 6 January 2023).
- National League for Nursing** (2005b) Simulation Design Scale® (Student Version). <https://www.nln.org/education/teaching-resources/tools-and-instruments> (accessed 6 January 2023).
- National League for Nursing** (2019) Descriptions of available instruments. <https://www.nln.org/education/teaching-resources/tools-and-instruments> (accessed 6 January 2023).
- Olaussen C, Heggdal K and Tvedt C** (2020) Elements in scenario-based simulation associated with nursing students' self-confidence and satisfaction: A cross-sectional study. *Nursing Open* **7**(1), 170–179. doi:10.1002/nop.2.375
- Paal P, Brandstotter C, Lorenzi S, et al.** (2019) Postgraduate palliative care education for all healthcare providers in Europe: Results from an EAPC survey. *Palliative and Supportive Care* **17**, 495–506. doi:10.1017/S1478951518000986
- Palliative Care Curriculum for Undergraduates** (2022) About PCC4U. Brisbane: QUT. <https://pcc4u.org.au/about/> (accessed 25 October 2022).
- International Nursing Association for Clinical Simulation and Learning Standards Committee, Persico L, Belle A, DiGregorio H, et al.** (2021) Healthcare Simulation Standards of Best Practice™ facilitation. *Clinical Simulation in Nursing* **58**, 22–26. doi:10.1016/j.ecns.2021.08.010
- Risteovski E, Leach M, Bolton E, et al.** (2022) Rural community-based nurses' self-reported knowledge and skills in the provision of psychosocial care to palliative and end-of-life clients and carers. *Palliative and Supportive Care* **20**, 823–831. doi:10.1017/S1478951521001644
- Rosenzweig M, Hravnak M, Magdic K, et al.** (2008) Patient communication simulation laboratory for students in an acute care nurse practitioner program. *American Journal of Critical Care* **17**(4), 364–372. doi:10.4037/ajcc2008.17.4.364
- Sarabia-Cobo C, Alconero-Camarero A, Lavin-Alconero L, et al.** (2016) Assessment of a learning intervention in palliative care based on clinical simulations for nursing students. *Nurse Education Today* **45**, 219–224. doi:10.1016/j.nedt.2016.08.014
- Sarmasoglu S, Dinc L and Elcin M** (2016) Using standardised patients in nursing education: Effects on students' psychomotor skill development. *Nurse Educator* **41**(2), E1–E5. doi:10.1097/NNE.0000000000000188
- So H, Chen P, Wong G, et al.** (2019) Simulation in medical education. *Journal of the Royal College of Physicians of Edinburgh* **49**(1), 52–57. doi:10.4997/JRCPE.2019.112
- Tamaki T, Inumaru A, Yokoi Y, et al.** (2019) The effectiveness of end-of-life care simulation in undergraduate nursing education: A randomised controlled trial. *Nurse Education Today* **76**, 1–7. doi:10.1016/j.nedt.2019.01.005
- Thidemann I and Söderhamn O** (2013) High-fidelity simulation among bachelor students in simulation groups and use of different roles. *Nurse Education Today* **33**(12), 1599–1604. doi:10.1016/j.nedt.2012.12.004
- Zapko K, Ferranto M, Blasiman R, et al.** (2018) Evaluating best educational practices, student satisfaction, and self-confidence in simulation: A descriptive study. *Nurse Education Today* **60**, 28–34. doi:10.1016/j.nedt.2017.09.006
- Zulkosky K, Minchhoff D, Dommel L, et al.** (2021) Effect of repeating simulation scenarios on student knowledge, performance, satisfaction and self-confidence. *Clinical Simulation in Nursing* **55**, 27–36. doi:10.1016/j.ecns.2021.03.004