

In this paper we will present investigation of therapeutic factors by Yalom's questionnaire in different phases of group psychodynamic oriented psychotherapy with psychotic patients, respectively in patients who have participated in group psychotherapy for less than six months and in those that have participated in group psychotherapy for more than five years. We expect the results of this study to contribute to better understanding of group process with psychotics and to the development of guidelines for the definition of more specific therapeutic strategies.

P0201

Early intervention in psychosis: A comprehensive hospital outpatient programme for the first episodes during the critical period of illness

B. Restek-Petrovic, M. Mihanovic, M. Grah, A. Bogovic, S. Molnar, V. Grosic, S. Kezic. *Psychiatric hospital, Zagreb, Croatia*

Early intervention in first episodes of psychotic disorders is very important in therapeutic treatment because first three to five years is considered to be critical period in the course of illness (Birchwood, 2002), because it significantly determinates patient's future cognitive, social and emotional functioning and also reflects on their family functioning.

Psychiatric hospital «Sveti Ivan» in Zagreb has started the programme which is consisted of psychoeducation for patients and their family members, and psychodynamic group psychotherapy for patients and group psychotherapy for family members.

In order to improve patient's follow-up we also investigated cognitive functioning in different phases of treatment, object relations, separation and individuation and empathy.

Considering that stigma related to mental disease leads to decrease of self-esteem, loneliness and reduction of quality of life, we applied self-esteem questionnaire, loneliness scale, SF 36 and perceived social support questionnaire.

In this paper partial preliminary results of our programme with group psychotherapy after first six months-period by using Scale of emotional empathy, Rosenberg Self-Esteem Scale and UCLA Loneliness Scale (short version) will be presented.

P0202

Evaluation of basic symptoms in schizophrenia in Iran, Ardabil

P. Molavi¹, E. Rezaei Ghalechi². ¹ *Fatemi Hospital, Ardabil, Iran*
² *Islamic Azad Medical University, Ardabil, Iran*

Background: The objective of this research is determines the most frequent basic symptom & the most frequent type of schizophrenia in Ardabil hospitals patients.& providing a new classification of basic symptoms in these patients, so that psychiatrics of Ardabil city can use the results of this study to achieve an accurate & also rapid diagnosis of schizophrenia.

Methods: This study is a descriptive – analytic one. A sample of 100 patients of both sexes from 17 up to 40 years old was selected. To study the basic symptoms of schizophrenia in them, we used a 44 item questionnaire which was provided through a combination of PANSS & Kitamura questionnaires. To screen the patients with schizophrenia, we used MMPI test. The findings was analyzed by descriptive statistics & through the computer software SPSS. Also a PCA method of factor analysis was used to the hypotheses.

Results: single (62%=49 patients), unemployed (52%=52), with no education (31%=31), negative family history (91%=91), with frequent admission (47%=47). The most frequent basic symptom in

these patients was social withdrawal was at second place. the most frequent type was paranoid schizophrenia (44%=44).

Conclusion: The findings have a concordance with results of the last studies about the basic symptoms of schizophrenia & also with DSM-IV & Bleuler& Kraepelin symptoms all of these systems introduce the different fields of a single diagnostic structure rather than antagonist ones.

Anyway, two important factors that were recognized were cognitive disorder. This study suggests that if we recognized emotional symptoms

P0203

Follow and study basic symptoms & problems in schizophrenia patients in north west of Iran

P. Molavi¹, H. Ghamari Ghivi², E. Rezaei Ghale Chi³, R. Arab¹.
¹ *Medicine University, Ardebil, Iran* ² *Mohaghegh University, Ardebil, Iran* ³ *Azad Medicine University, Ardebil, Iran*

Background: The objective of this research is studying the basic symptoms in schizophrenic patients and determines the most frequent basic symptom and the most frequent type of schizophrenia in them. Also, providing a new classification of basic symptoms

Methods: This study is a descriptive – analytic one. A sample of 100 patients of both sexes(age: 17-40)was selected. To study the basic symptoms of schizophrenia in them, we used a 44 item questionnaire which was provided through a combination of PANSS (Emsley, 2003) and Kitamura questionnaires. we used MMPI test. The findings were analyzed by descriptive statistics and through the computer software SPSS. Also a PCA method of factor analysis was used to the hypotheses.

Results: single (62%=49 patients), unemployed (52%=52 patients), with no education (31%=31 patients), negative family history (91%=91 patients), with frequent admission (47%=47 patients). The most frequent basic symptom in these patients was attention deficit, and social withdrawal was at second place. Also, the highest frequent type of schizophrenia was paranoid schizophrenia (44%=44 patients).

Conclusion: This study's findings are similar to with the results of the last studies about the basic symptoms of schizophrenia and also with DSM-IV and Bleulerand Kraepelin symptoms . it seems that all of these systems introduce the different fields of a single diagnostic structure rather than antagonist ones.

However, two important recognized factors were cognitive disorder and relationship disorder. This study suggests that if we recognized emotional symptoms and restlessness, we should think of other psychotic disorders.

P0204

The incidence of risk factors for metabolic syndrome in patients with mental illness

N. Robins, M. Selvaraj, J. Scaria. *Moorside Unit, Trafford General Hospital, Manchester, UK*

Patients with schizophrenia are twice as likely to have coronary heart disease and their overall life expectancy is reduced by approximately 10 years. Metabolic syndrome is a cluster of risk factors known to increase the risk of cardiovascular disease and diabetes. This study aims

to identify at-risk patients by offering a physical health screening to all patients on antipsychotics.

A total of 79 patients have been screened, with the majority of patients were being treated with an oral atypical antipsychotic (88.6%).

Most patients (81%) were found to be overweight (BMI \geq 25) and 58.2% of patients had a waist circumference above the recommended limit. 40.5% of patients had a blood pressure reading above 130/85.

Cholesterol testing found that 54.9% of patients had a total cholesterol $>$ 5.0mmol/l, 58.8% had an LDL of \geq 3mmol/l, 44.8% had a triglyceride level \geq 1.7mmol/l and 30.6% had a HDL below the recommended limit.

Random blood glucose (RBG) testing identified five patients with a RBG $>$ 11.1mmol/l which is indicative of diabetes. A further 17 patients (25%) were in the range 6–11mmol/l and therefore require further tests.

Overall, 38.7% of patients met the criteria for metabolic syndrome. This is significantly higher than that found in the general population with European studies suggesting the prevalence to be 12–25%.

These findings suggest that a significant number of patients with serious mental illness are at risk of cardiovascular disease. Patients should be offered lifestyle advice, further tests and appropriate treatment. A follow-up study should address the impact of these interventions.

P0205

An audit of antipsychotic depot: Patients' perspective

N.M. Robins, M. Selvaraj, J. Scaria. *North Trafford Sector Moorside Unit, Trafford General Hospital, Manchester, UK*

The aim of the study was to assess patient perspectives of conventional depots and risperidone long-acting injection (RLAI). All adult patients receiving treatment with a depot were invited to complete a patient survey.

70 patients completed the survey, of which 20 were on RLAI. The mean duration of treatment with conventional depot was 12.1 years compared to 1.5 years for RLAI. Patients on conventional depot had an average of 0.58 admissions each over the past two years compared to 0.8 for RLAI patients. 72% of conventional depot patients had been relapse-free during the past two years compared to 45% of RLAI patients.

33.3% of patients on RLAI reported no side-effects compared to 10.2% on conventional depots. 77.6% of conventional depot patients reported that their treatment was effective, compared to 61.9% with RLAI. 61.2% of conventional patients reported that they were satisfied with their treatment compared to 47.6% with RLAI.

Although RLAI was generally better tolerated than conventional depots, more patients on conventional depots rated their treatment as effective, and satisfaction levels were also higher. However, other studies have found high patient satisfaction with RLAI, and a review of the literature² found that side-effects are one of the main drivers of patient satisfaction.

A limitation of this study is the differences between the two treatment groups in baseline demographics and this may have impacted on the results. Patient satisfaction with treatment is based on many variables and further research into this is needed.

P0206

Empirical evidence of residential skills therapy for schizophrenic patients: A controlled study including a 5-year follow-up

V. Roder, D.R. Mueller, U. Keppeler. *University Psychiatric Services Bern, Bern, Switzerland*

Background: Against the background of a paradigm shift in the locus of treatment from the hospital to the community, successful housing and re-establishing residential skills therefore are among the main preconditions for the independent living of schizophrenia outpatients. But so far only a few empirically evaluated treatment approaches exist, which specifically focus on improving these skills.

Method: During the last years, we developed a new manualised cognitive-behavioural group therapy program for residential skills (experimental group, EG).

In an international controlled study, the EG was compared to a conventional, unspecific therapy for social skills (control group, CG). EG and CG were conducted in groups consisting of 5–8 patients over a period of 6 months. Assessments were applied before and after therapy and at a 1-year follow-up. Additionally, relapse rates and rehabilitation status were assessed over 5 years after the beginning of therapy. Altogether 54 patients participated in this study (EG: N=26; CG: N=28).

Results: EG patients displayed a significantly higher frequency of change from a high structured in-patient setting to a lower structured outpatient setting and unassisted living compared to CG patients during 5 years of assessment (odds ratio=12.5). Moreover, lower relapse rates, significantly reduced psychopathology and higher treatment motivation of EG patients compared to controls were found. Both treatment groups significantly improved in cognitive and general social functioning.

Conclusions: Due to these results, the newly developed residential skills program could constitute an additional treatment option in the context of multimodal psychiatric rehabilitation.

P0207

Characteristics of patients at first contact for psychosis in Bologna

E. Rossi, D. Pecile, E. Mori, N. Cevenini, R. Michetti, D. Berardi. *Institute of Psychiatry 'P.Ottonello', Bologna, Italy*

Introduction: International literature has recently given great attention to risk factors and early detection of psychosis. According to this research line we have investigated the main characteristics of subjects at first contact for psychotic disorder with one of the three Community Mental Health Centers (CMHC) of Bologna Ovest.

Method: A retrospective study was conducted recruiting subjects with ICD-10 F2 diagnosis (schizophrenia, schizotypal disorder and delusional disorder), psychosis onset between 18–30 years and at first contact with a Bologna Ovest CMHC over a five-year period (January 2002–December 2006).

The identified sample was evaluated on socio-demographic characteristics, history of substance abuse, history of hospital admissions and pathways to CMHCs.

Results: The sample consists of 56 subjects. They are predominantly male (70%), unmarried (95%) and the most of them (79%) lives with their parents. Twenty-four patients (43%) have an eight year education, whereas 22 (39%) a thirteen year one. During the assessed period 24 (43%) subjects are unemployed and 15 (27%) are studying.

The median age of psychosis onset is 23,3 years and a concurrent substance abuse concerns 26 subjects (46%). Twenty-one (37,5%) patients present an history of hospital admissions.