

HOMOTOPY GROUPS OF COMPACT LIE GROUPS

E₆, E₇ AND E₈

HIDEYUKI KACHI

§ 1. Introduction

Let G be a simple, connected, compact and simply-connected Lie group. If k is the field with characteristic zero, then the algebra of cohomology $H^*(G; k)$ is the exterior algebra generated by the elements x_1, \dots, x_l of the odd dimension n_1, \dots, n_l ; the integer l is the rank of G and $n = \sum_{i=1}^l n_i$ is the dimension of G . Let X be the direct product of spheres of dimension n_1, \dots, n_l , then there exists a continuous map $f: G \rightarrow X$ which induces isomorphisms of $H^i(X; k)$ to $H^i(G; k)$ for all i (cf. [8]). From this we deduce by Serre's C -theory [8] that $f_*: \pi_i(G) \rightarrow \pi_i(X)$ are C -isomorphisms for all i , where C is the class of finite abelian groups. Therefore the rank of $\pi_q(G)$ is equal to the number of such i that n_i is equal to q , and particularly if q is even, then $\pi_q(G)$ is finite. It is a classical fact that $\pi_2(G) = 0$ and $\pi_3(G) = Z$.

According to Bott-Samelson [6];

$$\begin{aligned} \pi_i(E_6) &= 0 & \text{for } 4 \leq i \leq 8, & & \pi_9(E_6) &= Z, \\ \pi_i(E_7) &= 0 & \text{for } 4 \leq i \leq 10, & & \pi_{11}(E_7) &= Z, \\ \pi_i(E_8) &= 0 & \text{for } 4 \leq i \leq 14, & & \pi_{15}(E_8) &= Z. \end{aligned}$$

where E_6, E_7 and E_8 are compact exceptional Lie groups.

In this paper, using the killing method we compute the 2-components of homotopy group $\pi_j(G)$, where $G = E_6, E_7$ and E_8 . The results are stated as follows;

j	$4 \leq j \leq 14$	15	16	17	18	19	20	21	22	23
$\pi_j(E_8 : 2)$	0	Z	Z_2	Z_2	Z_8	0	0	Z_2	0	$Z + Z_2$

j	24	25	26	27	28
$\pi_j(E_8 : 2)$	$Z_2 + Z_2$	Z_2	0	Z	0

Received April 3, 1967.

j	$4 \leq j \leq 10$	11	12	13	14	15	16	17	18	19
$\pi_j(E_7 : 2)$	0	Z	Z_2	Z_2	0	Z	Z_2	Z_2	Z_4	$Z + Z_2$

j	20	21	22	23	24	25
$\pi_j(E_7 : 2)$	Z_2	Z_2	Z_4	$Z_2 + Z_2 + Z_2$	$Z_2 + Z_2 + Z_2$	$Z_2 + Z_2$

j	$4 \leq j \leq 8$	9	10	11	12	13	14	15	16	17
$\pi_j(E_6 : 2)$	0	Z	0	Z	Z_4	0	0	Z	0	$Z + Z_2$

j	18	19	20	21	22
$\pi_j(E_6 : 2)$	$Z_{16} + Z_2$	0	Z_8	0	0

All spaces that we consider in this paper are those which have the homotopy groups of finite type. Let G be such a space, then $\pi_i(G)$ is isomorphic to the direct sum of a free part F and the p -components of $\pi_i(G)$ for every prime p . We denote by $\pi_i(G : p)$ the direct sum of a certain subgroup F' of F and the p -component of $\pi_i(G)$, where the index $[F : F']$ is prime to p .

Given an exact sequence for such A, B and C

$$\dots \longrightarrow \pi_i(A) \longrightarrow \pi_i(B) \longrightarrow \pi_i(C) \longrightarrow \dots,$$

then we can form the following exact one in our case

$$\dots \longrightarrow \pi_i(A : p) \longrightarrow \pi_i(B : p) \longrightarrow \pi_i(C : p) \longrightarrow \dots.$$

The author is indebted to Professor H. Toda for his advice during the preparation of the paper.

§ 2. The cohomology of the 3-connective fibre spaces of E_6, E_7 and E_8 .

H. Cartan and J.P. Serre introduced a method to calculate the homotopy group in [7]. Let $K(\pi, n)$ be an Eilenberg-Mac-Lane space of type (π, n) .

THEOREM 2. 1. *Let X be an arcwise connected topological space, then there exists a sequence of $(n - 1)$ -connected spaces (X, n) ($n = 1, 2, \dots$, and $(X, 1) = X$) and continuous maps $f_n : (X, n + 1) \longrightarrow (X, n)$ such that:*

- (I) the triple $((X, n + 1), f_n, (X, n))$ is a fibre space with a fibre $K(\pi_n(X), n - 1)$.
- (II) there exists a fibre space X'_n over the base space $K(\pi_n(X), n)$, where X'_n and (X, n) are of the same homotopy type, such that the fibre is $(X, n + 1)$.

Hence $f_1 \circ f_2 \circ \dots \circ f_{n-1}$ defines the isomorphism of $\pi_i(X, n)$ to $\pi_i(X)$ for $i \geq n$.

LEMMA 2.2. Let X be a 2-connected topological space. Assume that X satisfies the following conditions,

- (1) $\pi_3(X)$ is isomorphic to an infinite cyclic group,
- (2) $H^*(X; Z_2) = A_0 \otimes A_1 \otimes \dots \otimes A_r \otimes B$

where x_3 is a generator of $H^3(X; Z_2) \approx Z_2, A_0 = Z_2[x_3]/(x_3)^{s_0}, A_i = Z_2[Sq^{2^i}Sq^{2^{i-1}} \dots Sq^{2^2}x_3]/(Sq^{2^i}Sq^{2^{i-1}} \dots Sq^{2^2}x_3)^{2^{s_i}}$ ($s_i \geq 1$) $1 \leq i \leq r$, and $Sq^{2^{r+1}}Sq^{2^r} \dots Sq^{2^2}x_3 = 0$, then

$$H^*((X, 4); Z_2) = Z_2[w] \otimes A(a_0, a_1, \dots, a_r) \otimes B'$$

where the $deg. a_i = (2^{i+1} + 1)(2^{s_i} - 1) + 2^{2^i}$, $deg. w = 2^{2^{r+1}}$, $A(a_0, a_1, \dots, a_r)$ indicates a submodule having a_0, \dots, a_r as a simple system of generators and B' is isomorphic to B by $(f_1 \circ f_2 \circ f_3)^* : H^*(X; Z_2) \rightarrow H^*((X, 4); Z_2)$.

Proof. From the above theorem, there exists a fibre space $((X, 4), f_1 \circ f_2 \circ f_3, X)$ with a fibre $K(Z, 2)$. Since $K(Z, 2)$ is the infinite dimensional complex projective space, its mod 2 cohomology structure is $H^*(Z, 2; Z_2) \approx Z_2[u]$, where u is a generator of $H^2(Z, 2; Z_2)$. Let $\{E_r^{**}\}$ be the mod 2 spectral sequence associated to the above fibration $((X, 4), X, K(Z, 2))$, then

$$E_2^{**} = A_0 \otimes A_1 \otimes \dots \otimes A_r \otimes B \otimes Z_2[u].$$

Clearly we have $d_3(1 \otimes u) = x_3 \otimes 1$. Hence if n is even, $d_3(1 \otimes u^n) = 0$, if n is odd, $d_3(1 \otimes u^n) = x_3 \otimes u^{n-1}$, and $d_3(x_3^{s_0-1} \otimes u^n) = 0$ for all $n > 0$. Thus we obtain

$$E_4^{**} = A(\bar{a}_0) \otimes A_1 \otimes A_2 \otimes \dots \otimes A_r \otimes B \otimes Z_2[u^n]$$

where $\bar{a}_0 = (x_3)^{s_0-1} \otimes u$.

Let τ be the transgression, $\tau(u^2) = Sq^2x_3$, since the transgression commutes the Steenrod operation. Thus $d_5(1 \otimes u^2) = Sq^2x_3 \otimes 1$. Since d_t is derivative, $d_5(1 \otimes u^{2n}) = 0$ if n is even, $d_5(1 \otimes u^{2n}) = Sq^2x_3 \otimes u^{2(n-1)}$ if n is odd, and $d_5((Sq^2x_3)^{2^{s_i-1}} \otimes u^{2^n}) = 0$ for all $n \geq 1$. Thus

$$E_6^{**} = A(\bar{a}_0, \bar{a}_1) \otimes A_2 \otimes A_3 \otimes \cdots \otimes A_r \otimes B \otimes Z_2[u^4]$$

where $\bar{a}_1 = (Sq^2 x_3)^{2^{s_1}-1} \otimes u^2$.

Carrying on similarly, we have

$$E_{2^{r+1}+2}^{**} = A(\bar{a}_0, \bar{a}_1, \dots, \bar{a}_r) \otimes B \otimes Z_2[u^{2^{r+1}}]$$

where $\bar{a}_i = (Sq^{2^i} Sq^{2^{i-1}} \cdots Sq^2 x_3)^{2^{s_i}-1} \otimes u^{2^i}$, $i = 0, 1, \dots, r$, and $s_i \geq 1$. Clearly $d_t = 0$ for all $t \geq 2^{r+1} + 2$. Thus we obtain

$$E_\infty^{**} = A(a_0, a_1, \dots, a_r) \otimes B \otimes Z_2[u^{2^{r+1}}].$$

Since E_∞^{**} is the graded algebra associated to $H^*((X, 4); Z_2)$, assume that a_i, w, B' correspond to $\bar{a}_i, u^{2^{r+1}}, B$ respectively. We have the lemma.

Particularly, we can assume that B is mapped isomorphically onto B' by the homomorphism $(f_1 \circ f_2 \circ f_3)^*$; $H^*(X; Z_2) \rightarrow H^*((X, 4); Z_2)$. Thus the relation of B are arranged in B' .

The mod 2 cohomology algebra of the exceptional Lie groups have been determined by S. Araki [2] and S. Araki-Y. Shikata [3]. These algebra are as follow.

- (2. 1) $H^*(F_4; Z_2) = Z_2[x_3]/(x_3^4) \otimes A(Sq^2 x_3, x_{15}, Sq^8 x_{15}),$
- (2. 2) $H^*(E_6; Z_2) = Z_2[x_3]/(x_3^4) \otimes A(Sq^2 x_3, Sq^4 Sq^2 x_3, x_{15}, Sq^8 Sq^4 Sq^2 x_3, Sq^8 x_{15}),$
- (2. 3) $H^*(E_7; Z_2) = Z_2[x_3, Sq^2 x_3, Sq^4 Sq^2 x_3]/(x_3^4, (Sq^2 x_3)^4, (Sq^4 Sq^2 x_3)^4) \otimes A(x_{15}, Sq^8 Sq^4 Sq^2 x_3, Sq^8 x_{15}, Sq^4 Sq^8 x_{15}),$
- (2. 4) $H^*(E_8; Z_2) = Z_2[x_3, Sq^2 x_3, Sq^4 Sq^2 x_3, x_{15}]/(x_3^{16}, (Sq^2 x_3)^8, (Sq^4 Sq^2 x_3)^4, x_{15}^4) \otimes A(Sq^8 Sq^4 Sq^2 x_3, Sq^8 x_{15}, Sq^4 Sq^8 x_{15}, Sq^2 Sq^4 Sq^8 x_{15})$

where x_i denotes a generator of degree i .

(2. 5) In the inclusion $F_4 \subset E_6 \subset E_7 \subset E_8$, every subgroup is totally non-homologous to zero mod 2 in any bigger group containing it, where each exceptional group denotes simply-connected one. (See, S. Araki and Y. Shikata [3], Theorem 3).

If $Sq^{16} Sq^8 Sq^4 Sq^2 x_3 = 0$ in E_8 , then this is a primitive element. By (2. 4), there is no primitive element of degree 33. Thus $Sq^{16} Sq^8 Sq^4 Sq^2 x_3 = 0$ in E_8 . Similarly we have $Sq^{16} Sq^8 Sq^4 Sq^2 x_3 = 0$ in E_6, E_7 and $Sq^4 Sq^2 x_3 = 0$ in F_4 .

COROLLARY 2. 3. *Let \tilde{G} be the 3-connective fibre space over G : where $G = F_4, E_6, E_7, E_8$, then*

$$\begin{aligned}
 H^*(\tilde{F}_4; Z_2) &= Z_2[y_8] \otimes \mathcal{A}(y_9, y_{11}, y_{15}, y_{23}), \\
 H^*(\tilde{E}_6; Z_2) &= Z_2[y_{32}] \otimes \mathcal{A}(y_9, y_{11}, y_{15}, y_{17}, y_{23}, y_{33}), \\
 H^*(\tilde{E}_7; Z_2) &= Z_2[y_{32}] \otimes \mathcal{A}(y_{11}, y_{15}, y_{19}, y_{23}, y_{27}, y_{33}, y_{35}), \\
 H^*(\tilde{E}_8; Z_2) &= Z_2[y_{15}, y_{32}]/(y_{15}^4) \otimes \mathcal{A}(y_{23}, y_{27}, y_{29}, y_{33}, y_{35}, y_{39}, y_{47}),
 \end{aligned}$$

where y_i denotes a generator of degree i . By the naturality of the homomorphism $p^* = (f_1 f_2 f_3)^*$, we have

$$\begin{aligned}
 Sq^8 y_{15} &= y_{23} && \text{in } \tilde{E}_6, \tilde{E}_7, \tilde{E}_8 \text{ and } \tilde{F}_4, \\
 Sq^4 y_{23} &= y_{27} && \text{in } \tilde{E}_7, \tilde{E}_8, \\
 Sq^2 y_{27} &= y_{29} && \text{in } \tilde{E}_8.
 \end{aligned}$$

LEMMA 2.4. We have the following relations,

- (i) $Sq^1 y_8 = y_9, Sq^2 y_9 = y_{11}$ in \tilde{F}_4 ,
- (ii) $Sq^2 y_9 = y_{11}, Sq^8 y_9 = y_{17}$ in \tilde{E}_6 ,
- (iii) $Sq^8 y_{11} = y_{19}$ in \tilde{E}_7 .

Proof. (i) From Theorem 2.1, there exists a fibration $(\tilde{F}_4, K(Z, 3), \tilde{F}_4)$, where \tilde{F}_4 denotes the space which has same homotopy type as F_4 . We consider the spectral sequence $\{E_r^{**}\}$ over Z_2 associated with the above fibration. Then

$$E_2^{**} = H^*(Z, 3; Z_2) \otimes H^*(\tilde{F}_4; Z_2).$$

It is known that

$$H^*(Z, 3; Z_2) = Z_2[v, Sq^2 v, Sq^4 Sq^2 v, \dots]$$

where v is a fundamental class of $H^3(Z, 3; Z_2)$. From the mod 2 cohomology algebra of $F_4, Sq^4 v \otimes 1, (Sq^2 v)^2 \otimes 1$ and $v^4 \otimes 1$ must be d_r -images for some r . If $p \neq 0$ and $0 < q < 8$, or $q \neq 0$ and $0 < p < 3$, then $E_r^{p, q} = 0$ for all r . Since $E_r^{0, 8}$ has only one element $1 \otimes y_8$ for $r \leq 9$, $Sq^4 Sq^2 v \otimes 1$ is not a d_r -image for $r \leq 8$. Thus τ be the transgression, we have $\tau(y_8) = Sq^4 Sq^2 v$. Since $E_r^{0, 9}$ has only one generator $1 \otimes y_9$ and $(Sq^2 v)^2 \otimes 1$ is not a d_r -image for $r \leq 10$, we have that $\tau(y_9) = (Sq^2 v)^2$. Consider

$$d_r : E_r^{p, q} \longrightarrow E_r^{1+2, 0} \quad \text{for } p + q = 11 \text{ and } r = q + 1.$$

From Corollary 2.3, we have $E_r^{p, q} = 0$ for $q \neq 8, 9$. But $E_7^{2, 9} = 0$. $E_7^{3, 8}$ has one generator $v \otimes y_8$ and $d_9(v \otimes y_8) = v Sq^4 Sq^2 v \otimes 1 \neq 0$, for $d_9(1 \otimes y_8)$

$= Sq^4Sq^2v \otimes 1$. Thus $E_{12}^{0,11}$ has only one generator $1 \otimes y_{11}$ and $v^4 \otimes 1$ is not a d_r -image for $r \leq 11$. Therefore we have that $\tau(y_{11}) = v^4$. Using Adem's relation, from $Sq^1Sq^4Sq^2v = Sq^5Sq^2v = (Sq^2v)^2$, $Sq^2(Sq^2v)^2 = Sq^2Sq^5Sq^2v = Sq^6Sq^3v = v^4$, we obtain $Sq^1y_8 = y_9$, and $Sq^2y_9 = y_{11}$.

(ii) From Theorem 2.1, there exists a fibration $(\bar{E}_6, K(Z, 3), \tilde{E}_6)$ where \bar{E}_6 denotes the space which has the same homotopy type as E_6 . Let τ be the transgression associated with this fibration. Let $\{E_r^{p,q}\}$ be the mod 2 spectral sequence associated with this fibration. Then

$$E_2^{**} = H^*(Z, 3; Z_2) \otimes H^*(\tilde{E}_6; Z_2).$$

By the same argument as in \tilde{F}_4 , we have that $\tau(y_9) = (Sq^2v)^2$ and $\tau(y_{11}) = v^4$. Consider

$$d_r; E_r^{p,q} \longrightarrow E_r^{18,0} \text{ for } p + q = 17 \text{ and } r = q + 1.$$

From Corollary 2.3, we have $E_r^{p,q} = 0$ for $q \neq 9, 11, 15$ and 17 ($q \leq 22$). But $E_r^{2,15} = 0$. $E_{10}^{8,9}$ has one generator $(vSq^2v) \otimes y_9$ and $d_{10}((vSq^2v) \otimes y_9) = v(Sq^2v)^3 \otimes 1 \neq 0$, for $d_{10}(1 \otimes y_9) = (Sq^2v)^2 \otimes 1$. $E_{12}^{6,11}$ has one generator $v^2 \otimes y_{11}$ and $d_{12}(v^2 \otimes y_{11}) = v^6 \otimes 1 \neq 0$ for $d_{12}(1 \otimes y_{11}) = v^4 \otimes 1$. Thus, since $E_{17}^{0,17}$ has one generator y_{17} and $(Sq^4Sq^2v)^2 \otimes 1$ is not a d_r -image for $r \leq 16$, $d_{18}(1 \otimes y_{17}) = (Sq^4Sq^2v)^2 \otimes 1$, i.e. $\tau(y_{17}) = (Sq^4Sq^2v)^2$. Using Adem's relation, $Sq^2(Sq^2v)^2 = Sq^2Sq^5Sq^2v = Sq^6Sq^3v = v^4$ and $Sq^8(Sq^2v)^2 = Sq^8Sq^5Sq^2v = Sq^9Sq^4Sq^2v = (Sq^4Sq^2v)^2$. From the commutativity of the Steenrod operation and the transgression, we obtain $Sq^2y_9 = y_{11}$ and $Sq^8y_9 = y_{17}$.

(iii) Consider the fibration $(\bar{E}_7, K(Z, 3), \tilde{E}_7)$ of theorem 2.1 (II), where \bar{E}_7 has the same homotopy type as E_7 . Let $\{E_r^{p,q}\}$ be the mod 2 spectral sequence associated with this fibration. Then

$$E_r^{**} = H^*(Z, 3; Z_2) \otimes H^*(\tilde{E}_7; Z_2).$$

From the mod 2 cohomology algebra of E_7 , $v^4 \otimes 1$ and $(Sq^2v)^4 \otimes 1$ must be the d_r -images for some r . Since $H^*(\bar{E}_7; Z_2) = 0$ for degree ≤ 10 , we have $E_r^{p,q} = 0$ for $p \neq 0$ and $0 < q < 10$. Thus we have that $\tau(y_{11}) = v^4$, where τ is the transgression. Consider

$$d_r; E_r^{p,q} \longrightarrow E_r^{20,0} \text{ for } p + q = 19 \text{ and } r = q + 1.$$

From $H^i(\tilde{E}_7; Z_2) = 0$ for $i \neq 11, 15$ and $i < 19$, it follows that $E_r^{p,q} = 0$ for $(p, q) \neq (4, 11)$ and $(2, 15)$. On the other hand $H^i(Z, 3; Z_2) = 0$ for $i = 2, 4$

and $i \leq 4$. Thus $E_r^{p,q} = 0$ for $(p, q) = (4, 11)$ and $(2, 15)$. From this we obtain $\tau(y_{19}) = (Sq^2v)^4$. By Adem's relation $Sq^8v^4 = Sq^8Sq^6Sq^3v = Sq^{10}Sq^4Sq^3v + Sq^{11}Sq^3Sq^3v = Sq^{10}Sq^5Sq^2 + Sq^{11}Sq^5Sq^1v = (Sq^2v)^4$. Thus we obtain $Sq^8y_{11} = y_{19}$.

LEMMA 2.5. *Let a topological space X be 2-connected and the homology of finite type. Assume that $H^*(X; Z_2)$ has the additive basis a_1, \dots, a_s for $\dim. < N$. Then there exist a finite cell complex $K = * \cup e_1 \cup e_2 \cup \dots \cup e_s$, where $\dim.e_i = \text{degree } a_i = n_i$ and a continuous map $f; K \rightarrow X$ such that f induces isomorphism of $H^*(X; Z_2)$ onto $H^*(K; Z_2)$ for $\dim. < N$.*

Particularly if $\pi_{n_i-1}(K^{n_i-1})$ is finite, then we can assume that the class of attaching map of e_i belong to the 2-components. Here $$ denotes a vertex and K^n the n -skelton of K .*

Proof. We prove this by induction on dimension N . Suppose that there exist a finite cell complex $K_0 = K^{N-1}$ and a continuous map $f_0; K_0 \rightarrow X$ satisfying lemma 2.5 for $\dim. < N$. Here we may assume that $f_0; K_0 \rightarrow X$ is the injection by the mapping-cylinder argument. Suppose that $H^N(X; Z_2)$ has generator a_{s+1}, \dots, a_r .

From the cohomology exact sequence for pair (X, K_0) and the assumption of the induction, we have

$$H^i(X, K_0; Z_2) = 0 \quad \text{for } i < N,$$

$$H^N(X, K_0; Z_2) \approx H^N(X; Z_2).$$

By the duality, we obtain

$$H_i(X, K_0; Z_2) = 0 \quad \text{for } i < N$$

and

$$H_N(X, K_0; Z_2) \text{ has the generators } \bar{a}_{s+1}, \dots, \bar{a}_r.$$

By Serre's C -theory [8], we have that $\pi_N(X, K_0) \otimes Z_2 \rightarrow H_N(X, K_0) \otimes Z_2$ is an isomorphism. Let $f_i: (E^N, S^{N-1}) \rightarrow (X, K_0)$ ($i = 1, 2, \dots, r - s$) be the generators of $\pi_N(X, K_0)$ such that they correspond to \bar{a}_{s+i} by the above isomorphism and construct a cell complex K which is obtained from the disjoint union of $C(S_1^{N-1} \vee \dots \vee S_{r-s}^{N-1})$ and K_0 by identifying $S_1^{N-1} \vee \dots \vee S_{r-s}^{N-1}$ with its image under a map $(f_1|S_1^{N-1}) \vee \dots \vee (f_{r-s}|S_{r-s}^{N-1}); S_1^{N-1} \vee \dots \vee S_{r-s}^{N-1} \rightarrow K_0$, where CY is a cone over the space Y and S_i^{N-1} is a $(N - 1)$ -sphere. Using the map f_i the inclusion map $f_0; K_0 \rightarrow X$ has

an extension over K and we denote this extension by $g : K \rightarrow X$. Then $g : K \rightarrow X$ induce an isomorphism $H_N(K, K_0; Z_2)$ onto $H_N(X, K_0; Z_2)$ and from the duality between homology and cohomology, it follows that $g^* : H^N(X, K_0; Z_2) \rightarrow H^N(K, K_0; Z_2)$ is an isomorphism onto.

Applying the five lemma to the diagram

$$\begin{array}{ccccccc} H^{N-1}(K_0; Z_2) & \longrightarrow & H^N(X, K_0; Z_2) & \longrightarrow & H^N(X; Z_2) & \longrightarrow & H^N(K_0; Z_2) = 0 \\ \downarrow \approx & & \downarrow g^* & & \downarrow g^{**} & & \downarrow \approx \\ H^{N-1}(K_0; Z_2) & \longrightarrow & H^N(K, K_0; Z_2) & \longrightarrow & H^N(K; Z_2) & \longrightarrow & H^N(K_0; Z_2) = 0, \end{array}$$

we obtain that

$$g^* : H^N(X; Z_2) \longrightarrow H^N(K; Z_2)$$

is an isomorphism.

Particularly if $\pi_{N-1}(K_0)$ is finite, then there exists an odd integer q such that $q\{f_i | S^{N-1}\}$ belongs to the 2-component of $\pi_{N-1}(K_0)$. Displacing f_i by qf_i , it is sufficient for the last statement that we construct a cell complex K from K_0 . Consequently the lemma is proved.

Let α be an element of $\pi_{n+i-1}(S^n)$ and consider a cell complex $K_\alpha = S^n \cup e^{n+i}$ which is uniquely determined by α up to homotopy type.

THEOREM 2.6. *Let $n > i$ and $i = 2$ (4 or 8 respectively), then $Sq^i : H^n(K_\alpha; Z_2) \rightarrow H^{n+i}(K_\alpha; Z_2)$ is an isomorphism onto if and only if $\alpha \equiv \eta_n$, (ν_n or σ_n respectively) mod $2\pi_{n+i-1}(S^n)$. (For the proof see H. Tada; [11] Proposition 8.1)*

From Lemma 2.5 and Corollary 2.3, there exist a cell complex $M = S^8 \cup e^9 \cup e^{11} \cup e^{15}$ and a continuous map $f : M \rightarrow \tilde{F}_4$ such that f induces an C_2 -isomorphisms $\pi_i(M)$ onto $\pi_i(\tilde{F}_4)$ for $i \leq 14$, where C_2 is the classes of finite abelian group whose 2-primary components are zero. Since $Sq^1 y_8 = y_9$ in \tilde{F}_4 , we may assume that e^9 is attached to S^8 by a map of degree two. Then we have

$$(2.6) \quad \begin{aligned} \pi_{13}(S^8 \cup_2 e^9 : 2) &= 0, \\ \pi_{14}(S^8 \cup_2 e^9 : 2) &\approx \pi_{14}(S^8 : 2) = Z_2 \quad \text{generated by } \nu_8^2, \end{aligned}$$

we denote by ν_8^2 a generator of $\pi_{14}(S^8 \cup_2 e^9 : 2)$ identifying with that of $\pi_{14}(S^8 : 2)$ by the inclusion $S^8 \subset S^8 \cup_2 e^9$.

Consider the following exact sequence

$$\pi_i(S^8 : 2) \longrightarrow \pi_i(S^8 : 2) \longrightarrow \pi_i(S^8 \cup e^9 : 2) \longrightarrow \pi_i(S^9 : 2) \longrightarrow \pi_i(S^9 : 2)$$

for $i \leq 15$. From $\pi_{12}(S^8) = \pi_{13}(S^9) = \pi_{14}(S^9) = 0$ and $\pi_{14}(S^8) = \{\nu_8^2\} = Z_2$, (2. 6) is obtained.

Consider the exact sequence

$$\begin{aligned} \pi_{14}(S^{10} : 2) &\longrightarrow \pi_{14}(S^8 \cup_2 e^9 : 2) \xrightarrow{i^*} \pi_{14}(S^8 \cup_2 e^9 \cup e^{11} : 2) \xrightarrow{j_*} \pi_{14}(S^{11} : 2) \\ &\longrightarrow \pi_{14}(S^9 \cup_2 e^{10} : 2) \end{aligned}$$

where i is the inclusion $S^8 \cup_2 e^9 \subset S^8 \cup_2 e^9 \cup e^{11}$, and $j : S^8 \cup_2 e^9 \cup e^{11} \longrightarrow S^{11}$ is the projection. From (2. 6), we have the following exact sequence

$$(2. 7) \quad 0 \longrightarrow \pi_{14}(S^8 \cup_2 e^9 : 2) \xrightarrow{i^*} \pi_{14}(S^8 \cup_2 e^9 \cup e^{11} : 2) \xrightarrow{j_*} \pi_{14}(S^{11} : 2) \longrightarrow 0.$$

Then there exists a coextension (in the sense of [11]) $\tilde{\nu}_{10}$ of ν_{10} and $j_*\tilde{\nu}_{10} = \nu_{11}$. Assume that $8\tilde{\nu}_{10} = 0$, then $-i_*\nu_8^2 = i_*\nu_8^2 = 8\tilde{\nu}_{10}$. Let $f : S^{14} \vee S^{11} \longrightarrow S^8 \cup_2 e^9 \cup e^{11}$ be a map such that $f|_{S^{14}}$ and $f|_{S^{11}}$ representative of $8i_{14} \oplus \nu_{11}$, then $f \circ g : S^{14} \longrightarrow S^8 \cup_2 e^9 \cup e^{11}$ is homotopic to zero. Consider a mapping cone C_f of f , then there exists a coextension $G : S^{15} \longrightarrow C_f$ of g . Let K be a mapping cone of G , then we have a complex

$$K = S^8 \cup e_6 \cup e^{11} \cup e^{12} \cup e^{15} \cup e^{16}$$

and $Sq^4u_8 = u_{12}$, $Sq^4u_{12} = u_{16}$, where u_8, u_{12} and u_{16} are cohomology classes mod 2 which are represented by S^8, e^{12} and e^{16} respectively. Thus it is verified that $Sq^4Sq^4u_8 \neq 0$ in K . By use of Adem's relation

$$Sq^4Sq^4u_8 = Sq^5Sq^2u_8 + Sq^2Sq^6u_8.$$

Since there is no cell of dimension 10 or 14 in K , the right side of the above equation vanishes in K , but this is a contradiction. Thus we have proved that $8\tilde{\nu}_{10} = 0$. Therefore, from the exact sequence (2. 7), we obtain

$$\pi_{14}(S^8 \cup e^9 \cup e^{11} : 2) = \{i_*\nu_8^2\} + \{\tilde{\nu}_{10}\} \approx Z_2 + Z_8.$$

In the complex $M = S^8 \cup_2 e^9 \cup e^{11} \cup e^{15}$, let e^{15} be attached to $S^8 \cup_2 e^9 \cup e^{11}$ by a map $h : S^{14} \longrightarrow S^8 \cup_2 e^9 \cup e^{11}$, then we have the sequence

$$\pi_{14}(S^{14} : 2) \xrightarrow{h_*} \pi_{14}(S^8 \cup_2 e^9 \cup e^{11} : 2) \longrightarrow \pi_{14}(M : 2) \longrightarrow \pi_{14}(S^{15} : 2) = 0$$

is exact. By Lemma 5.5 of [10], $\pi_{14}(F_4) = Z_2$. Thus $\pi_{14}(M : 2) \approx Z_2$ and

$$h_*\iota_{14} = b\bar{\nu}_{10} + a(i_*\nu_8^2) \quad \text{where } a = 0 \text{ or } 1,$$

for an odd integer b . Thus

$$j_*h_*\iota_{14} = \nu_{11} \pmod{2\pi_{14}(S^{11})}.$$

By theorem 2.6, we have the following important lemma.

LEMMA 2.7. $Sq^4y_{11} = y_{15}$ in \tilde{F}_4 .

Considering the natural inclusions $\tilde{F}_4 \subset \tilde{E}_6 \subset \tilde{E}_7$, we have

COROLLARY 2.8. $Sq^4y_{11} = y_{15}$ in \tilde{E}_6 and \tilde{E}_7 .

§ 3. Homotopy group of some cell complexes.

Let X be an m -connected CW-complex and let α be an element of $\pi_{n-1}(X)$ ($n > m$). Consider a CW-complex $K_\alpha = X \cup_{\alpha} e^n$.

LEMMA 3.1. Let i be an injection $X \rightarrow K_\alpha$ and let $p : K_\alpha \rightarrow S^n$ be a mapping which shrinks X to a point. Then the following sequence is exact for $j \leq m + n - 1$

$$(3.1) \quad \dots \rightarrow \pi_j(S^{n-1}) \xrightarrow{\alpha_*} \pi_j(X) \xrightarrow{i_*} \pi_j(K_\alpha) \rightarrow \pi_{j-1}(S^{n-1}) \xrightarrow{\alpha_*} \pi_{j-1}(X) \rightarrow \dots$$

Here ∂ is a composition $E^{-1} \circ p_* : \pi_j(K_\alpha) \rightarrow \pi_{j-1}(S^{n-1})$, and $E : \pi_{j-1}(S^{n-1}) \rightarrow \pi_j(S^n)$ is the suspension homomorphism. If α is of order a power of 2, then the above sequence is exact for the 2-primary components.

Proof. See Blakers-Massey [4].

We introduce necessary results on the homotopy group of spheres. According to [11], the results are listed in the following table;

(i) $n > k + 1$

(3.2)

$k =$	0	1	2	3	4	5	6	7	8
$\pi_{n+k}(S^n : 2)$	Z	Z_2	Z_2	Z_8	0	0	Z_2	Z_{16}	$Z_2 + Z_2$
Generator	ι_n	η_n	η_n^2	ν_n			ν_n^2	σ_n	$\bar{\nu}_n, \varepsilon_n$

$k =$	9	10	11	12	13
$\pi_{n+k}(S^n : 2)$	$Z_2 + Z_2 + Z_2$	Z_2	Z_8	0	0
Generator	$\nu_n^3, \eta_n \varepsilon_{n+1}, \mu_n$	$\eta_n \mu_{n+1}$	ζ_n		

(ii) $n \leq k + 1$ $n = 9, 10, 11, 13, 14.$

(3. 3)

$k =$	8	9	10	11
$\pi_{k+9}(S^9 : 2)$	$Z_2 + Z_2 + Z_2$	$Z_2 + Z_2 + Z_2 + Z_2$	$Z_8 + Z_2$	$Z_8 + Z_2$
Generator	$\sigma_9 \eta_{16}, \bar{\nu}_9, \epsilon_9$	$\sigma_9 \eta_{16}^2, \nu_9^3, \mu_9, \eta_9 \epsilon_{10}$	$\sigma_9 \nu_{16}, \eta_9 \mu_{10}$	$\zeta_9, \bar{\nu}_9 \nu_{17}$
$\pi_{k+10}(S^{10} : 2)$		$Z + Z_2 + Z_2 + Z_2$	$Z_4 + Z_2$	Z_8
Generator		$\Delta(\ell_{21}), \nu_{10}^3, \mu_{10}, \eta_{10} \epsilon_{11}$	$\sigma_{10} \nu_{17}, \eta_{10} \mu_{11}$	ζ_{10}
$\pi_{k+11}(S^{11} : 2)$			$Z_2 + Z_2$	Z_8
Generator			$\sigma_{11} \nu_{18}, \eta_{11} \mu_{12}$	ζ_{11}
$\pi_{k+13}(S^{13} : 2)$				
Generator				
$\pi_{k+14}(S^{14} : 2)$				
Generator				

$k =$	12	13	14
$\pi_{k+9}(S^9 : 2)$	0	Z_2	$Z_{16} + Z_4$
Generator		$\sigma_9 \nu_{16}^2$	σ_9^2, κ_9
$\pi_{k+10}(S^{10} : 2)$	Z_4	Z_2	$Z_{16} + Z_2$
Generator	$\Delta(\nu_{21})$	$\sigma_{10} \nu_{17}^2$	$\sigma_{10}^2, \kappa_{10}$
$\pi_{k+11}(S^{11} : 2)$	Z_2	$Z_2 + Z_2$	$Z_{16} + Z_2$
Generator	θ'	$\theta' \eta_{23}, \sigma_{11} \nu_{18}^2$	$\sigma_{11}^2, \kappa_{11}$
$\pi_{k+13}(S^{13} : 2)$	Z_2	Z_2	$Z_{16} + Z_2$
Generator	$E\theta$	$E\theta \eta_{25}$	$\sigma_{13}^2, \kappa_{13}$
$\pi_{k+14}(S^{14} : 2)$		Z	$Z_8 + Z_2$
Generator		$\Delta(\ell_{29})$	$\sigma_{14}^2, \kappa_{14}$

We shall use the following relations;

(3. 4) $\sigma_n \circ \mu_{n+7} = \eta_n \circ \sigma_{n+1} = \bar{\nu}_n + \epsilon_n$ for $n \geq 10$
 by Lemma 6. 4 of [11],

- (3. 5) $\sigma_n \circ \eta_{n+7}^2 = \eta_n^2 \circ \sigma_{n+2} = \nu_n^3 + \eta_n \circ \varepsilon_{n+1}$ for $n \geq 10$
by Lemma 6. 3 of [11],
- (3. 6) $\sigma_n \circ \nu_{n+7} = 0$ for $n \geq 12$
 $\nu_n \circ \sigma_{n+3} = 0$ for $n \geq 11$,
 $2\sigma_{10} \circ \nu_{17} = \nu_{10} \circ \sigma_{13}$ by (7. 20) of [11],
 $\varepsilon_n \circ \eta_{n+8}^2 = \eta_n^2 \circ \varepsilon_{n+2} = 0$ for $n \geq 9$ by (7. 10) and (7. 20) of [11],
- (3. 7) $\sigma_n \circ \bar{\nu}_{n+7} = 0$ for $n \geq 11$ by (10. 8) of [11],
 $\sigma_n \circ \varepsilon_{n+7} = 0$ for $n \geq 6$ by Lemma 10. 7 of [11],
- (3. 8) $\nu_n \circ \varepsilon_{n+3} = \nu_n \circ \nu_{n+3} = 0$ for $n \geq 7$ by (7. 17) of [11],
 $\nu_n \circ \eta_{n+3} = \eta_n \circ \nu_{n+1} = 0$ for $n \geq 6$ by (5. 9) of [11],
- (3. 9) $\nu_n \circ \mu_{n+3} = 0$ for $n \geq 7$ by Theorem 7. 6 of [11],
- (3. 10) $A(\iota_{21}) \circ \eta_{19} = 2\sigma_{10} \circ \nu_{17}$ by (7. 21) of [11].

Consider a generator σ_n of $\pi_{n+7}(S^n : 2) \approx Z_{16}$ for $n \geq 9$ and a cell complex $K_{\sigma_n} = S^n \cup_{\sigma_n} e^{n+8}$. Let $i : S^n \rightarrow K_{\sigma_n}$ be the injection.

PROPOSITION 3. 2. *We have the following tables of the homotopy groups $\pi_j(K_{\sigma_n} : 2)$ for $n = 9, 10, 11, 14$ and 15, and generator of their 2-primary components.*

(3. 11)

j	$j \leq 8$	9	10	11	12	13	14	15	16
$\pi_j(K_{\sigma_9} : 2)$	0	Z	Z_2	Z_2	Z_8	0	0	Z_2	0
Generator		$i_*\iota_9$	$i_*\eta_9$	$i_*\eta_9^2$	$i_*\nu_9$			$i_*\nu_9^2$	

j	17	18	19	20	21	22
$\pi_k(K_{\sigma_9} : 2)$	$Z + Z_2 + Z_2$	$Z_2 + Z_2 + Z_2$	Z_2	$Z_8 + Z_2$	0	0
Generator	$\widetilde{16}\iota_{16}, i_*\varepsilon_9, i_*\bar{\nu}_9$	$i_*\eta_9\varepsilon_{10}, i_*\nu_9^8, i_*\mu_9$	$i_*\eta_9\mu_{10}$	$i_*\zeta_9, i_*\bar{\nu}_9\nu_{17}$		

(3. 12)

j	$j \leq 9$	10	11	12	13	14	15	16	17
$\pi_j(K_{\sigma_{10}} : 2)$	0	Z	Z_2	Z_2	Z_8	0	0	Z_2	0
Generator		$i_*\iota_{10}$	$i_*\eta_{10}$	$i_*\eta_{10}^2$	$i_*\nu_{10}$			$i_*\nu_{10}^2$	

j	18	19	20	21	22	23
$\pi_j(K_{\sigma_{10}} : 2)$	$Z + Z_2$	$Z + Z_2 + Z_2$	Z_2	Z_{16}	Z_4	0
Generator	$\widetilde{16\epsilon_{17}}, i_*\epsilon_{10}$	$i_*\Delta(\epsilon_{21}), i_*\eta_{10}\epsilon_{11}, i_*\mu_{10}$	$i_*\eta_{10}\mu_{11}$	$\widetilde{4\nu_{17}}$	$i_*\Delta(\nu_{21})$	

(3. 13)

j	$j \leq 9$	11	12	13	14	15	16	17	18
$\pi_j(K_{\sigma_{11}} : 2)$	0	Z	Z_2	Z_2	Z_8	0	0	Z_2	0
Generator		$i_*\epsilon_{11}$	$i_*\eta_{11}$	$i_*\eta_{11}^2$	$i_*\nu_{11}$			$i_*\nu_{11}^2$	

j	19	20	21	22	23	24	25
$\pi_j(K_{\sigma_{11}} : 2)$	$Z_2 + Z$	$Z_2 + Z_2$	Z_2	Z_{32}	Z_2	Z_2	Z_2
Generator	$i_*\epsilon_{14}, \widetilde{16\epsilon_{18}}$	$i_*\mu_{11}, i_*\eta_{11}\epsilon_{12}$	$i_*\eta_{11}\mu_{12}$	$\widetilde{2\nu_{18}}$	$i_*\theta'$	$i_*\theta'\eta_{23}$	$i_*\kappa_{11}$

(3. 14)

j	$j \leq 13$	14	15	16	17	18	19	20	21
$\pi_j(K_{\sigma_{14}} : 2)$	0	Z	Z_2	Z_2	Z_8	0	0	Z_2	0
Generator		$i_*\epsilon_{14}$	$i_*\eta_{14}$	$i_*\eta_{14}^2$	$i_*\nu_{14}$			$i_*\nu_{14}^2$	

j	22	23	24	25	26	27
$\pi_j(K_{\sigma_{14}} : 2)$	$Z + Z_2$	$Z_2 + Z_2$	Z_2	Z_{64}	0	Z
Generator	$\widetilde{16\epsilon_{21}}, i_*\epsilon_{14}$	$i_*\mu_{14}, i_*\eta_{14}\epsilon_{15}$	$i_*\eta_{14}\mu_{15}$	$\widetilde{\nu_{21}}$		$i_*\Delta(\epsilon_{29})$

(3. 15)

j		15	16	17	18	19	20	21	22
$\pi_j(K_{\sigma_{15}} : 2)$	0	Z	Z_2	Z_2	Z_8	0	0	Z_2	0
Generator		$i_*\epsilon_{15}$	$i_*\eta_{15}$	$i_*\eta_{15}^2$	$i_*\nu_{15}$			$i_*\nu_{15}^2$	

j	23	24	25	26	27	28
$\pi_j(K_{\sigma_{15}} : 2)$	$Z + Z_2$	$Z_2 + Z_2$	Z_2	Z_{64}	0	0
Generator	$\widetilde{16\epsilon_{22}}, i_*\epsilon_{15}$	$i_*\mu_{15}, i_*\eta_{15}\epsilon_{16}$	$i_*\eta_{15}\mu_{16}$	$\widetilde{\nu_{22}}$		

Here we denote by $\tilde{\beta}$ an element of $\pi_i(K_{\sigma_n} : 2)$ such that $\partial\tilde{\beta} = \beta \in \pi_{i-1}(S^{n+7} : 2)$ i.e. we may consider that $\tilde{\beta}$ is a coextension of β .

Proof. Consider the exact sequence

$$\begin{aligned} \dots \longrightarrow \pi_j(S^{n+7} : 2) &\xrightarrow{\sigma_{n*}} \pi_j(S^n : 2) \xrightarrow{i_*} \pi_j(K_{\sigma_n} : 2) \xrightarrow{\partial} \pi_{j-1}(S^{n+7} : 2) \\ &\xrightarrow{\sigma_{n*}} \pi_{j-1}(S^n : 2) \longrightarrow \dots \end{aligned}$$

of (3. 1) for $j \leq 2n + 5$. From $\pi_j(S^{n+7} : 2) = 0$ for $j \leq n + 6$ and from the exactness of the above sequence, it follows that

$$i_* : \pi_j(S^n : 2) \longrightarrow \pi_j(K_{\sigma_n} : 2)$$

are isomorphisms onto for $j \leq n + 6$, and $n = 9, 10, 11, 14, 15$.

It follows from (3. 1) that the sequence

$$\pi_{n+7}(S^{n+7} : 2) \xrightarrow{\sigma_{n*}} \pi_{n+7}(S^n : 2) \xrightarrow{i_*} \pi_{n+7}(K_{\sigma_n} : 2) \xrightarrow{\partial} \pi_{n+6}(S^{n+7} : 2) = 0$$

is exact for $n \geq 9$. From $\pi_{n+7}(S^n : 2) \approx \{\sigma_n\} \approx Z_{16}$, we have that

$$(3. 16) \quad \sigma_{n*} : \pi_{n+7}(S^{n+7} : 2) \longrightarrow \pi_{n+7}(S^n : 2)$$

is an epimorphism. Thus we obtain $\pi_{n+7}(K_{\sigma_n} : 2) = 0$ for $n = 9, 10, 11, 14$ and 15.

Consider the exact sequence

$$\pi_{n+8}(S^{n+7} : 2) \xrightarrow{\sigma_n} \pi_{n+8}(S^n : 2) \xrightarrow{i_*} \pi_{n+8}(K_{\sigma_n} : 2) \xrightarrow{\partial} Z = \{16\epsilon_{n+7}\} \longrightarrow 0$$

of (3. 1) for $n \geq 9$. From (3. 2), (3. 3) and (3. 4) we have that

$$(3. 17) \quad \sigma_{n*} : \pi_{n+8}(S^{n+7} : 2) \longrightarrow \pi_{n+8}(S^n : 2)$$

are monomorphisms for $n \geq 9$. Thus it follows from the exactness of the above sequence that the table is true for $\pi_{n+8}(K_{\sigma_n} : 2)$, $n = 9, 10, 11, 14, 15$.

From (3. 17) and the exact sequence (3. 1), it follows that the sequence

$$\pi_{n+9}(S^{n+7} : 2) \xrightarrow{\sigma_{n*}} \pi_{n+9}(S^n : 2) \xrightarrow{i_*} \pi_{n+9}(K_{\sigma_n} : 2) \longrightarrow 0$$

is exact for $n \geq 9$. From (3. 5), (3. 2) and (3. 3), we have that

$$(3. 18) \quad \sigma_{n*} : \pi_{n+9}(S^{n+7} : 2) \longrightarrow \pi_{n+9}(S^n : 2)$$

is monomorphisms for $n \geq 9$. Thus we obtain that

$$\pi_{n+9}(K_{\sigma_n} : 2) \approx \pi_{n+9}(S^n : 2) / \{\sigma_n \circ \eta_{n+7}\}.$$

From (3. 18) and the exact sequence (3. 1), it follows that the sequence

$$\pi_{n+10}(S^{n+7} : 2) \xrightarrow{\sigma_{n*}} \pi_{n+10}(S^n : 2) \xrightarrow{i_*} \pi_{n+10}(K_{\sigma_n} : 2) \longrightarrow 0$$

is exact for $n \geq 9$. From (3. 2), (3. 3) and (3. 6), it follows that

(3. 19) $\sigma_{9*} : \pi_{19}(S^{16} : 2) \longrightarrow \pi_{19}(S^9 : 2)$ is a monomorphism,
 $\sigma_{n*} : \pi_{n+10}(S^{n+7} : 2) \longrightarrow \pi_{n+10}(S^n : 2)$ is trivial for $n = 14, 15$,
 the kernel of $\sigma_{10*} : \pi_{20}(S^{17} : 2) \longrightarrow \pi_{20}(S^{10} : 2)$ is

generated by $\{4\nu_{17}\}$, and

$$\text{the kernel of } \sigma_{11*} : \pi_{21}(S^{18} : 2) \longrightarrow \pi_{21}(S^{11} : 2) \text{ is}$$

generated by $\{2\nu_{18}\}$.

Thus it follows that the table is true for $\pi_{n+10}(K_{\sigma_n} : 2)$ $n = 9, 10, 11, 14$ and 15 .

In the stable rangs, we have the exact sequence

$$0 \longrightarrow \pi_{n+11}(S^n : 2) \xrightarrow{i_*} \pi_{n+11}(K_{\sigma_n} : 2) \xrightarrow{\partial} \pi_{n+10}(S^{n+7} : 2) \longrightarrow 0$$

of (3. 1) for $n \geq 13$. Moreover we have the following relation in the stable secondary compositions

$$\begin{aligned} \zeta \in \langle \sigma, 4\nu, 2\iota \rangle \pmod{2G_{11}} & \quad \text{from Lemma 9. 1 of [11],} \\ \supset \langle \sigma, \nu, 8\iota \rangle & \quad \text{from Proposition 1. 2 of [11],} \end{aligned}$$

and $\langle \sigma, \nu, 8\iota \rangle$ is a coset of the subgroup $\sigma \circ G_4 + 8G_{11} = 8G_{11}$. Thus

$$\zeta \equiv \langle \sigma, \nu, 8\iota \rangle \pmod{2 G_{11}}$$

where G_n is the n -th stable homotopy group of the sphere and ζ is a generator of the 2-components of G_{11} .

From Proposition 1. 8 of [11], we obtain

$$\begin{aligned} i_*\xi &= i_* \langle \sigma, \nu, 8\iota \rangle \pmod{2 i_*G_{11}} \\ &= -8\bar{\nu} \end{aligned}$$

where $\bar{\alpha} \in \pi_i(K_{\sigma_n} : 2)$ is a coextension of $\alpha \in \pi_{i-1}(S^{n+7} : 2)$. Thus, from this and from the exactness of the above sequence it follows that

(3. 20) $\pi_{n+11}(K_{\sigma_n} : 2) = \{\bar{\nu}\} = Z_{64}$

for $n \geq 13$

From (3. 1), (3. 19) and from $\pi_{n+11}(S^{n+7} : 2) = 0$ for $n \geq 0$, it follows the next four exact sequences and the commutative diagram

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \pi_{20}(S^9 : 2) & \xrightarrow{i_*} & \pi_{20}(K_{\sigma_9} : 2) & \longrightarrow & 0 \\
 & & \downarrow E & & \downarrow E & & \\
 0 & \longrightarrow & \pi_{21}(S^{10} : 2) & \xrightarrow{i_*} & \pi_{21}(K_{\sigma_{10}} : 2) & \xrightarrow{\theta} & \{4\nu_{17}\} \longrightarrow 0 \\
 & & \downarrow E & & \downarrow E & & \downarrow E \\
 0 & \longrightarrow & \pi_{22}(S^{11} : 2) & \xrightarrow{i_*} & \pi_{22}(K_{\sigma_{11}} : 2) & \xrightarrow{\theta} & \{2\nu_{18}\} \longrightarrow 0 \\
 & & \downarrow E^{n-11} & & \downarrow E^{n-11} & & \downarrow E^{n-11} \\
 0 & \longrightarrow & \pi_{n+11}(S^n : 2) & \xrightarrow{i_*} & \pi_{n+11}(K_{\sigma_n} : 2) & \xrightarrow{\theta} & \{\nu_{n+7}\} \longrightarrow 0
 \end{array}$$

for $n \geq 13$, where $E : \pi_{21}(S^{10} : 2) \longrightarrow \pi_{22}(S^{11} : 2)$ and $E^{n-11} : \pi_{22}(S^{11} : 2) \longrightarrow \pi_{n+11}(S^n : 2)$ are isomorphisms. From (3. 20) and the above diagram, we obtain that

$$\begin{aligned}
 \pi_{20}(K_{\sigma_9} : 2) &= \{i_*\zeta_9\} + \{i_*\bar{\nu}_9 \circ \nu_{17}\} \approx Z_8 + Z_2, \\
 \pi_{21}(K_{\sigma_{11}} : 2) &= \{\tilde{4}\nu_{17}\} \approx Z_{16}, \\
 \pi_{22}(K_{\sigma_{11}} : 2) &= \{\tilde{2}\nu_{18}\} \approx Z_{32}, \\
 \pi_{n+11}(K_{\sigma_n} : 2) &= \{\tilde{\nu}_{n+7}\} \approx Z_{64} \quad \text{for } n \geq 13.
 \end{aligned}$$

It is easily seen the results of $\pi_{n+12}(K_{\sigma_n} : 2)$ and $\pi_{n+13}(K_{\sigma_n} : 2)$ from the exact sequence of (3. 1), the table (3. 2), (3. 3) and the relation (3. 6).

Consider the exact sequence

$$\pi_{25}(S^{18} : 2) \xrightarrow{\sigma_{11}^*} \pi_{25}(S^{11} : 2) \xrightarrow{i^*} \pi_{25}(K_{\sigma_{11}} : 2) \xrightarrow{\theta} \pi_{24}(S^{18} : 2) \xrightarrow{\sigma_n^*} \pi_{24}(S^{11} : 2)$$

of (3. 1). From (3. 2), (3. 3) it follows that

$$(3. 21) \quad \sigma_{11}^* : \pi_j(S^{18} : 2) \longrightarrow \pi_j(S^{11} : 2) \quad \text{for } j = 24, 25$$

are monomorphisms. Thus from the exactness of the above sequence we have

$$\pi_{25}(K_{\sigma_{11}} : 2) \approx \pi_{25}(S^{11} : 2) / \{\sigma_{11}^2\} = \{\kappa_{11}\} \approx Z_2$$

From (3. 1) and (3. 2), we have the exact sequence

$$\pi_{26}(S^{18} : 2) \xrightarrow{\sigma_{11}^*} \pi_{26}(S^{11} : 2) \xrightarrow{i^*} \pi_{26}(K_{\sigma_{11}} : 2) \longrightarrow 0.$$

From (3. 7) and (3. 2), we have that

3. 22)
$$i_* : \pi_{26}(S^{11} : 2) \longrightarrow \pi_{26}(K_{\sigma_{11}} : 2)$$

an isomorphism onto.

Next consider a generator ν_{10} of $\pi_{13}(S^{10} : 2)$ of order 8 and an element $= d(\iota_{21}) + \gamma$ of $\pi_{19}(S^{10} : 2)$ of order infinite order, where γ is an element $\eta_{20} \circ \varepsilon_{11} + b\nu_{10}^3$ of $\pi_{19}(S^{10} : 2)$ with the order at most 2 ($a, b = 0$ or 1). Let a cell complex $K = S^{11} \cup C(S^{13} \vee S^{19})$ be obtained by attaching $C(S^{13} \vee S^{19})$ to S^{10} by $\nu_{10} \vee \beta : S^{13} \vee S^{19} \longrightarrow S^{10}$. Then we have the following lemma.

LEMMA 3. 3. *We have the following table of homotopy group $\pi_j(K : 2)$ for $j \leq 21$;*

j	$j \leq 9$	10	11	12	13	14	15	16
$\pi_j(K : 2)$	0	Z	Z_2	Z_2	0	Z	Z_2	Z_2
Generator		$i_*\iota_{10}$	$i_*\eta_{10}$	$i_*\eta_{10}^2$		$\widetilde{8\iota_{13}}$	$\widetilde{\eta_{13}}$	$\widetilde{\eta_{13}^2}$
j	17	18	19	20	21			
$\pi_j(K : 2)$	$Z_{16} + Z_4$	$Z_2 + Z_2$	$Z_2 + Z_2$	$Z_2 + Z_2$	Z_{128}			
Generator	$i_*\sigma_{10}, \widetilde{2\nu_{13}}$	$i_*\nu_{10}, i_*\varepsilon_{10}$	$i_*\eta_{10}\varepsilon_{11}, i_*\mu_{10}$	$i_*\sigma_{10}\nu_{17}, i_*\eta_{10}\mu_{11}$	$\sigma_{13} \oplus \eta_{19}$			

Here $i : S^{10} \longrightarrow K$ is an injection and we denote by $\bar{\alpha}$ an element of $\pi_j(K : 2)$ such that $\bar{\alpha}$ is a coextension of $\alpha \in \pi_{j-1}(S^{13} \vee S^{19} : 2)$.

Proof. By (3. 1), we have an exact sequence

3. 23)
$$\begin{aligned} \dots &\longrightarrow \pi_j(S^{13} \vee S^{19} : 2) \xrightarrow{(\nu_{10} \vee \beta)_*} \pi_j(S^{10} : 2) \xrightarrow{i_*} \pi_j(K : 2) \\ &\xrightarrow{\partial} \pi_{j-1}(S^{13} \vee S^{19} : 2) \xrightarrow{(\nu_{10} \vee \beta)_*} \pi_{j-1}(S^{10} : 2) \longrightarrow \dots \end{aligned}$$

or $j \leq 21$. We can identify $\pi_j(S^{13} \vee S^{19} : 2)$ ($(\nu_{10} \vee \beta)_*$ respectively) with $\pi_j(S^{13} : 2) \oplus \pi_j(S^{19} : 2)$ ($\nu_{10*} + \beta_*$ respectively) for $j \leq 21$ and we shall use the notation $\alpha = \nu_{10*} + \beta_*$.

From the tables (3. 2), (3. 3), the relations (3. 6), (3. 8) and the exact sequence (3. 23), it is easy to see the results of $\pi_j(K : 2)$ for $j \neq 17, 21$.

Consider the exact sequence

$$\begin{aligned} \pi_{17}(S^{13} : 2) \oplus \pi_{17}(S^{19} : 2) &\xrightarrow{\alpha} \pi_{17}(S^{10} : 2) \xrightarrow{i_*} \pi_{17}(K : 2) \\ &\xrightarrow{\partial} \pi_{16}(S^{13} : 2) \oplus \pi_{16}(S^{19} : 2) \xrightarrow{\alpha} \pi_{16}(S^{10} : 2) \end{aligned}$$

of (3. 23), where $\pi_{16}(S^{13} : 2) \oplus \pi_{16}(S^{19} : 2) = \pi_{16}(S^{13} : 2) = \{\nu_{13}\} \approx Z_8$ and $\pi_{17}(S^{13} : 2) \oplus \pi_{17}(S^{19} : 2) = 0$ by (3. 2). We have that the homomorphism $\alpha : \pi_{16}(S^{13} : 2) \oplus \pi_{16}(S^{19} : 2) \longrightarrow \pi_{16}(S^{10} : 2)$ is an epimorphism and its kernel is generated by $\{2\nu_{13}\}$. Thus we obtain the following sequence

$$(3. 24) \quad 0 \longrightarrow \{\sigma_{10}\} \xrightarrow{i_*} \pi_{17}(K : 2) \xrightarrow{\partial} \{2\nu_{13}\} \longrightarrow 0.$$

By Adams [1],

$$\{\nu_{10}, 2\nu_{13}, 4\epsilon_{16}\} \equiv 0 \pmod{4\pi_{17}(S^{10} : 2)}$$

and we have, by Proposition 1. 8 of [11], $4\widetilde{2\nu_{13}} = -i_*\{\nu_{10}, 2\nu_{13}, 4\epsilon_{16}\} \in 4i_*\pi_{17}(S^{10} : 2)$. Thus $4(\widetilde{2\nu_{13}} + i_*\alpha) = 0$ for some $\alpha \in \pi_{17}(S^{10} : 2)$. We may replace $\widetilde{2\nu_{13}} + i_*\alpha$ by $\widetilde{2\nu_{13}}$. Thus, from (3. 24), follows that

$$\pi_{17}(K : 2) = \{i_*\sigma_{10}\} + \{\widetilde{2\nu_{13}}\} \approx Z_{16} + Z_4.$$

From (3. 23), we have the exact sequence

$$\begin{aligned} \pi_{21}(S^{13} : 2) \oplus \pi_{21}(S^{19} : 2) &\xrightarrow{\alpha} \pi_{21}(S^{10} : 2) \xrightarrow{i_*} \pi_{21}(K : 2) \\ &\xrightarrow{\partial} \pi_{20}(S^{13} : 2) \oplus \pi_{20}(S^{19} : 2) \xrightarrow{\alpha} \pi_{20}(S^{10} : 2). \end{aligned}$$

By (3. 6), (3. 10) and the diagram (3. 2), (3. 3), we have

$$\alpha\{\sigma_{13}\} = \nu_{10} \circ \sigma_{13} = 2\sigma_{10} \circ \nu_{17} = \mathcal{A}(\epsilon_{21}) \circ \eta_{19} = \alpha\{\eta_{19}\}.$$

Thus we obtain that

$$(3. 25) \quad \text{the kernel of } \alpha : \pi_{20}(S^{13} : 2) \oplus \pi_{20}(S^{19} : 2) \longrightarrow \pi_{20}(S^{10} : 2)$$

is generated by $\{\sigma_{13} \oplus \eta_{19}\} \approx Z_{16}$.

By (3. 8), (3. 10) and the diagram (3. 2),

$$(3. 26) \quad \begin{aligned} \alpha\{\bar{\nu}_{13}\} &= \nu_{10} \circ \bar{\nu}_{13} = 0, \\ \alpha\{\epsilon_{13}\} &= \nu_{10} \circ \epsilon_{13} = 0, \\ \alpha\{\eta_{19}^2\} &= \beta\{\eta_{19}^2\} = \mathcal{A}(\epsilon_{21}) \circ \eta_{19}^2 + a\eta_{10} \circ \epsilon_{11} \circ \eta_{19}^2 + b\nu_{10}^3 \circ \eta_{19}^2 \\ &= 2\sigma_{10} \circ \nu_{18} \circ \eta_{19}^2 + 4a\nu_{10} \circ \epsilon_{13} \\ &= 0. \end{aligned}$$

Thus, from (3. 25), (3. 26) and the from above sequence, it follows that the sequence

$$0 \longrightarrow \{\zeta_{10}\} \xrightarrow{i_*} \pi_{21}(K : 2) \xrightarrow{\theta} \{\sigma_{13} \oplus \eta_{19}\} \longrightarrow 0$$

is exact. By (9. 3) of [11],

$$\zeta_{10} \in \{\nu_{10}, 2\sigma_{13}, 8\iota_{20}\} \pmod{8\pi_{21}(S^{10} : 2)}$$

and by Proposition 1. 3 of [11]

$$\begin{aligned} i_*\zeta_{10} &\in i_*\{\nu_{10}, 2\sigma_{13}, 8\iota_{20}\} \\ &= -8 \widetilde{2\sigma_{13}} \\ &= -16 \widetilde{\sigma_{13} \oplus \eta_{19}}. \end{aligned}$$

Thus we obtain that

$$\pi_{21}(K : 2) = \widetilde{\{\sigma_{13} \oplus \eta_{19}\}} \approx Z_{128}.$$

§ 4. Homotopy groups of exceptional Lie groups E_6, E_7 and E_8 .

(I) HOMOTOPY GROUPS $\pi_j(E_8 : 2)$ for $j \leq 28$.

From Corollary 2. 3, Lemma 2. 5, there exist a cell complex $K_{\tilde{E}_8} = S^{15} \cup_{\sigma_{15}} e^{23} \cup e^{27} \cup e^{29}$ and a continuous map $f : K_{\tilde{E}_8} \longrightarrow \tilde{E}_8$, from which the following isomorphism f_* , induced by a map f , is obtained;

$$(4. 1) \quad f_* : \pi_j(S^{15} \cup_{\sigma_{15}} e^{23} \cup e^{27} \cup e^{29} : 2) \approx \pi_j(\tilde{E}_8 : 2) \quad \text{for } j \leq 28.$$

Let e^{27} be attached to $K_{\sigma_{15}} = S^{15} \cup_{\sigma_{15}} e^{23}$ by a map $g : S^{26} \longrightarrow K_{\sigma_{15}}$ and e^{29} be attached to $S^{15} \cup_{\sigma_{15}} e^{23} \cup e^{27}$ by a map $h : S^{28} \longrightarrow S^{15} \cup_{\sigma_{15}} e^{23} \cup e^{27}$, then, from Corollary 2. 3 and Theorem 2. 6, it follows that the next diagrams are commutative

$$(4. 2) \quad \begin{array}{ccc} (i) & \begin{array}{ccc} S^{26} & \xrightarrow{g} & K_{\sigma_{15}} \\ & \searrow \nu_{23} & \downarrow p \\ & & S^{23} \end{array} & (ii) \quad \begin{array}{ccc} S^{28} & \xrightarrow{h} & S^{15} \cup_{\sigma_{15}} e^{23} \cup e^{27} \\ & \searrow \eta_{27} & \downarrow p' \\ & & S^{27} \end{array} \end{array}$$

where p, p' are the maps which shrink $S^{15}, S^{15} \cup_{\sigma_{15}} e^{23}$ are respectively to a point. From (4. 1),

$$\pi_j(\tilde{E}_8 : 2) \approx \pi_j(S^{15} \cup_{\sigma_{15}} e^{23} \cup e^{27} : 2) \quad \text{for } j \leq 27.$$

Consider the exact sequence

$$\pi_{26}(S^{26} : 2) \xrightarrow{g_*} \pi_{26}(K_{\sigma_{15}} : 2) \xrightarrow{i'_*} \pi_{26}(S^{15} \cup_{\sigma_{15}} e^{23} \cup e^{27} : 2) \xrightarrow{\partial} \pi_{25}(S^{26} : 2)$$

of (3. 1), where $i' : K_{\sigma_{15}} \rightarrow S^{15} \cup_{\sigma_{15}} e^{23} \cup e^{27}$ is the inclusion map. From (i) of (4. 2) and the table (3. 15), we have that

$$(4. 3) \quad g_* : \pi_{26}(S^{26} : 2) \rightarrow \pi_{26}(K_{\sigma_{15}} : 2)$$

is an epimorphism. Thus, from the exactness of the above sequence, we obtain

$$(4. 4) \quad \pi_{26}(S^{15} \cup_{\sigma_{15}} e^{23} \cup e^{27} : 2) = 0.$$

It follows from (3. 1), (3. 15) and (4. 3) that the sequence

$$0 = \pi_{27}(K_{\sigma_{15}} : 2) \xrightarrow{i'_*} \pi_{27}(S^{15} \cup_{\sigma_{15}} e^{23} \cup e^{27} : 2) \xrightarrow{\partial} \pi_{26}(S^{26} : 2) \xrightarrow{g_*} \pi_{26}(K_{\sigma_{15}} : 2) \rightarrow 0$$

is exact. Thus we obtain

$$(4. 5) \quad \pi_{27}(S^{15} \cup_{\sigma_{15}} e^{23} \cup e^{27} : 2) = Z.$$

Next consider the diagram;

$$\begin{array}{ccccccc} & & \pi_{28}(S^{15} \cup_{\sigma_{15}} e^{23} : 2) = 0 & & & & \\ & & \downarrow i'_* & & & & \\ \pi_{28}(S^{28} : 2) & \xrightarrow{h_*} & \pi_{28}(S^{15} \cup e^{23} \cup e^{27} : 2) & \xrightarrow{i''_*} & \pi_{28}(S^{15} \cup e^{23} \cup e^{27} \cup e^{29} : 2) & \xrightarrow{\partial} & \pi_{27}(S^{28} : 2) = 0 \\ \downarrow \eta_{27*} & \swarrow \eta'_* & \downarrow \partial & & & & \\ \pi_{28}(S^{27} : 2) & \xleftarrow{E} & \pi_{27}(S^{26} : 2) & & & & \\ & & \downarrow g_* & & & & \\ & & \pi_{27}(S^{15} \cup_{\sigma_{15}} e^{23} : 2) = 0 & & & & \end{array}$$

where i'' is a inclusion map. From (3. 1) the row and column sequences are exact, and from (ii) of (4. 2) and from the definition of ∂ , it follows that the diagram is commutative. By (3. 15), $\partial : \pi_{28}(S^{15} \cup e^{23} \cup e^{27} : 2) \rightarrow \pi_{27}(S^{26} : 2)$ is an isomorphism, and $E : \pi_{27}(S^{26} : 2) \rightarrow \pi_{28}(S^{27} : 2)$ is an isomorphism. Thus, from the commutativity of the above diagram, it follows that

$$h_* : \pi_{28}(S^{28} : 2) \rightarrow \pi_{28}(S^{15} \cup_{\sigma_{15}} e^{23} \cup e^{27} : 2)$$

is epimorphic. Thus, from the exactness of the column sequence, we obtain

$$(4.6) \quad \pi_{28}(S^{15} \cup_{\sigma_{15}} e^{23} \cup e^{27} \cup e^{29} : 2) = 0.$$

From (4.1), (3.15) and (4.4)–(4.9), it follows the next table of the homotopy groups of exceptional Lie group E_8 .

PROPOSITION 4.1.

j	1, 2	3	$4 \leq j \leq 14$	15	16	17	18	19	20
$\pi_j(E_8 : 2)$	0	Z	0	Z	Z_2	Z_2	Z_8	0	0

j	21	22	23	24	25	26	27	28
$\pi_j(E_8 : 2)$	Z_2	0	$Z + Z_2$	$Z_2 + Z_2$	Z_2	0	Z	0

(II) HOMOTOPY GROUPS $\pi_j(E_7 : 2)$ for $j \leq 25$.

From Lemma 2.5, there exist a cell complex $K_{\tilde{E}_7} = S^{11} \cup e^{15} \cup e^{19} \cup e^{23} \cup e^{26} \cup e^{27}$ and a continuous map $k : K_{\tilde{E}_7} \rightarrow \tilde{E}_7$ such that $k_* : \pi_j(K_{\tilde{E}_7}) \rightarrow \pi_j(\tilde{E}_7)$ are C_2 -isomorphism onto for $j \leq 28$. By Corollary 2.8 and Lemma 2.4, e^{15} is attached to S^{11} by a representative of $\nu_{11} \in \pi_{14}(S^{11} : 2)$.

Consider the diagram

$$\begin{array}{ccc}
 S^{11} \cup e^{15} \cup e^{19} & \longrightarrow & S^{15} \cup e^{19} \\
 \nu_{11} \downarrow k & & \downarrow \bar{k} \swarrow f \\
 \tilde{E}_7 & \subset & \tilde{E}_8 \quad S^{15}
 \end{array}$$

where p is a map which shrinks S^{11} to a point and $\tilde{E}_7 \subset \tilde{E}_8$ is the natural inclusion. Since $\pi_i(\tilde{E}_8) = 0$ for $i \leq 14$, $k|S^{11} \simeq 0$ in E_8 . Thus there exists a map $\bar{k} : S^{15} \cup e^{19} \rightarrow \tilde{E}_8$ such that the above diagram is homotopy commutative. A generator $x_{15} \in H^{15}(\tilde{E}_7 : Z_2)$ corresponds to a generator $x_{15} \in H^{15}(\tilde{E}_8 : Z_2)$ by the natural inclusion $\tilde{E}_7 \subset \tilde{E}_8$. Thus, from the commutativity of the above diagram, $x_{15} \in H^{15}(\tilde{E}_8 : Z_2)$ corresponds to a generator of $H^{15}(S^{15} \cup e^{19} : Z_2)$ by \bar{k}^* . Let $f : S^{15} \rightarrow \tilde{E}_8$ be a representative of a generator $\{f\}$ of $\pi_{15}(\tilde{E}_8) = Z$, then $\bar{k}|S^{15}$ is homotopic to $x\{f\}$ for some odd integer x . Let e^{19} be attached to S^{15} by $\beta : S^{15} \rightarrow S^{15}$ for a cell complex $S^{15} \cup e^{19}$ of the above diagram.

Since \bar{k} is extended over e^{19} , we have

$$0 = (\bar{k}|S^{15})_*\beta = x(f_*\beta) \quad \text{in 2-component.}$$

By (4. 1), $f_* : \pi_j(S^{15}) \longrightarrow \pi_j(\tilde{E}_8)$ are C_2 -isomorphism onto for $j \leq 21$. Thus it follows $\beta = 0$. From this we have that $S^{11} \cup e^{19}$ is a subcomplex of $K_{\tilde{E}_7}$, and e^{19} is attached to S^{11} by σ_{11} .

LEMMA 4. 2. *We may regard the inclusion $j : K_{\sigma_{11}} = S^{11} \cup_{\sigma_{11}} e^{19} \subset K_{\tilde{E}_7}$ as the fibre map. Let F be the fibre, then $H^*(F ; Z_2)$ has additive basis $\{1, a_{14}, a_{22}, a_{26}\}$ for degree < 29 , where a_i denote a generator of degree i .*

Proof. From lemma 2. 5, $H^*(K_{\tilde{E}_7} ; Z_2) = \Delta(x_{11}, x_{15}, x_{19}, x_{23}, x_{27})$ for degree < 30 and $Sq^4 x_{11} = x_{15}$, $Sq^8 x_{15} = x_{23}$, $Sq^4 x_{23} = x_{27}$, $Sq^8 x_{11} = x_{19}$. Let $\{E_r^{**}\}$ be the mod 2 spectral sequence associated with the above fibering, then we have

$$E_2^{**} = H^*(K_{\tilde{E}_7} ; Z_2) \otimes H^*(F ; Z_2)$$

and

$$E_\infty^{**} = \Delta(x_{11}, x_{19}) \text{ for degree } < 30.$$

Clearly $K_{\tilde{E}_7}$ and F are 10-and 13-connected respectively. We have the following cohomology exact sequence $\cdots \longrightarrow H^*(K_{\tilde{E}_7} ; Z_2) \xrightarrow{j^*} H^*(K_{\sigma_{11}} ; Z_2) \longrightarrow H^*(F ; Z_2) \xrightarrow{\tau} H^*(K_{\tilde{E}_7} ; Z_2) \longrightarrow \cdots$ for degree ≤ 24 . It follows that $H^*(F ; Z_2) = \{1, a_{14}, a_{22}\}$ for degree < 24 where $\tau(a_{14}) = x_{15}$ and $\tau(a_{22}) = x_{23}$, i.e., $d_{15}(1 \otimes a_{14}) = x_{15} \otimes 1$ and $d_{23}(1 \otimes a_{22}) = x_{23} \otimes 1$. For $24 \leq q \leq 29$, any non-zero element of $E_2^{0, q}$ must be cancelled by d_r with some element of $E_r^{r, q-r+1}$. By the dimensional reason, the only possibilities of such q are $q = 24, 25, 26$ corresponding to $x_{11} \otimes a_{14}$, $x_{11}x_{15} \otimes 1$ and $x_{27} \otimes 1$ respectively. Thus $H^q(F ; Z_2) = 0$ for $q = 27, 28, 29$. Since $d_{15}(x_{11} \otimes a_{14}) = x_{11}x_{15} \otimes 1 \neq 0$, $x_{11} \otimes a_{14}$ is not a d_{15} -image, hence $H^{24}(F ; Z_2) = 0$. We have also $H^{25}(F ; Z_2) = 0$ since $x_{11}x_{15} \otimes 1 = 0$ in $E_6^{26, 0}$. By the dimensional reason, we see that $x_{27} \otimes 1 \neq 0$ in $E_{27}^{27, 0}$, hence there exists an element a_{26} such that $d_{28}(1 \otimes a_{26}) = x_{27} \otimes 1$ and a_{26} generates $H^{26}(F ; Z_2) \approx Z_2$.

From the proof of this lemma, we have that a_{14}, a_{22}, a_{26} are transgressive elements. Since $Sq^8 x_{15} = x_{23}$, $Sq^4 x_{23} = x_{27}$, it follows, from the commutativity of the Steenrod operation and the transgression, that

$$(4. 7) \quad Sq^8 a_{14} = a_{22}, \quad Sq^4 a_{22} = a_{26}.$$

By Lemma 2. 5 and Theorem 2. 6, there exists a cell complex $K_F = S^{14} \cup e^{22} \cup e^{26}$ and a continuous map from K_F to F which induces

isomorphisms from $\pi_j(K_F : 2)$ onto $\pi_j(F : 2)$ for $j \leq 26$. Let $f : K_F \rightarrow K_{\sigma_{11}} = S^{11} \cup e^{19}$ be the mapping from a fibre to the total space identifying F with K_F for dimension ≤ 26 . Then $f|S^{14}$ is a representative of ν_{11} .

Consider the exact sequence

$$(4.8) \quad \begin{array}{ccccccc} \cdots & \longrightarrow & \pi_j(K_F : 2) & \xrightarrow{f_*} & \pi_j(K_{\sigma_{11}} : 2) & \xrightarrow{j_*} & \pi_j(K_{E_7} : 2) \\ & & & & \xrightarrow{\partial} & & \\ & & \pi_{j-1}(K_F : 2) & \xrightarrow{f_*} & \pi_{j-1}(K_{\sigma_{11}} : 2) & \longrightarrow & \cdots \end{array}$$

associated with the above fibering for $j \leq 26$ and the following homotopy commutative diagram

$$(4.9) \quad \begin{array}{ccc} S^{14} & \xrightarrow{i} & K_F \\ \downarrow \nu_{11} & & \downarrow f \\ S^{11} & \xrightarrow{i} & K_{\sigma_{11}}. \end{array}$$

From (3. 1), (3. 14) and from the fact that e^{26} is attached to $K_{\sigma_{14}}$ by a coextension of ν_{22} , we have the next table ;

(4. 10)

j	$j \leq 13$	14	15	16	17	18	19	20
$\pi_j(K_F : 2)$	0	Z	Z_2	Z_2	Z_8	0	0	Z_2
Generator		$i_*\iota_{14}$	$i_*\eta_{14}$	$i_*\eta_{14}^2$	$i_*\nu_{14}$			$i_*\nu_{14}^2$

j	21	22	23	24	25	26
$\pi_j(K_F : 2)$	0	$Z + Z_2$	$Z_2 + Z_2$	Z_2	0	Z
Generator		$\widetilde{16}\iota_{21}, i_*\epsilon_{14}$	$i_*\mu_{14}, i_*\eta_{14}\epsilon_{15}$	$i_*\eta_{14}\mu_{15}$		$\widetilde{64}\iota_{25}$

LEMMA 4. 3. For the homomorphism $f_* : \pi_j(K_F : 2) \rightarrow \pi_j(K_{\sigma_{11}} : 2)$, we have the following table ;

(4. 11)


$\alpha =$	$i_*\iota_{14}$	$i_*\eta_{14}$	$i_*\eta_{14}^2$	$i_*\nu_{14}$	$i_*\nu_{14}^2$	$i_*\widetilde{16}\iota_{21}$	$i_*\epsilon_{14}$	$i_*\mu_{14}$	$i_*\eta_{14}\epsilon_{15}$	$i_*\eta_{14}\mu_{15}$
$f_*\alpha =$	$i_*\nu_{14}$	0	0	$i_*\nu_{14}^2$	$i_*\eta_{11}\epsilon_{12}$	$4\widetilde{2}\nu_{18}$	0	0	0	0

Proof. From (4. 9), (3. 8), (3. 9), it follows that the table is true excepting for $\alpha = i_*\widetilde{16}\iota_{21}, i_*\nu_{14}^2$.

The relation $i_*\eta_{11} \circ \varepsilon_{12} = i_*\nu_{11}^3$ in $\pi_{20}(K\sigma_{11} : 2)$ imply the formula

$$f_*(i_*\nu_{14}^2) = i_*\eta_{11} \circ \varepsilon_{12}.$$

Consider the following commutative diagram



where $\widetilde{16\iota_{21}}$ is a coextension of $16\iota_{21}$ and $\bar{\nu}_{11}$ is an extension of ν_{11} . We have

$$\begin{aligned}
 f_*\widetilde{16\iota_{21}} &= i_*\bar{\nu}_{11} \circ \widetilde{16\iota_{21}} \\
 &= -i_*\{\nu_{11}, \sigma_{14}, 16\iota_{21}\} \text{ by Proposition 1. 8 of [11],} \\
 &= -i_*\zeta_{11} \text{ by (9. 3) of [11],} \\
 &= -4 \widetilde{2\nu_{21}}.
 \end{aligned}$$

PROPOSITION 4. 4. *The homotopy groups $\pi_j(E_7 : 2)$ for $j \leq 25$ are listed in the following table;*

j	1, 2	3	$4 \leq j \leq 10$	11	12	13	14	15
$\pi_j(E_7 : 2)$	0	Z	0	Z	Z_2	Z_2	0	Z

j	16	17	18	19	20	21	22	23
$\pi_j(E_7 : 2)$	Z_2	Z_2	Z_4	$Z + Z_2$	Z_2	Z_2	Z_4	$Z + Z_2 + Z_2$

j	24	25
$\pi_j(E_7 : 2)$	$Z_2 + Z_2 + Z_2$	$Z_2 + Z_2$

Proof. The results of $\pi_j(E_7 : 2)$ for $j \leq 22$ follow immediately from the tables (4. 10), (3. 13), (4. 11) and from the exactness of the sequence of (4. 8).

$$\{\nu_{11}, \varepsilon_{14}, 2\iota_{22}\} \supset E^4\{\nu_7, \varepsilon_{10}, 2\iota_{18}\} \subset E^4\pi_{19}(S^7) = 0.$$

Thus we have

$$(4.12) \quad \{\nu_{11}, \varepsilon_{14}, 2\iota_{22}\} \equiv 0 \pmod{2\pi_{23}(S^{11})}.$$

Similarly we have

$$(4.13) \quad \{\nu_{11}, \mu_{14}, 2\iota_{23}\} \equiv 0 \pmod{2\pi_{24}(S^{11})}.$$

$\{\nu_{11}, \eta_{14} \circ \varepsilon_{15}, 2\iota_{23}\} \supset \{\nu_{11} \circ \eta_{14}, \varepsilon_{15}, 2\iota_{23}\} = \{0, \varepsilon_{15}, 2\iota_{23}\} \equiv 0$ by Proposition 1. 2 of [11]. Thus we have

$$(4.14) \quad \{\nu_{11}, \eta_{14} \circ \varepsilon_{15}, 2\iota_{23}\} \equiv 0 \pmod{2\pi_{24}(S^{11} : 2)}.$$

Similarly,

$$(4.15) \quad \{\nu_{11}, \eta_{14} \circ \mu_{15}, 2\iota_{24}\} \equiv 0 \pmod{2\pi_{25}(S^{11} : 2)}.$$

Consider the commutative diagram

$$(4.16) \quad \begin{array}{ccccccccc} \pi_j(K_F : 2) & \xrightarrow{f_*} & \pi_j(K_{\sigma_{11}} : 2) & \xrightarrow{j_*} & \pi_j(K_{\tilde{E}_7} : 2) & \xrightarrow{\partial} & \pi_{j-1}(K_F : 2) & \xrightarrow{f_*} & \pi_{j-1}(K_{\sigma_{11}} : 2) \\ \uparrow i_* & & \uparrow i_* & & \uparrow i_* & & \uparrow i_* & & \uparrow i_* \\ \pi_j(S^{14} : 2) & \xrightarrow{\nu_{11,*}} & \pi_j(S^{11} : 2) & \xrightarrow{i_*} & \pi_j(S^{11} \cup e^{15} : 2) & \xrightarrow{\partial} & \pi_{j-1}(S^{14} : 2) & \xrightarrow{\nu_{11,*}} & \pi_{j-1}(S^{11} : 2) \end{array}$$

where i, j are inclusions.

From Proposition 1. 8 of [11] and the above secondary composition, coextension $\tilde{\varepsilon}_{14}, \tilde{\mu}_{14}, \widetilde{\eta_{14} \circ \varepsilon_{15}}$ and $\widetilde{\eta_{14} \circ \mu_{15}}$ of $\varepsilon_{14}, \mu_{14}, \eta_{14} \circ \varepsilon_{15}$ and $\eta_{14} \circ \mu_{15}$ respectively are elements of order 2. Thus from the commutativity and the exactness of the above diagram. (4. 16), the results of $\pi_j(K_{\tilde{E}_7} : 2)$ for $j = 23, 24, 25$, are obtained.

(III) HOMOTOPY GROUPS $\pi_j(E_6 : 2)$ for $j \leq 22$.

By Corollary 2. 3,

$$H^*(\tilde{E}_6 ; Z_2) = Z_2[y_{32}] \otimes \mathcal{A}(y_9, y_{11}, y_{15}, y_{17}, y_{23}, y_{33})$$

and

$$Sq^2 y_9 = y_{11}, Sq^8 y_9 = y_{17}, Sq^4 y_{11} = y_{15}, Sq^8 y_{15} = y_{23}.$$

From Lemma 2. 5, there exists a cell complex $K_{\tilde{E}_6}$ and a continuous

map $l : K\tilde{E}_6 \rightarrow \tilde{E}_6$ such that $l_* : \pi_j(K\tilde{E}_6) \rightarrow \pi_j(\tilde{E}_6)$ are C_2 -isomorphism onto for $j \leq 24$, i.e, $K\tilde{E}_6 = S^9 \cup e^{11} \cup e^{15} \cup e^{17} \cup e^{20} \cup e^{23} \cup e^{24}$.

By Corollary 2. 8, e^{11} is attached to S^9 by γ_9 .

LEMMA 4. 5. $K\sigma_9 = S^9 \cup_{\sigma_9} e^{17}$ is a subcomplex of $K\tilde{E}_6$. Exchanging an inclusion map $K\sigma_9 \rightarrow K\tilde{E}_6$ by a fibre map, we denote by F the fibre of this fibering. Then $H^*(F ; Z_2)$ has the additive basis $\{1, a_{10}, a_{14}, a_{20}, a_{22}\}$ for degree ≤ 25 such that $Sq^4 a_{10} = a_{14}$, $Sq^8 a_{14} = a_{22}$, where a_i denotes a generator of degree i .

Proof. From Lemma 2. 5, $H^*(K\tilde{E}_6 ; Z_2) = \mathcal{A}(x_9, x_{11}, x_{15}, x_{17}, x_{23})$ for degree < 32 and $Sq^2 x_9 = x_{11}$, $Sq^4 x_{11} = x_{15}$, $Sq^8 x_{15} = x_{23}$, $Sq^8 x_9 = x_{17}$.

By use of Adem's relation we have relations

$$\begin{aligned} Sq^6 x_{11} &= Sq^6 Sq^2 x_9 = Sq^4 Sq^4 x_9 + Sq^7 Sq^1 x_9, \\ Sq^2 x_{15} &= Sq^2 Sq^4 x_{11} = Sq^5 Sq^1 x_{11} + Sq^6 x_{11}. \end{aligned}$$

Since there is no cell of dimension 10 and 13, $Sq^6 x_{11} = 0$ in $K\tilde{E}_6$. Since there is no cell of dimension 12 and $Sq^6 x_{11} = 0$, $Sq^2 x_{15} = 0$ in $K\tilde{E}_6$. Then e^{17} is inessential to e^{15} , that is, up to homotopy type $S^9 \cup e^{11} \cup e^{17}$ is a subcomplex. Since $\pi_{16}(S^9 \cup e^{11}, S^9) \approx \pi_{16}(S^{11}) = 0$, we have that $S^9 \cup e^{17}$ is a subcomplex. Then, by Theorem 2. 6, we may consider that $S^9 \cup e^{17} = K\sigma_9$ is a subcomplex of $K\tilde{E}_6$.

Let $\{E_r^{**}\}$ be the mod 2 spectral sequence associated with a fibering $\{K\sigma_9, i, K\tilde{E}_6\}$ with the fibre F , then

$$E_2^{**} = H^*(K\tilde{E}_6 ; Z_2) \otimes H^*(F ; Z_2)$$

and

$$E_\infty^{**} = \wedge (x_9, x_{17}) \quad \text{for degree } \leq 25.$$

By concerning the cohomology exact sequence associated with this fibering, we have $H^*(F ; Z_2) = \{1, a_{10}, a_{14}\}$ for degree < 18 with generator a_{10}, a_{14} such that $d_{11}(1 \otimes a_{10}) = x_{11} \otimes 1$ and $d_{15}(1 \otimes a_{14}) = x_{15} \otimes 1$. For the total degree < 27 , E_2^{**} is the sum of $'E_2^{**} = H^*(K\tilde{E}_6 ; Z_2) \otimes \{1, a_{10}, a_{14}\}$ and $\sum_{q \geq 18} 1 \otimes H^q(F ; Z_2)$. From $'E_2^{**}$ we compute $'E_r^{**}$ giving d_r trivially except $d_r(b \otimes a_{10}) = b x_{11} \otimes 1$ and $d_r(b \otimes a_{14}) = b x_{15} \otimes 1$, $b \in H^*(K\tilde{E}_6 ; Z_2)$. Then we have for the total degree < 30 , $'E_\infty^{**} = \mathcal{A}(x_9, x_{17}, x_{23}) \otimes 1 + \{x_{11} \otimes a_{10}, x_{15} \otimes a_{14}\}$, where we use the fact $x_{11}^2 = x_{15}^2 = 0$. Compare this with E_∞^{**} , we conclude that $x_{23} \otimes 1, x_{11} \otimes a_{10}$ must be cancelled by some elements a_{22}, a_{20} , i.e,

$d_{23}(1 \otimes a_{22}) = x_{23} \otimes 1$ and $d_{11}(1 \otimes a_{20}) = x_{11} \otimes a_{10}$. Moreover, no other non-zero elements exists in $H^*(F; Z_2)$ for degree ≤ 25 . Thus $H_*(F; Z_2) = \{1, a_{10}; a_{14}, a_{20}, a_{22}\}$ for degree ≤ 25 .

From the above proof, a_{10}, a_{14} and a_{22} are transgressive element. Since $Sq^4 a_{11} = x_{15}$ and $Sq^8 x_{15} = x_{23}$, using the commutativity of Steenrod operation and transgression we have $Sq^4 a_{10} = a_{14}$ and $Sq^8 a_{14} = a_{22}$.

By Lemma 2.5, there exists a cell complex $K_F = S^{10} \cup e^{14} \cup e^{20} \cup e^{22}$ and a continuous map which induce C_2 -isomorphisms from $\pi_j(K_F)$ to $\pi_j(F)$ for $j \leq 24$. We identify the fiber to the total space, then we have a commutative diagram

$$(4.17) \quad \begin{array}{ccc} S^{10} & \xrightarrow{i} & K_F \\ \downarrow \eta_9 & & \downarrow f \\ S^9 & \xrightarrow{i} & K_{\sigma_9} \end{array}$$

where i is inclusion map, and the exact sequence

$$(4.18) \quad \begin{array}{ccccccc} \dots & \longrightarrow & \pi_j(K_F : 2) & \longrightarrow & \pi_j(K_{\sigma_9} : 2) & \longrightarrow & \pi_j(K\tilde{E}_6 : 2) \\ & & \longrightarrow & \longrightarrow & \longrightarrow & \longrightarrow & \dots \end{array}$$

Consider the cell complex $K_F = S^{10} \cup e^{14} \cup e^{20} \cup e^{22}$. Since $Sq^4 a_{10} = a_{14}$, e^{14} is attached to S^{10} by a representative of ν_{10} .

From $\pi_{19}(S^{10} \cup e^{14}, S^{10}) \approx \pi_{18}(S^{13}) = 0$, we may assume that $K_F = S^{10} \cup C(S^{13} \vee S^{19}) \cup e^{22}$.

Let $\alpha : S^{21} \longrightarrow S^{10} \cup C(S^{13} \vee S^{19})$ be the attaching map of e^{22} and e^{20} be attached to S^{10} by $\beta : S^{19} \longrightarrow S^{10}$. Consider the exact sequence

$$\pi_{21}(S^{10} : 2) \longrightarrow \pi_{21}(S^{10} \cup e^{14} \cup e^{20} : 2) \xrightarrow{\partial} \pi_{20}(S^{13} \vee S^{19} : 2) \xrightarrow{(\nu_{10} \vee \beta)_*} \pi_{20}(S^{10} : 2).$$

From the definition of ∂ , we have the commutative diagram

$$\begin{array}{ccc} \pi_{21}(S^{10} \cup e^{14} \cup e^{20} : 2) & \xrightarrow{\partial} & \pi_{20}(S^{13} \vee S^{19} : 2) \\ \searrow p_* & & \swarrow E \\ \pi_{21}(S^{14} \vee S^{20} : 2) & = & \pi_{21}(S^{14} : 2) + \pi_{21}(S^{20} : 2) \end{array}$$

where p is a map which shrinks S^{10} to a point. Since $Sq^8 a_{14} = a_{22}$, $p_* \alpha = \sigma_{14} + x \eta_{20}$ for $x = 1$ or 0 . From the exactness of the above sequence, $0 = (\nu_{10} \vee \beta)_* \circ \partial \alpha = \nu_{10} \circ \sigma_{13} + x(\beta \circ \eta_{19})$. Thus we have $x(\beta \circ \eta_{19}) = \nu_{10} \circ \sigma_{13} \neq 0$ and $x = 1$.

Put $\beta = a\mathcal{A}(\iota_{21}) + b\eta_{10} \circ \varepsilon_{11} + c\nu_{11}^3 + d\mu_{10}$ for some integers a, b, c, d , then we have

$$\begin{aligned} \nu_{10} \circ \sigma_{13} &= \beta \circ \eta_{19} \\ &= a(\mathcal{A}(\iota_{21})) \circ \eta_{19} + b\eta_{10}^2 \circ \varepsilon_{12} + c\nu_{10}^3 \circ \eta_{19} + d\eta_{10} \circ \mu_{11} \\ &= a\nu_{10} \circ \sigma_{13} + 0 + 0 + d\eta_{10} \circ \mu_{11} \quad \text{by (3. 6) and (3. 10).} \end{aligned}$$

Thus by (3. 3) $a = 1$ and $d = 0$. Therefore

$$(4. 19) \quad \begin{aligned} \beta &= \mathcal{A}(\iota_{21}) + b\eta_{10} \circ \varepsilon_{11} + c\nu_{10}^3 \quad \text{where } b, c = 0 \text{ or } 1. \\ \partial\alpha &= \sigma_{13} + \eta_{19} \end{aligned}$$

From (4. 19), Lemma 3. 3 and from the exact sequence

$$\dots \longrightarrow \pi_j(S^{21} : 2) \xrightarrow{\alpha_*} \pi_j(S^{10} \cup e^{14} \cup e^{20} : 2) \longrightarrow \pi_j(K_F : 2) \longrightarrow \pi_{j-1}(S^{21} : 2) \xrightarrow{\alpha_*} \dots$$

of (3. 1), we have the next table;

(4. 20)

j	$j \leq 9$	10	11	12	13	14	15	16	17
$\pi_j(K_F : 2)$	0	Z	Z_2	Z_2	0	Z	Z_2	Z_2	$Z_{16} + Z_4$
Generator		$i_*\iota_{10}$	$i_*\eta_{10}$	$i_*\eta_{10}^2$		$\widetilde{8\iota_{13}}$	$\widetilde{\eta_{13}}$	$\widetilde{\eta_{13}^2}$	$i_*\sigma_{10}, \widetilde{2\nu_{13}}$

j	18	19	20	21
$\pi_j(K_F : 2)$	$Z_2 + Z_2$	$Z_2 + Z_2$	$Z_2 + Z_2$	0
Generator	$i_*\overline{\nu_{10}}, i_*\varepsilon_{10}$	$i_*\eta_{10}\varepsilon_{11}, i_*\mu_{10}$	$i_*\sigma_{10}\nu_{17}, i_*\eta_{10}\mu_{11}$	

LEMMA 4. 6. For the homomorphism $f_* : \pi_j(K_F : 2) \longrightarrow \pi_F(K_{\sigma_9} : 2)$, we have the following table;

(4. 21)

$\alpha =$	$i_*\iota_{10}$	$i_*\eta_{10}$	$i_*\eta_{10}^2$	η_{13}	$i_*\sigma_{10}$	$2\nu_{13}$	$i_*\nu_{10}$	$i_*\varepsilon_{10}$
$f_*\alpha =$	$i_*\eta_9$	$i_*\eta_9^2$	$4i_*\nu_9$	$i_*\nu_9^2$	$i_*\varepsilon_9 + i_*\overline{\nu_9}$	$i_*\varepsilon_9$	$i_*\nu_9^3$	$i_*\eta_9\varepsilon_{10}$


$\alpha =$	$i_*\eta_{10} \circ \varepsilon_{11}$	$i_*\mu_{10}$	$i_*\sigma_{10}\nu_{17}$	$i_*\eta_{10}\mu_{11}$
$f_*\alpha =$	0	$i_*\eta_9\mu_{10}$	$i_*\nu_9\nu_{17}$	$4i_*\zeta_9$

Proof. We shall use the next relations

$$\begin{aligned}
 \eta_n^3 &= 4\nu_n & \text{for } n \geq 5 & & \text{by (5. 5) of [11],} \\
 \eta_n \circ \bar{\nu}_{n+1} &= \nu_n^3 & \text{for } n \geq 6 & & \text{by Lemma 6. 3 of [11],} \\
 \eta_9 \circ \sigma_{10} &= \bar{\nu}_9 + \varepsilon_9 & & & \text{by Lemma 6. 4 of [11],} \\
 \eta_n^2 \circ \varepsilon_{n+2} &= 0 & \text{for } n \geq 9 & & \text{by (7. 10), (7. 20) of [11],} \\
 4\zeta_n &= \eta_n^2 \circ \mu_{n+2} & \text{for } n \geq 5 & & \text{by Lemma 6. 7 of [11].}
 \end{aligned}$$

From (4. 17), (4. 22), it follows that the table is true except for $\alpha = \widetilde{\eta}_{13}$ and $\widetilde{2\nu}_{13}$.


From the definition of $\widetilde{\eta}_{13}$ and (4. 17), we have the commutative diagram



where p is the mapping which shrinks S^{10} to a point and $\overline{\eta_9}$ is a extension of η_9 . Thus we have

$$f_*\widetilde{\eta}_{13} = i_*\overline{\eta_9} \circ \widetilde{\eta}_{13} = i_*\{\eta_9, \nu_{10}, \eta_{13}\} \ni i_*\nu_9^2 \quad \text{by Lemma 5. 5 of [11].}$$

Consider the commutative diagram



then we have

$$\begin{aligned}
 f_*\widetilde{2\nu}_{13} &= i_*\overline{\eta_9} \circ \widetilde{2\nu}_{13} \in i_*\{\eta_9, \nu_{10}, 2\nu_{13}\} & \text{by Proposition 1. 7 of [11],} \\
 &\in i_*\varepsilon_9 & \text{by (6. 1) of [11].}
 \end{aligned}$$

PROPOSITION 4. 7. *The homotopy groups $\pi_j(E_6 : 2)$ for $j \leq 22$ are listed in the following table;*

j	1, 2	3	$4 \leq j \leq 8$	9	10	11	12	13	14
$\pi_j(E_6 : 2)$	0	Z	0	Z	0	Z	Z_4	0	0

j	15	16	17	18	19	20	21	22
$\pi_j(E_6 : 2)$	Z	0	$Z + Z_2$	$Z_{16} + Z_2$	0	Z_8	0	0

Proof. The results of $\pi_j(E_6 : 2)$ for $j \neq 18, 20$, follow immediately from the table the (3. 11), (4. 20), (4. 21) and from the exact sequence (4. 18).

By (3. 9) and Proposition 1. 2 of [11], $\mu \in \langle \eta, 8\iota, 2\sigma \rangle \cong \langle \eta, 2\sigma, 8\iota \rangle + \langle 2\sigma, \eta, 8\iota \rangle$ and $\langle 2\sigma, \eta, 8\iota \rangle \cong \langle \sigma, 2\eta, 8\iota \rangle \cong 0$. Then, by concerning the suspension homomorphism, we obtain

$$\{\eta_9, 2\sigma_{10}, 8\iota_{17}\} \ni \mu_9.$$

By Lemma 9. 1 of [11], we have

$$\{\eta_9, \eta_{10} \circ \varepsilon_{11}, 2\iota_{19}\} \ni \zeta_9.$$

Consider the commutative diagram

$$\begin{array}{ccccccccc}
 \pi_{18}(K_F : 2) & \xrightarrow{f_*} & \pi_{18}(K_{\sigma_9} : 2) & \xrightarrow{j_*} & \pi_{18}(KE_6 : 2) & \xrightarrow{\partial} & \pi_{17}(K_F : 2) & \xrightarrow{f_*} & \pi_{17}(K_{\sigma_9} : 2) \\
 \uparrow i_* & & \uparrow i_* & & \uparrow i_* & & \uparrow i_* & & \uparrow i_* \\
 \pi_{18}(S^{10} : 2) & \xrightarrow{\eta_{9*}} & \pi_{18}(S^9 : 2) & \xrightarrow{j_*} & \pi_{18}(S^9 \cup e^{11} : 2) & \xrightarrow{\partial} & \pi_{17}(S^{10} : 2) & \xrightarrow{\eta_{9*}} & \pi_{17}(S^9 : 2)
 \end{array}$$

where j is a inclusion map $S^9 \longrightarrow S^9 \cup e^{11}$.

By Proposition 1. 8 of [11], we have

$$j_*\mu_9 \in j_*\{\eta_9, 2\sigma_{10}, 8\iota_{17}\} = -8 \widetilde{2\sigma_{10}}.$$

From the above commutative diagram and from the tables (3. 11), (4. 20), (4. 21), we obtain

$$\pi_{18}(K_{\widetilde{E}_6} : 2) \approx Z_{16} + Z_2.$$

We have the following commutative diagram

$$\begin{array}{ccccccccc}
 \pi_{20}(K_F : 2) & \xrightarrow{f_*} & \pi_{20}(K_{\sigma_9} : 2) & \longrightarrow & \pi_{20}(K_{E_6} : 2) & \longrightarrow & \pi_{19}(K_F : 2) & \xrightarrow{f_*} & \pi_{19}(K_{\sigma_9} : 2) \\
 \uparrow \tilde{i}_* & & \uparrow i_* & & \uparrow i_* & & \uparrow i_* & & \uparrow i_* \\
 \pi_{20}(S^{10} : 2) & \xrightarrow{\eta_{9^*}} & \pi_{20}(S^9 : 2) & \xrightarrow{j_*} & \pi_{20}(S^9 \cup e^{11} : 2) & \xrightarrow{\partial} & \pi_{19}(S^{10} : 2) & \xrightarrow{\eta_{9^*}} & \pi_{19}(S^9 : 2)
 \end{array}$$

and from Proposition 1. 7 of [11]

$$j_*\zeta_9 \in j_*\{\eta_9, \eta_{10} \circ \varepsilon_{11}, 2\iota_{19}\} = -2\widetilde{\eta_{10} \circ \varepsilon_{11}}.$$

From the exact sequence (4. 18) and from the table (3. 8), (4. 10), (4. 21), we obtain

$$\pi_{20}(K_{\widetilde{E}_6} : 2) \approx Z_8.$$

BIBLIOGRAPHY

- [1] J. F. Adams: On the group $J(X)$ IV, *Topology*, **5-1** (1966), 21-71.
- [2] S. Araki: Cohomology modulo 2 of the compact exceptional groups E_6 and E_7 , *J. of Math. Osaka C.V.*, Vol. **12** (1961), 43-65.
- [3] S. Araki and Y. Shikata: Cohomology mod 2 of the compact exceptional group E_8 , *Proc. Japan Acad.*, **37** (1961), 619-622.
- [4] A.L. Blakers and W.S. Massey: The homotopy groups of a triad II, *Ann. of Math.*, **55** (1952), 192-201.
- [5] R. Bott: The stable homotopy of the classical groups, *Ann. of Math.*, **70** (1959), 313-337.
- [6] R. Bott and H. Samelson: Application of the theory of Morse to symmetric spaces, *Amer. J. Math.*, **80** (1958), 964-1029.
- [7] H. Cartan and J.P. Serre: Espaces fibrés et groupes d'homotopie I, II, *C.R. Acad. Sci. Paris.*, **234** (1952), 288-290, 393-395.
- [8] J.P. Serre: Groupes d'homotopie et classes de groupes abélien, *Ann. of Math.*, **58** (1953), 258-294.
- [9] J.P. Serre: Cohomologie modulo 2 des complexes d'Eilenberg Mac-Lane, *Comm. Math. Helv.*, **27** (1953), 198-231.
- [10] M. Mimura: The homotopy group of Lie groups of low rank, *J. Math. Kyoto Univ.*, **6-2** (1967), 131-176.
- [11] H. Toda: Composition methods in homotopy groups of spheres, *Ann. of Math. Studies.*, (1962).

*Mathematical Institute
Nagoya University*