




# Complete mitochondrial genome of *Penicillidia jenynsii* (Diptera: Hippoboscoidea: Nycteribiidae) and phylogenetic relationship

Jinting Yang<sup>1</sup> , Xiaobin Huang<sup>1</sup>, Yujuan Wang<sup>2</sup>, Huijuan Yang<sup>1</sup>, Xianzheng Zhang<sup>1</sup> and Xiaoyan Zheng<sup>1</sup>

<sup>1</sup>Institute of Pathogens and Vectors, Yunnan Provincial Key Laboratory for Zoonosis Control and Prevention, Dali University, Dali, Yunnan, China and <sup>2</sup>Jilin Provincial Key Laboratory of Animal Resource Conservation and Utilization, Northeast Normal University, Changchun, China

## Research Article

**Cite this article:** Yang J, Huang X, Wang Y, Yang H, Zhang X, Zheng X (2023). Complete mitochondrial genome of *Penicillidia jenynsii* (Diptera: Hippoboscoidea: Nycteribiidae) and phylogenetic relationship. *Parasitology* **150**, 623–630. <https://doi.org/10.1017/S003118202300032X>

Received: 4 February 2023

Revised: 18 March 2023

Accepted: 20 March 2023

First published online: 27 March 2023

### Keywords:

Bat fly; mitochondrial genome; Nycteribiidae; *Penicillidia jenynsii*

### Corresponding author:

Xiaobin Huang,

E-mail: [huangxb633@nenu.edu.cn](mailto:huangxb633@nenu.edu.cn)

## Abstract

In recent years, bat-associated pathogens, such as 2019 novel coronavirus, have been ravaging the world, and ectoparasites of bats have received increasing attention. *Penicillidia jenynsii* is a member of the family Nycteribiidae which is a group of specialized ectoparasites of bats. In this study, the complete mitochondrial genome of *P. jenynsii* was sequenced for the first time and a comprehensive phylogenetic analysis of the superfamily Hippoboscoidea was conducted. The complete mitochondrial genome of *P. jenynsii* is 16 165 base pairs (bp) in size, including 13 protein-coding genes (PCGs), 22 transfer RNA genes, 2 ribosomal RNA genes and 1 control region. The phylogenetic analysis based on 13 PCGs of the superfamily Hippoboscoidea known from the NCBI supported the monophyly of the family Nycteribiidae, and the family Nycteribiidae was a sister group with the family Streblidae. This study not only provided molecular data for the identification of *P. jenynsii*, but also provided a reference for the phylogenetic analysis of the superfamily Hippoboscoidea.

## Introduction

Nycteribiidae, belonging to the Dipteran superfamily Hippoboscoidea, is a specialized parasitic insect on the body surface of bats (Szentivanyi *et al.*, 2016). More than 270 species of Nycteribiidae are known to exist (Dick and Gracioli, 2018). Together with the family Streblidae, they are called bat flies which feed on the blood of bats. As a member of the superfamily Hippoboscoidea, the Nycteribiidae also had a unique life history process that involves reproducing *via* viviparous puparity. Eggs are fertilized internally and all larval stages develop within the female, nourished by intrauterine “milk” glands. The larvae moult twice within the female and gravid females deposit a single terminal (third instar) larva on the roost substrate. Once deposited, the larva (called a prepupa) immediately forms a puparium. After a pupal stage of 3–4 weeks, an adult emerges and must find and colonize a host (Dick and Patterson, 2006). In recent years, bats and their surface parasites have received a great deal of attention as bat pathogens, such as 2019 novel coronavirus (Fan *et al.*, 2019), have become widespread worldwide. Numerous studies have found zoonotic pathogens in bat flies, e.g. Bartonella (Low *et al.*, 2022), narnaviruses, reoviruses (Xu *et al.*, 2022). Therefore, it is hypothesized that bat flies are potential vectors for pathogen transmission. The monitoring of Nycteribiidae relies heavily on basic research on its biology, genetics, genomics, etc.

The mitochondrial genome is the only extra-nuclear genetic information carrier in animals, and it has the characteristics of lack of extensive recombination, conserved gene content, matrilineal inheritance, small molecular weight (approximately 14–21 kb), high mutation rate and a fast evolutionary rate (Yang *et al.*, 2022). Therefore, mitochondrial DNA is often considered a useful molecular marker for species identification, phylogeography, analysis of population structure and dynamics and molecular evolution. Currently, only 4 species of Nycteribiidae in the Genbank database have undergone complete mitochondrial genome sequencing, which is grossly inconsistent with the rich species diversity of Nycteribiidae. Therefore, in order to understand the phylogenetic relationships of the superfamily Hippoboscoidea and further study the population genetics of Nycteribiidae, we sequenced the complete mitochondrial genome of *Penicillidia jenynsii* and analysed its characteristics and evolutionary relationships among the superfamily Hippoboscoidea.

## Materials and methods

### Sample collection and identification of morphological characteristics

Bat flies were collected from the body surface of captured *Miniopterus fuliginosus* (Chiroptera: Miniopterus) in July 2022 in Binchuan (100.58°E, 25.83°N), Yunnan Province, China, placed in sample tubes containing 95% ethanol and stored in a –20°C refrigerator until sample processing. In the laboratory, the collected bat flies were placed directly under the SZ2-ILST dissection microscope (Olympus, Tokyo, Japan) for species identification. One sample was mounted onto

© The Author(s), 2023. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

**Table 1.** Mitochondrial genome information used in this study

Family	Species	Length (bp)	GenBank accession No.	References
Glossinidae	<i>Glossina austeni</i>	17 449	MZ826152	Porter <i>et al.</i> (2022)
Glossinidae	<i>Glossina brevipalpis</i>	17 751	MZ826153	Porter <i>et al.</i> (2022)
Hippoboscidae	<i>Melophagus ovinus</i>	15 573	KX870852	Liu <i>et al.</i> (2017)
Hippoboscidae	<i>Lipoptena</i> sp.	16 953	MT679542	Wang <i>et al.</i> (2021)
Hippoboscidae	<i>Ornithomya biloba</i>	18 654	MZ379837	Li <i>et al.</i> (2022)
Streblidae	<i>Paradyschiria parvula</i>	14 588	MK896865	Trevisan <i>et al.</i> (2019)
Streblidae	<i>Paratrichobius longicrus</i>	16 296	MK896866	Trevisan <i>et al.</i> (2019)
Nycteribiidae	<i>Basilina ansifera</i>	16 964	MZ826150	Porter <i>et al.</i> (2022)
Nycteribiidae	<i>Dipseliopoda setosa</i>	19 164	MZ826151	Porter <i>et al.</i> (2022)
Nycteribiidae	<i>Nycteribia parvula</i>	16 060	OP442519	Unpublished
Nycteribiidae	<i>Phthiridium szechuanum</i>	14 896	OP459298	Unpublished
Limulidae	<i>Carcinoscorpius rotundicauda</i>	15 037	JX437074	Baek <i>et al.</i> (2014)
Limulidae	<i>Limulus polyphemus</i>	14 985	NC_003057	Lavrov <i>et al.</i> (2000)

glass slides with Hoyer's solution. After dehydration, drying and transparency, the mounted specimen was photographed under a Leica DM 3000 LED microscope (Leica, Weztlar, Germany) for pictures of the morphological features of the sample.

#### DNA extraction, library preparation and mitogenome sequencing

The total DNA was extracted from the insects' whole body tissue using the Tissue DNA Kit (Omega Georgia, Connecticut, USA) according to the manufacturer's instructions. The DNA content was quantified using a NanoDrop One Microvolume UV-Vis Spectrophotometer (Thermo Scientific, Massachusetts, USA). A measure of 0.2 µg DNA was used as the input material for DNA library preparation. The sequencing library was prepared using the NEBNext UltraTM DNA Library Prep Kit (New England Biolabs, New York, USA) for Illumina according to the manufacturer's recommendations, and index codes were added. DNA fragments were then end-polished, A-tailed and ligated with the full-length adapter for Illumina sequencing, followed by further polymerase chain reaction (PCR) amplification. After PCR, the products were purified using the AMPure XP system (Beverly, Los Angeles, USA). The quality of the libraries was then assessed using the Agilent 5400 system (Agilent, Palo Alto, USA) and quantified by quantitative PCR (1.5 nM). The qualified libraries were pooled and sequenced on Illumina platforms with the PE150 strategy at Novogene Bioinformatics Technology Co., Ltd. (Beijing, China) according to the effective library concentration and the required amount of data.


#### Assembly, annotation and sequence analysis

This project used MitoZ 2.3 (<https://doi.org/10.1101/489955>) to assemble the mitochondrial genome. The 13 protein-coding genes (PCGs) and 2 ribosomal RNA genes (rRNA) of mitochondrial genome were annotated by MITOS (Donath *et al.*, 2019), and manually compared with the known Nycteribiidae mitochondrial sequences. The 22 transfer RNA genes (tRNA) of the genome were annotated by tRNAscan-SE (Chan *et al.*, 2021) and second structures of tRNAs were predicted by MITOS. Finally, the circular mitochondrial genome map was drawn using GENOMEVX (Conant and Wolfe, 2008). Codon W 1.4.2 (<https://sourceforge.net/projects/codonw/>) was used to calculate

the composition of the base and the relative synonymous codon usage (RSCU). To calculate GC- and AT- skews, the following formulas were used: AT-skew =  $(A - T)/(A + T)$  and GC-skew =  $(G - C)/(G + C)$  (Perna and Kocher, 1995). MEGA 11 was used to estimate the genetic distances using the Kimura-2 parameter (K2P) (Tamura *et al.*, 2021). The non-synonymous (Ka) to synonymous rate (Ks) ratio test of 13 PCGs was conducted in DnaSP 6 (Rozas *et al.*, 2017).

#### Phylogenetic analysis

The phylogenetic relationships of *P. jenynsii* with other species of the superfamily Hippoboscoidea, which could be found in the NCBI (Table 1), were constructed on the basis of 13 PCGs, selecting *Carcinoscorpius rotundicauda* and *Limulus polyphemus* as outgroups. Since tRNA genes are highly conserved, they were not considered when constructing the phylogenetic tree (Gray *et al.*, 1984). Thirteen PCGs from *P. jenynsii* and other species available in GenBank were aligned using MAFFT 7, using default

**Figure 1.** Dorsal side of female *Penicillidia jenynsii*.


**Table 2.** Mitochondrial genome organization of *Penicillidia jenynsii*

Gene	Coding strand	Position and length (bp)	Nucleic acid		Codon		AT-skew	GC-skew	Anticodon	Intergenic nucleotides
			AT%	GC%	Start	Stop				
<i>trnI</i>	J	1–67 (67)	80.60	19.40					GAT	–2
<i>trnQ</i>	N	65–133 (69)	82.61	17.39					TTG	–1
<i>trnM</i>	J	133–200 (68)	75.00	25.00					CAT	0
<i>nad2</i>	J	201–1196 (996)	85.04	14.96	ATT	TAA	–0.18	–0.23		–2
<i>trnW</i>	J	1195–1262 (68)	73.53	26.47					TCA	–8
<i>trnC</i>	N	1255–1314 (60)	68.33	31.67					GCA	0
<i>trnY</i>	N	1315–1378 (64)	76.56	23.44					GTA	–2
<i>cox1</i>	J	1377–2910 (1534)	73.60	26.40	TCG	T	–0.15	–0.09		1
<i>trnL2</i>	J	2912–2974 (63)	77.78	22.22					TAA	2
<i>cox2</i>	J	2977–3655 (679)	77.32	22.68	ATG	T	–0.09	–0.14		0
<i>trnK</i>	J	3656–3726 (71)	70.42	29.58					CTT	–1
<i>trnD</i>	J	3726–3791 (66)	89.39	10.61					GTC	0
<i>atp8</i>	J	3792–3953 (162)	88.89	11.11	ATT	TAA	0.03	–0.56		–7
<i>atp6</i>	J	3947–4621 (675)	79.11	20.89	ATG	TAA	–0.19	–0.34		–1
<i>cox3</i>	J	4621–5409 (789)	78.71	21.29	ATG	TAA	–0.12	–0.13		2
<i>trnG</i>	J	5412–5476 (65)	89.23	10.77					TCC	0
<i>nad3</i>	J	5477–5827 (351)	82.91	17.09	ATT	TAG	–0.18	–0.13		–2
<i>trnA</i>	J	5826–5888 (63)	73.02	26.98					TGC	–1
<i>trnR</i>	J	5888–5948 (61)	77.05	22.95					TCG	–3
<i>trnN</i>	J	5946–6009 (64)	75.00	25.00					GTT	0
<i>trnS1</i>	J	6010–6077 (68)	76.47	23.53					GCT	3
<i>trnE</i>	J	6081–6147 (67)	94.03	5.97					TTC	136
<i>trnF</i>	N	6284–6337 (54)	85.19	14.81					GAA	0
<i>nad5</i>	N	6338–8075 (1738)	83.43	16.57	TTG	T	–0.12	–0.27		0
<i>trnH</i>	N	8076–8139 (64)	82.81	17.19					GTG	0
<i>nad4</i>	N	8137–9469 (1330)	82.41	17.59	ATG	T	–0.18	–0.37		–1
<i>nad4L</i>	N	9469–9747 (279)	85.30	14.70	ATG	TAA	0.15	–0.66		2
<i>trnT</i>	J	9750–9815 (66)	86.36	13.64					TGT	0
<i>trnP</i>	N	9815–9880 (65)	87.69	12.31					TGG	5
<i>nad6</i>	J	9886–10 389 (504)	88.10	11.90	ATT	TAA	–0.14	–0.47		–1
<i>Cytb</i>	J	10 389–11 516 (1128)	78.10	21.90	ATG	TAA	–0.11	–0.27		1
<i>trnS2</i>	J	11 518–11 584 (67)	82.09	17.91					TGA	16
<i>nad1</i>	N	11 601–12 536 (936)	79.49	20.51	ATG	TAG	0.18	–0.32		4
<i>trnL1</i>	N	12 541–12 601 (61)	80.33	19.67					TAG	0
<i>16S rRNA</i>	N	12 602–13 874 (1273)	83.82	16.18	TAA	T				71
<i>trnV</i>	N	13 946–14 016 (71)	81.69	18.31					TAC	0
<i>12S rRNA</i>	N	14 017–14 799 (783)	81.10	18.99	ATA	ATG				0

parameters (Kato and Standley, 2013). Gblocks 0.91b software was used to remove the intergenic gaps and ambiguous sites (Castresana, 2000). The best-fit model was selected using PartitionFinder 2 (Lanfear *et al.*, 2017). The PhyloSuite 1.2.2 was used to perform phylogenetic analyses using maximum likelihood (ML) and Bayesian inference (BI) methods (Zhang *et al.*, 2020). For the ML analysis, an ML phylogenetic tree was

constructed using IQ-TREE 2.1.2 with 1000 ultra-fast bootstrapping under the GTR + F + I + G4 model (Minh *et al.*, 2020). The construction of Bayesian phylogenetic trees was performed using the MRBayes 3.2 software based on the best-fit partitioning scheme and corresponding models in Table S1 (Ronquist *et al.*, 2012), in which the initial 25% of the sampled data were discarded as aged data. When the mean standard deviation of the split





**Figure 4.** Ka, Ks and Ka/Ks of 13 PCGs of *Penicillidia jenynsii*.

standard ATN codons (4 ATT and 7 ATG). Except for 4 PCGs (*cox1*, *cox2*, *nad4* and *nad5*) that ended with incomplete stop codons T, the remaining 11 PCGs (7 TAA and 2 TAG) have complete stop codons.

PCGs' codon usage and RSCU were calculated (Fig. 3). The results show that UUA, AUU, UUU and AUA were the most frequently used codons among the *P. jenynsii*'s mitochondrial PCGs, and GCG, CCG and AGG were less frequently used codons.

The Ka, Ks and Ka/Ks were computed for each PCG, and details are shown in Fig. 4. The values of Ka/Ks ranged from 0.07 to 2.52. The evolution rate of 13 PCGs was sequentially *nad1* > *nad2* > *nad4L* > *nd6* > *nad4* > *nad5* > *nad3* > *atp8* > *atp6* > *cox3* > *cox2* > *cytb* > *cox1*. Among them, *nad1* was positively selected (Ka/Ks > 1), while other genes were purified (Ka/Ks < 1).

### tRNA, rRNA genes and control region

The length of the 22 tRNAs ranges from 54 to 71 bp. All tRNAs fold into a typical cloverleaf structure except *trnF* that lacked the TΨC arms and *trnS1* that lacked the dihydrouracil (DHU) arm (Fig. S1). The 16S rRNA was 1273 bp in length with 83.82% AT content, which was located between *trnL1* and *trnV*. The 12S rRNA was 789 bp in length with 81.10% AT content, which was located between *trnV* and the control region. The control region was located between *trnV* and *trnI*, with a sequence length of 1717 bp. Furthermore, the AT content was 90.62%.

## Phylogenetic analysis of *P. jenynsii*

### Genetic distance

The analysis of corrected genetic distances was performed to investigate the phylogenetic relationships of the superfamily Hippoboscoidea (Table 3). The range of genetic distance is from 0.114 to 0.298. The average genetic distance is 0.270. Among the genetic distances of *P. jenynsii* and others, the genetic distance between *P. jenynsii* and *Phthiridium szechuanum* is the smallest (0.138).

### Phylogenetic analysis


The topologies of the phylogenetic tree completed by the 2 methods were identical, with high support for each node (Figs 5 and 6). The results demonstrate the superfamily Hippoboscoidea is divided into 2 branches: one is Nycteribiida and Streblidae, and the other is Glossinidae and Hippoboscidae. The family Nycteribiidae is a monophyletic and sister group with the family Streblidae. This is consistent with the morphological classification. *Penicillidia jenynsii* has the closest affinity to *P. szechuanum* and *Nycteribia parvula*.

## Discussion

In this study, we sequenced the complete mitochondrial genome of *P. jenynsii* for the first time and conducted a comprehensive phylogenetic analysis based on 13 PCGs of *P. jenynsii* and the other 11 species. The sequence of the mitochondrial genome of *P. jenynsii* is the same as that known in the superfamily Hippoboscoidea and is identical to the sequence of *Drosophila melanogaster* which is the classical structure of Diptera (Lewis *et al.*, 1995). Mitochondrial genome rearrangements as evolutionary events are relatively rare; therefore, this is an important tool for assessing phylogenetic relationships between species. Eight of the 13 PCGs have the classical ATN as the start codon, while *cox1* has TCG as the start codon, which is common in species of the superfamily Hippoboscoidea (Tang *et al.*, 2018; Porter *et al.*, 2022). Nine of the 13 PCGs end with complete stop codons, while 4 genes (*cox1*, *cox2*, *nad4*, *nad5*) utilize incomplete stop codon T, which could be completed as TAA via posttranscriptional polyadenylation (Ojala *et al.*, 1981). The tRNA can form a cloverleaf secondary structure except for the absence of the DHU arm of *trnS1* and the absence of the TΨC arms of *trnF*. The absence of the DHU arm of *trnS1* is a common phenomenon

**Table 3.** Estimation of pairwise genetic distances (%) among the species of the superfamily Hippoboscoidea

	1	2	3	4	5	6	7	8	9	10	11
1. <i>Penicillidia jenynsii</i>											
2. <i>Basilina ansifera</i>	0.175										
3. <i>Dipseliopoda setosa</i>	0.211	0.219									
4. <i>Glossina austeni</i>	0.294	0.298	0.295								
5. <i>Glossina brevivalpis</i>	0.273	0.275	0.274	0.177							
6. <i>Lipoptena</i> sp	0.268	0.270	0.267	0.224	0.208						
7. <i>Melophagus ovinus</i>	0.274	0.269	0.272	0.236	0.216	0.138					
8. <i>Nycteribia parvula</i>	0.141	0.172	0.210	0.296	0.274	0.263	0.266				
9. <i>Ornithomya biloba</i>	0.282	0.286	0.285	0.209	0.200	0.189	0.202	0.283			
10. <i>Paradyschiria parvula</i>	0.287	0.281	0.282	0.273	0.260	0.260	0.263	0.280	0.270		
11. <i>Paratrachobius longicrus</i>	0.268	0.274	0.266	0.255	0.241	0.233	0.238	0.264	0.243	0.204	
12. <i>Phthiridium szechuanum</i>	0.138	0.168	0.207	0.292	0.269	0.268	0.270	0.114	0.282	0.281	0.262




**Figure 5.** Phylogenetic relationships of the superfamily Hippoboscoidea using ML analyses based on 13 PCGs of mitogenomes.

in Metazoa, and tRNA genes with incomplete secondary structures can maintain their normal functions through post-transcriptional modifications (Negrisolo *et al.*, 2011). The positions of these 2 rRNAs are more conserved as in the other Dipteran mitochondrial genomes. The control region, also known as the AT-enriched region, has an AT content of 90.62% in *P. jenynsii* and is thought to control the replication and transcription of the mitochondrial genome (Boore, 1767). The length variation among insect mitochondrial genomes depends mainly on the variation in the A + T-rich region, which ranges from 70 to 13 kb in length (Zhang and Hewitt, 1997).

The base composition of the mitochondrial genome shows a severe bias, with 82.15% AT content, similar to other species in the superfamily Hippoboscoidea and it has been suggested that

the species with higher AT content in mitochondrial genes are considered to be more evolutionarily advantageous (Vrijenhoek, 1994). The results of AT-skew and GC-skew could reflect the preference for A and G bases in the mitochondrial genome (Lindahl, 1993). The main causes of base skewness are asymmetric mutation of bases during replication and transcription, as well as selection pressure.

Ka/Ks values of PCGs can reflect the selective pressure on the gene and, to some extent, the conservativeness of the gene; Ka/Ks < 1 is considered purifying selection; Ka/Ks > 1 is considered positive selection. The smaller the value of Ka/Ks, the greater the selective pressure on the gene, and the more conservative the gene; the larger the value of Ka/Ks, the faster the evolutionary rate of the gene, and the less conservative the gene (Nekrutenko



**Figure 6.** Phylogenetic relationships of the superfamily Hippoboscoidea using BI analyses based on 13 PCGs of mitogenomes.

*et al.*, 2002). In the mitochondrial genome, the Ka/Ks values of all PCGs except *nad1* are less than 1. It is evident that most of the PCGs are conserved, among which the Ka/Ks values of *cox1* and *cytb* are significantly smaller than those of other genes, indicating that they are under strong selection pressure and the genes are more conserved. On the contrary, the Ka/Ks values of *nad1* and *nad2* were relatively larger, and these genes were under weaker selection pressure and were relatively less conservative. Other studies have also shown that the *cox1* and *cytb* genes are the most conserved in the mitochondrial genome, which are now widely used for molecular markers and phylogenetic studies (Park *et al.*, 2010; Tobe *et al.*, 2010).

Two methods of completing the phylogenetic trees of the superfamily Hippoboscoidea revealed that the families Nycteribiidae, Hippoboscidae, Glossinidae and Streblidae are monophyletic, and the species within these families clustered together with high confidence. The families Hippoboscidae, Nycteribiidae, Streblidae and Glossinidae have been combined into 1 superfamily because of their unique reproductive mode, adenotrophic viviparity (Petersen *et al.*, 2007). Except for family Glossinidae which is free-living, the other 3 families are all true ectoparasites, spending all or most of their adult lives within the fur or feathers of their mammalian and bird hosts, and thus undergo parasitic adaptations (McAlpine, 1989). This, together with the relatively small amount of molecular data on Hippoboscoidea species, has led to disagreement in the classification of Hippoboscoidea. Initially, only 3 families of species, Hippoboscidae, Nycteribiidae and Streblidae, were considered to belong to the superfamily Hippoboscoidea, and some experts would classify Nycteribiidae and Streblidae into Hippoboscidae (Petersen *et al.*, 2007). In this study, all PCGs of the mitochondrial genes were analysed phylogenetically to provide definitive evidence for the classification of the superfamily Hippoboscoidea.

**Supplementary material.** The supplementary material for this article can be found at <https://doi.org/10.1017/S003118202300032X>.

**Data availability.** The genome sequence data that support the findings of this study are openly available in GenBank of NCBI at <https://www.ncbi.nlm.nih.gov/>, reference number OQ127278. The associated SRA number is SRR23882031.

**Author's contribution.** The research was designed by J. Yang, X. Huang, Y. Wang, H. Yang, X. Zhang and X. Zheng. J. Yang performed the research. Material preparation and data collection were performed by J. Yang, X. Huang, X. Zhang and Y. Wang. J. Yang, H. Yang and X. Huang analysed the data. The first draft of the manuscript was written by J. Yang. X. Huang critically revised the content of the paper. All authors have read and approved the final version of the manuscript to be published; and all authors agree to be accountable for all aspects of the work.

**Financial support.** This work was supported by the National Natural Science Foundation of China (No. 32001096 to Xiaobin Huang) and the Science and Technology Planning Project of the Yunnan Provincial Department of Science and Technology (No. 202001AT070025 to Xiaobin Huang).

**Competing interests.** None.

**Ethical standards.** No specific permits were required for the insects collected for this study. The bats were captured according to the standards and procedures set by the Animal Ethics Committee of Dali University (name: Dali University Ethics Committee; approval ID: MECDU-202104-27).

## References

Baek SY, Choi EH, Jang KH, Ryu SH, Park SM, Suk HY, Chang CY and Hwang UW (2014) Complete mitochondrial genomes of *Carcinoscorpis rotundicauda* and *Tachypleus tridentatus* (Xiphosura, Arthropoda) and

- implications for chelicerate phylogenetic studies. *International Journal of Biological Sciences* **10**, 479.
- Boore JL (1967) Animal mitochondrial genomes. *Nucleic Acids Research* **27**, 1767–1780.
- Castresana J (2000) Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Molecular Biology and Evolution* **17**, 540–552.
- Chan PP, Lin BY, Mak AJ and Lowe TM (2021) tRNAscan-SE 2.0: improved detection and functional classification of transfer RNA genes. *Nucleic Acids Research* **49**, 9077–9096.
- Conant GC and Wolfe KH (2008) GenomeVx: simple web-based creation of editable circular chromosome maps. *Bioinformatics* **24**, 861–862.
- Dick CW and Gracioli G (2018) *Checklist of World Streblidae (Diptera: Hippoboscoidea)*. Chicago, IL: Center for Integrative Research, The Field Museum of Natural History. Available from [https://www.researchgate.net/publication/322578987\\_CHECKLIST\\_OF\\_WORLD\\_STREBLIDAE\\_DIPTERA\\_HIPPOBOSCOIDEA](https://www.researchgate.net/publication/322578987_CHECKLIST_OF_WORLD_STREBLIDAE_DIPTERA_HIPPOBOSCOIDEA) (accessed 4 April 2019).
- Dick CW and Patterson BD (2006) *Micromammals and Macroparasites: from Evolutionary Ecology to Management*. Berlin, Germany: Springer, pp. 179–194.
- Donath A, Jühling F, Al-Arab M, Bernhart SH, Reinhardt F, Stadler PF, Middendorf M and Bernt M (2019) Improved annotation of protein-coding genes boundaries in metazoan mitochondrial genomes. *Nucleic Acids Research* **47**, 10543–10552.
- Fan Y, Zhao K, Shi ZL and Zhou P (2019) Bat coronaviruses in China. *Viruses* **11**, 210.
- Gray MW, Sankoff D and Cedergren RJ (1984) On the evolutionary descent of organisms and organelles: a global phylogeny based on a highly conserved structural core in small subunit ribosomal RNA. *Nucleic Acids Research* **12**, 5837–5852.
- Katoh K and Standley DM (2013) MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution* **30**, 772–780.
- Lanfear R, Frandsen PB, Wright AM, Senfeld T and Calcott B (2017) PartitionFinder 2: new methods for selecting partitioned models of evolution for molecular and morphological phylogenetic analyses. *Molecular Biology and Evolution* **34**, 772–773.
- Lavrov DV, Boore JL and Brown WM (2000) The complete mitochondrial DNA sequence of the horseshoe crab *Limulus polyphemus*. *Molecular Biology and Evolution* **17**, 813–824.
- Lewis O, Farr C and Kaguni L (1995) *Drosophila melanogaster* mitochondrial DNA: completion of the nucleotide sequence and evolutionary comparisons. *Insect Molecular Biology* **4**, 263–278.
- Li X, Wang L and Yang D (2022) The complete mitochondrial genome of *Ornithomya biloba* (Diptera, Hippoboscidae). *Mitochondrial DNA Part B* **7**, 856–858.
- Lindahl T (1993) Instability and decay of the primary structure of DNA. *Nature* **362**, 709–715.
- Liu ZQ, Kuermanali N, Li Z, Chen SJ, Wang YZ, Tao H and Chen CF (2017) The complete mitochondrial genome of the parasitic sheep ked *Melophagus ovinus* (Diptera: Hippoboscidae). *Mitochondrial DNA Part B* **2**, 432–434.
- Low VL, Tan TK, Tohiran KA, Lim YAL, AbuBakar S and Nasir DM (2022) A novel clade of bat-associated Bartonella detected in the bat fly *Leptocyclopodia ferrari* (Diptera: Nycteribiidae) parasitizing *Cynopterus brachyotis* (Chiroptera: Pteropodidae). *Veterinary Microbiology* **264**, 109284.
- McAlpine J (1989) Phylogeny and classification of the Muscomorpha. *Manual of Nearctic Diptera* **3**, 1397–1518.
- Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, Von HA and Lanfear R (2020) IQ-TREE 2: new models and efficient methods for phylogenetic inference in the genomic era. *Molecular Biology and Evolution* **37**, 1530–1534.
- Negriso E, Babbucci M and Patarnello T (2011) The mitochondrial genome of the ascalaphid owlfly *Libelloides macaronius* and comparative evolutionary mitochondrial genomics of neuropterid insects. *BMC Genomics* **12**, 1–26.
- Nekrutenko A, Makova KD and Li WH (2002) The KA/KS ratio test for assessing the protein-coding potential of genomic regions: an empirical and simulation study. *Genome Research* **12**, 198–202.
- Ojala D, Montoya J and Attardi G (1981) tRNA punctuation model of RNA processing in human mitochondria. *Nature* **290**, 470–474.

- Park DS, Suh SJ, Oh HW and Hebert PD** (2010) Recovery of the mitochondrial COI barcode region in diverse Hexapoda through tRNA-based primers. *BMC Genomics* **11**, 1–7.
- Perna NT and Kocher TD** (1995) Patterns of nucleotide composition at four-fold degenerate sites of animal mitochondrial genomes. *Journal of Molecular Evolution* **41**, 353–358.
- Petersen FT, Meier R, Kuttly SN and Wiegmann BM** (2007) The phylogeny and evolution of host choice in the Hippoboscoidea (Diptera) as reconstructed using four molecular markers. *Molecular Phylogenetics and Evolution* **45**, 111–122.
- Porter ML, Lutz H, Steck M and Chong RA** (2022) The complete mitochondrial genomes and phylogenetic analysis of two Nycteribiidae bat flies (Diptera: Hippoboscoidea). *Mitochondrial DNA Part B* **7**, 1486–1488.
- Ronquist F, Teslenko M, Van Der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA and Huelsenbeck JP** (2012) MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* **61**, 539–542.
- Rothschild M and Theodor O** (1967) *An Illustrated Catalogue of the Rothschild Collection of Nycteribiidae (Diptera) in the British Museum (Natural History)*. London and Tonbridge, UK: Whitefriars Press.
- Rozas J, Ferrer-Mata A, Sánchez-DelBarrio JC, Guirao-Rico S, Librado P, Ramos-Onsins SE and Sánchez-Gracia A** (2017) DnaSP 6: DNA sequence polymorphism analysis of large data sets. *Molecular Biology and Evolution* **34**, 3299–3302.
- Szentivanyi T, Estók P and Földvári M** (2016) Checklist of host associations of European bat flies (Diptera: Nycteribiidae, Streblidae). *Zootaxa* **4205**, 101–126.
- Tamura K, Stecher G and Kumar S** (2021) MEGA11: molecular evolutionary genetics analysis version 11. *Molecular Biology and Evolution* **38**, 3022–3027.
- Tang JM, Li F, Cheng TY, Duan DY and Liu GH** (2018) Comparative analyses of the mitochondrial genome of the sheep ked *Melophagus ovinus* (Diptera: Hippoboscoidea) from different geographical origins in China. *Parasitology Research* **117**, 2677–2683.
- Tobe SS, Kitchener AC and Linacre AM** (2010) Reconstructing mammalian phylogenies: a detailed comparison of the cytochrome b and cytochrome oxidase subunit I mitochondrial genes. *PLoS ONE* **5**, e14156.
- Trevisan B, Alcantara DM, Machado DJ, Marques FP and Lahr DJ** (2019) Genome skimming is a low-cost and robust strategy to assemble complete mitochondrial genomes from ethanol preserved specimens in biodiversity studies. *PeerJ* **7**, e7543.
- Vrijenhoek R** (1994) DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* **3**, 294–299.
- Wang M, Wang JS, Guo YY, Zheng QL, Nouhoum DB and Meng FM** (2021) Complete mitochondrial genome of a potential vector louse fly, *Lipoptena grahami* (Diptera, Hippoboscoidea). *Mitochondrial DNA Part B* **6**, 1752–1753.
- Xu Z, Feng Y, Chen X, Shi M, Fu S, Yang W, Liu WJ, Gao GF and Liang G** (2022) Virome of bat-infesting arthropods: highly divergent viruses in different vectors. *Journal of Virology* **96**, e01464–e01421.
- Yang HJ, Yang ZH and Dong WG** (2022) Morphological identification and phylogenetic analysis of Laelapin mite species (Acari: Mesostigmata: Laelapidae) from China. *The Korean Journal of Parasitology* **60**, 273–279.
- Zhang DX and Hewitt GM** (1997) Insect mitochondrial control region: a review of its structure, evolution and usefulness in evolutionary studies. *Biochemical Systematics and Ecology* **25**, 99–120.
- Zhang D, Gao F, Jakovlić I, Zou H, Zhang J, Li WX and Wang GT** (2020) PhyloSuite: an integrated and scalable desktop platform for streamlined molecular sequence data management and evolutionary phylogenetics studies. *Molecular Ecology Resources* **20**, 348–355.