

exposure to the HCW while she was symptomatic. The four HCWs were contacted, and chemoprophylaxis initiated for them and for the two hospitalized patients.

DISCUSSION

Nosocomial pertussis has been reported previously, most commonly in pediatric wards of hospitals or in residential facilities for the disabled.^{10,11} The index case, as in this report, usually has been a patient with undiagnosed infection. Infected employees, although rarely the source case, may be involved in propagating infection within the health-care institution¹² or within their own households. Thus, diagnosis of a healthcare worker with pertussis, particularly in an acute-care pediatric institution, is of utmost importance.

Although the diagnosis of pertussis in our index case was delayed, PCR was able to diagnose infection in both the patient and HCW within 24 hours of processing the specimen. Prolonged contact between the HCW and the patient confirms the potential for nosocomial acquisition of infection, particularly when high-risk activities such as suctioning are performed. The infection control service should work together with the Occupational Health Department to identify all HCWs with substantial exposure to an infective case.

Current recommendations for chemoprophylaxis of pertussis after substantial exposure are the same as those for treatment of infection.¹³ The value of confirming the diagnosis in the HCW is to restrict them from work during the most contagious catarrhal phase of infection and thus from the potential of further spread within the institution. In autumn and winter months, when respiratory symptoms due to other infectious agents may be prevalent, a confirmed diagnosis of pertussis is useful to advise the symptomatic HCW to refrain from working, as has been recommended.¹⁴ Polymerase chain reaction has proven to be a useful and rapid test in this regard; the high sensitivity and rapid turnaround time makes this test useful for the diagnosis of nosocomially acquired pertussis.

This case report supports the utility of pertussis PCR in the diagnosis of nosocomially acquired infection.

96-CC-238. Address reprint requests to Anne G. Matlow, MD, Division of Microbiology/Infectious Diseases, The Hospital for Sick Children, 555 University Ave, Toronto, Ontario M5G 1X8, Canada.

REFERENCES

- Hallander HO, Storsaeter J, Mollby R. Evaluation of serology and nasopharyngeal cultures for diagnosis of pertussis in a vaccine efficacy trial. *J Infect Dis* 1991;163:1046-1054.
- Reizenstein E, Johansson B, Mardin L, Abens J, Mollby R, Hallander HO. Diagnostic evaluation of polymerase chain reaction discriminative for *Bordetella pertussis*, *B paraptussis* and *Bronchiseptica*. *Diagn Microbiol Infect Dis* 1993;17:185-191.
- van der Zee A, Agterberg C, Peeters M, Schellekens J, Mooi F. Polymerase chain reaction assay for pertussis: simultaneous detection and discrimination of *Bordetella pertussis* and *Bordetella paraptussis*. *J Clin Microbiol* 1993;31:2134-2140.
- Glare E, Paton J, Premier R, Lawrence A, Nisbet I. Analysis of a repetitive DNA sequence from *Bordetella pertussis* and its application to the diagnosis of pertussis using the polymerase chain reaction. *J Clin Microbiol* 1990;28:1982-1987.
- Grimpel E, Begue P, Anjak I, Betsou F, Guiso N. Comparison of polymerase chain reaction, culture and Western immunoblot serology for diagnosis of *Bordetella pertussis* infection. *J Clin Microbiol* 1993;31:2745-2750.
- He Q, Mertsola J, Soini H, Viljanen MK. Sensitive and specific polymerase chain reaction assays for detection of *Bordetella pertussis* in nasopharyngeal specimens. *J Pediatr* 1994;124:421-426.
- He Q, Mertsola J, Soini H, Skurnik M, Ruuskanen O, Viljanen M. Comparison of polymerase chain reaction with culture and enzyme immunoassay for diagnosis of pertussis. *J Clin Microbiol* 1993;31:642-645.
- Nelson S, Matlow A, McDowell C, et al. Detection of *Bordetella pertussis* in clinical specimens using the polymerase chain reaction and a micro titre-based DNA hybridization assay. *J Clin Microbiol* 1997;35:642-645.
- McLafferty M, Harcus D, Hewlett E. Nucleotide sequence and characterization of a repetitive DNA element from the genome of *Bordetella pertussis* with characteristics of an insertion sequence. *J Gen Microbiol* 1988;134:2297-2306.
- Steketee RW, Wassilak GF, Adkins WN Jr, et al. Evidence for a high attack rate and efficacy of erythromycin prophylaxis in a pertussis outbreak in a facility for the developmentally disabled. *J Infect Dis* 1988;157:434-440.
- Fisher MC, Long SS, McGowan KL, Kaselis E, Smith DG. Outbreak of pertussis in a residential facility for handicapped people. *J Pediatr* 1989;114:934-939.
- Kurt TL, Yeager AS, Guenette S, Dunlop S. Spread of pertussis by hospital staff. *JAMA* 1972;221:264-267.
- Committee on Infectious Diseases. *Red Book*. Elk Grove Village, IL: American Academy of Pediatrics; 1994:355-367.
- Weber DJ, Rutala WA. Management of health care workers exposed to pertussis. *Infect Control Hosp Epidemiol* 1994;15:411-415.

Calendar

March 18-21, 1998. The Second Australian Wound Management Association Conference, Wound Management Toward 2000—A Blossoming Paradigm, will be held in Brisbane and hosted by the Queensland Wound Care Association, Inc, on behalf of the Australian Wound Management Association. Proposed topics include, but are not limited to, the following: chronic wound healing; chronic venous ulcers; commercialization of medical interventions; molecu-

lar mechanisms of normal and aberrant wound repair; proto-oncogene control of fetal wounds; skin-graft storage analyses; cell biology of animal models of impaired wound healing; mechanisms of cell death by apoptosis in vascularized skin flaps; in vitro models of wound fibroblasts; acute, chronic, and burn wounds; optimizing scar outcome using dermal substitutes and cultured epithelial autograft; comorbidities of attendees of wound clinic—co-existing diseases, dependencies,

and cognitive status; hyperbaric oxygen therapy.

October 15, 1997, is the deadline for submission of abstracts. Faxed abstracts will not be considered.

To request further information, contact the Second AWMA Conference, PO Box 1280, Unit 11, 97 Castlemaine St, Milton QLD 4064, Australia; e-mail, wm98@im.com.au; telephone, 61 (0)7 3369 0477; fax, 61 (0)7 3369 1512.