

ing habits, as presented at the 1967 investigation. It is concluded that none of these factors seemed to influence the future development of CHD in twins apparently tainted with a heredity for this disease.

Ingvar Liljefors, M.D., St Görans Hospital, Box 12500, 11281 Stockholm, Sweden

## METABOLIC RESEARCH IN MZ TWINS WITH DIABETES MELLITUS

### Progress Report

T. GOECKE, W. GROTE

*Department of Human Genetics and Anthropology, University of Düsseldorf, German Federal Republic*

An intravenous and oral glucose-tolerance test and an intravenous tolbutamid-test have been carried out in a sample of 12 MZ twin pairs discordant for diabetes mellitus. Blood sugar, insulin, and free fatty acids were determined. The research aimed at finding out whether prediabetic subjects may show any characteristic feature in the metabolism of sugars and fats, that may reveal a diabetic ground. Preliminary results show that, in MZ twins with discordant juvenile diabetes mellitus, metabolic values may remain discordant for quite a number of years.

Dr. T. Goecke, Institut für Humangenetik und Anthropologie, Ulenbergstrasse 127/129, Düsseldorf, German Federal Republic

## A GENETIC STUDY ON DEAF TWINS

M. HORIUCHI

*Keio University, Tokyo, Japan*

A study was carried out on 33 cases of twins with early total deafness from the Japanese young population. Complete physical and otological signs of all subjects were examined by otolaryngologists. Twenty-five cases

were accepted as index cases for genetic analysis. They had no evidence of the known exogenous causes of deafness. Family and developmental history were taken. The sample included 17 MZ and 8 DZ pairs. Cousin-mating rate was 32%, i.e., five times the average population rate in Japan.

Audiometric tests were analysed. Nonsignificant differences were found between right and left ear in zygosity group, but intrapair differences of the hearing impairment were larger in DZ than MZ pairs. The deafness concordance rates for DZ and MZ pairs were respectively assessed as 63 and 88%. Penetrance was established as 0.94. The average intrapair difference of the hearing loss was 8 dB in the MZ and 30 dB in the DZ group, and a dissolution indicated remarkable small value in the MZ group. It is suggested that the degree of hearing impairment is strongly influenced by heredity.

Dr. M. Horiuchi, 1-35-16 Shimizu Suginami-ku, Tokyo, Japan

## GENETIC FACTORS IN MYOPIA

JOHN L. KARLSSON

*Napa State Hospital, Napa, California, USA*

The present study is concerned primarily with the nature of the well documented relationship between nearsightedness and enhanced performance on scholastic or intelligence tests. Myopia is established to be a familial disorder, and there are great variations in its frequency between different ethnic or national groups. In some populations, such as those of Jewish, Japanese, or West-European origin, the rate of myopia is in the order of 25%, while primitive societies are almost free of myopia.

Many eye specialists favor the view that nearsightedness may result from excessive close work, often referring to the condition arising during the period of education as "school myopia". Recent surveys which indicate that an increase may be occurring in myopia in Eskimos and North-Canadian Indians have also been interpreted in terms of external factors. Some centers are engaged in experimental work with monkeys,