

# Review: Importance of colostrum supply and milk feeding intensity on gastrointestinal and systemic development in calves

H. M. Hammon<sup>1†</sup> , W. Liermann<sup>1</sup>, D. Frieten<sup>2</sup> and C. Koch<sup>3</sup>

<sup>1</sup>Institute of Nutritional Physiology 'Oskar Kellner', Leibniz Institute for Farm Animal Biology (FBN), Wilhelm-Stahl-Allee 2, Dummerstorf 18196, Germany; <sup>2</sup>Department of Life Sciences and Engineering, University of Applied Sciences Bingen, Berlinstrasse 109, Bingen am Rhein 55411, Germany; <sup>3</sup>Educational and Research Centre for Animal Husbandry, Hofgut Neumuehle, Münchweiler an der Alsenz 67728, Germany

(Received 28 March 2019; Accepted 21 August 2019)

---

*Feeding management of the postnatal and preweaning calf has an important impact on calf growth and development during this critical period and affects the health and well-being of the calves. After birth, an immediate and sufficient colostrum supply is a prerequisite for successful calf rearing. Colostrum provides high amounts of nutrient as well as non-nutrient factors that promote the immune system and intestinal maturation of the calf. The maturation and function of the neonatal intestine enable the calf to digest and absorb the nutrients provided by colostrum and milk. Therefore, colostrum intake supports the start of anabolic processes in several tissues, stimulating postnatal body growth and organ development. After the colostrum feeding period, an intensive milk feeding protocol, that is, at least 20% of BW milk intake/day, is required to realise the calf potential for growth and organ development during the preweaning period. Insufficient milk intake delays postnatal growth and may have detrimental effects on organ development, for example, the intestine and the mammary gland. The somatotrophic axis as the main postnatal endocrine regulatory system for body growth is stimulated by the intake of high amounts of colostrum and milk and indicates the promotion of anabolic metabolism in calves. The development of the forestomach is an important issue during the preweaning period in calves, and forestomach maturation is best achieved by solid feed intake. Unfortunately, intensive milk-feeding programmes compromise solid feed intake during the first weeks of life. In the more natural situation for beef calves, when milk and solid feed intake occurs at the same time, calves benefit from the high milk intake as evidenced by enhanced body growth and organ maturation without impaired forestomach development during weaning. To realise an intensive milk-feeding programme, it is recommended that the weaning process should not start too early and that solid feed intake should be at a high extent despite intensive milk feeding. A feeding concept based on intensive milk feeding prevents hunger and abnormal behaviour of the calves and fits the principles of animal welfare during preweaning calf rearing. Studies on milk performance in dairy cows indicate that feeding management during early calf rearing influences lifetime performance. Therefore, an intensive milk-feeding programme affects immediate as well as long-term performance, probably by programming metabolic pathways during the preweaning period.*

---

**Keywords:** preweaning calf, somatotrophic axis, nutrient intake, organ maturation, growth

## Implications

Successful calf rearing depends on an enhanced colostrum management and is improved by an intensive milk-feeding programme that allows calves to develop their potential for growth and organ maturation. Feeding regimes with a restricted milk supply of 4 to 6 kg milk or milk replacer/day focus on forestomach development and may disregard body growth and the maturation of other visceral organs beyond the rumen. An *ad libitum* milk-feeding programme stimulates calf growth and organ development and is

consistent with calf well-being and avoids hunger. In addition, intensive milk feeding may have a strong impact on calf health and long-life performance in cattle.

## Introduction

Mortality and morbidity rates are still unacceptably high during early calf rearing. The incidence for mortality in the perinatal period, defined as the duration from birth to 48 h after birth, ranges in dairy herds worldwide between 3% and 9% (Compton *et al.*, 2017). A recent survey on mortality

---

† E-mail: [hammon@fbn-dummerstorf.de](mailto:hammon@fbn-dummerstorf.de)

rates in Germany revealed up to 17% calf losses (calf losses after birth up to 6 months of age) in dairy farms (Tautenhahn, 2017). In US dairy herds, current mortality rates of 5% and morbidity rates of 34% were published for preweaning calves (Urie *et al.*, 2018). The UK Department of the Environment, Food and Rural Affairs reported that economic losses from calf mortality were around £60 million/year (DEFRA, 2003). It is obvious that the high mortality and morbidity rates contradict the aim of increased animal welfare for farm animals and compromise the breeding of robust animals (Huber, 2018). In dairy farming, calves usually do not grow up with their dam, and calves are immediately separated from their dams after birth. Thus, farmers are highly responsible for the colostrum and milk feeding management and can significantly contribute to an improved calf health and the reduction of calf losses during the postnatal period.

Feeding management during the neonatal and preweaning period has a great impact on the success of calf rearing and, in addition, affects health and performance in later life (Khan *et al.*, 2011; Ballou, 2012; Van Amburgh and Soberon, 2013). Because severe diarrhoea is a main reason for neonatal calf losses, the management of milk feeding and especially colostrum supply in the first days of life is of particular importance for the success of calf rearing (DEFRA, 2003; Tautenhahn, 2017; Urie *et al.*, 2018). An adequate and immediate (within 2 to 3 h after birth) colostrum supply is important for establishing passive immunity in calves, and the amount of colostrum fed to newborn calves directly correlates with the prevention of illness and calf losses (Godden, 2008; Mee, 2008).

Furthermore, there is increasing evidence that an enhanced milk or milk replacer (MR) feeding schedule during the preweaning period not only affects growth but also promotes organ development and well-being (Geiger *et al.*, 2016; Schäff *et al.*, 2016; Rosenberger *et al.*, 2017). This review aims to summarise the research on the impact of colostrum supply and subsequent intensive milk feeding on the gastrointestinal and systemic development and maturation of the preweaning calf. An intensive milk feeding schedule orientates on a daily milk intake of 20% instead of 10% of BW (Khan *et al.*, 2011), which is closely related to *ad libitum* milk (Jasper and Weary, 2002; Schiessler *et al.*, 2002; Maccari *et al.*, 2015) or MR feeding (Schäff *et al.*, 2016; Frieten *et al.*, 2017) in preweaning calves.

## Impact of colostrum supply on postnatal maturation

### *Colostrum supply and intestinal development and maturation*

Bovine colostrum provides newborn calves with high amounts of nutrient and non-nutrient biologically active substances (Blum and Baumrucker, 2008; Nissen *et al.*, 2017). In addition to the great importance of colostral immunoglobulins for the passive immunity of neonatal calves (Barrington and Parish, 2001; Godden, 2008), colostrum contains a large number of immunomodulatory peptides that may also affect

neonatal immune response (Chase *et al.*, 2008; Stelwagen *et al.*, 2009; Nissen *et al.*, 2017). Some of these factors are provided by colostral immune cells that are involved in the establishment of local and systemic neonatal immunity (Liebler-Tenorio *et al.*, 2002; Stelwagen *et al.*, 2009; Langel *et al.*, 2015). In addition, potential effects of colostrum on the neonatal microbiome in the gut are likely and become more important in the research of calf nutrition (Malmuthuge and Guan, 2017). The importance of the colostrum supply for the development and maturation of the immune system of the newborn calf is far beyond the provision of immunoglobulins. Recent studies in humans, investigating the effects of breastmilk feeding on neonatal intestinal development, illustrate the significance of colostral immune cells and the intestinal microbiome on the maturation of the neonatal immune response in the gut (Molès *et al.*, 2018). A comparable function of bovine colostrum in the neonatal intestine of the calf is conceivable but requires more investigations.

Bovine colostrum has an overall importance for the postnatal development of the gut (Blum, 2006). The high concentrations of hormones, growth factors and cell-modulating factors in colostrum (Blum and Baumrucker, 2008; Nissen *et al.*, 2017) stimulate villus growth of the small intestinal mucosa in calves (Blum, 2006; Steinhoff-Wagner *et al.*, 2014). Colostrum feeding promotes mucosal cell growth and protein synthesis in the enterocytes of neonatal mammals (Donovan and Odle, 1994; Burrin *et al.*, 1995; Xu, 1996). The amount of overall ingested colostrum corresponds to the villus size in the intestinal mucosa, leading to a greater villus size in repeatedly colostrum-fed calves (Blum, 2006). When feeding a colostrum extract, that is, a fraction originating from first colostrum including most of the growth-promoting peptides, together with a milk-based formula, the villus size is stimulated when compared to a milk-based formula feeding with similar protein and energy as in colostrum but no growth-stimulating peptides (Roffler *et al.*, 2003). This finding supports the general assumption that colostral peptides, such as IGF-I, or hormones, such as insulin, are involved in the growth-stimulating effect on the intestinal mucosa of neonatal calves (Blum, 2006).

In general, the proliferation rate of intestinal crypt cells depends on feeding (Johnson 1988; Mathers, 1998). Colostrum or colostral components stimulate crypt cell proliferation in the intestinal mucosa of neonatal calves (Blum, 2006). When comparing colostrum feeding and milk-based formula feeding (same nutrient content but no growth-promoting bioactive factors as colostrum during the first 3 days after birth), the greater stimulation of cell proliferation corresponded to the greater villus growth in colostrum than formula-fed calves on day 8 of life (Blum, 2006). The cell turnover of the intestinal mucosa depends on cell proliferation and programmed cell death (apoptosis; mainly seen at the villus tips) (Ramachandran *et al.*, 2000). Colostrum intake reduces apoptosis of epithelial cells and therefore prolongs the lifespan of the epithelial cells (Blum, 2006).

Milk-borne factors such as IGF-I are known for their stimulation of cell proliferation (Burrin *et al.*, 1996; Hammon *et al.*, 2013; Ontsouka *et al.*, 2016) as well as inhibition of cell death due to apoptosis or inflammation in the intestinal mucosa (Mylonas *et al.*, 2000; Blum, 2006). Recombinant human IGF-I, fed together with MR in neonatal calves, increases intestinal cell proliferation (Blum, 2006). A more distinct stimulation of mucosal cell proliferation is observed when feeding a colostrum extract (see above) instead of a single growth-promoting peptide (Roffler *et al.*, 2003). This finding indicates that not a single factor but the interaction of the large amount of growth-stimulating substances in the colostrum promotes intestinal cell proliferation and growth (Hammon *et al.*, 2012). Receptors for IGF-I, IGF-II and insulin (IGF1R, IGF2R and InsR, respectively) are present in the intestinal mucosa throughout the total gut in neonatal calves, and their expression and/or binding capacities are modified by colostrum feeding and orally administered rhIGF-I (Blum, 2006; Hammon *et al.*, 2013; Ontsouka *et al.*, 2016). The density of IGF1R and InsR, but not IGF2R, in the intestinal mucosa seems to be associated with crypt cell proliferation (Georgiev *et al.*, 2003).

Most biologically active factors in colostrum, such as IGF-I, IGF-II and insulin, are barely absorbed and therefore likely have no systemic function (Blum, 2006; Hammon *et al.*, 2013). Therefore, local effects of colostrum factors on crypt cell proliferation, intestinal epithelial growth and intestinal maturation may dominate in neonatal farm animals (Donovan and Odle, 1994; Xu, 1996; Blum, 2006). However, recent findings in calves indicate the absorption of colostrum adiponectin in neonatal calves (Kesser *et al.*, 2015), an adipokine involved in the regulation of insulin sensitivity (Kadowaki *et al.*, 2006). With respect to the lactocrine signalling theory described in pigs (Bartol *et al.*, 2013), the intestinal absorption of adiponectin may contribute to the stimulation of anabolic metabolism in neonatal calves after colostrum feeding (Hammon *et al.*, 2012). Colostrum feeding supports protein synthesis in the skeletal muscle of piglets (Burrin *et al.*, 1995), and the present research suggests a similar effect in neonatal calves, indicating enhanced protein synthesis in skeletal muscle due to insulin action after colostrum feeding (Hammon *et al.*, 2012; Sadri *et al.*, 2017). The impact of adiponectin on this finding, however, is not clear and needs further investigation.

#### *Colostrum supply and glucose metabolism*

Due to its growth-stimulating effect in the small intestine, colostrum intake promotes the absorptive capacity of the small intestine. Measurements of xylose and glucose absorption in neonatal calves clearly indicate a greater absorption after feeding with colostrum instead of formula or MR (Blum, 2006; Steinhoff-Wagner *et al.*, 2011; Gruse *et al.*, 2015). Xylose absorption on day 5 of age after feeding colostrum only once was similar to that after feeding colostrum for the first 3 days after birth (Hammon *et al.*, 2013). Therefore, the intake of first colostrum during the first hours after birth is of great importance for glucose absorption and

the postnatal glucose status in neonatal calves. In contrast, digestive enzymes and mucosal transporters with respect to carbohydrate digestion, such as lactase and SGLT1 and GLUT2, seem to be less affected by colostrum feeding. A distinct stimulation of lactase activity and the glucose transporter when feeding colostrum instead of a milk-based formula was barely observed in neonatal calves (Sauter *et al.*, 2004; Steinhoff-Wagner *et al.*, 2014). For more details concerning digestive enzymes in neonatal calves, readers are referred to Guilloteau *et al.* (2009a). Recent studies using metabolomics approaches in neonatal calves indicate that the uptake and metabolism of other nutrients (e.g. amino acids) are also influenced by colostrum feeding (Qi *et al.*, 2018; Zhao *et al.*, 2018).

First-pass glucose uptake in the splanchnic tissue on days 2 and 7 of life is greater in formula-fed than colostrum-fed calves, indicating a greater glucose utilisation in the splanchnic tissue (gastrointestinal tract and liver) of calves not fed with colostrum (Steinhoff-Wagner *et al.*, 2011; Gruse *et al.*, 2015). Possibly, nutrient absorption is generally impaired in formula-fed calves, leading to increased glucose utilisation in the splanchnic tissue, whereas colostrum-fed calves are able to use greater amounts of digested fat and protein as energy fuel in the splanchnic tissue. This hypothesis is supported by the finding that oral fat absorption is greater in colostrum- than in formula-fed or MR-fed calves, providing more fat (i.e. medium-chain fatty acids from colostrum) as energy fuel in the splanchnic tissue that partly can replace glucose utilisation (Hammon *et al.*, 2012). In contrast to intestinal glucose absorption, a stimulating influence of colostrum intake on endogenous glucose production, as supposed to be the case in piglets (Lepine *et al.*, 1991), does not occur in bovine neonates. Thus, growth-promoting substances of ingested colostrum do not affect endogenous glucose production in neonatal calves (Steinhoff-Wagner *et al.*, 2011). Nevertheless, the increased plasma glucose concentration and the greater hepatic glycogen content in colostrum-fed calves indicate an improved glucose status by colostrum feeding (Steinhoff-Wagner *et al.*, 2011; Hammon *et al.*, 2012 and 2013). These findings lead to the conclusion that the improved glucose status in calves fed colostrum immediately after birth and for 3 days is a result of enhanced glucose absorption and probably of less glucose utilisation in the splanchnic tissue but is not a result of increased endogenous glucose production.

#### *Colostrum supply and maturation in the somatotrophic axis*

The elevated glucose availability and the improved insulin status in colostrum-fed calves are important prerequisites for the accelerated maturation of the somatotrophic axis, as indicated by several studies in neonatal calves that were previously summarised (Blum, 2006; Hammon *et al.*, 2012). The stimulation of gastrointestinal hormones due to colostrum feeding may contribute to the elevated insulin secretion in the calves (Hadorn *et al.*, 1997; Inabu *et al.*, 2019). The elevated insulin status due to colostrum feeding in neonatal calves is probably the trigger for stimulating endogenous

IGF-I and the postnatal somatotrophic axis (Hammon *et al.*, 2012) because glucose and insulin stimulate the hepatic gene expression of the growth hormone receptor and IGF-I as well as IGF-I secretion (Brameld *et al.*, 1999; Butler *et al.*, 2003). On the other hand and as discussed earlier, studies in neonatal calves and piglets indicate no intestinal absorption of colostral IGF-I or insulin (Donovan *et al.*, 1997; Blum, 2006). Thus, the endogenously produced IGF-I determines the IGF-I status of the calf. Therefore, the nutrient supply is responsible for the maturation of the neonatal somatotrophic axis, and the colostral IGF-I promotes intestinal development of the neonatal calf but does not contribute to systemic IGF-I availability (Blum, 2006; Hammon *et al.*, 2012; Ontsouka *et al.*, 2016).

In summary, the postnatal maturation of the neonatal intestine is enhanced due to colostrum intake, and the improved intestinal maturation results in a greater nutrient absorption and stimulation of anabolic processes that are a prerequisite for accelerated postnatal growth.

### Development of the preweaning calf due to intensive milk feeding

#### *Definition of intensive milk feeding*

After the colostrum period, the calf depends on the intake of liquid feed in the form of milk or high-quality MR for nutrient supply. Although it is a common feeding strategy to increase solid feed intake as soon as possible in the preweaning period by reducing milk feeding (Huber, 1969; Gelsing *et al.*, 2016; Kertz *et al.*, 2017), solid feed intake during the first 3 weeks of age is low, and the digestion of solid feed is impaired due to the immature forestomach in the postnatal period (Drackley, 2008; Khan *et al.*, 2016; Frieten *et al.*, 2017). Thus, sufficient milk or MR supply during the first weeks of life is a prerequisite for calf growth and development. The World Organisation for Animal Health (OIE) defines animal welfare in the Terrestrial Animal Health Code as a state where the animal is healthy, comfortable, well nourished, safe, able to express natural behaviour and not suffering from pain, fear and distress (OIE, 2017). Feeding calves limited amounts of liquid feed (i.e. 4 to 6 kg/day) during the first weeks of life results in a lack of expression in natural suckling behaviour (Schuessler *et al.*, 2002; Miller-Cushon and DeVries, 2015) followed by hunger (Jensen and Holm, 2003; De Paula Vieira *et al.*, 2008; Borderas *et al.*, 2009; Gerbert *et al.*, 2018) and stress for the calves. Allowing calves to drink unlimited amounts of milk or MR for several weeks during the preweaning period more than doubles liquid feed intake compared with restricted amounts of 4 to 6 kg/day of MR or milk (Hammon *et al.*, 2002; Maccari *et al.*, 2015; Schäff *et al.*, 2016; Frieten *et al.*, 2017). Therefore, an intensive milk-feeding programme contributes to the overall well-being of preweaning calves (Von Keyserlingk *et al.*, 2009; FAWC, 2015; OIE, 2017).

An intensive milk feeding provides milk or MR in unrestricted amounts all day long for 24 h. The calves have *ad*

*libitum* access to milk (Jasper and Weary, 2002; Schuessler *et al.*, 2002; Maccari *et al.*, 2015) or MR feeding (Schäff *et al.*, 2016; Frieten *et al.*, 2017; Korst *et al.*, 2017) for several weeks. *Ad libitum* MR feeding (125 g powder/l, 21.7% CP and 18.3 MJ metabolisable energy (ME) per kg DM) provides on average 1.6 kg DM/day, 35 MJ ME/day and 347 g protein/day to the calves during the intensive milk-feeding period. In this context, daily peaks of more than 2 kg DM intake were observed in previous studies (Schäff *et al.*, 2016; Frieten *et al.*, 2017). Such feeding schedules provide much more protein and energy than commonly used milk-feeding programmes of 4 to 6 kg milk/day. *Ad libitum* milk-feeding programmes are comparable to early calf rearing in beef production where calves live together with their dams and have free access to milk all day long (Egli and Blum, 1998; Schuessler *et al.*, 2002). Main findings of an intensive milk feeding protocol compared to restricted milk feeding on calf growth, organ development, metabolic and endocrine changes, feeding behaviour and immune response have been summarised in Table 1.

#### *Stimulation of growth and endocrine growth regulation by intensive milk feeding*

*Ad libitum* milk or MR feeding (Maccari *et al.*, 2015; Schäff *et al.*, 2016; Korst *et al.*, 2017) or enhanced milk-feeding programmes using MR with a greater protein content (up to 30% CP in DM) (Smith *et al.*, 2002; Geiger *et al.*, 2016) resulted in an elevated body growth during the preweaning period when compared to restricted milk or MR feeding (4 to 6 kg milk or MR/day). In addition to stimulating muscle and fat growth (Bartlett *et al.*, 2006; Geiger *et al.*, 2016; Schäff *et al.*, 2016; Koch *et al.*, 2019), intensive milk or MR intake accelerates organ growth, for example, small intestine, mammary gland, thymus and endocrine pancreas (Prokop *et al.*, 2015; Geiger *et al.*, 2016; Soberon and Van Amburgh, 2017; Koch *et al.*, 2019). As discussed later in this review, the velocity of body growth could temporarily decrease in intensive milk-fed calves during the weaning process due to the adaptation to solid feed intake. However, BW at the end of the weaning process is still greater in intensively than restrictively milk-fed calves (Maccari *et al.*, 2015; Schäff *et al.*, 2016; Frieten *et al.*, 2017).

Studies on the hepatic transcriptome and proteome in lambs and metabolome in blood plasma of calves reveal marked changes with respect to protein and energy metabolism when animals receive MR *ad libitum* instead of in restricted amounts (Kenéz *et al.*, 2018; Santos *et al.*, 2019). In lambs, restricted MR feeding stimulates hepatic pathways involved in gluconeogenesis, amino acid degradation and hepatic fatty acid oxidation, which points at changes in energy utilisation to stabilise glucose homeostasis as compared to *ad libitum* MR-fed lambs (Santos *et al.*, 2019). In calves, *ad libitum* instead of restricted MR feeding seems to increase the capacity for mitochondrial transport of fatty acids and probably affects fatty acid oxidation as well (Kenéz *et al.*, 2018). In addition, an enhanced MR-feeding programme leads to a greater metabolic activity in muscle

**Table 1** Effects of intensive milk feeding immediately after birth on growth, development, behaviour and immune response in preweaning calves<sup>1</sup>

Item effect due to intensive feeding when compared to restricted milk feeding	Type of intensive milk or milk replacer feeding <sup>2</sup>	Duration in weeks	Reference
Increased BW and body growth	<i>Ad libitum</i> feeding of milk powder	4	Hammon <i>et al.</i> (2002)
	<i>Ad libitum</i> feeding of whole milk	5	Jasper and Weary (2002)
	20% of BW/day of whole milk	3.5	Khan <i>et al.</i> (2007b)
	1.4 kg DM/day milk replacer with 27% CP and 28% crude fat	9	Daniels <i>et al.</i> (2009)
	2.1% of BW DM/day milk replacer with 30.6% CP and 16.1% crude fat	4	Davis Rincker <i>et al.</i> (2011)
	<i>Ad libitum</i> feeding of whole milk	3	Maccari <i>et al.</i> (2015)
	1.08 kg DM/day milk replacer with 28.9% CP and 26.2% crude fat	8	Geiger <i>et al.</i> (2016)
	<i>Ad libitum</i> feeding of milk replacer	5	Schäff <i>et al.</i> (2016)
	<i>Ad libitum</i> feeding of milk replacer	8	Frieten <i>et al.</i> (2017)
	<i>Ad libitum</i> feeding of milk replacer or whole milk	3.5	Korst <i>et al.</i> (2017)
Organ growth and development			
Elevated forestomach and rumen papillae growth postweaning	20% of BW/day of whole milk	3.5	Khan <i>et al.</i> (2007a)
	2% of BW DM/day milk replacer with 28.5% CP and 15% crude fat	5	Naeem <i>et al.</i> (2012)
No impairment on rumen development postweaning	<i>Ad libitum</i> feeding of milk replacer	5	Schäff <i>et al.</i> (2018)
	<i>Ad libitum</i> feeding of milk replacer	8	Koch <i>et al.</i> (2019)
Enhanced muscle and body fat growth	2% of BW DM/day milk replacer with 28.5% CP and 15% crude fat	5	Wang <i>et al.</i> (2014)
	<i>Ad libitum</i> feeding of milk replacer	5	Schäff <i>et al.</i> (2016)
Enhanced growth of the small intestine	<i>Ad libitum</i> feeding of milk replacer	8	Koch <i>et al.</i> (2019)
Enhanced growth and development of the mammary gland	1.4 kg DM/day milk replacer with 27% CP and 28% crude fat	9	Daniels <i>et al.</i> (2009)
	1.08 kg DM/day milk replacer with 28.9% CP and 26.2% crude fat	8	Geiger <i>et al.</i> (2016)
	1.24 MJ/kg BW <sup>0.75</sup> per day milk replacer with 28.5% CP and 15% crude fat	8	Soberon and Van Amburgh (2017)
Stimulation of the endocrine pancreas secretion	<i>Ad libitum</i> feeding of whole milk	3	Prokop <i>et al.</i> (2015)
Metabolic and endocrine changes			
Systemic metabolic changes indicating greater metabolic activity	<i>Ad libitum</i> feeding of milk replacer	5	Schäff <i>et al.</i> (2016)
	<i>Ad libitum</i> feeding of milk replacer	8	Frieten <i>et al.</i> (2017)
	<i>Ad libitum</i> feeding of milk replacer or whole milk	3.5	Kenéz <i>et al.</i> (2018)
Elevated plasma insulin and stimulation of the somatotrophic axis	20% of BW/day of whole milk	3.5	Khan <i>et al.</i> (2007a)
	1.4 kg DM/day milk replacer with 27% CP and 28% crude fat	9	Daniels <i>et al.</i> (2008)
	<i>Ad libitum</i> feeding of whole milk	3	Maccari <i>et al.</i> (2015)
	<i>Ad libitum</i> feeding of milk replacer	5	Schäff <i>et al.</i> (2016)
	<i>Ad libitum</i> feeding of milk replacer	8	Frieten <i>et al.</i> (2018)
	10 kg/day of whole milk	7	Haisan <i>et al.</i> (2018)
Elevated metabolic activity in the ruminal epithelium	2% of BW DM/day milk replacer with 28.5% CP and 15% crude fat	5	Naeem <i>et al.</i> (2014)
Elevated metabolic activity in the omental adipose tissue	1.26 MJ/kg BW <sup>0.75</sup> per day milk replacer with 28.5% CP and 15% crude fat	8	Leal <i>et al.</i> (2018)

Table 1 (Continued)

Item effect due to intensive feeding when compared to restricted milk feeding	Type of intensive milk or milk replacer feeding <sup>2</sup>	Duration in weeks	Reference
Feeding behaviour			
Elevated milk/milk replacer intake and less sign of hunger	<i>Ad libitum</i> feeding of milk powder	4	Hammon <i>et al.</i> (2002)
	<i>Ad libitum</i> feeding of whole milk	5	Jasper and Weary (2002)
	8 kg/day of milk and milk replacer/day <sup>3</sup>	7	Jensen and Holm (2003)
	<i>Ad libitum</i> feeding of whole milk	1	De Paula Vieira <i>et al.</i> (2008)
	<i>Ad libitum</i> feeding of milk replacer; 12 kg/day of whole milk	6	Borderas <i>et al.</i> (2009) <sup>4</sup>
		6	
	<i>Ad libitum</i> feeding of milk replacer	6	Miller-Cushon <i>et al.</i> (2013)
	<i>Ad libitum</i> feeding of milk replacer	5	Schäff <i>et al.</i> (2016)
	<i>Ad libitum</i> feeding of milk replacer	8	Gerbert <i>et al.</i> (2018)
Immune response and health			
Altered interferon- $\gamma$ and <i>no</i> production of peripheral blood mononuclear cells <i>ex vivo</i>	2.5% of BW DM milk replacer with 30% CP and 20% crude fat	8.5	Nonnecke <i>et al.</i> (2003)
Greater plasma haptoglobin concentration 24 h after LPS challenge; greater neutrophil oxidative burst intensity after <i>Escherichia coli</i> co-culture	810 and 1180 g DM/day of milk replacer with 28% CP and 20% crude fat	6	Ballou (2012)
Faster improvement of faecal scores after infection with <i>Cryptosporidium parvum</i>	0.95 to 1.24 MJ/kg BW <sup>0.75</sup> per day milk replacer with 28% CP and 20% crude fat	3	Ollivett <i>et al.</i> (2012)
Decreased systemic inflammation after oral <i>Salmonella typhimurium</i> challenge in postweaning calves	610 and 735 g DM/day of milk replacer with 28% CP and 25% crude fat	6	Ballou <i>et al.</i> (2015)
Reduced plasma fibrinogen concentration in blood	<i>Ad libitum</i> feeding of milk replacer	5	Schäff <i>et al.</i> (2016)
Enhanced activation of pathways related to the intestinal immune system	<i>Ad libitum</i> feeding of milk replacer	8	Hammon <i>et al.</i> (2018)

<sup>1</sup>Intensive milk feeding is defined as daily milk or milk replacer intake of 20% of BW, *ad libitum* milk or milk replacer feeding or feeding enhanced amounts of milk replacer with elevated CP and fat content.

<sup>2</sup>When not stated in the table, milk replacer contained 21% to 23% of CP and 17% to 20% of crude fat based on DM. Whole milk contained 320 to 350 g CP and 370 to 400 g crude fat/kg milk.

<sup>3</sup>Data apply only for Holstein calves.

<sup>4</sup>Two studies in the reference.

and fat tissue as well as the ruminal epithelium (Naeem *et al.*, 2012 and 2014; Wang *et al.*, 2014; Leal *et al.*, 2018).

The improved growth development and protein accretion in calves fed intensively with milk or MR are confirmed by the stimulation of the somatotrophic axis (Maccari *et al.*, 2015; Schäff *et al.*, 2016; Frieten *et al.*, 2018; Haisan *et al.*, 2018). Important elements of the somatotrophic axis are growth hormone (GH), IGF-I and several IGF-binding proteins (IGFBPs). The postnatal interaction of GH, IGF-I and IGFBP affects body growth and organ development in mammals (Breier *et al.*, 2000), including the development of the mammary gland (Akers, 2006) and immune function (Clark, 1997). The stimulation of the postnatal somatotrophic axis depends on the nutrient supply and reflects the glucose and insulin status of the animal (Brameld *et al.*, 1999; Renaville *et al.*, 2000; Smith *et al.*, 2002). Plasma IGF-I and IGFBP-3 concentrations are elevated, and the IGFBP-2 concentration is decreased during growth in well-nourished animals as compared to animals of same age with restricted feed intake (Breier *et al.*, 2000; Renaville *et al.*, 2000). A key factor in maturation of the somatotrophic axis is the increased expression of the GH receptor, particularly in the liver, with age (Breier *et al.*, 2000; Hammon *et al.*, 2012). The GH receptor mediates GH action on IGF-I synthesis and secretion and is stimulated by insulin (Breier *et al.*, 2000; Butler *et al.*, 2003). The glucose, insulin, IGF-I and IGFBP-3 plasma concentrations are much greater, and hepatic gene expression of the GH receptor and IGF-I is higher in intensively milk-fed calves than in calves with restricted milk intake (e.g., 6 l MR/day; Maccari *et al.*, 2015; Schäff *et al.*, 2016; Frieten *et al.*, 2017 and 2018). The IGFBP-2 plasma concentration and hepatic gene expression behave the other way round, as expected from the literature (Renaville *et al.*, 2000; Schäff *et al.*, 2016; Frieten *et al.*, 2018). No signs of impaired insulin response are seen during enhanced milk feeding in calves (MacPherson *et al.*, 2019).

During the first weeks of life, the elevated concentrate intake in restrictively milk-fed calves cannot compensate for impaired nutrient intake due to reduced milk feeding, and consequently, the somatotrophic axis is not stimulated during early postnatal life when concentrate and forage feeding are favoured instead of milk feeding. In particular, the elevated IGFBP-2 plasma concentration in milk-restricted calves indicates an impaired nutrient intake (Renaville *et al.*, 2000; Schäff *et al.*, 2016; Frieten *et al.*, 2018). An impaired nutrient supply with decreased IGF-I and IGFBP-3 and increased IGFBP-2 plasma concentrations also occurs during weaning when milk feeding is reduced too quickly and the solid feed intake does not meet nutrition requirements (Schäff *et al.*, 2016; Frieten *et al.*, 2018). These changes in the somatotrophic axis are reflected by a depressed growth rate during the weaning process (Schäff *et al.*, 2016; Frieten *et al.*, 2017). To prevent a growth depression during weaning in calves with an intensified milk-feeding programme, a delayed weaning age or individual weaning based on solid feed intake is recommended (Eckert *et al.*, 2015; de Passillé and Rushen, 2016; Welboren *et al.*, 2019).

Parameters of the somatotrophic axis may provide useful information on the metabolic status and may help to avoid detrimental weaning programmes in calves.

#### *Development and maturation of the gastrointestinal tract and immune response by intensive milk feeding*

An intensive milk feeding strategy affects intestinal development. The size and the absorptive capacity increase in pre-weaning calves with an enhanced MR-feeding programme (Geiger *et al.*, 2016; Koch *et al.*, 2019). In addition, an intensive milk or MR-feeding programme seems to stimulate the expression of long non-coding RNA involved in the regulation of tight-junction protein synthesis (Weikard *et al.*, 2018). It is well established in ruminants that the diet affects tight-junction protein expression (Steele *et al.*, 2016). According to the upregulation of tight-junction protein-encoding genes, it is suggested that the first week of life is crucial for the development of the intestinal epithelium and the intestinal barrier of the mucosal immune system in calves (Malmuthuge and Guan, 2017). In piglets, specific amino acids can affect intestinal permeability and integrity, protein synthesis, intestinal repair after injury and cell proliferation in the gastrointestinal tract (Jacobi and Odle, 2012). Because of its composition and ingredients, milk provide the best conditions for nutrient supply in the postnatal and preweaning period to promote intestinal integrity in the bovine (Steele *et al.*, 2016).

There is growing evidence that an adequate nutrient supply is important for the maturation of the intestinal immune system and for successful defence against pathogens (Khan *et al.*, 2011; Hammon *et al.*, 2018). A greater nutrient supply may have beneficial effects on intestinal maturation, including the generation of a proper adaptive immune system and a stable microbiota, which may protect against diarrhoeal diseases in the neonatal and preweaning period. Common feeding schedules of not more than 6 kg of milk per day may delay the establishment of a proper immune response and microbiota in the intestine. Studies in neonatal calves have investigated the diet-dependent establishment of the intestinal microbiome, but the impact of the milk amount on the intestinal microbiome is still unclear (Malmuthuge and Guan, 2017). A higher plane (considering the amount of MR, concentration of MR and protein and fat concentration) of nutrition seems to protect the intestine against pathogenic infections and promote overcoming of pathogenic infections. Calves with a higher plane of nutrition (intake energy: 1.3 MJ/kg metabolic BW instead of 0.5 MJ/kg metabolic BW by MR feeding) indicate a faster resolution from diarrhoea caused by infection with *Cryptosporidium parvum* (Ollivett *et al.*, 2012). Ballou *et al.* (2015) showed that a higher plane of nutrition (610 and 735 g/day DM MR during week 1 and weeks 2 to 6, respectively, of a 28% CP and 25% fat MR instead of 409 g/day DM MR of a 20% CP and 20% fat MR) results in a higher resistance against *Salmonella typhimurium* in postweaning calves. Even there is first evidence that an intensive milk or MR feeding stimulates intestinal development and maturation and the intestinal immune

response (Hammon *et al.*, 2018), more studies are needed to investigate the interaction between the changes of the intestinal microbiome and immune response due to an intensive milk-feeding programme.

An intensive milk feeding may delay rumen development by reducing solid feed intake during the early preweaning period (Baldwin *et al.*, 2004; Khan *et al.*, 2011 and 2016). However, solid feed intake and rumen development accelerate during the weaning process, and rumen function is not impaired at the end of the weaning process when calves received 20% instead of 10% of BW milk per day (Khan *et al.*, 2011). Rumen papilla growth and concentrations of volatile fatty acids are the same when calves are fed MR *ad libitum* for 5 and 8 weeks after birth, respectively, compared to 6 kg/day MR intake (Schäff *et al.*, 2018; Koch *et al.*, 2019). These findings are supported by the fact that *ad libitum* milk-fed calves immediately increase their solid feed intake and plasma  $\beta$ -hydroxybutyrate concentration in blood when MR intake is reduced (Schäff *et al.*, 2016; Frieten *et al.*, 2017; Welboren *et al.*, 2019). Plasma  $\beta$ -hydroxybutyrate results from ketogenesis from butyrate in the rumen epithelial cells and is an indicator for maturation of the rumen function (Baldwin *et al.*, 2004). Interestingly, the metabolic activity in ruminal epithelium is enhanced in calves fed elevated amounts of MR (Naeem *et al.*, 2012 and 2014).

In summary, an intensive milk-feeding regime is required to realise the potential for growth and development in preweaning calves. Body growth and organ maturation are improved with an intensive milk-feeding programme, and calves are less hungry and probably more resilient to infectious diseases during the preweaning period. The development of the forestomach could be delayed during the intensive milk-feeding period, but applying an adapted weaning protocol for intensive milk-fed calves avoids an impaired rumen development and body growth depression during the postweaning period.

### Impact of preweaning growth and development on lifetime performance and health

Colostrum and milk feeding not only influence the postnatal and preweaning development and growth of the calves but also influence performance and health in later life (Van Amburgh and Soberon, 2013; Huber, 2018). The improved mammary gland development during the preweaning rearing period is an example of the importance of the nutrient supply during the preweaning period for organ development (Geiger *et al.*, 2016; Soberon and Van Amburgh, 2017), with consequences for lifetime performance (Van Amburgh and Soberon, 2013). However, presently it is not known whether early calf nutrition has long-lasting effects on other organ systems, for example, the liver or the immune system. Culling rates are still high in dairy production, and metabolic diseases and immune suppression around calving are heavily involved in this unfavourable situation (Hare *et al.*, 2006;


Ingvartsen and Moyes, 2015; Probo *et al.*, 2018; Gross and Bruckmaier, 2019). Interestingly, different patterns of metabolic parameters due to variable milk or MR feeding in preweaning calves seem to be maintained when calves become dairy cows, and epigenetic effects due to different milk-feeding programmes during the preweaning period have been assumed (Kenéz *et al.*, 2018). Postnatal nutritional programming is well known from human studies and in other species (Guilloteau *et al.*, 2009b), but it still remains unclear whether variable metabolic profiles of young individuals are conserved and are the basis for the different metabolic types of the adults in their productive life span. Thus, more research is needed to investigate the impact of early calf nutrition on metabolic performance in later life and whether early calf nutrition may improve robustness and resilience in dairy cows.

### Conclusions

An intensive milk-feeding programme, starting immediately after birth, with an enhanced colostrum intake and subsequent intensive milk feeding supports postnatal growth and development of dairy calves, prevents behavioural anomalies and promotes the raising of robust young animals. Providing only 4 to 6 kg milk/day to the preweaning calves is not consistent with animal welfare principles (FAWC, 2015; OIE, 2017; Huber, 2018). Thus, a change of the early calf management is needed to follow the natural processes of preweaning calf rearing, for example, as known from beef calf management (Egli and Blum, 1998; Schiessler *et al.*, 2002). Research will continue to investigate the impact of an intensive milk-feeding regime on raising robust and well-performing dairy cows and bulls.

### Acknowledgements

This review is based on an invited presentation at the 13th International Symposium on Ruminant Physiology (ISRP 2019) held in Leipzig, Germany, September 2019.

 H. M. Hammon 0000-0001-8698-1257

### Declaration of interest

The authors declare no conflicts of interest.

### Ethics statement

None.

### Software and data repository resources

None.

### References

- Akers RM 2006. Major advances associated with hormone and growth factor regulation of mammary growth and lactation in dairy cows. *Journal of Dairy Science* 89, 1222–1234.
- Baldwin RL, McLeod KR, Klotz JL and Heitmann RN 2004. Rumen development, intestinal growth and hepatic metabolism in the pre- and postweaning ruminant. *Journal of Dairy Science* 87, E55–E65.



- Ballou MA 2012. Immune responses of Holstein and Jersey calves during the preweaning and immediate postweaned periods when fed varying planes of milk replacer. *Journal of Dairy Science* 95, 7319–7330.
- Ballou MA, Hanson DL, Cobb CJ, Obeidat BS, Sellers MD, Pepper-Yowell AR, Carroll JA, Earleywine TJ and Lawhon SD 2015. Plane of nutrition influences the performance, innate leukocyte responses, and resistance to an oral *Salmonella enterica* serotype Typhimurium challenge in Jersey calves. *Journal of Dairy Science* 98, 1972–1982.
- Barrington GM and Parish SM 2001. Bovine neonatal immunology. *Veterinary Clinics of North America: Food Animal Practice* 17, 463–476.
- Bartlett KS, McKeith FK, VandeHaar MJ, Dahl GE and Drackley JK 2006. Growth and body composition of dairy calves fed milk replacers containing different amounts of protein at two feeding rates. *Journal of Animal Science* 84, 1454–1467.
- Bartol FF, Wiley AA, Miller DJ, Silva AJ, Roberts KE, Davolt ML, Chen JC, Frankshun AL, Camp ME, Rahman KM, Vallet JL and Bagnell CA 2013. Lactation biology symposium: lactocrine signaling and developmental programming. *Journal of Animal Science* 91, 696–705.
- Blum JW 2006. Nutritional physiology of neonatal calves. *Journal of Animal Physiology and Animal Nutrition* 90, 1–11.
- Blum JW and Baumrucker CR 2008. Insulin-like growth factors (IGFs), IGF binding proteins, and other endocrine factors in milk: role in the newborn. *Advances in Experimental Medicine and Biology* 606, 397–422.
- Borderas TF, de Passille AMB and Rushen J 2009. Feeding behavior of calves fed small or large amounts of milk. *Journal of Dairy Science* 92, 2843–2852.
- Brameld JM, Gilmour RS and Buttery PJ 1999. Glucose and amino acids interact with hormones to control expression of insulin-like growth factor-I and growth hormone receptor mRNA in cultured pig hepatocytes. *Journal of Nutrition* 129, 1298–1306.
- Breier BH, Oliver MH and Gallaher BW 2000. Regulation of growth and metabolism during postnatal development. In *Ruminant physiology: digestion, metabolism, growth and reproduction* (ed. PB Cronjé 2000). Regulation of growth and metabolism during postnatal development. In *Ruminant physiology: digestion, metabolism, growth and reproduction* (ed. ), pp. 187–204. CABI Publishing, New York, NY, USA.
- Burrin DG, Davis TA, Ebner S, Schoknecht PA, Fiorotto ML, Reeds PJ and McAvoy S 1995. Nutrient-independent and nutrient-dependent factors stimulate protein synthesis in colostrum-fed newborn pigs. *Pediatric Research* 37, 593–599.
- Burrin DG, Wester TJ, Davis TA, Amick S and Heath JP 1996. Orally administered IGF-I increases intestinal mucosal growth in formula-fed neonatal pigs. *American Journal of Physiology* 270, R1085–1091.
- Butler ST, Marr AL, Pelton SH, Radcliff RP, Lucy MC and Butler WR 2003. Insulin restores GH responsiveness during lactation-induced negative energy balance in dairy cattle: effects on expression of IGF-I and GH receptor 1A. *Journal of Endocrinology* 176, 205–217.
- Chase CCL, Hurley DJ and Reber AJ 2008. Neonatal immune development in the calf and its impact on vaccine response. *Veterinary Clinics of North America: Food Animal Practice* 24, 87–104.
- Clark R 1997. The somatogenic hormones and insulin-like growth factor-1: stimulators of lymphopoiesis and immune function. *Endocrine Reviews* 18, 157–179.
- Compton CWR, Heuer C, Thomsen PT, Carpenter TE, Phyn CVC and McDougall S 2017. Invited review: a systematic literature review and meta-analysis of mortality and culling in dairy cattle. *Journal of Dairy Science* 100, 1–16.
- Daniels KM, Capuco AV, McGilliard ML, James RE and Akers RM 2009. Effects of milk replacer formulation on measures of mammary growth and composition in Holstein heifers. *Journal of Dairy Science* 92, 5937–5950.
- Daniels KM, Hill SR, Knowlton KF, James RE, McGilliard ML and Akers RM 2008. Effects of milk replacer composition on selected blood metabolites and hormones in preweaned Holstein heifers. *Journal of Dairy Science* 91, 2628–2640.
- Davis Rincker LE, Vandehaar MJ, Wolf CA, Liesman JS, Chapin LT and Weber Nielsen MS 2011. Effect of intensified feeding of heifer calves on growth, pubertal age, calving age, milk yield, and economics. *Journal of Dairy Science* 94, 3554–3567.
- de Passillé AM and Rushen J 2016. Using automated feeders to wean calves fed large amounts of milk according to their ability to eat solid feed. *Journal of Dairy Science* 99, 3578–3583.
- Department for Environment, Food and Rural Affairs (DEFRA) 2003. Improving calf survival. Retrieved on 20 June 2019 from <http://adlib.everysite.co.uk/resources/000/020/709/calfsurvival.pdf>
- De Paula Vieira A, Guesdon V, De Passille AM, von Keyserlingk MAG and Weary DM 2008. Behavioural indicators of hunger in dairy calves. *Applied Animal Behaviour Science* 109, 180–189.
- Donovan SM, Chao J, Zijlstra RT and Odle J 1997. Orally administered iodinated recombinant human insulin-like growth factor-I (I-125-rhIGF-I) is poorly absorbed by the newborn piglet. *Journal of Pediatric Gastroenterology and Nutrition* 24, 174–182.
- Donovan SM and Odle J 1994. Growth factors in milk as mediators of infant development. *Annual Review of Nutrition* 14, 147–167.
- Drackley JK 2008. Calf nutrition from birth to breeding. *Veterinary Clinics of North America: Food Animal Practice* 24, 55–86.
- Eckert E, Brown HE, Leslie KE, DeVries TJ and Steele MA 2015. Weaning age affects growth, feed intake, gastrointestinal development, and behavior in Holstein calves fed an elevated plane of nutrition during the preweaning stage. *Journal of Dairy Science* 98, 6315–6326.
- Egli CP and Blum JW 1998. Clinical, haematological, metabolic and endocrine traits during the first three months of life of suckling simmental calves held in a cow-calf operation. *Zentralblatt für Veterinärmedizin. Reihe A* 45, 99–118.
- Farm Animal Welfare Committee (FAWC) 2015. Opinion on the welfare implications of nutritional management strategies for artificially-reared calves from birth to weaning. FAWC, University of Bristol, Bristol, UK.
- Frieten D, Gerbert C, Koch C, Dusel G, Eder K, Kanitz E, Weitzel JM and Hammon HM 2017. Ad libitum milk replacer feeding, but not butyrate supplementation, affects growth performance as well as metabolic and endocrine traits in Holstein calves. *Journal of Dairy Science* 100, 6648–6661.
- Frieten D, Gerbert C, Koch C, Dusel G, Eder K, Hoeflich A, Mielenz B and Hammon HM 2018. Influence of ad libitum milk replacer feeding and butyrate supplementation on the systemic and hepatic insulin-like growth factor I and its binding proteins in Holstein calves. *Journal of Dairy Science* 101, 1661–1672.
- Geiger AJ, Parsons CLM, James RE and Akers RM 2016. Growth, intake, and health of Holstein heifer calves fed an enhanced preweaning diet with or without postweaning exogenous estrogen. *Journal of Dairy Science* 99, 3995–4004.
- Gelsing SL, Heinrichs AJ and Jones CM 2016. A meta-analysis of the effects of preweaned calf nutrition and growth on first-lactation performance. *Journal of Dairy Science* 99, 6206–6214.
- Georgiev IP, Georgieva TM, Pfaffl M, Hammon HM and Blum JW 2003. Insulin-like growth factor and insulin receptors in intestinal mucosa of neonatal calves. *Journal of Endocrinology* 176, 121–132.
- Gerbert C, Frieten D, Koch C, Dusel G, Eder K, Stefaniak T, Bajzert J, Jawor P, Tuchscherer A and Hammon HM 2018. Effects of ad libitum milk replacer feeding and butyrate supplementation on behavior, immune status, and health of Holstein calves in the postnatal period. *Journal of Dairy Science* 101, 7348–7360.
- Godden S 2008. Colostrum management for dairy calves. *Veterinary Clinics of North America: Food Animal Practice* 24, 19–39.
- Gross JJ and Bruckmaier RM 2019. Invited review: metabolic challenges and adaptation during different functional stages of the mammary gland in dairy cows: perspectives for sustainable milk production. *Journal of Dairy Science* 102, 2828–2843.
- Gruse J, Görs S, Tuchscherer A, Otten W, Weitzel JM, Metges CC, Wolfram S and Hammon HM 2015. The effects of oral quercetin supplementation on splanchnic glucose metabolism in 1-week-old calves depend on diet after birth. *Journal of Nutrition* 145, 2486–2495.
- Guilloteau P, Zabielski R and Blum JW 2009a. Gastrointestinal tract and digestion in the young ruminant: ontogenesis, adaptations, consequences and manipulations. *Journal of Physiology and Pharmacology* 60 (suppl. 3), 37–46.
- Guilloteau P, Zabielski R, Hammon HM and Metges CC 2009b. Adverse effects of nutritional programming during prenatal and early postnatal life, some aspects of regulation and potential prevention and treatments. *Journal of Physiology and Pharmacology* 60 (suppl. 3), 17–35.
- Hadorn U, Hammon H, Bruckmaier RM and Blum JW 1997. Delaying colostrum intake by one day has important effects on metabolic traits and on gastrointestinal and metabolic hormones in neonatal calves. *Journal of Nutrition* 127, 2011–2023.
- Haisan J, Oba M, Ambrose DJ and Steele MA 2018. Short communication: the effects of offering a high or low plane of milk preweaning on insulin-like growth

- factor and insulin-like growth factor binding proteins in dairy heifer calves. *Journal of Dairy Science* 101, 11441–11446.
- Hammon HM, Frieten D, Gerbert C, Koch C, Dusel G, Weikard R and Kühn C 2018. Different milk diets have substantial effects on the jejunal mucosal immune system of pre-weaning calves, as demonstrated by whole transcriptome sequencing. *Scientific Reports* 8, 1693.
- Hammon HM, Schiessler G, Nussbaum A and Blum JW 2002. Feed intake patterns, growth performance, and metabolic and endocrine traits in calves fed unlimited amounts of colostrum and milk by automate, starting in the neonatal period. *Journal of Dairy Science* 85, 3352–3362.
- Hammon HM, Steinhoff-Wagner J, Flor J, Schönhusen U and Metges CC 2013. Lactation biology symposium: role of colostrum and colostrum components on glucose metabolism in neonatal calves. *Journal of Animal Science* 91, 685–695.
- Hammon HM, Steinhoff-Wagner J, Schönhusen U, Metges CC and Blum JW 2012. Energy metabolism in the newborn farm animal with emphasis on the calf: endocrine changes and responses to milk-borne and systemic hormones. *Domestic Animal Endocrinology* 43, 171–185.
- Hare E, Norman HD and Wright JR 2006. Survival rates and productive herd life of dairy cattle in the United States. *Journal of Dairy Science* 89, 3713–3720.
- Huber JT 1969. Development of the digestive and metabolic apparatus of the calf. *Journal of Dairy Science* 52, 1303–1315.
- Huber K 2018. Invited review: resource allocation mismatch as pathway to disproportionate growth in farm animals—prerequisite for a disturbed health. *Animal* 12, 528–536.
- Inabu Y, Pyo J, Pletts S, Guan LL, Steele MA and Sugino T 2019. Effect of extended colostrum feeding on plasma glucagon-like peptide-1 concentration in newborn calves. *Journal of Dairy Science* 102, 4619–4627.
- Ingvartsen KL and Moyes KM 2015. Factors contributing to immunosuppression in the dairy cow during the periparturient period. *Japanese Journal of Veterinary Research* 63 (suppl. 1), S15–24.
- Jacobi SK and Odle J 2012. Nutritional factors influencing intestinal health of the neonate. *Advances in Nutrition* 3, 687–696.
- Jasper J and Weary DM 2002. Effects of ad libitum milk intake on dairy calves. *Journal of Dairy Science* 85, 3054–3058.
- Jensen MB and Holm L 2003. The effect of milk flow rate and milk allowance on feeding related behaviour in dairy calves fed by computer controlled milk feeders. *Applied Animal Behaviour Science* 82, 87–100.
- Johnson LR 1988. Regulation of gastrointestinal mucosal growth. *Physiological Reviews* 68, 456–502.
- Kadowaki T, Yamauchi T, Kubota N, Hara K, Ueki K and Tobe K 2006. Adiponectin and adiponectin receptors in insulin resistance, diabetes, and the metabolic syndrome. *Journal of Clinical Investigation* 116, 1784–1792.
- Kenéz Á, Koch C, Korst M, Kesser J, Eder K, Sauerwein H and Huber K 2018. Different milk feeding intensities during the first 4 weeks of rearing dairy calves: part 3: plasma metabolomics analysis reveals long-term metabolic imprinting in Holstein heifers. *Journal of Dairy Science* 101, 8446–8460.
- Kertz AF, Hill TM, Quigley JD 3rd, Heinrichs AJ, Linn JG and Drackley JK 2017. A 100-year review: calf nutrition and management. *Journal of Dairy Science* 100, 10151–10172.
- Kesser J, Hill M, Heinz JF, Koch C, Rehage J, Steinhoff-Wagner J, Hammon HM, Mielenz B, Sauerwein H and Sadri H 2015. The rapid increase of circulating adiponectin in neonatal calves depends on colostrum intake. *Journal of Dairy Science* 98, 7044–7051.
- Khan MA, Bach A, Weary DM and von Keyserlingk MAG 2016. Invited review: transitioning from milk to solid feed in dairy heifers. *Journal of Dairy Science* 99, 885–902.
- Khan MA, Weary DM and von Keyserlingk MA 2011. Invited review: effects of milk ration on solid feed intake, weaning, and performance in dairy heifers. *Journal of Dairy Science* 94, 1071–1081.
- Khan MA, Lee HJ, Lee WS, Kim HS, Ki KS, Hur TY, Suh GH, Kang SJ and Choi YJ 2007a. Structural growth, rumen development, and metabolic and immune responses of Holstein male calves fed milk through step-down and conventional methods. *Journal of Dairy Science* 90, 3376–3387.
- Khan MA, Lee HJ, Lee WS, Kim HS, Kim SB, Ki KS, Ha JK, Lee HG and Choi YJ 2007b. Pre- and postweaning performance of Holstein female calves fed milk through step-down and conventional methods. *Journal of Dairy Science* 90, 876–885.
- Koch C, Gerbert C, Frieten D, Dusel G, Eder K, Zitnan R and Hammon HM 2019. Effects of ad libitum milk replacer feeding and butyrate supplementation on the epithelial growth and development of the gastrointestinal tract in Holstein calves. *Journal of Dairy Science* 102, 8513–8526.
- Korst M, Koch C, Kesser J, Müller U, Romberg FJ, Rehage J, Eder K and Sauerwein H 2017. Different milk feeding intensities during the first 4 weeks of rearing in dairy calves: part 1: effects on performance and production from birth over the first lactation. *Journal of Dairy Science* 100, 3096–3108.
- Langel SN, Wark WA, Garst SN, James RE, McGilliard ML, Petersson-Wolfe CS and Kanevsky-Mullarky I 2015. Effect of feeding whole compared with cell-free colostrum on calf immune status: the neonatal period. *Journal of Dairy Science* 98, 3729–3740.
- Leal LN, Romao JM, Hooiveld GJ, Soberon F, Berends H, Boekshoten MV, Van Amburgh ME, Martin-Tereso J and Steele MA 2018. Nutrient supply alters transcriptome regulation in adipose tissue of pre-weaning Holstein calves. *PLoS ONE* 13, e0201929.
- Lepine AJ, Boyd RD and Whitehead DM 1991. Effect of colostrum intake on hepatic gluconeogenesis and fatty acid oxidation in the neonatal pig. *Journal of Animal Science* 69, 1966–1974.
- Liebler-Tenorio EM, Riedel-Caspari G and Pohlenz JF 2002. Uptake of colostrum leukocytes in the intestinal tract of newborn calves. *Veterinary Immunology and Immunopathology* 85, 33–40.
- Maccari P, Wiedemann S, Kunz HJ, Piechotta M, Sanftleben P and Kaske M 2015. Effects of two different rearing protocols for Holstein bull calves in the first 3 weeks of life on health status, metabolism and subsequent performance. *Journal of Animal Physiology and Animal Nutrition* 99, 737–746.
- MacPherson J, Meale SJ, Macmillan K, Haisan J, Bench CJ, Oba M and Steele MA 2019. Effects of feeding frequency of an elevated plane of milk replacer and calf age on behavior, and glucose and insulin kinetics in male Holstein calves. *Animal* 13, 1385–1393.
- Malmuthuge N and Guan LL 2017. Understanding the gut microbiome of dairy calves: opportunities to improve early-life gut health. *Journal of Dairy Science* 100, 5996–6005.
- Mathers JC 1998. Nutrient regulation of intestinal proliferation and apoptosis. *Proceedings of the Nutrition Society* 57, 219–223.
- Mee JF 2008. Newborn dairy calf management. *Veterinary Clinics of North America: Food Animal Practice* 24, 1–17.
- Miller-Cushon EK, Bergeron R, Leslie KE and DeVries TJ 2013. Effect of milk feeding level on development of feeding behavior in dairy calves. *Journal of Dairy Science* 96, 551–564.
- Miller-Cushon EK and DeVries TJ 2015. Invited review: development and expression of dairy calf feeding behaviour. *Canadian Journal of Animal Science* 95, 341–350.
- Molès JP, Tuillon E, Kankasa C, Bedin AS, Nagot N, Marchant A, McDermid JM and Van de Perre P 2018. Breastmilk cell trafficking induces microchimerism-mediated immune system maturation in the infant. *Pediatric Allergy and Immunology* 29, 133–143.
- Mylonas PG, Matsouka PT, Papandoniou EV, Vagianos C, Kalfarentzos F and Alexandrides TK 2000. Growth hormone and insulin-like growth factor I protect intestinal cells from radiation induced apoptosis. *Molecular and Cellular Endocrinology* 160, 115–122.
- Naeem A, Drackley JK, Stamey J and Looor JJ 2012. Role of metabolic and cellular proliferation genes in ruminal development in response to enhanced plane of nutrition in neonatal Holstein calves. *Journal of Dairy Science* 95, 1807–1820.
- Naeem A, Drackley JK, Lanier JS, Everts RE, Rodriguez-Zas SL and Looor JJ 2014. Ruminal epithelium transcriptome dynamics in response to plane of nutrition and age in young Holstein calves. *Functional & Integrative Genomics* 14, 261–273.
- Nissen A, Andersen PH, Bendixen E, Ingvartsen KL and Rontved CM 2017. Colostrum and milk protein rankings and ratios of importance to neonatal calf health using a proteomics approach. *Journal of Dairy Science* 100, 2711–2728.
- Nonnecke BJ, Foote MR, Smith JM, Pesch BA and Van Amburgh ME 2003. Composition and functional capacity of blood mononuclear leukocyte populations from neonatal calves on standard and intensified milk replacer diets. *Journal of Dairy Science* 86, 3592–3604.
- OIE 2017. Terrestrial Animal Health Code. Section 7. Animal Welfare. Retrieved on 3 July 2019 from <http://oie.int/international-standard-setting/terrestrial-code>
- Ollivett TL, Nydam DV, Linden TC, Bowman DD and Van Amburgh ME 2012. Effect of nutritional plane on health and performance in dairy calves after experimental infection with *Cryptosporidium parvum*. *Journal of the American Veterinary Medical Association* 241, 1514–1520.

- Ontsouka EC, Albrecht C and Bruckmaier RM 2016. Invited review: growth-promoting effects of colostrum in calves based on interaction with intestinal cell surface receptors and receptor-like transporters. *Journal of Dairy Science* 99, 4111–4123.
- Probo M, Pascottini OB, LeBlanc S, Opsomer G and Hostens M 2018. Association between metabolic diseases and the culling risk of high-yielding dairy cows in a transition management facility using survival and decision tree analysis. *Journal of Dairy Science* 101, 9419–9429.
- Prokop L, Kaske M, Maccari P, Lucius R, Kunz HJ and Wiedemann S 2015. Intensive rearing of male calves during the first three weeks of life has long-term effects on number of islets of Langerhans and insulin stained area in the pancreas. *Journal of Animal Science* 93, 988–998.
- Qi Y, Zhao X, Huang D, Pan X, Yang Y, Zhao H, Hu H and Cheng G 2018. Exploration of the relationship between intestinal colostrum or milk, and serum metabolites in neonatal calves by metabolomics analysis. *Journal of Agricultural and Food Chemistry* 66, 7200–7208.
- Ramachandran A, Madesh M and Balasubramanian KA 2000. Apoptosis in the intestinal epithelium: its relevance in normal and pathophysiological conditions. *Journal of Gastroenterology and Hepatology* 15, 109–120.
- Renaville R, Van Eenaeme C, Breier BH, Vleurick L, Bertozzi C, Gengler N, Hornick JL, Parmentier I, Istasse L, Haezebroeck V, Massart S and Portetelle D 2000. Feed restriction in young bulls alters the onset of puberty in relationship with plasma insulin-like growth factor-I (IGF-I) and IGF-binding proteins. *Domestic Animal Endocrinology* 18, 165–176.
- Roffler B, Föh A, Sauter SN, Hammon HM, Gallmann P, Brem G and Blum JW 2003. Intestinal morphology, epithelial cell proliferation, and absorptive capacity in neonatal calves fed milk-borne insulin-like growth factor-I or a colostrum extract. *Journal of Dairy Science* 86, 1797–1806.
- Rosenberger K, Costa JHC, Neave HW, von Keyserlingk MAG and Weary DM 2017. The effect of milk allowance on behavior and weight gains in dairy calves. *Journal of Dairy Science* 100, 504–512.
- Sadri H, Steinhoff-Wagner J, Hammon HM, Bruckmaier RM, Gors S and Sauerwein H 2017. Mammalian target of rapamycin signaling and ubiquitin proteasome-related gene expression in 3 different skeletal muscles of colostrum-versus formula-fed calves. *Journal of Dairy Science* 100, 9428–9441.
- Santos A, Giraldez FJ, Frutos J and Andres S 2019. Liver transcriptomic and proteomic profiles of preweaning lambs are modified by milk replacer restriction. *Journal of Dairy Science* 102, 1194–1204.
- Sauter SN, Roffler B, Philipona C, Morel C, Rome V, Guilloteau P, Blum JW and Hammon HM 2004. Intestinal development in neonatal calves: effects of glucocorticoids and dependence of colostrum feeding. *Biology of the Neonate* 85, 94–104.
- Schäff CT, Gruse J, Maciej J, Mielenz M, Wirthgen E, Hoefflich A, Schmicke M, Pfuhr R, Jawor P, Stefaniak T and Hammon HM 2016. Effects of feeding milk replacer ad libitum or in restricted amounts for the first five weeks of life on the growth, metabolic adaptation, and immune status of newborn calves. *PLoS ONE* 11, e0168974.
- Schäff CT, Gruse J, Maciej J, Pfuhr R, Zitnan R, Rajskey M and Hammon HM 2018. Effects of feeding unlimited amounts of milk replacer for the first 5 weeks of age on rumen and small intestinal growth and development in dairy calves. *Journal of Dairy Science* 101, 783–793.
- Schiessler G, Nussbaum A, Hammon HM and Blum JW 2002. Calves sucking, colostrum and milk from their dams or from an automatic feeding station starting in the neonatal period: metabolic and endocrine traits and growth performance. *Animal Science* 74, 431–444.
- Smith JM, Van Amburgh ME, Diaz MC, Lucy MC and Bauman DE 2002. Effect of nutrient intake on the development of the somatotrophic axis and its responsiveness to GH in Holstein bull calves. *Journal of Animal Science* 80, 1528–1537.
- Soberon F and Van Amburgh ME 2017. Effects of preweaning nutrient intake in the developing mammary parenchymal tissue. *Journal of Dairy Science* 100, 4996–5004.
- Steele MA, Penner GB, Chaucheyras-Durand F and Guan LL 2016. Development and physiology of the rumen and the lower gut: targets for improving gut health. *Journal of Dairy Science* 99, 4955–4966.
- Steinhoff-Wagner J, Gors S, Junghans P, Bruckmaier RM, Kanitz E, Metges CC and Hammon HM 2011. Intestinal glucose absorption but not endogenous glucose production differs between colostrum- and formula-fed neonatal calves. *Journal of Nutrition* 141, 48–55.
- Steinhoff-Wagner J, Zitnan R, Schönhusen U, Pfannkuche H, Hudakova M, Metges CC and Hammon HM 2014. Diet effects on glucose absorption in the small intestine of neonatal calves: importance of intestinal mucosal growth, lactase activity, and glucose transporters. *Journal of Dairy Science* 97, 6358–6369.
- Stelwagen K, Carpenter E, Haigh B, Hodgkinson A and Wheeler TT 2009. Immune components of bovine colostrum and milk. *Journal of Animal Science* 87, 3–9.
- Tautenhahn, A. 2017. Risikofaktoren für eine erhöhte Kälbersterblichkeit und geringe Tageszunahmen von Aufzuchtältern in norddeutschen Milchkuhhaltungen. Dissertation for Doctor of Veterinary Medicine. Free University of Berlin, Berlin, Germany.
- Urie NJ, Lombard JE, Shivley CB, Koprak CA, Adams AE, Earleywine TJ, Olson JD and Garry FB 2018. Preweaned heifer management on US dairy operations: part V. Factors associated with morbidity and mortality in preweaned dairy heifer calves. *Journal of Dairy Science* 101, 9229–9244.
- Van Amburgh ME and Soberon F 2013. The role of calf nutrition and management on lifetime productivity of dairy cattle. In *Proceedings of the Cow Longevity Conference*, 28–29 August 2013, Hamra Farm, Tumba, Sweden, pp. 178–197.
- Von Keyserlingk MAG, Rushen J, de Passille AM and Weary DM 2009. The welfare of dairy cattle—key concepts and the role of science. *Journal of Dairy Science* 92, 4101–4111.
- Wang P, Drackley JK, Stamey-Lanier JA, Keisler D and Looor JJ 2014. Effects of level of nutrient intake and age on mammalian target of rapamycin, insulin, and insulin-like growth factor-1 gene network expression in skeletal muscle of young Holstein calves. *Journal of Dairy Science* 97, 383–391.
- Weikard R, Hadlich F, Hammon HM, Frieren D, Gerbert C, Koch C, Dusel G and Kuehn C 2018. Long noncoding RNAs are associated with metabolic and cellular processes in the jejunum mucosa of pre-weaning calves in response to different diets. *Oncotarget* 9, 21052–21069.
- Welboren AC, Leal LN, Steele MA, Khan MA and Martin-Tereso J 2019. Performance of ad libitum fed dairy calves weaned using fixed and individual methods. *Animal* 13, 1891–1898.
- Xu RJ 1996. Development of the newborn GI tract and its relation to colostrum milk intake: a review. *Reproduction Fertility and Development* 8, 35–48.
- Zhao XW, Qi YX, Huang DW, Pan XC, Cheng GL, Zhao HL and Yang YX 2018. Changes in serum metabolites in response to ingested colostrum and milk in neonatal calves, measured by nuclear magnetic resonance-based metabolomics analysis. *Journal of Dairy Science* 101, 7168–7181.