

EV0365

Validation of the Tunisian version of the patient health questionnaire (PHQ-9)H. Belhadj^{1,*}, R. Jomli², U. Ouali², Y. Zgueb², F. Nacef²¹ Military hospital, psychiatry, Tunis, Tunisia² Razi Hospital, psychiatry, Manouba, Tunisia

* Corresponding author.

Introduction The PHQ-9 has been recommended as the best available screening and case-finding instrument for primary care based on its brevity, and ability to inform the clinicians on both depression severity and diagnostic criteria.

Objective Our study evaluated the reliability and the validity of the Tunisian version of the PHQ-9 in detecting major depression in general population.

Method We undertook a cross-sectional and analytical study. A total of 134 participants, representative of the Tunisian general population, were enrolled. The PHQ-9 was validated against the HAD reference standard. The types of validity determined for the PHQ-9 in this study were: translation validity, internal reliability and criterion validity.

Results Test-Retest reliability was determined by intraclass correlation. This scale is stable over 2 weeks (ICC = 0.97). The Tunisian version of the PHQ-9 was found to have good internal reliability (Cronbach's alpha = 0.84). As for criterion validity of the PHQ-9, the Pearson's correlation coefficient between the PHQ-9 and HAD was 0.94 and the Spearman's correlation coefficient was 0.81. This indicated a positive association of good strength between the two instruments. A cut-off score of 10 or higher on the PHQ-9 had a sensitivity of 86.2 and a specificity of 83.8. The VPP was 0.6 and the VPV was 0.9.

Conclusion The Tunisian version of the PHQ-9 has several potential advantages. It was found to be a valid and reliable casefinding instrument for detecting depression in general population.

Disclosure of interest The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.01.695>

EV0366

Prevalence of depression in Tunisian general populationH. Belhadj^{1,*}, R. Jomli²¹ Military hospital, psychiatry, Tunis, Tunisia² Razi Hospital, psychiatry, Mannouba, Tunisia

* Corresponding author.

Introduction Depression is a major burden for the health-care system worldwide.

Objective To identify prevalence and severity of depression in Tunisian general population and define socio-demographic characteristics of screened positive to depression participants.

Method We undertook a cross-sectional and descriptive study. A total of 134 participants, representative of the Tunisian general population, were enrolled. Age, gender, and educational level were the major criteria for representativeness. Depression was assessed with the Patient Health Questionnaire (PHQ-9).

Results The cut-off score was 10. The prevalence of depression was 13.4%. There were no statistical difference in gender, education and age for the prevalence of depression.

Conclusion The World Health Organization ranks depression as the fourth leading cause of disability worldwide. Thus, the detection of depression and the dissemination of treatment in the general population are very important to reduce the burden of the disease.

Disclosure of interest The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.01.696>

EV0367

Anxiety and depression at the medical students in post-examination

S. Ben Saadi*, O. Moula, J. Oueslati, I. Jelalia, R. Ghachem

Razi Hospital, Department B, Mannouba, Tunisia

* Corresponding author.

Introduction The course of medical studies is a long route asking for a lot of breath. It is enclosed by the national examination for specialisation, which allows to access professional training. This examination requires a diligent preparation over several months in difficult conditions for the most part of the candidates who are in practical training.

Objectives and aims We suggest estimating the degree of anxiety and depression at the candidates in the week following the national examination for specialization of 2016.

Methods We have put online on the pages of the various Tunisian Faculties of Medicine an anonymous questionnaire intended for the candidates who took the national examination for specialization of 2016. We used the scale HAD (Hospital Anxiety and depression scale).

Results We recruited 220 participants. More or less half of the participants declare to be rather often in a good mood, that they take little only of the pleasure in the same things as before, that they laugh and see the highly-rated voucher of things really less than before, that they are made of the concern very often and that they sometimes experience sensations of fear. A third of the participants feel tense or irritated most of the time feel sudden sensations of panic rather often and have the impression to work in slow motion rather often.

Conclusion Taking the national examination of specialization rhymes with stress and anxiety. It leads us to question: is it necessary to assure a psychological coverage during the preparation and upstream of the competition?

Disclosure of interest The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.01.697>

EV0368

Depression, anxiety and stress [DASS21] symptoms in menopausal Arab women: Shedding more light on a complex relationship

A. Bener

Istanbul University, Biostatistics and Public Health, Istanbul, Turkey

Aim The objective was to determine correlation between depression, anxiety and stress in menopausal and post-menopausal women and shedding more light on a complex relationship.

Methods A cross-sectional based on Arabian women at the Primary Health Care (PHC) Centers in Qatar during July 2012 and May 2014 and 1101 women agreed to participate and responded to the study. Depression, anxiety and stress were measured using the Depression Anxiety Stress Scales (DASS-21). Data on body mass index (BMI), clinical and other parameters were used.

Results The mean age and SD of the menopausal age was 49.55 ± 3.12, and postmenopausal age was 58.08 ± 3.26 (P < 0.001). There were statistically significant differences between menopausal stages with regards to age, ethnicity, educational status, occupation status, and place of living. Also, there were statistically significant differences between menopausal stages with regards to BMI, systolic and diastolic blood pressure, vitamin D deficiency, and diseases. Depression and anxiety were more common among postmenopause women. The multivariate regression analyses revealed that age in years, diastolic BP, consanguinity, regular exercise were predictor for depression. Meanwhile, diastolic BP, occupation and physical activity considered the main risk factors for anxiety. Furthermore, age in years, occupation and