

1. Emergency Medicine, Komfo Anokye Teaching Hospital, Kumasi, Ghana
2. Oxford University, Oxford, United Kingdom
3. CEO, Tiny David, Ghana
4. Trauma and Orthopedic, Komfo Anokye Teaching Hospital, Kumasi, Ghana

Introduction: The continuous development of the knowledge and skill of the emergency medical technicians (EMTs) in Ghana is important for the success of the pre-hospital system. Due to distance and time constraints, an online e-learning platform is a good way to educate the Emergency Medicine Technicians in Ghana.

Aim: The study looked at the feasibility of developing a distant learning module for the training and continuous medical education of EMTs.

Methods: EMTs in the Ashanti Region were randomly selected to be part of the study. They received online lectures and notes that were accessible by their mobile phones. They all received a test at the end of each model. The study measured their willingness to participate, average attendance for each model, and the scores for each model test. The study also measured the overall feasibility of the distant learning program.

Results: The study developed a training course comprised of 7 modules: trauma and surgical emergencies, obstetric emergencies, pediatric emergencies, disaster management, medical emergencies, basic ultrasound, and medical research. Tests and quizzes were electronically sent to EMTs over the course of the research period, with an average test score of 70.14% (low: 35%, high: 95%) for the cohort. Feedback from participants showed gains in knowledge and skill delivery. The average attendance for all model was 56.6% ranging from 47.37%–63.16% for the models. Challenges for attendance included internet access, heavy duties, and other personal reasons. The post-training interview showed 100% willingness to participate in future online programs with the most common reasons stated as low cost, ease of attendance for models, and reduced expense.

Discussion: The study concluded that online, distant learning models can be used in Ghana for training and continuous medical education for EMTs. It is an easy and cost-effective model compared to a face-to-face model.

Prehosp Disaster Med 2019;34(Suppl. 1):s119–s120

doi:10.1017/S1049023X19002565

Development of Trauma and Disaster Response in Togo, Africa

Dr. Odeda Benin-Goren¹, Dr. Nimrod Aviran², Dr. Iris Adler¹, Dr. Oran Zlotnik², Dr. Yossi Baratz³

1. ODRON, Tel Aviv, Israel
2. Rabin Medical Center, Petach Tiqva, Israel
3. Division of International Development Cooperation (MASHAV) Jerusalem, Israel

Introduction: The project was provided under the auspice and support of the Israel Agency for International Development Cooperation (MASHAV) at the Ministry of Foreign Affairs (MFA). Togo, one of the smallest and least developed countries in West Africa, has a population of ~7.9 million. About 65% of its population lives in rural areas. Due to the lack of medical

resources, Togo suffers from health problems including those related to trauma and mass events. In May 2017, a trauma and disaster team came to Togo to train the medical team in the new trauma unit, donated and built by the MFA. The unit was built in the Atakpame Regional Hospital (ARH), located 160km north of the capital, Lomé. ARH serves one million inhabitants, mostly from rural areas.

Methods: The training included lectures, simulations, drills, case studies, bedside teaching, and operation of medical technologies.

Results: Following the training, it was recommended to continue the program and to move forward with advanced training. Following the team's recommendations, MASHAV decided to expand the program and to provide a multilateral project to Togo and ten other West African countries within five months after the first training ended. Twenty participants (mostly senior doctors) were chosen from ten Western African countries and brought to Lomé. The participants joined a two-day Trauma and Disaster Preparedness seminar. Following the seminar, they were moved to Atakpame to join the local team and the facilitators, to visit the trauma unit, and to learn about it as a model for trauma care that can be modified to the capabilities of the local facility.

Discussion: Lessons learned and recommendations from those two projects were brought to the MFA that will try to develop more training and cooperation models to help and establish better trauma care and disaster response, supported by the Israeli team.

Prehosp Disaster Med 2019;34(Suppl. 1):s120

doi:10.1017/S1049023X19002577

The Diagnostic Value of Ultrasonic Measurement of Inferior Vena Cava Diameter Respiratory Variability on Volume Response Evaluation of Geriatric Hip Fracture

Dr. Yang Li, Ms. Yuan Cen

Army Medical Center of PLA, Chongqing, China

Aim: To evaluate the value of ultrasonic measurement of the respiratory variability of inferior vena cava (IVC) in the preoperative volume-response evaluation of elderly hip fracture patients.

Methods: Volume-loading tests were carried out in elderly patients with hip fractures requiring surgical treatment from August 2017 to February 2018. The maximum diameter (IVCe) and minimum diameter (IVCi) of the IVC were measured by ultrasound, and the variation of IVC (IVC-CI) was calculated before surgery. SV was monitored by a FloTrac/Vigileo system, and positive volume responsiveness was defined as ΔSV increasing by more than 15%. The sensitivity and specificity of IVC-CI to volume responsiveness evaluation was analyzed by ROC, and the correlation between IVC-CI and ΔSV was analyzed by Spearman correlation analysis.

Results: Ultrasound measurements and volume-loading tests were successfully performed in 39 of the 44 patients. Among them 21 cases were volume responsiveness positive (group R) and 18 cases were volume responsiveness negative (group N). Before the volume-loading test, IVCi in group R was significantly smaller than group N and IVC-CI was significantly larger than group N ($P < 0.05$), while the difference between IVCe and group N was not statistically significant ($P > 0.05$). After the volume-loading test, the differences between IVCe,