
INSTRUCTIONS TO CONTRIBUTORS

The *Journal of Helminthology* publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance.

Manuscripts, which must be in English or French (with an English summary), should be addressed to:

The Editor, Journal of Helminthology
London School of Hygiene and Tropical Medicine
Keppel Street, London WC1E 7HT
England

Two copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief introduction (unheaded); Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration may be printed is 13.5 × 18 cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space should be indicated. Some indication of scale should normally be given on the figure. Photocopies of illustrations should also be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints, preferably of the same size as they are to appear in the Journal (maximum size 13.5 × 18 cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

DUKE, B. O. L. (1971) The ecology of onchocerciasis in man and animals. In *Ecology and physiology of parasites* (ed A. M. Fallis) pp. 213-222. Adam Hilger Ltd.: London.

JAMES, C. and WEBBE, G. (1973) A comparison of Egyptian and East African strains of *Schistosoma haematobium*. *Journal of Helminthology*, **47**, 49-59.

50 offprints are provided free of charge; additional copies may be ordered at the proof stage.

Contents

	Pages
Studies on the mechanism of long-term survival of <i>Taenia taeniaeformis</i> in rats. B. A. KWA and F. Y. LIEW	1-6
Intrauterine development of the microfilariae of <i>Dipetalonema viteae</i> . D. S. ELLIS, R. ROGERS, A. E. BIANCO and D. A. DENHAM	7-10
Hybridization between different strains of <i>Schistosoma intercalatum</i> Fisher, 1934 from Cameroun and Zaire. F. FRANSEN	11-22
Egg-laying and hatching rhythms in the monogenean <i>Diplozoon homoion gracile</i> from the southern barbel (<i>Barbus meridionalis</i>). S. MACDONALD and A. JONES	23-28
Allometric growth in three species of digenetic trematodes of marine fishes from Belize. J. H. FISCHTHAL	29-39
New schistorophine nematodes (Acuariidae) from New Zealand shore birds. W. C. CLARK	41-50
Life history of <i>Allocreadium fasciatusi</i> Kakaji, 1969 (Trematoda: Allocreadiidae) from the freshwater fish <i>Aplocheilus melastigma</i> McClelland. R. MADHAVI	51-59
The influence of some physico-chemical factors on the host-finding capacity of <i>Fasciola hepatica</i> miracidia. N. Ø. CHRISTENSEN, P. NANSEN and F. FRANSEN	61-67
Gastro-intestinal nematodes of British police dogs. Research Note. E. J. PEGG	68-70
Prevalence of hydatid disease in food animals reported at meat inspection in Nigeria, 1971-1975. Research Note. B. J. O. DADA	70-72
<i>Schistosoma margrebowiei</i> of cobs in Chad. Research Note. M. GRABER	72-74
<i>Onchocerca sweetae</i> (Nematoda: Filarioidea): notes on the intermediate host. D. M. SPRATT, A. L. DYCE and H. A. STANDFAST	75-81
Activity of disaccharidases in the intestinal contents of piglets infected with gastrointestinal nematodes and activity in these nematodes. K. ENIGK and A. DEY-HAZRA	83-89
SUPPLEMENTARY REVIEW ARTICLE	
Ascaridoid nematodes of amphibians and reptiles: <i>Goezia</i> . J. F. A. SPRENT	91-98