

Recombination Line Abundances in the Winds of the [WC] Wolf-Rayet Stars

R. L. Kingsburgh¹, I. Dashevsky¹ and M. J. Barlow²

¹York University, Dept. of Physics and Astronomy;

²University College London, Dept. of Physics and Astronomy

Abstract: We present a preliminary abundance analysis, utilising recombination theory, for the [WC] Wolf-Rayet central stars of the planetary nebulae NGC 6751 and NGC 6905. This analysis is based on optical spectrophotometry of a sample of [WC] stars which show a strong O VI 3811, 34 Å feature in emission. We have performed a recombination line analysis of stellar wind emission lines which are judged to be optically thin, in order to derive relative C, O and He abundances. We also present a comparison of the derived wind abundances with those of Population I WO stars. For the [WC 4] central star of NGC 6751, we have derived C/He = 0.35 and C/O = 4.2, by number. And for the [WC 3] central star of NGC 6905, we have derived C/He = 0.36 and C/O < 12. These surface abundances are comparable to the abundances that have been derived for early type [WC] stars via more complex NLTE modelling (e.g. Koesterke & Hamann 1996).

1. Introduction

The [WC] Wolf-Rayet stars are a hydrogen-deficient sub-group of the central stars of PN. Their spectra exhibit strong broad emission line features, indicative of stellar winds similar to those found in the Population I WC and WO stars. Of particular interest are the members of the O VI sequence defined by Smith & Aller (1969) which show the O VI 3p–3s doublet at 3811, 34 Å strongly in emission, since the spectra of these stars shows the highest degree of ionization amongst all WR stars. Their spectra are dominated by lines of He, C and O. Unlike the Pop. I WR stars, there are no [WN] counterparts to the [WC] central stars, although occasionally the N V 1240Å feature is seen, indicating that although both Pop. I and II stellar winds reveal He-processed material, different mixing histories would have been experienced by the very different progenitor stars, allowing for trace amounts of nitrogen to be present in the Pop. II [WC] stars.

We have initiated a study of the O VI-sequence [WC] stars, aimed at deriving global parameters including their surface abundances and radiative energy distributions, in order to obtain insight into their evolutionary status. The current work presents a preliminary analysis for NGC 6751 and NGC 6905.

2. Observations

Both narrow- and wide-slit spectrophotometry were obtained using the RGO Spectrograph at the 3.0 m AAT, with an IPCS as detector. Data reduction was performed as described by Kingsburgh & Barlow (1994).

Equivalent widths, EW, were measured from the narrow-slit spectra, while the absolute level of the underlying continuum, $F_c(\lambda)$, was determined from the wide-slit spectra. Multiplication of the EW's by the appropriate values of $F_c(\lambda)$ yielded absolute line fluxes. These were dereddened by adopting the extinction coefficient derived from the Balmer decrement of the nebular spectrum (Kingsburgh & Barlow, 1994) together with the Galactic reddening law of Howarth (1983).

III. Central Stars

3. Recombination line analysis

A Case B recombination line analysis was performed for lines which were judged to be optically thin. The method is described in detail by Kingsburgh, Barlow & Storey (1995). The adopted wind temperature and density were 50000K and 10^{11}cm^{-3} respectively. Relative carbon, oxygen and helium abundances were derived for NGC 6751 and NGC 6905, and are presented in Table 1, along with the abundances derived for the Population I WO stars by the same method.

The oxygen abundance derived for NGC 6905 is currently a lower limit, based only on the $\text{O}^{6+}/\text{He}^{2+}$ abundance, as the $\text{OV } 5590\text{\AA}$ line is predicted to be partially optically thick, and cannot be used to estimate the O^{5+} abundance. The accuracy of the oxygen abundances for both objects will improve as UV spectra are incorporated into the analysis. The C/He and C/O ratios for NGC 6751 and 6905 are within 30% of the values found for the two galactic population I WO stars, Sand 4 and Sand 5, although these stars have had completely different evolutionary histories; the pop I WC/WO stars originating from $\sim 40\text{--}60M_{\odot}$ stars, and the Pop. II [WC] stars from $\sim 1M_{\odot}$ stars.

NGC 6751 has recently been analyzed by Koesterke and Hamann (1996; KH) using a detailed non-LTE model of the stellar wind. They derive the following mass fractions: $X(\text{C})=0.31$, $X(\text{O})=0.15$, $X(\text{He})=0.54$. From Table 1, we derive the following mass fractions for NGC 6751: $X(\text{C})=0.44$, $X(\text{O})=0.14$, $X(\text{He})=0.42$. Our $X(\text{O})$ is in agreement with that of KH, however our $X(\text{C})$ is a factor of 1.4 higher. The C IV 5470\AA line is best suited for deriving carbon abundances (Hillier 1989); however KH's models underestimate this line's strength, and they conclude that a value of $X(\text{C})$ as high as ~ 0.5 (or $X(\text{C})/X(\text{O})=1.0$) cannot be ruled out.

Table 1. A Comparison Between Population I and II Wolf-Rayet Wind Abundances By Number

	Object	class	C/He	O/He	C/O	Ref
pop I	NGC 6751	[WC4]	0.35	0.086	4.2	1
	NGC 6905	[WC3]	0.36	>0.029	<12	1
pop II	Sand 1	WO4	0.81	0.30	2.7	2
	Sand 2	WO4	0.52	0.11	4.6	2
	Sand 4	WO1	0.51	0.11	4.6	2
	Sand 5	WO2	0.52	0.10	5.2	2

Notes: 1—this work; 2—Kingsburgh, Barlow & Storey 1995.

REFERENCES

- Howarth, I. D., 1983, MNRAS, 203, 301.
Hillier, J. D., 1989, ApJ, 347, 392
Kingsburgh, R. L. & Barlow, M. J., 1994, MNRAS.
Kingsburgh, R. L., Barlow, M. J. & Storey, P. J., 1995, A&A, 295, 75.
Koesterke, L. & Hamann, W.-R., 1996, A&A, in press. (KH)
Smith, L. & Aller, L. H., 1969, ApJ, 157, 1245.