



# The Bochner-flat cone of a CR manifold

Liana David

## ABSTRACT

We construct a Kähler structure (which we call a generalised Kähler cone) on an open subset of the cone of a strongly pseudo-convex CR manifold endowed with a one-parameter family of compatible Sasaki structures. We determine those generalised Kähler cones which are Bochner-flat and we study their local geometry. We prove that any Bochner-flat Kähler manifold of complex dimension bigger than two is locally isomorphic to a generalised Kähler cone.

## 1. Introduction

The Bochner tensor of a Kähler manifold is the biggest irreducible component of the curvature tensor under the action of the unitary group. In complex dimension two, the Bochner tensor coincides with the anti-self-dual Weyl tensor. A Kähler manifold is Bochner-flat if its Bochner tensor vanishes. Bochner-flat Kähler manifolds represent an important class of Kähler manifolds and have been intensively studied: the local geometry of Bochner-flat Kähler manifolds and its interactions with Sasaki geometry has been studied, using the Webster's correspondence, in [DG06]; complete Bochner-flat Kähler structures on simply connected manifolds have been classified in [Bry01]; generalisations of Bochner-flat Kähler manifolds (like weakly Bochner-flat Kähler manifolds and Kähler manifolds with a hamiltonian 2-form) have also been developed (see, for example, [ACG04, ACG06, Gau01]).

An important class of Kähler manifolds is represented by the Kähler cones of Sasaki manifolds. Unfortunately, except when the Sasaki manifold is an open subset of the standard CR sphere with its standard metric as the Sasaki metric, the Kähler cones are not Bochner-flat. In this paper we propose an alternative construction, which is a natural generalisation of the Kähler cone construction and which produces, locally, all Bochner-flat Kähler structures of complex dimension bigger than two. More precisely, we consider, on a fixed CR manifold  $(N, H, I)$ , a one-parameter family of Sasaki Reeb vector fields  $\{T_r, r \in \mathcal{J}\}$  (with  $\mathcal{J} \subset \mathbb{R}^{>0}$  an open connected interval). On the cone manifold  $N \times \mathcal{J}$  we define an almost complex structure  $J$ , which on  $H \subset T(N \times \mathcal{J})$  coincides with  $I$  and which sends the radial vector field  $V = r \partial/\partial r$  to the vector field  $T$ , defined by  $T(p, r) := T_r(p)$ , for any  $(p, r) \in N \times \mathcal{J}$ . It turns out that  $J$  is integrable and that the pair  $(\omega := \frac{1}{4} dd^J r^2, J)$  is a Kähler structure on the open subset of  $N \times \mathcal{J}$  where  $\omega(V, T) > 0$ . Such a Kähler structure will be called a generalised Kähler cone and coincides with the usual Kähler cone of a Sasaki manifold when the family of Reeb vector fields is constant. A strong motivation for this construction comes from the fact that the Bryant family of Bochner-flat Kähler structures (which were discovered by Bryant in his classification theorem of complete Bochner-flat Kähler structures on simply connected manifolds [Bry01] and have been further studied in [DG06]) are generalised Kähler cones. Our main result is the following.

---

Received 9 January 2006, accepted in final form 1 October 2007.

*2000 Mathematics Subject Classification* 53C55, 53C21, 53D10.

*Keywords:* Bochner-flat Kähler manifolds, cone manifolds, CR and Sasaki manifolds.

Research supported by Centro di Ricerca Matematica 'Ennio de Giorgi', Scuola Normale Superiore, Pisa, Italy, and partially supported by CNCSIS grant no. GR 54/2007 code 811.

This journal is © [Foundation Compositio Mathematica](#) 2008.

**THEOREM 1.** *Any Bochner-flat Kähler manifold of complex dimension bigger than two is locally isomorphic to a generalised Kähler cone.*

The plan of the paper is the following. In § 2 we review the theory of Kähler and Sasaki manifolds, which will be useful later on in our study of generalised Kähler cones. In § 3 we determine the generalised Kähler cones which are Bochner-flat and in § 4 we study their local geometry. This study will readily imply Theorem 1. The last section is dedicated to examples. We explain how Kähler manifolds with constant holomorphic sectional curvature, weighted projective spaces and Bryant family of Bochner-flat Kähler structures fit into our formalism of generalised Kähler cones. We also look at Bochner-flat generalised Kähler cones of order one and at those which are of Tachibana and Liu type [TL70].

## 2. Notation and earlier results

### 2.1 The Bochner tensor of a Kähler manifold

In this section we recall the definition of the Bochner tensor of a Kähler manifold. We use the formalism developed in [ACG06, Gau01].

Let  $(V, g, J)$  be a real vector space together with a complex structure  $J$  and a  $J$ -invariant positive definite metric  $g$ . We shall identify vectors and covectors of  $V$  using the metric  $g$ . Let  $\omega := g(J\cdot, \cdot)$  be the Kähler form. Recall that the space  $\mathcal{K}(V)$  of Kähler curvature tensors of  $(V, g, J)$ , defined as those curvature tensors which annihilate all  $J$ -anti-invariant 2-forms on  $V$ , decomposes into a  $g$ -orthogonal sum

$$\mathcal{K}(V) := c_{\mathcal{K}}^*(\text{Sym}^{1,1}(V)) \oplus \mathcal{W}(V). \tag{1}$$

Here  $c_{\mathcal{K}}^* : \text{Sym}^{1,1}(V) \rightarrow \mathcal{K}(V)$  is the adjoint of the Ricci contraction

$$c_{\mathcal{K}} : \mathcal{K}(V) \rightarrow \text{Sym}^{1,1}(V), \quad c_{\mathcal{K}}(R)(v, w) := \text{trace } R(v, \cdot, w, \cdot), \quad v, w \in V,$$

and has the following expression [Gau01]:

$$c_{\mathcal{K}}^*(S) = \frac{1}{2} \left[ \frac{S \wedge \text{Id} + (J \circ S) \wedge J}{2} + \omega \otimes S + \beta \otimes J \right], \tag{2}$$

where  $S \in \text{Sym}^{1,1}(V)$  is a symmetric  $J$ -invariant endomorphism of  $V$ , ‘Id’ is the identity endomorphism,  $\beta \in \Lambda^{1,1}(V)$  is the  $J$ -invariant 2-form on  $V$ , related to  $S$  by  $\beta(v, w) := g(SJv, w)$ , and, for two endomorphisms  $S$  and  $T$  of  $V$ ,  $S \wedge T$  is the endomorphism of  $\Lambda^2(V)$  defined by the formula

$$(S \wedge T)(v \wedge w) := S(v) \wedge T(w) - S(w) \wedge T(v), \quad v, w \in V.$$

According to the decomposition (1), a Kähler curvature tensor  $R \in \mathcal{K}(V)$  decomposes into the sum

$$R = c_{\mathcal{K}}^*(S) + W^K,$$

where  $W^K \in \mathcal{W}(V)$  is its principal part (or the Bochner tensor of  $R$ ) and  $S \in \text{Sym}^{1,1}(V)$  is a modified Ricci tensor.

Consider now a Kähler manifold  $(M, g, J)$ . The curvature  $R^g$  of the Kähler metric  $g$  is, at any point  $p \in M$ , a Kähler curvature tensor of the tangent space  $(T_p M, g_p, J_p)$ . The principal part of  $R^g$  is called the Bochner tensor of  $(M, g, J)$  and is a section of the symmetric product  $\Lambda^{1,1}(M) \odot \Lambda^{1,1}(M)$ . The Kähler manifold  $(M, g, J)$  is Bochner-flat if its Bochner tensor vanishes.

### 2.2 Review of CR and Sasaki manifolds

Recall that an oriented (strongly pseudo-convex) CR manifold  $(N, H, I)$  has a codimension one oriented subbundle  $H$  of the tangent bundle  $TN$ , called the contact bundle, and a bundle homomorphism  $I : H \rightarrow H$  with  $I^2 = -\text{Id}$ , such that, for every smooth section  $X, Y \in \Gamma(H)$ ,

$[IX, IY] - [X, Y]$  is also a section of  $H$  and the integrability condition

$$[IX, IY] - [X, Y] = I([IX, Y] + [X, IY]) \tag{3}$$

is satisfied. Since  $N$  and  $H$  are oriented, the co-contact line bundle  $L := TN/H$  is also oriented, hence trivialisable. A positive section  $\mu$  of  $L$  defines a contact form  $\theta := \eta\mu^{-1}$  on  $M$ , where  $\eta : TN \rightarrow L$  is the natural projection and  $\mu^{-1} \in \Gamma(L^*)$  is the dual section of  $\mu$ , i.e. the natural contraction between  $\mu$  and  $\mu^{-1}$  is the function on  $N$  identically equal to one. The bilinear form  $g(X, Y) := \omega(X, IY) := \frac{1}{2} d\theta(X, IY)$  of the bundle  $H$  is independent, up to a positive multiplicative function, of the choice of the contact form and is positive definite – the strongly pseudo-convexity condition. The contact form  $\theta$  determines a Reeb vector field  $T$ , uniquely defined by the conditions  $\theta(T) = 1$  and  $i_T d\theta = 0$ . Note that the Reeb vector field preserves the bundle  $H$ , i.e.  $[T, X] \in \Gamma(H)$  when  $X \in \Gamma(H)$  and hence  $\mathcal{L}_T(I)$  is a well-defined endomorphism of  $H$ . We associate to  $\theta$  a Riemannian metric  $g$  on the manifold  $N$ , which on  $H$  is defined above and such that  $T$  is of norm one and  $g$ -orthogonal to  $H$ . Finally, we need to recall the definition of the Tanaka connection [Tan75] associated to  $\theta$ . It is the unique linear connection  $\nabla$  on  $N$  with the following three properties: (i) it preserves the bundle  $H$ ; (ii)  $I, g$  and  $T$  are  $\nabla$ -parallel; and (iii) the torsion  $T^\nabla$  of  $\nabla$  has the expression

$$\begin{aligned} T^\nabla(X, Y) &= 2\omega(X, Y), \\ T^\nabla(T, X) &= -\frac{1}{2}I\mathcal{L}_T(I)(X), \end{aligned}$$

for every  $X, Y \in \Gamma(H)$ . It turns out that, on  $H$ ,  $\nabla$  is determined by a Koszul-type formula

$$\begin{aligned} 2g(\nabla_X Y, Z) &= X(g(Y, Z)) + Y(g(X, Z)) - Z(g(X, Y)) + g([X, Y]^H, Z) \\ &\quad - g([X, Z]^H, Y) - g([Y, Z]^H, X), \end{aligned}$$

where  $X, Y, Z \in \Gamma(H)$  and, for a vector field  $W$  of  $N$ ,  $W^H := W - \theta(W)T$  is its  $g$ -orthogonal projection on the bundle  $H$ . The metric  $g$  is called Sasaki if  $T$  is a Killing vector field for the metric  $g$  or, equivalently, if  $\mathcal{L}_T(I) = 0$ . In this case, the curvature  $R^\nabla$  of the Tanaka connection on the bundle  $H$  is an element of the tensor product  $\Lambda^2(N) \otimes \Lambda^{1,1}(H)$  and its restriction to bivectors of  $H$  belongs to the symmetric product  $\Lambda^{1,1}(H) \odot \Lambda^{1,1}(H)$  and is a Kähler curvature tensor of the complex Riemannian vector bundle  $(H, g, I)$ . Its Bochner part (called the Chern–Moser tensor [CM74, Dav04] of the CR manifold  $(N, H, I)$ ) is independent of the choice of the compatible Sasaki structure on  $(N, H, I)$ . A CR manifold with vanishing Chern–Moser tensor is called flat. The importance of the Chern–Moser tensor comes from the fact that if the CR manifold is flat and of dimension bigger than three, then it is locally isomorphic with a sphere with its standard CR structure [CM74, BS76]. On the other hand, if  $g$  is Sasaki, the complex structure  $I$  and the metric  $g$  of the bundle  $H$  descend on the quotient  $N/T$  and determine a Kähler structure on this quotient. (In our conventions, the quotient  $N/T$  denotes the space of leaves of the foliation generated by  $T$  in a sufficiently small open subset of  $N$ , so that  $N/T$  is a manifold.) Moreover, the Bochner tensor of the Kähler manifold  $N/T$  becomes identified with the Chern–Moser tensor of the CR manifold  $(N, H, I)$  (see [Web77, Dav04]). Since any Kähler manifold can be locally written as a quotient of a Sasaki manifold under its Reeb vector field, by means of a choice of a local primitive of the Kähler form, it follows that a Bochner-flat Kähler manifold of complex dimension  $m \geq 2$  is locally isomorphic with the quotient of a standard CR sphere  $S^{2m+1}$  under the Reeb vector field of a compatible Sasaki structure.

### 2.3 The local type of Bochner-flat Kähler structures

The local geometry of Bochner-flat Kähler structures, of complex dimension  $m \geq 2$ , is of four types [Bry01, Cal02, DG06]. This follows from the fact that the compatible Sasaki structures on the CR sphere  $S^{2m+1}$  are determined by elements of the Lie algebra  $su(m + 1, 1)$ , and that there

are four types of conjugacy classes in this Lie algebra (elliptic, hyperbolic, one-step and two-step parabolic). In order to explain this, it is convenient to identify  $S^{2m+1}$  with the hermitian sphere  $\Sigma^{2m+1}$  of all complex null lines in a hermitian complex vector space  $W$  of signature  $(m + 1, 1)$ , with hermitian metric  $(\cdot, \cdot)$ , by fixing an orthonormal basis of  $W$ , i.e. a basis  $\{e_0, \dots, e_{m+1}\}$  with  $(e_0, e_0) = -1$ ,  $(e_j, e_j) = 1$ , for  $j \in \{1, \dots, m + 1\}$ , and  $(e_i, e_j) = 0$  for  $i \neq j$ , and by associating to a complex null line  $x$  of  $W$  its unique representative of the form  $e_0 + u$ , where  $u \in S^{2m+1}$  belongs to the unit sphere of the positive definite hermitian vector space  $\text{Span}\{e_1, \dots, e_{m+1}\}$ . Let  $\eta$  be the natural (line bundle valued) contact form of  $\Sigma^{2m+1}$ :

$$\eta(X) := \text{Im}(\hat{X}w, w), \quad X \in T_x\Sigma^{2m+1}, \quad 0 \neq w \in x, \quad x \in \Sigma^{2m+1},$$

where  $\hat{X} \in \text{Hom}_{\mathbb{C}}(x, W)$  is a representative of  $X \in \text{Hom}_{\mathbb{C}}(x, W/x)$ . A hermitian trace-free endomorphism  $A$  of  $W$  determines a Reeb vector field  $T^A$  of a Sasaki structure on the open subset

$$\Sigma_A^{2m+1} := \{x \in \Sigma^{2m+1} : (Aw, w) > 0, \quad w \in x, \quad w \neq 0\},$$

defined in the following way. At a point  $x \in \Sigma_A^{2m+1}$ ,  $T_x^A \in \text{Hom}_{\mathbb{C}}(x, W/x)$  associates to a non-zero vector  $w \in x$ , the class of  $iAw$  in the quotient  $W/x$ . The contact form of  $T^A$  is  $\eta_A := \eta/(Aw, w)$ , i.e.

$$\eta_A(X) = \frac{\text{Im}(\hat{X}w, w)}{(Aw, w)}, \quad X \in T_x\Sigma^{2m+1}, \quad 0 \neq w \in x, \quad x \in \Sigma_A^{2m+1}.$$

Employing the notation of [DG06], we shall denote by  $M_A$  the induced Kähler structure on the quotient  $\Sigma_A^{2m+1}/T^A$  and by  $g_A$  its Kähler metric.

We end this section with a simple lemma on hermitian operators which will play an important role in our treatment. For completeness of the exposition, we include its proof.

LEMMA 2. *Let  $A : W \rightarrow W$  be a hermitian operator on a complex vector space  $W$  with a hermitian metric  $(\cdot, \cdot)$  of signature  $(m + 1, 1)$ . Suppose that  $A$  satisfies  $(Aw, w) = 0$ , for any null vector  $w$  which belongs to a non-empty open subset  $D$  of  $W$ . Then  $A = \lambda \text{Id}$ , for  $\lambda \in \mathbb{R}$ . If, moreover,  $A$  is trace-free, then  $A = 0$ .*

*Proof.* Let  $w = w_t$  be a curve in  $D$ , with  $w_t$  null for any  $t$ ,  $w_0 = w \in D$  and  $\dot{w}_0 = X$ . Taking the derivative at  $t = 0$  of the equality  $(Aw_t, w_t) = 0$  and using the fact that  $A$  is hermitian, we get  $\text{Re}(Aw, X) = 0$ . In particular, we deduce that  $(Aw, X) = 0$ , for any null vector  $w \in D$  and any  $X \in W$ , which is hermitian orthogonal to  $w$ . This implies that  $Aw = \lambda w$ , where  $\lambda \in \mathbb{R}$  depends *a priori* on  $w$ . It follows that the map

$$W \ni u \rightarrow Au \wedge u \in \Lambda^2(W) \tag{4}$$

vanishes when  $u \in D$  is null. Being holomorphic, the map (4) must be identically zero. We deduce that, for any  $u \in W$ ,  $Au$  is a multiple of  $u$  which clearly implies the first claim. The second claim is trivial. □

### 2.4 The Bryant minimal and characteristic polynomials

The local type of a Bochner-flat Kähler manifold  $(M, g, J)$  is encoded into the Bryant minimal and characteristic polynomials, which can be defined as follows. Let  $S$  be the modified Ricci operator which satisfies  $c_{\mathcal{K}}^*(S) = R^g$  (where  $R^g$  is the curvature of  $g$ ) and  $P(t)$  be the characteristic polynomial of a new modified Ricci operator  $\Theta$ , related to  $S$  by

$$\Theta := \frac{1}{4} \left( S - \frac{\text{trace}_{\mathbb{R}}(S)}{2(m + 2)} \text{Id} \right), \tag{5}$$

where  $m$  is the complex dimension of  $M$ . The Ricci operator  $\Theta$  has been introduced in [Bry01]. It will be considered as a complex linear operator on the complex vector bundle  $(TM, J)$ . Its trace

is called the modified scalar curvature of  $(M, g, J)$ . Denote by  $\xi_1, \dots, \xi_l$  the non-constant roots of  $P$  and by  $P_n$  its non-constant part, defined by  $P_n(t) := (t - \xi_1) \cdots (t - \xi_l)$ . The number  $l$  is called the order of  $(M, g, J)$ . On a dense open subset  $M^0$  of  $M$ , the eigenvalues  $\xi_j$  (for any  $j \in \{1, \dots, l\}$ ) are simple, different from each other at any point and different, at any point, from any constant eigenvalue of  $\Theta$ ; the functions  $\xi_1, \dots, \xi_l$  are functionally independent on  $M^0$  and

$$|\text{grad}_g(\xi_j)|^2 = -4 \frac{p_m(\xi_j)}{P'_n(\xi_j)}, \quad j \in \{1, \dots, l\}, \tag{6}$$

where  $p_m$  is a monic polynomial of degree  $l + 2$ , with constant coefficients, independent of  $j$ , called the Bryant minimal polynomial of  $(M, g, J)$ . The Bryant characteristic polynomial  $p_c$  of  $(M, g, J)$  is by definition the product of  $p_m$  with the constant part  $P_c := P/P_n$  of  $P$ .

Suppose now that  $(M, g, J) \cong M_A = \Sigma_A^{2m+1}/T^A$ , for a hermitian operator  $A$  of  $W$ . Denote by  $\tilde{a}$  the reduced adjoint operator of  $A$ , defined by

$$\tilde{a}(t) = t^{l+2} + a_1 t^{l+1} + \cdots + a_{l+2},$$

where

$$a_k := A^k - \sigma_1(q_A)A^{k-1} + \cdots + (-1)^k \sigma_k(q_A)$$

and  $\sigma_k(q_A)$  is the  $k$ -elementary symmetric function of the roots of the minimal polynomial  $q_A$  of  $A$ . The reduced adjoint operator  $\tilde{a}$  satisfies  $(t \text{Id} - A)\tilde{a}(t) = q_A(t)\text{Id}$ , for any  $t \in \mathbb{R}$ . The eigenspace of  $\Theta$  corresponding to a non-constant eigenvalue  $\xi_j$  is spanned by the vector field  $L_j$  which, viewed as a section of  $H$ , is equal, at a point  $x \in \Sigma_A^{2m+1}$ , to

$$L_j(w) = \tilde{a}(\xi_j)w \pmod w.$$

The non-constant part  $P_n(t)$  of the modified Ricci operator  $\Theta$  of  $(M, g, J)$ , viewed as a polynomial with function coefficients defined on  $\Sigma_A^{2m+1}$ , is equal, at a point  $x \in \Sigma_A^{2m+1}$ , to

$$p_{A,x}(t) := \frac{(\tilde{a}(t)w, w)}{(Aw, w)}, \quad w \in x, \quad w \neq 0.$$

The constant eigenvalues of  $\Theta$  can also be described in terms of  $A$ : if  $\lambda$  is a constant eigenvalue of  $\Theta$  of multiplicity  $n$ , then it is a multiple eigenvalue of  $A$ , of multiplicity  $n + 1$ , and the eigenspace of  $\Theta$ , at a point  $x \in \Sigma_A^{2m+1}$ , corresponding to  $\lambda$ , can be identified with the intersection of the hermitian orthogonal  $x^\perp \subset W$  with the eigenspace of  $A$  corresponding to  $\lambda$  (see [DG06]). The Bryant minimal and characteristic polynomials  $p_m$  and  $p_c$  coincide with the minimal polynomial  $q_A$ , respectively, with the characteristic polynomial  $Q_A$  of  $A$ , (see [Cal02, DG06]). The modified scalar curvature of  $M_A$ , viewed as a function on  $\Sigma_A^{2m+1}$ , is equal, at  $x \in \Sigma_A^{2m+1}$ , to  $-(A^2w, w)/(Aw, w)$ , where  $w \in x$  is non-zero.

The following lemma will be useful in our study of generalised Kähler cones and is an easy consequence of the theory developed in [DG06]. For completeness of the exposition, we include its proof.

LEMMA 3. For any  $t \in \mathbb{R}$ ,  $x \in \Sigma_A^{2m+1}$  and  $w \in x$  non-zero,

$$g_A(dp_{A,x}(t), dp_{A,x}(t)) = 4 \left( q'_A(t)p_{A,x}(t) - q_A(t)p'_{A,x}(t) - 2tp_{A,x}^2(t) + p_{A,x}^2(t) \frac{(A^2w, w)}{(Aw, w)} \right).$$

*Proof.* Via the metric  $g_A$ , the 1-form  $dp_A(t)$  corresponds to the vector field  $L_t$ , which, viewed as a section of  $H$ , is equal, at  $x \in \Sigma_A^{2m+1}$ , to the endomorphism

$$L_t(w) := 2(\tilde{a}(t)w - p_{A,x}(t)Aw) \pmod w.$$

Its square norm is equal to

$$g_A(L_t, L_t) = 4 \frac{(\tilde{a}(t)w, \tilde{a}(t)w) - 2p_{A,x}(t)(\tilde{a}(t)w, Aw) + p_{A,x}^2(t)(A^2w, w)}{(Aw, w)} \\ = 4 \left( q'_A(t)p_{A,x}(t) - q_A(t)p'_{A,x}(t) - 2tp_{A,x}^2(t) + \frac{p_{A,x}^2(t)(A^2w, w)}{(Aw, w)} \right),$$

where we have used  $(A\tilde{a}(t)w, w) = t(\tilde{a}(t)w, w)$  (which follows from  $(t\text{Id} - A)\tilde{a}(t) = q_A(t)\text{Id}$  and  $(w, w) = 0$ ) and

$$\frac{(\tilde{a}(t)w, \tilde{a}(t)w)}{(Aw, w)} = q'_A(t)p_{A,x}(t) - q_A(t)p'_{A,x}(t), \tag{7}$$

which has been proved in [DG06]. □

### 3. Definition of generalised Kähler cones; the Bochner-flatness condition

Let  $(N, H, I)$  be an oriented CR manifold and  $\{T_r\}_{r \in \mathcal{J}}$ , with  $\mathcal{J} \subseteq \mathbb{R}^{>0}$  a connected open interval, a family of Reeb vector fields of Sasaki structures on  $(N, H, I)$ , with contact forms  $\{\theta\}_{r \in \mathcal{J}}$ . Let  $\omega_r := \frac{1}{2}d\theta_r \in \Omega^2(N)$  and  $g_r := \omega_r(\cdot, I\cdot)$  be the corresponding (positive definite) metrics of the contact bundle  $H$ . On the cone manifold  $N \times \mathcal{J}$  define the vector fields  $T, V$ , a complex structure  $J$  and a 2-form  $\omega$  as in the Introduction, and also a symmetric bilinear form  $g := \omega(\cdot, J\cdot)$ .

Let  $M \subset N \times \mathcal{J}$  be the open subset where  $g(T, T)$  is positive. Define a positive function  $f : M \rightarrow \mathbb{R}^{>0}$  by  $g(T, T) = r^2f$ . Note that the restriction  $f_r := f(\cdot, r)$  of  $f$  to  $N_r := M \cap (N \times \{r\})$  is positive. We introduce a new family of contact forms  $\tilde{\theta}_r = (1/f_r)\theta_r$ ; for any  $r$ , the contact form  $\tilde{\theta}_r$  is defined on  $N_r$  (viewed as an open subset of  $N$ ).

*Conventions of notation.* For a function  $h : M \rightarrow \mathbb{R}$ , we shall denote by  $\dot{h} : M \rightarrow \mathbb{R}$  its derivative with respect to  $r$  and by  $h_r : N_r \rightarrow \mathbb{R}$  the restriction of  $h$  to a level set  $N_r$ .

LEMMA 4. *The following equalities hold:*

$$f = 1 + \frac{r\dot{\theta}_r(T_r)}{2} \tag{8}$$

and

$$\dot{\tilde{\theta}}_r = -\frac{2G}{r}\tilde{\theta}_r, \tag{9}$$

where

$$G : M \rightarrow \mathbb{R}, \quad G := \frac{rf}{2f} - f + 1.$$

*Proof.* Note that  $\omega = \frac{1}{2}(dr \wedge d^Jr + rdd^Jr)$ . It is straightforward to see that  $d^Jr = r\theta$ , where  $\theta \in \Omega^1(N \times \mathcal{J})$  is defined by  $\theta(Z) := \theta_r(\pi_*Z)$ , for a tangent vector  $Z \in T_{(p,r)}(N \times \mathcal{J})$ , where  $\pi : N \times \mathcal{J} \rightarrow N$  is the natural projection. At a tangent space  $T_{(p,r)}(N \times \mathcal{J}) = T_pN \times \mathbb{R}$ ,

$$d\theta = d\theta_r + dr \wedge \dot{\theta}_r,$$

and then, restricted to the same tangent space,

$$\omega = r \left( dr \wedge \left( \theta_r + \frac{r}{2}\dot{\theta}_r \right) + \frac{r}{2}d\theta_r \right). \tag{10}$$

It follows that

$$r^2f = \omega(V, T) = r^2 \left( 1 + \frac{r\dot{\theta}_r(T_r)}{2} \right),$$

which implies (8). To prove (9), we take the derivative with respect to  $r$  of the equality  $\tilde{\theta}_r = (1/f_r)\theta_r$  and we use the fact that  $\dot{\theta}_r = \dot{\theta}_r(T_r)\theta_r$ . We get

$$\dot{\tilde{\theta}}_r = -\frac{\dot{f}_r}{f_r}\tilde{\theta}_r + \frac{1}{f_r}\dot{\theta}_r = \left(-\frac{\dot{f}_r}{f_r} + \dot{\theta}_r(T_r)\right)\tilde{\theta}_r = -\frac{2G}{r}\tilde{\theta}_r,$$

which proves our lemma. □

LEMMA 5. *The pair  $(\omega, J)$  defines a Kähler structure on  $M$ .*

*Proof.* From relation (3), it is clear that the integrability tensor  $N^J$  of the almost complex structure  $J$ , applied to a pair of sections  $(X, Y)$  of  $H$ , vanishes. On the other hand,  $N^J(X, V)$ , restricted to a level set  $N \times \{r\}$ , is equal to  $-L_{T_r}(I)(X)$ , which is zero, because  $T_r$  is the Reeb vector field of a Sasaki structure. It follows that  $J$  is integrable. From (10) it is easy to see that  $T$  is hermitian orthogonal to  $H$  and that the restriction of  $g$  to  $H_{(p,r)} \subset T_{(p,r)}(N \times \mathcal{J})$  coincides with  $r^2g_r$ , which is positive definite. We deduce that  $g$  is positive definite on the subset  $M$  of  $N \times \mathcal{J}$  where  $g(T, T) > 0$ , and that  $(M, \omega, J)$  is a Kähler manifold (the 2-form  $\omega$  being closed). □

DEFINITION 6. The Kähler manifold  $(M, \omega, J)$  is a generalised Kähler cone over the CR manifold  $(N, H, I)$ . It is a restricted generalised Kähler cone if the function  $f$  is constant along the trajectories of the vector field  $T$ .

*Convention.* For simplicity, in this paper we will consider only restricted generalised Kähler cones; when we refer to a generalised Kähler cone, we will actually mean restricted generalised Kähler cone; this is true also for the statement of Theorem 1.

*Remark 1* (Main class of generalised Kähler cones). We shall be mainly concerned with generalised Kähler cones over (open subsets) of hermitian CR spheres. Suppose that  $N \subset \Sigma^{2m+1}$  is an open subset of the hermitian CR sphere of complex null lines in  $W = \mathbb{C}^{m+1,1}$ . Then

$$\theta_r = \frac{\eta}{(B_r w, w)}, \quad \tilde{\theta}_r = \frac{\eta}{(A_r w, w)}$$

for some hermitian trace-free operators  $A_r, B_r$  of  $W$ . The condition  $T(f) = 0$  is equivalent to  $[A_r, B_r] = 0$  for any  $r$ , as the following lemma shows.

LEMMA 7.

(i) *The operators  $A_r$  and  $B_r$  are related in the following way:*

$$A_r = B_r - \frac{r}{2}\dot{B}_r. \tag{11}$$

(ii) *The functions  $f$  and  $G$  have the following expression: for any  $(x, r) \in M$ ,*

$$f_r(x) = \frac{(A_r w, w)}{(B_r w, w)}, \quad G(x, r) = \frac{r(\dot{A}_r w, w)}{2(A_r w, w)}, \quad w \in x, \quad w \neq 0.$$

(iii) *The condition  $T(f) = 0$  is equivalent to  $[A_r, B_r] = 0$  for any  $r$ .*

*Proof.* Note that

$$\dot{\theta}_r = -\frac{(\dot{B}_r w, w)}{(B_r w, w)}\theta_r \quad \text{and} \quad \dot{\theta}_r(T_r) = -\frac{(\dot{B}_r w, w)}{(B_r w, w)}.$$

It follows that

$$f = 1 + \frac{r\dot{\theta}_r(T_r)}{2} = 1 - \frac{r(\dot{B}_r w, w)}{2(B_r w, w)} = \frac{(B_r w - \frac{1}{2}r\dot{B}_r w, w)}{(B_r w, w)}. \tag{12}$$

We deduce that

$$\tilde{\theta}_r = \frac{1}{f_r}\theta_r = \frac{(B_r w, w)}{(B_r w - \frac{1}{2}r\dot{B}_r w, w)} \cdot \frac{\eta}{(B_r w, w)} = \frac{\eta}{(B_r w - \frac{1}{2}r\dot{B}_r w, w)}. \tag{13}$$

Relation (11) follows from (13),  $\tilde{\theta}_r = \eta/(A_r w, w)$  and Lemma 2. From (11) and (12) we get the expression of  $f$ . On the other hand, from Lemma 4 and  $\tilde{\theta}_r = \eta/(A_r w, w)$ , we have

$$\dot{\tilde{\theta}}_r = -\frac{2G}{r}\tilde{\theta}_r = -\frac{(\dot{A}_r w, w)}{(A_r w, w)}\tilde{\theta}_r,$$

which implies that  $G$  is of the required form. To prove the last statement, note that

$$\begin{aligned} T_r(f_r) &= \frac{[(A_r T_r w, w) + (A_r w, T_r w)](B_r w, w) - [(B_r T_r w, w) + (B_r w, T_r w)](A_r w, w)}{(B_r w, w)^2} \\ &= \frac{i([A_r, B_r]w, w)}{(B_r w, w)}, \end{aligned}$$

since, at a point  $x \in N$ ,  $T_r(x) \in \text{Hom}(x, W/x)$  is the homomorphism  $T_r w = iB_r w \pmod{w}$  and the operators  $A_r$  and  $B_r$  are hermitian. We conclude from Lemma 2.  $\square$

LEMMA 8. *The Levi-Civita connection  $D^g$  of a generalised Kähler cone  $(M, \omega, J)$  has the following expression:*

$$\begin{aligned} D_X^g Y &= \nabla_X^r Y - \omega_r(X, Y)T_r - g_r(X, Y)V, \\ D_V^g Y &= fY + \frac{Y(f)}{2f}V - \frac{(JY)(f)}{2f}T, \\ D_T^g Y &= \mathcal{L}_T(Y) + fJY + \frac{Y(f)}{2f}T + \frac{(JY)(f)}{2f}V, \\ D_X^g V &= fX + \frac{X(f)}{2f}V - \frac{(JX)(f)}{2f}T, \\ D_V^g V &= -\frac{1}{2}v + (G + f)V, \\ D_T^g V &= \frac{1}{2}Jv + (G + 3f - 2)T. \end{aligned}$$

Here  $X, Y \in \Gamma(H)$ , the vector field  $D_X^g Y$  is restricted to a level set  $N_r$ ,  $\nabla^r$  is the Tanaka connection of the contact form  $\theta_r$  of the CR manifold  $(N, H, I)$  and  $v$  is a vector field on  $M$  which belongs, at any point  $(p, r) \in M$ , to  $H_p \subset T_{(p,r)}M$  and is determined by the condition:

$$g_r(X, v_{(p,r)}) = df(X), \quad \text{for all } X \in H_p \subset T_{(p,r)}M. \tag{14}$$

*Proof.* The proof is a straightforward computation based on the Koszul formula. It uses the expression of the Tanaka connection on the contact bundle  $H$ , mentioned in §2.  $\square$

LEMMA 9. *The curvature  $R^g$  of a generalised Kähler cone  $(M, \omega, J)$  has the following expression:*

$$\begin{aligned} g(R_{X,T}^g Y, Z) &= -\frac{Y(f)}{2}\omega(X, Z) - \frac{(JY)(f)}{2}g(X, Z) + \frac{Z(f)}{2}\omega(X, Y) \\ &\quad + \frac{(JZ)(f)}{2}g(X, Y) - X(f)\omega(Y, Z), \\ g(R_{X_1, X_2}^g Y, Z) &= g(R_{X_1, X_2}^{\nabla^r} Y, Z) - \frac{f}{r^2}\{g(X_1, Y)g(X_2, Z) - g(X_2, Y)g(X_1, Z)\} \\ &\quad + \frac{f}{r^2}\{-\omega(X_1, Y)\omega(X_2, Z) + \omega(X_2, Y)\omega(X_1, Z) - 2\omega(X_1, X_2)\omega(Y, Z)\}, \\ g(R_{X,T}^g V, Y) &= \frac{r^2}{2}(\nabla^r df)^{J,-}(X, JY) + \frac{r^2 f}{2}\left(\nabla^r\left(\frac{df}{f}\right)\right)^{J,+}(X, JY) \\ &\quad + f(2 - G - 2f)\omega(X, Y), \\ g(R_{T,V}^g V, T) &= g(v, v) + r^2 f((G - 2)(8f - 2) + r\dot{G} + 12f^2), \\ g(R_{T,V}^g V, Z) &= -\frac{r^3}{2}df(JZ) + r^2(G - 1)df(JZ), \end{aligned}$$



where  $X, X_1, X_2, Y, Z \in \Gamma(H)$ ,  $g(R_{X_1, X_2}^g Y, Z)$  is restricted to a level set  $N_r$ ,  $v$  is the vector field defined by (14) and the superscripts  $J, + (J, -)$  denote the  $J$ -invariant part (respectively the  $J$ -anti-invariant part) of a bilinear form.

*Proof.* The proof is a long but straightforward computation. □

PROPOSITION 10. *The generalised Kähler cone  $(M, \omega, J)$  is Bochner-flat if and only if the CR manifold  $(N, H, I)$  is flat and the following two conditions hold.*

- (i) *The function  $G$  depends only on  $r$ .*
- (ii) *For every  $r \in \mathcal{J}$ , the contact form  $\tilde{\theta}_r = (1/f_r)\theta_r$  is the contact form of a Sasaki structure on  $(N_r, H, I)$ , which determines an Einstein Kähler structure on the quotient  $N_r/\tilde{T}_r$  (where  $\tilde{T}_r$  is the Reeb vector field of  $\tilde{\theta}_r$ ), with modified Ricci tensor*

$$\tilde{S}_r = \left( \frac{r\dot{G}}{2} - G + 2 \right) \text{Id}.$$

*Proof.* Suppose first that  $(M, \omega, J)$  is Bochner flat, i.e.

$$R^g = c_{\mathcal{K}}^*(S), \tag{15}$$

for a tensor field  $S \in \text{Sym}^{1,1}(M)$ . Plugging into (15) the arguments  $(T, V, V, T)$  and  $(T, V, V, Z)$ , for  $Z \in \Gamma(H)$ , and using formula (2) for the adjoint of the Ricci contraction, we readily deduce that  $S(T, T)$  and  $S(Z, T)$  are related to the curvature  $R^g$  as follows:

$$S(T, T) = -\frac{1}{2r^2 f} g(R_{T, V}^g V, T), \quad S(Z, T) = -\frac{1}{r^2 f} g(R_{T, V}^g V, Z). \tag{16}$$

On the other hand, from Lemma 9 we know that

$$g(R_{X, V}^g Y, Z) = \frac{Y(f)g(X, Z)}{2} - \frac{Z(f)g(X, Y)}{2} - \frac{(JY)(f)\omega(X, Z)}{2} + \frac{(JZ)(f)\omega(X, Y)}{2} - (JX)(f)\omega(Y, Z),$$

for every  $X, Y, Z \in \Gamma(H)$ , and, since  $c_{\mathcal{K}}^*(S)(X, V, Y, Z) = g(R_{X, V}^g Y, Z)$ , we readily get  $S(Z, T) = 2(JZ)(f)$ . Combining this with the second relation (16), we deduce that

$$-\frac{1}{r^2 f} g(R_{T, V}^g V, Z) = 2(JZ)(f),$$

which is equivalent, using the expression of  $g(R_{T, V}^g V, Z)$  provided by Lemma 9, to  $(JZ)(G) = 0$ . Since  $Z \in \Gamma(H)$  is arbitrary, we obtain the first condition of the proposition (since  $X(G) = Y(G) = 0$ , for  $X, Y \in \Gamma(H)$ , also  $[X, Y](G) = 0$ ; recall now that vector fields of the form  $\{X, [X, Y], X, Y \in \Gamma(H)\}$  span the entire  $TN$ ). To obtain the second condition of the proposition, we notice that the expression of  $g(R_{X_1, X_2}^g Y, Z)$  found in Lemma 9, combined with (15), imply that the CR manifold  $(N, H, I)$  is flat and that, on the bundle  $H$  restricted to a level set  $N_r$ ,

$$S^\perp = \frac{1}{r^2} (S_r - 2f \text{Id}), \tag{17}$$

where  $S^\perp : H \rightarrow H$  is induced by  $S$  followed by  $g$ -orthogonal projection and  $S_r \in \text{End}(H)$  is the modified Ricci tensor of the Kähler curvature  $R^{\nabla^r}|_{H \times H} \in \Lambda^{1,1}(H) \odot \Lambda^{1,1}(H)$  of  $\nabla^r$ . Plugging into (15) the argument  $(X, T, V, Y)$  and using relation (17), we obtain

$$\begin{aligned} g(R_{X, T}^g V, Y) &= \frac{1}{4} (S(JX, Y)g(V, V) + \omega(X, Y)S(V, V)) \\ &= \frac{f}{4} g(S_r(JX), Y) - \frac{1}{4} \left( 2f^2 + \frac{g(R_{T, V}^g V, T)}{2r^2 f} \right) \omega(X, Y). \end{aligned}$$

Using the expression of  $g(R_{X,T}^g V, Y)$  provided by Lemma 9 we deduce that

$$g_r(S_r X, Y) = -\frac{2}{f} \nabla^r (df)^{J,-}(X, Y) - 2 \nabla^r \left( \frac{df}{f} \right)^{J,+}(X, Y) + \left( 4(2 - G) - 6f + \frac{g(R_{T,V}^g V, T)}{2f^2 r^2} \right) g_r(X, Y).$$

This relation clearly implies that  $\nabla^r (df)^{J,-}|_{H \times H} = 0$ , which means that  $\tilde{\theta}_r = (1/f_r)\theta_r$  (for any  $r$ ) is the contact form of a Sasaki structure on  $(N_r, H, I)$  (see [GO98]). Moreover, the modified Ricci tensor  $\tilde{S}_r$  of the Sasaki structure determined by  $\tilde{\theta}_r$  is related to  $S_r$  in the following way [Dav04]:

$$\frac{1}{f} g_r(\tilde{S}_r(X), Y) = g_r(S_r(X), Y) + 2 \nabla^r \left( \frac{df}{f} \right)^{J,+}(X, Y) - \frac{g_r(v, v)}{2f^2} g_r(X, Y). \tag{18}$$

We deduce, using the previous expression of  $g_r(S_r(X), Y)$ , that

$$\frac{1}{f} g_r(\tilde{S}_r(X), Y) + \left( \frac{g_r(v, v)}{2f^2} + 4(G - 2) + 6f - \frac{g(R_{T,V}^g V, T)}{2r^2 f^2} \right) g_r(X, Y) = 0.$$

Using again Lemma 9 for the expression of  $g(R_{T,V}^g V, T)$  we obtain the second condition of the proposition. Conversely, it is easy to check that the flatness of  $(N, H, I)$  and the two conditions of the proposition ensure the Bochner-flatness of  $(M, \omega, J)$ . □

The main result of this section is the following.

**THEOREM 11.** *Let  $(M, g, J)$  be a Bochner-flat generalised Kähler cone of complex dimension  $m + 1 \geq 3$ , defined by a family of Sasaki Reeb vector fields  $\{T_r\}$  over a CR manifold  $(N, H, I)$ . Then  $(N, H, I)$  is locally isomorphic to the standard CR sphere  $\Sigma^{2m+1}$  of complex null lines in a complex hermitian vector space  $W$  of signature  $(m + 1, 1)$ , and  $\{T_r\}$  is defined by one of the following families of hermitian operators  $B_r$  of  $W$ .*

- (i)  $B_r = r^2(B - \mu(r^2)A)$ . Here the real function  $\mu$  satisfies  $\mu' > 0$  and is a solution of the differential equation

$$\mu' = \frac{1}{2} \mu^2 + d, \tag{19}$$

where  $d \in \mathbb{R}$  is an arbitrary real number. The operator  $A$  is hermitian semi-simple, with a positive definite eigenspace, of dimension  $m + 1$ , which corresponds to the eigenvalue  $1/(2(m + 2))$  and a one-dimensional timelike eigenspace, which corresponds to the eigenvalue  $-(m + 1)/(2(m + 2))$ .

- (ii)  $B_r = r^2(B + \mu(r^2)A)$ , where  $\mu$  satisfies (19) and  $\mu' < 0$ . The operator  $A$  is hermitian semi-simple, with an eigenspace of signature  $(m, 1)$ , which corresponds to the eigenvalue  $-1/(2(m + 2))$ , and a one-dimensional spacelike eigenspace, which corresponds to the eigenvalue  $(m + 1)/(2(m + 2))$ .
- (iii)  $B_r = r^2(B - r^2A)$ , where  $A$  is one-step parabolic, with all eigenvalues equal to zero.
- (iv)  $B_r = r^2(B - (e^{\lambda r^2}/\lambda)A)$ , where  $\lambda \in \mathbb{R} \setminus \{0\}$  and  $A$  is one-step parabolic with all eigenvalues equal to zero.

In all these cases,  $B$  is any hermitian, trace-free operator of  $W$  which commutes with  $A$  (see Remark 2 following the proof, below).

*Proof.* Since  $(N, H, I)$  is flat (see Proposition 10) and of dimension bigger than three, we can assume, restricting  $N$  if necessary, that  $(N, H, I)$  is an open subset of the hermitian CR sphere  $\Sigma^{2m+1}$  of complex null lines in a complex hermitian vector space  $W$  of signature  $(m + 1, 1)$  (see [BS76]). As explained in Lemma 7, the two families of contact forms  $\{\theta_r\}_{r \in \mathcal{J}}$  and  $\{\tilde{\theta}_r\}_{r \in \mathcal{J}}$  are generated by two

families of hermitian trace-free endomorphisms  $\{B_r\}_{r \in \mathcal{J}}$  and  $\{A_r\}_{r \in \mathcal{J}}$  of  $W$  respectively, such that, for any  $r$ , the operators  $A_r$  and  $B_r$  commute. From Lemma 4 we know that  $\dot{\tilde{\theta}}_r = -(2G_r/r)\tilde{\theta}_r$ . Since  $G$  depends only on  $r$ , we get

$$\tilde{\theta}_r = e^{-\int_{r_0}^r (2G_q/q) dq} \tilde{\theta}_{r_0}$$

and we infer that the modified Ricci tensor  $\tilde{S}_r$  of  $\tilde{\theta}_r$  has the expression

$$\tilde{S}_r = e^{\int_{r_0}^r (2G_q/q) dq} \tilde{S}_{r_0}.$$

The second condition of Proposition 10 is equivalent with

$$\left(\frac{r\dot{G}}{2} - G + 2\right) \text{Id} = e^{\int_{r_0}^r (2G_s/s) ds} \tilde{S}_{r_0}$$

and implies that

$$\left(\frac{r\dot{G}}{2} - G + 2\right)' = \frac{2G}{r} \left(\frac{r\dot{G}}{2} - G + 2\right). \tag{20}$$

Equation (20) can be solved as follows. Define a real function  $\mu$  by

$$\mu(t) := \frac{G(\sqrt{t}) - 2}{t}. \tag{21}$$

We shall write (20) in terms of  $\mu$ . For this, we first take the derivative of  $r^2\mu(r^2) = G(r) - 2$  and we get:

$$\dot{G}(r) = 2r\mu(r^2) + 2r^3\mu'(r^2).$$

It easily follows that

$$\frac{r\dot{G}}{2} - G + 2 = r^4\mu'(r^2). \tag{22}$$

Equation (20) becomes  $\mu'' = \mu'\mu$ . Since  $\mathcal{J}$  is connected,  $\mu$  satisfies (19), for a constant  $d \in \mathbb{R}$ . We have the following three possibilities.

- (i)  $\mu' > 0$ . From Lemma 7 we deduce that

$$(\dot{A}_r w, w) = \frac{2}{r} G(r) (A_r w, w) = \frac{2}{r} (r^2\mu(r^2) + 2) (A_r w, w).$$

Since  $\int r\mu(r^2) dr = \frac{1}{2} \ln(\mu'(r^2))$  when  $\mu' > 0$  we get

$$(A_r w, w) = K(w)r^4\mu'(r^2),$$

where  $K = K(w)$  depends only on  $w$ . Equivalently,

$$A_r = \mu'(r^2)r^4 A, \tag{23}$$

where  $A \in \text{End}(W)$  is hermitian, trace-free, satisfies  $(Aw, w) > 0$ , for  $w \in x$  non-zero, when  $(x, r) \in M$ . Moreover, (22) together with the second condition of Proposition 10 imply that the modified Ricci tensor  $S_A$  of the Kähler–Einstein structure  $M_A$  is the identity endomorphism, from where we deduce that  $A$  is as in the statement of the theorem (see [DG06]). On the other hand, from Lemma 7,  $B_r$  must satisfy (11), with  $A_r = \mu'(r^2)r^4 A$ . It follows that

$$B_r = r^2(B - \mu(r^2)A),$$

where  $B \in \text{End}(W)$  is hermitian and trace-free.

- (ii)  $\mu' < 0$ . Then  $\int r\mu(r^2) dr = \frac{1}{2} \ln(-\mu'(r^2))$ . A similar argument shows that

$$\begin{aligned} A_r &= -r^4\mu'(r^2)A \\ B_r &= r^2(B + \mu(r^2)A), \end{aligned}$$

but in this case the Bochner-flat Kähler structure  $M_A$  has the modified Ricci operator  $S_A = -\text{Id}$ , which implies that  $A$  is as in the statement of the theorem (see [DG06]).

(iii) It remains to consider the case when the function  $\mu$  is constant. Then  $\mu(t) = \lambda$  for  $\lambda \in \mathbb{R}$ ,  $G(x, r) = \lambda r^2 + 2$  and

$$(\dot{A}_r w, w) = \frac{2G}{r}(A_r w, w) = \frac{2(\lambda r^2 + 2)}{r}(A_r w, w).$$

We distinguish two subcases: (i)  $\lambda = 0$ ; (ii)  $\lambda \neq 0$ . In subcase (i) we obtain

$$A_r = r^4 A, \quad B_r = r^2(B - r^2 A),$$

and in subcase (ii), we obtain

$$A_r = r^4 e^{\lambda r^2} A, \quad B_r = r^2 \left( B - \frac{e^{\lambda r^2}}{\lambda} A \right).$$

Since  $\frac{1}{2}r\dot{G} - G + 2 = 0$ , the Kähler structure  $M_A$  is flat and hence the endomorphism  $A$  is one-step parabolic, with all eigenvalues zero (see [DG06]).

In all cases described above, the generalised Kähler cone condition  $T(f) = 0$  becomes  $[A, B] = 0$  (see Lemma 7). □

*Remark 2.* The condition  $[A, B] = 0$  of Theorem 11 determines the operator  $B$  as follows.

(i) In case (i) of Theorem 11,  $B$  preserves, up to a multiplicative constant, a timelike eigenvector  $v$  (unique, up to a non-zero multiplicative constant) of  $A$ . On the hermitian orthogonal  $v^\perp$ , the hermitian metric  $(\cdot, \cdot)$  of  $W$  is positive definite,  $A : v^\perp \rightarrow v^\perp$  is a multiple of the identity endomorphism and  $B : v^\perp \rightarrow v^\perp$ , being hermitian, is diagonalisable. It follows that  $A$  and  $B$  are simultaneously diagonalisable.

(ii) In case (ii) of Theorem 11,  $B$  preserves, up to a multiplicative constant, a spacelike eigenvector  $v$  (unique, up to a non-zero multiplicative constant) of  $A$ , which corresponds to the eigenvalue  $(m + 1)/(2(m + 2))$ . On the hermitian orthogonal  $v^\perp$ , the hermitian metric  $(\cdot, \cdot)$  has signature  $(m, 1)$ ,  $A$  is a multiple of the identity endomorphism and  $B : v^\perp \rightarrow v^\perp$  can be elliptic, hyperbolic, one- or two-step parabolic.

(iii) Consider now cases (iii) and (iv) of Theorem 11. Note that  $A = 0$  on any positive definite eigenspace of  $B$  (since  $[A, B] = 0$ ,  $A$  preserves such an eigenspace, say  $W_1$ , of  $B$ ; because the hermitian metric  $(\cdot, \cdot)$  is positive definite on  $W_1$  and  $A$  is hermitian,  $A$  is diagonalisable on  $W_1$ ; this forces it to be zero, because  $A$  does not have non-zero eigenvalues). Let us denote by  $W_1, \dots, W_s$  the positive definite eigenspaces of  $B$  and by  $W_0$  the hermitian orthogonal of the direct sum  $\bigoplus_{j=1}^s W_j$ . The eigenspaces  $W_j$  (for  $j \in \{1, \dots, s\}$ ) correspond to eigenvalues, say  $\beta_j$ , of  $B$ , which can be any real numbers. It remains to study the restriction  $B_0$  of  $B$  to  $W_0$ . We notice first that  $B_0$  cannot be hyperbolic: if it was hyperbolic, then  $B$  would have two complex non-real eigenvalues, say  $\delta$  and  $\bar{\delta}$ , with one-dimensional eigenspaces, generated by two null independent vectors  $v_1$  and  $v_2$  respectively. However, since  $[A, B] = 0$ ,  $BAv_1 = \delta Av_1$  and  $BAv_2 = \bar{\delta} Av_2$ , which imply that  $Av_1 = Av_2 = 0$  (because  $A$  has no non-zero eigenvalues). But if we take an orthonormal basis  $\{e_0, \dots, e_n\}$  of  $W_0$  in which

$$A = \begin{pmatrix} -1 & 1 & 0 & \cdots & 0 \\ -1 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

(which is possible since  $A : W_0 \rightarrow W_0$  is one-step parabolic with all eigenvalues equal to zero) the conditions  $v_1, v_2$  null and  $Av_1 = Av_2 = 0$  would imply that  $v_1$  and  $v_2$  are multiples of  $e_0 + e_1$ .

TABLE 1. Local type of Bochner-flat generalised Kähler cones.

	$\mu' > 0$	$\mu' = 0$
$d > 0$	(all) hyperbolic	—
$d = 0$	(all) one-step parabolic	$\lambda = 0$ ; (all) two-step parabolic
$d < 0$	(all) elliptic	$\lambda = \pm\sqrt{-2d}$ , (all) one-, two-step parabolic, elliptic

In particular they would be dependent, which is a contradiction. We conclude that  $B_0$  can be elliptic or one- or two-step parabolic. Therefore,  $B_0 = \gamma I + N$ , for an endomorphism  $N$  of  $W_0$  which commutes with  $A$  and which satisfies  $N^3 = 0$ , and for  $\gamma \in \mathbb{R}$  which is different from all  $\beta_j$ . The endomorphism  $N^\perp$  of  $\widehat{W}_0 := \text{Span}\{e_2, \dots, e_n\}$  obtained from  $N$  by restriction and orthogonal projection is hermitian on  $\widehat{W}_0$ . Because the metric  $(\cdot, \cdot)$  is positive definite on  $\widehat{W}_0$ ,  $N^\perp$  is diagonalisable and hence there is a basis  $\{e'_2, \dots, e'_n\}$  of  $\widehat{W}_0$ , such that  $N^\perp$  is diagonal in this basis. If we consider now the basis  $\mathcal{B} := \{e_0, e_1, e'_2, \dots, e'_n\}$  of  $W_0$ , it is straightforward to see that  $[A, N] = 0$  and  $N$  hermitian imply that

$$N = \begin{pmatrix} \gamma_0 & \alpha & \mu_2 & \mu_3 & \cdots & \mu_n \\ -\alpha & \gamma_1 & \mu_2 & \mu_3 & \cdots & \mu_n \\ -\bar{\mu}_2 & \bar{\mu}_2 & \gamma_2 & 0 & \cdots & 0 \\ -\bar{\mu}_3 & \bar{\mu}_3 & 0 & \gamma_3 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ -\bar{\mu}_n & \bar{\mu}_n & 0 & 0 & \cdots & \gamma_n \end{pmatrix}$$

in the basis  $\mathcal{B}$ . Moreover,  $N^3 = 0$  if and only if  $\gamma_0 = -\gamma_1 = -\alpha$  and  $\gamma_k = 0$  for any  $k \in \{2, \dots, n\}$  and  $N^2 = 0$  if and only if  $\gamma_0 = -\gamma_1 = -\alpha$ ,  $\gamma_k = \mu_k = 0$ , for any  $k \in \{2, \dots, n\}$ . Since  $B$  is trace-free, the real constants  $\beta_j$  and  $\gamma$  must satisfy  $(n + 1)\gamma + \sum_{j=1}^s n_j \beta_j = 0$ , where  $n_j$  is the dimension of  $W_j$  (and  $n + 1$  is the dimension of  $W_0$ ).

#### 4. The local geometry of Bochner-flat generalised Kähler cones

Without further explanations, we shall continue to employ the notation of the previous section. The aim of this section is to prove our main Theorem 1. We will do this by showing that all local types of Bochner-flat Kähler manifolds (elliptic, hyperbolic, one- and two-step parabolic) are covered by the Bochner-flat generalised Kähler cones which belong to cases (i), (iii) and (iv) of Theorem 11. Therefore, for our purpose – to prove Theorem 1 – it is not necessary to go into the study of the local type of the Bochner-flat generalised Kähler cones which belong to the second case of Theorem 11. Formally, cases (i) and (ii) of Theorem 11 are similar. However, with respect to the first case, the second case is more difficult to analyse, due to the fact that there is no restriction on the operator  $B$ , except that it should be hermitian, trace-free and commute with  $A$ ; in particular,  $B$  is not necessarily diagonalisable, as it is in the first case (see Remark 2). We summarise the results we shall obtain in this section in Table 1.

##### 4.1 The case of Theorem 11(i)

In this subsection we analyse the first column of the above table.

PROPOSITION 12. *Let  $(M, g, J)$  be a Bochner-flat generalised Kähler cone which belongs to case (i) of Theorem 11. Then  $(M, g, J)$  is:*

- (i) *of hyperbolic type, if  $d > 0$ ;*

- (ii) of elliptic type, if  $d < 0$ ;
- (iii) of one-step parabolic type, if  $d = 0$ .

Conversely, any Bochner-flat Kähler manifold which is of elliptic, hyperbolic or one-step parabolic type can be locally realised as a generalised Kähler cone, which belongs to case (i) of Theorem 11.

*Proof.* In case (i) of Theorem 11,

$$M = \{(x, r) \in \Sigma^{2m+1} \times \mathbb{R}^{>0} : (Bw, w) > \frac{1}{2}\mu(r^2)(Aw, w), (Aw, w) > 0, \forall w \in x, w \neq 0\}.$$

From Remark 2, there is an orthonormal basis  $\mathcal{B} = \{e_0, e_1, \dots, e_{m+1}\}$  of  $W$  such that both operators  $A$  and  $B$  are diagonal in this basis:

$$B = \text{diag}(-k, k_1, \dots, k_{m+1}),$$

$$A = \frac{1}{2(m+2)} \text{diag}(-m-1, 1, \dots, 1).$$

Here  $k_j \in \mathbb{R}$ , for any  $j \in \{1, \dots, m+1\}$ , and  $k = k_1 + \dots + k_{m+1}$ . We shall identify  $\Sigma^{2m+1}$  with the unit sphere  $S^{2m+1}$  in  $\text{Span}\{e_1, \dots, e_{m+1}\}$  and  $S^{2m+1} \times \mathbb{R}^{>0}$  with  $\mathbb{C}^{m+1} \setminus \{0\}$ , by means of the diffeomorphism

$$h : S^{2m+1} \times \mathbb{R}^{>0} \rightarrow \mathbb{C}^{m+1} \setminus \{0\}, \quad f(z_1, \dots, z_{m+1}, r) := (rz_1, \dots, rz_{m+1}).$$

Read on the image

$$h(M) = \left\{ (z_1, \dots, z_{m+1}) \in \mathbb{C}^{m+1} \setminus \{0\}, \sum_{j=1}^{m+1} (k + k_j - \frac{1}{2}\mu(r^2))|z_j|^2 > 0 \right\},$$

the complex structure  $J$ , at a point  $z \in h(M)$ , satisfies

$$J(V) = r^2 \sum_{j=1}^{m+1} (k_j + k - \frac{1}{2}\mu(r^2)) \left( x_j \frac{\partial}{\partial y_j} - y_j \frac{\partial}{\partial x_j} \right),$$

$$J|_{z^\perp} = i,$$

and the Kähler form  $\omega$  is equal to  $\frac{1}{4}dd^J r^2$ . Here  $r^2 = |z_1|^2 + \dots + |z_{m+1}|^2$  and  $z_j = x_j + iy_j$  for any  $j \in \{1, \dots, m+1\}$ . For simplicity, we restrict the Kähler structure  $(\omega, J)$  to the subset

$$D := \{z \in \mathbb{C}^{m+1} \setminus \{0\}, \mu(r^2) < 2(k_j + k), \forall j \in 1, 2, \dots, m+1\} \tag{24}$$

of  $h(M)$ . We shall consider separately the three cases:  $d > 0$ ,  $d < 0$  and  $d = 0$ .

- (i) Suppose that  $d = \beta^2/2$ , with  $\beta > 0$ . Then  $\mu(t) = \beta \text{tg}(\beta t/2)$ . It can be checked that the map

$$F(z_1, \dots, z_{m+1}) := (w_1 = f_1(r^2)z_1, \dots, w_{m+1} = f_{m+1}(r^2)z_{m+1}), \tag{25}$$

where

$$f_j(t) = \left( \frac{\beta}{\sqrt{2}} \right)^{1/2} \frac{e^{(k_j+k)t/2}}{t^{1/2}\mu'(t)^{1/4}}, \tag{26}$$

is an isomorphism between the Kähler manifolds  $(D, \frac{1}{4}dd^J r^2, J)$  and  $(F(D), \frac{1}{4}dd^{J_0} x, J_0)$ . Here  $J_0$  is the standard complex structure of  $\mathbb{C}^{m+1}$  and the positive function  $x = x(w_1, \dots, w_{m+1})$  is defined by the implicit equation

$$\sum_{j=1}^{m+1} \frac{|w_j|^2}{f_j^2(x)} = x. \tag{27}$$

Let  $y = y(w_1, \dots, w_{m+1})$  be related to  $x$  by the formula

$$x = \frac{4}{\beta} \text{arctg}[(1+y)^{1/2}]$$

and notice that

$$\frac{e^{(k_j+k)x}}{\dot{\mu}(x)^{1/2}} = \frac{\sqrt{2}|y|}{\beta(2+y)} e^{2l_j \arctg[(1+y)^{1/2}]},$$

where  $l_j = (2/\beta)(k_j + k)$ , for  $j \in \{1, \dots, m + 1\}$ . For simplicity, we restrict to the set, say  $D' \subset F(D)$ , where  $y > 0$ . On this set,  $(\frac{1}{4}dd^{J_0}x, J_0)$  coincides with the Kähler structure  $(\frac{1}{\beta}dd^{J_0} \arctg[(1+y)^{1/2}], J_0)$ , where  $y$  is defined by the implicit equation

$$\sum_{j=1}^{m+1} \frac{|w_j|^2}{e^{2l_j \arctg[(1+y)^{1/2}]}} = \frac{y}{2+y}. \tag{28}$$

The Kähler structure  $((1/\beta)dd^{J_0} \arctg[(1+y)^{1/2}], J_0)$  is of hyperbolic type (communicated to the author by Paul Gauduchon), isomorphic with  $M_C$ , where  $C$  is a hyperbolic hermitian operator of  $\mathbb{C}^{m+2,1}$  with characteristic and minimal polynomials

$$Q_C = \left( \left( t + \frac{2(m+2)k}{\beta(m+3)} \right)^2 + 1 \right) \prod_{j=1}^{m+1} \left( t - \frac{2}{\beta} \left( k_j + \frac{k}{m+3} \right) \right),$$

$$q_C = \left( \left( t + \frac{2(m+2)k}{\beta(m+3)} \right)^2 + 1 \right) \prod_{i=1}^s \left( t - \frac{2}{\beta} \left( k_i + \frac{k}{m+3} \right) \right).$$

Here  $i \in \{1, \dots, s\}$  parametrises the distinct values of  $\{k_1, \dots, k_{m+1}\}$ . This proves our first claim.

(ii) Next, suppose that  $d := -\beta^2/2$ , where  $\beta > 0$ . Then

$$\mu(t) = \frac{\beta(1 + e^{t\beta+p})}{1 - e^{t\beta+p}},$$

where  $p \in \mathbb{R}$ . By means of the function (25) with

$$f_j(t) = \frac{\beta^{1/2} 2^{1/4} e^{\frac{1}{2}(k_j+k)(t+p/\beta)+p}}{t^{1/2} \mu'(t)^{1/4}}, \quad j \in \{1, \dots, m + 1\}, \tag{29}$$

the Kähler manifolds  $(D, \omega, J)$  and  $(F(D), \omega_0 := \frac{1}{4}dd^{J_0}x, J_0)$  are isomorphic, where  $x = x(w_1, \dots, w_{m+1})$  is defined by the implicit equation (27), with functions  $f_j$  defined in (29). Define a new function  $y = y(w_1, \dots, w_{m+1})$  by

$$y = e^{\beta x} - e^{-p}.$$

For simplicity, we restrict to the subset of  $F(D)$ , say  $D'$ , where  $y > 0$ . On  $D'$ , the function  $y$  satisfies the implicit equation

$$\sum_{j=1}^{m+1} \frac{|w_j|^2}{e^{p_j y} (e^{p_j y} + 1)^{\beta_j}} = 1,$$

with  $\beta_j := (1/\beta)(k_j + k - \beta/2)$ . In terms of  $y$ ,

$$\omega_0 = \frac{1}{4\beta} dd^{J_0} \log(e^{p_j y} + 1).$$

It follows that  $(\omega_0, J_0)$  is of elliptic type on  $D'$  (see [DG06, § 2.2]) and is isomorphic with the Kähler structure  $M_C$ , with  $C$  a semi-simple hermitian operator of  $\mathbb{C}^{m+2,1}$ , with eigenvalues

$$-\frac{\beta}{2} - \frac{(m+2)k}{m+3}, \quad k_1 + \frac{k}{m+3}, \quad \dots, \quad k_{m+1} + \frac{k}{m+3}, \quad \frac{\beta}{2} - \frac{(m+2)k}{m+3}.$$

Our second claim follows.

(iii) Finally, suppose that  $d = 0$ . Then  $\mu(t) = -2/(t + q)$ , for  $q \in \mathbb{R}$ . For simplicity, we assume that  $q \geq 0$ . The function (25), with

$$f_j(t) = \frac{e^{(k_j+k)t/2}(t + q)^{1/2}}{t^{1/2}},$$

defines an isomorphism between the Kähler manifolds  $(D, \omega, J)$  and  $(F(D), \omega_0 := \frac{1}{4}dd^J_0x, J_0)$  where  $x = x(w_1, \dots, w_{m+1})$  is a positive function defined implicitly by the equation

$$\sum_{j=1}^{m+1} \frac{|w_j|^2}{e^{(k_j+k)x}(x + q)} = 1.$$

Note that the function  $y := x + q > 0$  satisfies

$$\sum_{j=1}^{m+1} \frac{|w_j|^2}{e^{(k_j+k)(y-q)}} = y. \tag{30}$$

Moreover,  $\omega_0 = \frac{1}{4}dd^J_0y$ . The Kähler structure  $(\frac{1}{4}dd^J_0y, J_0)$  is of one-step parabolic type (see [DG06, § 3.1]). It is isomorphic to the Kähler structure  $M_C$ , where  $C$  is a one-step parabolic hermitian operator of  $\mathbb{C}^{m+2,1}$ , with characteristic and minimal polynomials

$$Q_C(t) = \left(t + \frac{(m + 2)k}{m + 3}\right)^2 \prod_{j=1}^{m+1} \left(t - k_j - \frac{k}{m + 3}\right),$$

$$q_C(t) = \left(t + \frac{(m + 2)k}{m + 3}\right)^2 \prod_{i=1, k_i \neq -k}^s \left(t - k_i - \frac{k}{m + 3}\right).$$

As before,  $i \in \{1, \dots, s\}$  parametrises the distinct values of  $\{k_1, \dots, k_{m+1}\}$ . Our third claim follows.

The last statement of the proposition follows by an examination of the minimal and characteristic polynomials of the operators  $C$  we found in each of the cases (i), (ii) and (iii) above. □

### 4.2 The cases of Theorem 11(iii) and (iv)

In this section we prove the following proposition.

PROPOSITION 13.

- (i) Let  $(M, g, J)$  be a Bochner-flat generalised Kähler cone, which belongs to case (iii) of Theorem 11. Then  $(M, g, J)$  is of two-step parabolic type.
- (ii) Let  $(M, g, J)$  be a Bochner-flat generalised Kähler cone, which belongs to case (iv) of Theorem 11. Then  $(M, g, J)$  is of two-step parabolic type, except when  $\mu_j = 0$  for any  $j \in \{2, \dots, n\}$ . In this case it is of one-step parabolic type if  $\alpha \neq 0$  and of elliptic type if  $\alpha = 0$ .
- (iii) Any two-step parabolic Bochner-flat Kähler structure can be locally realised as a Bochner-flat generalised Kähler cone which belongs to case (iii) of Theorem 11, and also as a Bochner-flat generalised Kähler cone which belongs to case (iv) of Theorem 11.

We divide the proof of Proposition 13 into several lemmas and propositions. Let  $(M, \omega, J)$  be a Bochner-flat generalised Kähler cone, which belongs to case (iii) or (iv) of Theorem 11. We continue to preserve the notation of Theorem 11 and Remark 2.



PROPOSITION 14. *The Bryant modified Ricci operator  $\Theta$  of  $(M, g, J)$  has the following expression:*

$$\begin{aligned} \Theta(L_j) &= \left(\eta_j - \frac{c}{m+3}\right)L_j - \frac{q_r(\xi_j)}{r^4(\eta_j - \gamma)}V, \quad j \in \{1, \dots, l\}, \\ \Theta(v_k) &= \left(\frac{\lambda_k}{r^2} - \frac{c}{m+3}\right)v_k, \quad k \in \{1, \dots, m-l\}, \\ \Theta(V) &= \frac{f}{r^4} \sum_{j=1}^l \frac{L_j}{p'_r(\xi_j)(\eta_j - \gamma)} - \left(\frac{f}{r^2} - \frac{(m+2)c}{m+3}\right)V. \end{aligned}$$

Here  $\xi_1, \dots, \xi_l$  (respectively,  $\lambda_1, \dots, \lambda_{m-l}$ ) are the non-constant (respectively, constant) eigenvalues of the Bryant modified Ricci operator  $\Theta_r$  of  $M_{B_r}$ ,  $L_1, \dots, L_l$  are vector fields on  $M$  which, at a point  $(x, r) \in M$ , belong to  $H_x = \text{Hom}_{\mathbb{C}}(x, W/x)$  and are the homomorphisms  $L_j(w) = \bar{b}_r(\xi_j)w \pmod{w}$ ,  $v_1, \dots, v_{m-l}$  are eigenvectors of  $\Theta$  which correspond to the eigenvalues  $\lambda_1, \dots, \lambda_{m-l}$ ,  $p_r(t) = \prod_{j=1}^l (t - \xi_j)$  is the non-constant part of the characteristic polynomial of  $\Theta_r$ ,  $\eta_1 := \xi_1/r^2, \dots, \eta_l := \xi_l/r^2$ ,  $q_r$  is the minimal polynomial of  $B_r$  and  $c = \gamma - \lambda$ .

*Proof.* Recall that  $\Theta$  is related to the modified Ricci tensor  $S$  from the proof of Proposition 10 by

$$\Theta = \frac{1}{4} \left( S - \frac{\text{trace}_{\mathbb{R}}(S)}{2(m+3)} \text{Id} \right).$$

We need to determine  $S(L_j)$ ,  $S(v_k)$ ,  $S(V)$  and  $\text{trace}_{\mathbb{R}}(S)$ . From the proof of Proposition 10 we know that, for any  $X \in H$ ,

$$S(X) = \frac{1}{r^2} (S_r X - 2fX) + \frac{2(JX)(f)}{r^2 f} T - \frac{2X(f)}{r^2 f} V. \tag{31}$$

It is easy to check the following equalities:

$$df(L_j) = \frac{2f q_r(\xi_j)}{r^2(\eta_j - \gamma)}, \quad df(JL_j) = 0, \quad df(v_k) = 0, \quad df(Jv_k) = 0, \tag{32}$$

which imply, using  $\Theta_r(L_j) = \xi_j L_j$ ,  $\Theta_r(v_k) = \lambda_k v_k$  and relation (5) applied to  $\Theta_r$ , that

$$\begin{aligned} S(L_j) &= \frac{2}{r^2} \left( 2\xi_j - \frac{(B_r^2 w, w)}{(B_r w, w)} - f \right) L_j - \frac{4q_r(\xi_j)}{r^4(\eta_j - \gamma)} V, \\ S(v_k) &= \frac{2}{r^2} \left( 2\lambda_k - \frac{(B_r^2 w, w)}{(B_r w, w)} - f \right) v_k. \end{aligned}$$

To evaluate  $S(V)$ , we write it as a sum of  $S^\perp(V)$ , the  $g$ -orthogonal projection of  $S(V)$  on  $H$ , and  $(S(V, V)/r^2 f)V$ . It is easy to check, using the fact that  $S$  is hermitian, relation (6) and  $L_j = \frac{1}{2} p'_r(\xi_j) \text{grad}_{g_r}(\xi_j)$  (see [DG06, Theorem 2]), that

$$S^\perp(V) = \frac{4f}{r^4} \sum_{j=1}^l \frac{L_j}{p'_r(\xi_j)(\eta_j - \gamma)}. \tag{33}$$

To evaluate  $S(V, V) = S(T, T)$ , we use the first equality of (16) and the expression of  $g(R_{T,V}^g V, T)$  provided by Lemma 9. Notice that the vector field  $v$  defined in Lemma 8 has the following expression: at a point  $(x, r) \in M$ ,  $v_{(x,r)} \in H_x = \text{Hom}(x, W/x)$  is equal to

$$v_{(x,r)}(w) = 2(A_r w - f_r(x) B_r w) \pmod{w},$$

and so

$$g_r(v_{(x,r)}, v_{(x,r)}) = 4f_r(x)^2 \left( \frac{(B_r^2 w, w)}{(B_r w, w)} - 2r^2 \gamma \right), \tag{34}$$

where  $w \in x$  is non-zero. It follows that

$$S(V, V) = -2f \left( \frac{(B_r^2 w, w)}{(B_r w, w)} - 2r^2 \gamma \right) - 4\lambda r^2 f - 6f^2. \tag{35}$$

Relation (33), together with (35), determine  $S(V)$ . It remains to calculate  $\text{trace}_{\mathbb{R}}(S)$ . Using (31) and (35), we have

$$\begin{aligned} \text{trace}_{\mathbb{R}}(S) &= \frac{1}{r^2}(\text{trace}_{\mathbb{R}}(S_r) - 4mf) + \frac{2}{r^2 f} S(V, V) \\ &= -\frac{4}{r^2} \left( (m+3) \left( \frac{(B_r^2 w, w)}{(B_r w, w)} + f \right) - 2r^2 c \right). \end{aligned}$$

Our claim follows now easily, combining the expressions of  $S(L_j)$ ,  $S(v_k)$ ,  $S(V)$  and  $\text{trace}_{\mathbb{R}}(S)$  determined above. □

We introduce a new family of hermitian operators  $\hat{B}_r := (1/r^2)B_r = B + \delta(r)A$ , with  $\delta(r) := -r^2$  when  $\lambda = 0$  (equivalently, when  $(M, \omega, J)$  belongs to case (iii) of Theorem 11) and  $\delta(r) = -e^{\lambda r^2} / \lambda$  when  $\lambda \neq 0$  (equivalently, when  $(M, \omega, J)$  belongs to case (iv) of Theorem 11). Let  $\hat{q}$ , respectively  $\hat{Q}$ , be the minimal and characteristic polynomials of  $\hat{B}_r$ , equal to

$$\hat{q}(t) = (t - \gamma)^3 \prod_{j=1}^s (t - \beta_j), \quad \hat{Q} := (t - \gamma)^{n+1} \prod_{j=1}^s (t - \beta_j)^{n_j},$$

if there is  $\mu_j \neq 0$  and to

$$\hat{q}(t) = (t - \gamma)^2 \prod_{j=1}^s (t - \beta_j), \quad \hat{Q}(t) = (t - \gamma)^{n+1} \prod_{j=1}^s (t - \beta_j)^{n_j},$$

otherwise. Let  $\hat{p}_r(t) := \prod_{j=1}^l (t - \eta_j)$  be the non-constant part of the characteristic polynomial of the Bryant modified Ricci operator  $\hat{\Theta}_r$  of the Kähler structure  $M_{\hat{B}_r}$ . It will be considered as a polynomial with function coefficients defined on  $\Sigma_{\hat{B}_r}^{2m+1}$ . We shall denote by  $\hat{p}_{r,x}$  its value at a null line  $x \in \Sigma_{\hat{B}_r}^{2m+1}$ , which is a polynomial with constant coefficients.

PROPOSITION 15. *Let  $\hat{q}_1$  be the constant polynomial equal to the quotient of  $\hat{q}$  by  $(t - \gamma)^2$ . Then the characteristic polynomial  $P(t)$  of the modified Ricci operator  $\Theta$  of  $(M, g, J)$  is equal to the product*

$$\frac{\hat{Q}(t + c/(m+3))}{\hat{q}(t + c/(m+3))} P_1(t),$$

where

$$P_1(t) := \left( t - \frac{(m+2)c}{m+3} \right) \hat{p}_r \left( t + \frac{c}{m+3} \right) + \frac{f}{r^2} \hat{q}_1 \left( t + \frac{c}{m+3} \right). \tag{36}$$

*Proof.* Using Proposition 14, together with the fact that the constant part of the characteristic polynomial of the modified Ricci operator  $\Theta_A$  of a Bochner-flat Kähler structure  $M_A$  is equal to the quotient of the characteristic polynomial by the minimal polynomial of  $A$  (see [DG06, § 1.5]), it is easy to see that

$$P(t) = \frac{\hat{Q}(t + c/(m+3))}{\hat{q}(t + c/(m+3))} P_1(t),$$

where

$$P_1(t) = \left( t - \frac{(m+2)c}{m+3} + \frac{f}{r^2} \right) \hat{p}_r \left( t + \frac{c}{m+3} \right) + \frac{f}{r^8} \sum_{j=1}^l \frac{q_r(\xi_j)}{(\eta_j - \gamma)^2 p'_r(\xi_j)} \prod_{i \neq j} \left( t - \eta_i + \frac{c}{m+3} \right).$$

We shall evaluate the expression

$$\mathcal{E}(t) := \frac{f}{r^8} \sum_{j=1}^l \frac{q_r(\xi_j)}{(\eta_j - \gamma)^2 p'_r(\xi_j)} \prod_{i \neq j} \left( t - \eta_i + \frac{c}{m+3} \right).$$

For this, let  $g_r, \hat{g}_r$  be the Bochner-flat Kähler metrics of  $M_{B_r}$ , respectively  $M_{\hat{B}_r}$  (viewed as metrics on  $H$ ). Then, from relation (10),  $\hat{g}_r = r^2 g_r = g$  on  $H$ . Using (6), we get

$$-4 \frac{q_r(\xi_j)}{p'_r(\xi_j)} = g_r(\text{grad}_{g_r}(\xi_j), \text{grad}_{g_r}(\xi_j)) = r^6 \hat{g}_r(\text{grad}_{\hat{g}_r}(\eta_j), \text{grad}_{\hat{g}_r}(\eta_j)) = -4r^6 \frac{\hat{q}(\eta_j)}{\hat{p}'_r(\eta_j)},$$

which implies that

$$\mathcal{E}(t) = \frac{f}{r^2} \sum_{j=1}^l \frac{\hat{q}_1(\eta_j)}{\hat{p}'_r(\eta_j)} \prod_{i \neq j} \left( t - \eta_i + \frac{c}{m+3} \right).$$

Note that  $\mathcal{E}_1(t) := (r^2/f)\mathcal{E}(t)$  is a polynomial of degree  $l - 1$  which satisfies

$$\mathcal{E}_1 \left( \eta_j - \frac{c}{m+3} \right) = \frac{\hat{q}_1(\eta_j)}{\hat{p}'_r(\eta_j)} \prod_{i \neq j} (\eta_j - \eta_i) = \hat{q}_1(\eta_j), \quad j \in \{1, \dots, l\}.$$

Since  $\hat{q}_1$  is a monic polynomial of degree  $l$  and  $\mathcal{E}_1$  is of degree  $l - 1$ , we deduce that  $\hat{q}_1(t) = \mathcal{E}_1(t - c/(m + 3)) + \hat{p}_r(t)$ , which implies that

$$\sum_{j=1}^l \frac{\hat{q}_1(\eta_j)}{\hat{p}'_r(\eta_j)} \prod_{i \neq j} \left( t - \eta_i + \frac{c}{m+3} \right) = \hat{q}_1 \left( t + \frac{c}{m+3} \right) - \hat{p}_r \left( t + \frac{c}{m+3} \right). \tag{37}$$

Our claim follows. □

The constant roots of the polynomial  $P_1$  (which are also constant eigenvalues of  $\Theta$ ) will be determined in Proposition 17. For the proof of this proposition we need the following additional lemma.

LEMMA 16. *The following equality holds:*

$$\frac{d}{dr} \hat{p}_r(t) = \frac{2f}{r} (\hat{p}_r(t) - \hat{q}_1(t)).$$

*Proof.* We take the derivative with respect to  $r$  of the equality

$$((\eta_j \text{Id} - B - \delta A)^{-1} w, w) = 0$$

(which follows from  $\hat{p}_r(\eta_j) = 0$ ) and we obtain

$$((\eta_j \text{Id} - \hat{B}_r)^{-1} (\dot{\eta}_j \text{Id} - \dot{\delta} A) (\eta_j \text{Id} - \hat{B}_r)^{-1} w, w) = 0. \tag{38}$$

On the other hand, since  $AB = \gamma A$ , it is easy to see that

$$(\eta_j \text{Id} - \hat{B}_r)^{-1} A (\eta_j \text{Id} - \hat{B}_r)^{-1} = \frac{A}{(\eta_j - \gamma)^2}. \tag{39}$$

Applying (7) to  $A := \hat{B}_r$  and using the fact that  $\eta_j \text{Id} - \hat{B}_r$  is invertible, we get

$$((\eta_j \text{Id} - \hat{B}_r)^{-2}w, w) = -\frac{\hat{p}'_r(\eta_j)}{\hat{q}(\eta_j)}(\hat{B}_r w, w). \tag{40}$$

Combining (38), (39) and (40), and using the fact that  $\delta(r) = -e^{\lambda r^2}/\lambda$  when  $\lambda \neq 0$  and  $\delta(r) = -r^2$  when  $\lambda = 0$ , we deduce the expressions of the derivatives  $\dot{\eta}_j$  as follows:

$$\dot{\eta}_j = \frac{2f}{r} \frac{\hat{q}_1(\eta_j)}{\hat{p}'_r(\eta_j)}. \tag{41}$$

Since

$$\frac{d}{dr}\hat{p}_r(t) = -\sum_{j=1}^l \dot{\eta}_j \prod_{i \neq j} (t - \eta_i)$$

we get, using (37), our claim. □

PROPOSITION 17. *The following statements hold:*

- (i) *Suppose that  $\hat{q}_1(c) \neq 0$ . Then the polynomial  $P_1$  defined in (36) does not have constant roots, except when  $\lambda \neq 0$  and  $\alpha = \mu_j = 0$  for any  $j$ . In this case,*

$$t := \frac{(m+2)c}{m+3} + \lambda$$

*is the unique constant root of  $P_1$  and is simple.*

- (ii) *Suppose that  $\hat{q}_1(c) = 0$ . Then*

$$t := \frac{(m+2)c}{m+3}$$

*is a simple root of  $P_1$ . The polynomial  $P_1$  has other constant roots if and only if  $\lambda \neq 0$  and  $\alpha = \mu_j = 0$  for any  $j$ . In this case, there is only one additional constant root of  $P_1$ , namely*

$$t := \lambda + \frac{(m+2)c}{m+3},$$

*which is simple.*

*Proof.* We first consider the case when  $\hat{q}_1(c) \neq 0$ . We claim that  $P_1$  has no multiple roots. Suppose, on the contrary, that  $t$  is a multiple (necessarily constant, because the non-constant eigenvalues of the Bryant modified Ricci operator are always simple) root of  $P_1$ . Since  $\hat{q}_1(c) \neq 0$ ,  $t$  cannot be equal to  $(m+2)c/(m+3)$  and so

$$\hat{q}_1\left(t + \frac{c}{m+3}\right) \neq 0$$

(because  $\hat{p}_r$  has no constant roots). The equalities  $P_1(t) = P'_1(t) = 0$  imply that

$$t_1 \hat{p}'_r\left(t + \frac{c}{m+3}\right) + t_2 \hat{p}_r\left(t + \frac{c}{m+3}\right) = 0, \tag{42}$$

where

$$t_1 := t - \frac{(m+2)c}{m+3} \in \mathbb{R} \setminus \{0\} \quad \text{and} \quad t_2 := -\frac{(m+2)c}{m+3} - \left(t - \frac{(m+2)c}{m+3}\right) \frac{\hat{q}'_1(t + c/(m+3))}{\hat{q}_1(t + c/(m+3))} \in \mathbb{R}.$$

But (42) cannot hold: if it did, it would imply that  $I, \hat{B}_r, \dots, \hat{B}_r^{l+1}$  were dependent, which contradicts the fact that the minimal polynomial of  $\hat{B}_r$  has degree  $l+2$ . We conclude that  $P_1$  cannot have multiple roots. We will now show that the only possible constant root of  $P_1$  is  $\lambda + (m+2)c/(m+3)$  and it is a root if and only if  $\lambda \neq 0$  and  $\alpha = \mu_j = 0$  for any  $j$ . For this, let  $t$  be a constant root

of  $P_1$ . Taking the derivative with respect to  $r$  of the equality  $P_1(t) = 0$  and using Lemma 16, we get

$$\left(t - \frac{(m+2)c}{m+3} - \lambda\right) \hat{q}_1\left(t + \frac{c}{m+3}\right) = 0,$$

from where we deduce that  $t = \lambda + (m+2)c/(m+3)$ , since

$$\hat{q}_1\left(t + \frac{c}{m+3}\right) \neq 0.$$

Moreover,  $P_1(t) = 0$  if and only if

$$\lambda \hat{p}_r(\gamma) + \frac{f}{r^2} \hat{q}_1(\gamma) = 0. \tag{43}$$

Equality (43) forces  $\lambda \neq 0$  (if  $\lambda = 0$ , then, from (43),  $\hat{q}_1(\gamma) = 0$ ; also,  $c = \gamma$ ; recall however that we are under the hypothesis  $\hat{q}_1(c) \neq 0$ ; we obtain a contradiction). Therefore,  $\lambda \neq 0$  and then  $A_r = r^4 e^{\lambda r^2} A$ . Relation (43) is equivalent with

$$\lambda \tilde{\hat{b}}_r(\gamma) w, w + \hat{q}_1(\gamma) e^{\lambda r^2} (Aw, w) = 0,$$

for any  $w \in W$  null, where  $\tilde{\hat{b}}_r$  denotes the reduced adjoint operator of  $\hat{B}_r$ . With the notation of Remark 2,  $\tilde{\hat{b}}_r(\gamma)$ , as well as  $A$ , act trivially on the subspaces  $W_j$  (for  $j \geq 1$ ) of  $W$  (here and below the reader is referred to [DG06, Lemma 3], which describes the action of the reduced adjoint operator of a  $k$ -step parabolic hermitian operator when applied to the parabolic eigenvalue). It follows that (43) is equivalent with

$$\lambda \tilde{\hat{b}}_r(\gamma) + \hat{q}_1(\gamma) e^{\lambda r^2} A = 0. \tag{44}$$

We claim that equality (44) holds if and only if  $\alpha = \mu_j = 0$  for any  $j$  (and  $\lambda \neq 0$ ). Notice first that if (44) holds then  $\hat{q}_1(\gamma) \neq 0$  (since  $\tilde{\hat{b}}_r(\gamma) \neq 0$ ), which implies that  $\hat{B}_r$  is one-step parabolic. We deduce that  $\mu_j = 0$  for any  $j$ . With the notation of Remark 2,  $\tilde{\hat{b}}_r(\gamma)$ , as well as  $A$ , act trivially on  $\widehat{W}_0$ ; on  $\text{Span}\{e_0, e_1\}$ ,  $\tilde{\hat{b}}_r(\gamma)$  acts by

$$\lambda \left(\alpha - \frac{e^{\lambda r^2}}{\lambda}\right) \prod_{j=1}^s (\gamma - \beta_j) A.$$

Relation (44) is equivalent to

$$(\alpha \lambda - e^{\lambda r^2}) \prod_{j=1}^s (\gamma - \beta_j) + \hat{q}_1(\gamma) e^{\lambda r^2} = 0;$$

since  $\hat{q}_1(\gamma) = \prod_{j=1}^s (\gamma - \beta_j)$  and  $\lambda \neq 0$ , it reduces to  $\alpha = 0$ . Since  $P_1$  does not have multiple roots,

$$\lambda + \frac{(m+2)c}{m+3}$$

is the (unique) simple root of  $P_1$ .

Now we consider the case when  $\hat{q}_1(c) = 0$ . Clearly,  $(m+2)c/(m+3)$  is a root in this case. Define the polynomial  $\hat{q}_2(t) := \hat{q}_1(t)/(t - c)$  and let  $t$  be a constant root of

$$\hat{p}_r\left(t + \frac{c}{m+3}\right) + \frac{f}{r^2} \hat{q}_2\left(t + \frac{c}{m+3}\right) = 0. \tag{45}$$

Taking the derivative with respect to  $r$  of (45) and using Lemma 16, we get

$$\left(t - \frac{(m+2)c}{m+3} - \lambda\right) \hat{q}_2\left(t + \frac{c}{m+3}\right) = 0.$$

This implies that

$$t = \lambda + \frac{(m + 2)c}{m + 3},$$

because

$$\hat{q}_2\left(t + \frac{c}{m + 3}\right) \neq 0$$

(because  $\hat{p}_r$  has no constant roots). As before,  $t$  is a root of (45) if and only if

$$\tilde{b}_r(\gamma) + \hat{q}_2(\gamma)e^{\lambda r^2}A = 0. \tag{46}$$

Notice first that if (46) holds, then  $\hat{q}_2(\gamma) \neq 0$ . Next, we prove that if (46) holds, then  $\hat{B}_r$  is one-step parabolic. The argument is the following. Suppose, on the contrary, that (46) holds and that  $\hat{B}_r$  is two-step parabolic; then  $\lambda = 0$ ; otherwise, since  $(t - \gamma)$  divides  $\hat{q}_1$  ( $\hat{B}_r$  being two-step parabolic),  $\hat{q}_2(\gamma) = 0$ , which is impossible. On the other hand,  $\tilde{b}_r(\gamma)$  acts as  $\sum_{k=2}^n |\mu_k|^2 \prod_{j=1}^s (\gamma - \beta_j)A$  on  $\text{Span}\{e_0, e_1\}$ , when  $\hat{B}_r$  is two-step parabolic; also  $\hat{q}_2(\gamma) = \prod_{j=1}^s (\gamma - \beta_j)$ , when  $\lambda = 0$ ; from (46) it follows that

$$\left(\sum_{k=2}^n |\mu_k|^2 + 1\right) \prod_{j=1}^s (\gamma - \beta_j)A = 0$$

on  $\text{Span}\{e_0, e_1\}$ , which cannot hold. We have proved that if  $\lambda + (m + 2)c/(m + 3)$  is a root, then  $\hat{B}_r$  is one-step parabolic. Moreover, in this case  $\lambda \neq 0$  (if  $\lambda = 0$  then  $c = \gamma$  and since  $\hat{q}_1(c) = 0$ , then  $\hat{q}_1(\gamma) = 0$  which is absurd because  $(t - \gamma)^3$  does not divide the minimal polynomial  $\hat{q}$  of  $\hat{B}_r$  when  $\hat{B}_r$  is one-step parabolic). Finally, when  $\hat{B}_r$  is one-step parabolic and  $\lambda \neq 0$ , relation (46) becomes

$$\left(\alpha - \frac{e^{\lambda r^2}}{\lambda}\right) \prod_{j=1}^s (\gamma - \beta_j) + \hat{q}_2(\gamma)e^{\lambda r^2} = 0,$$

which holds if and only if  $\alpha = 0$ , because

$$\hat{q}_2(\gamma) = \frac{1}{\gamma - c}\hat{q}_1(\gamma) = \frac{1}{\lambda}\hat{q}_1(\gamma) = \frac{1}{\lambda} \prod_{j=1}^s (\gamma - \beta_j).$$

We have proved that  $P_1$  has an additional constant root, besides  $(m + 2)c/(m + 3)$ , if and only if  $\hat{B}_r$  is one-step parabolic (i.e.  $\mu_j = 0$  for any  $j$ ),  $\lambda \neq 0$  and  $\alpha = 0$ . The additional constant root is

$$\lambda + \frac{(m + 2)c}{m + 3}.$$

It is easy to see that it is simple. □

In Proposition 19 we shall determine the Bryant minimal and characteristic polynomials of  $(M, \omega, J)$ . For the proof of this proposition we need the following additional lemma.

LEMMA 18. For any  $t \in \mathbb{R}$ ,  $(x, r) \in M$  and  $w \in x$  non-zero,

$$\begin{aligned} g_{(x,r)}(d^H \hat{p}_r(t), d^H \hat{p}_r(t)) &= 4\left(\hat{q}'(t)\hat{p}_{r,x}(t) - \hat{q}(t)\hat{p}'_{r,x}(t) - 2t\hat{p}_{r,x}^2(t) + \hat{p}_{r,x}^2(t)\frac{(B_r^2 w, w)}{r^2(B_r w, w)}\right), \\ g_{(x,r)}\left(d^H\left(\frac{f}{r^2}\right), d^H\left(\frac{f}{r^2}\right)\right) &= \frac{4f^2}{r^6}\left(\frac{(B_r^2 w, w)}{(B_r w, w)} - 2r^2\gamma\right), \\ g_{(x,r)}\left(d^H\left(\frac{f}{r^2}\right), d^H \hat{p}_r(t)\right) &= \frac{4f}{r^2}\left((t - \gamma)\hat{q}_1(t) - (t + \gamma)\hat{p}_{r,x}(t) + \frac{\hat{p}_{r,x}(t)(B_r^2 w, w)}{r^2(B_r w, w)}\right). \end{aligned}$$

*Proof.* The first equality follows from Lemma 3, applied to  $\hat{B}_r$ . To prove the second equality, we remark that the 1-form  $d^H(f/r^2)$  corresponds to the vector field  $(1/r^4)v$  by means of the metric  $g$ . Therefore,

$$g\left(d^H\left(\frac{f}{r^2}\right), d^H\left(\frac{f}{r^2}\right)\right) = \frac{1}{r^6}g_r(v, v).$$

The second equality of the lemma follows from (34). To prove the third equality we notice that

$$g\left(d^H\left(\frac{f}{r^2}\right), d^H\hat{p}_r(t)\right) = \frac{\hat{L}_t(f)}{r^2},$$

where  $\hat{L}_t$  is the vector field on  $M$  which, at a point  $(x, r) \in M$ , belongs to  $H_x = \text{Hom}_{\mathbb{C}}(x, W/x)$  and is the endomorphism

$$(\hat{L}_t)_{(x,r)}(w) = 2(\tilde{b}_r(t)w - \hat{p}_r(t)\hat{B}_r w) \pmod w, \quad w \in x.$$

This is true since  $\hat{L}_t$  corresponds to the 1-form  $d^H\hat{p}_r(t)$  by means of the metric  $g$  (which coincides with  $\hat{g}_r$  on the bundle  $H$ , restricted to a level set  $N_r$ ). It is straightforward to check that

$$\begin{aligned} (A_r\hat{L}_t w, w) &= 2((t - \gamma)\hat{q}_1(t) - \gamma\hat{p}_{r,x}(t))(A_r w, w), \\ (B_r\hat{L}_t w, w) &= 2\hat{p}_{r,x}(t)\left(t - \frac{(B_r^2 w, w)}{r^2(B_r w, w)}\right)(B_r w, w), \end{aligned}$$

so that

$$\hat{L}_t(f)_{(x,r)} = 4f\left((t - \gamma)\hat{q}_1(t) - (t + \gamma)\hat{p}_{r,x}(t) + \frac{\hat{p}_{r,x}(t)(B_r^2 w, w)}{r^2(B_r w, w)}\right),$$

which proves the third equality. □

Proposition 13 is now an easy consequence of the following proposition.

PROPOSITION 19. *The Bryant characteristic polynomial of  $(M, g, J)$  is equal to*

$$p_c(t) = \left(t - \frac{(m+2)c}{m+3}\right)\left(t + \frac{c}{m+3} - \gamma\right)^{n+1} \prod_{j=1}^s \left(t + \frac{c}{m+3} - \beta_j\right)^{n_j}.$$

The Bryant minimal polynomial  $p_m$  of  $(M, g, J)$  has the following expression:

(i) if there is  $\mu_k \neq 0$  and  $c$  is different from  $\beta_j$  (for any  $j$ ) and  $\gamma$ , then

$$p_m(t) = \left(t - \frac{(m+2)c}{m+3}\right)\left(t + \frac{c}{m+3} - \gamma\right)^3 \prod_{j=1}^s \left(t + \frac{c}{m+3} - \beta_j\right);$$

(ii) if there is  $\mu_k \neq 0$  and  $c$  is equal to  $\beta_{j_0}$  (for a certain  $j_0$ ) or to  $\gamma$ , then

$$p_m(t) = \left(t + \frac{c}{m+3} - \gamma\right)^3 \prod_{j=1}^s \left(t + \frac{c}{m+3} - \beta_j\right);$$

(iii) if all  $\mu_k = 0$  and  $c$  is different from  $\beta_j$  (for any  $j$ ), then

$$p_m(t) = \left(t - \frac{(m+2)c}{m+3}\right)\left(t + \frac{c}{m+3} - \gamma\right)^2 \prod_{j=1}^s \left(t + \frac{c}{m+3} - \beta_j\right),$$

except when  $\lambda \neq 0$  and  $\alpha = 0$ , in which case

$$p_m(t) = \left(t - \frac{(m+2)c}{m+3}\right)\left(t + \frac{c}{m+3} - \gamma\right) \prod_{j=1}^s \left(t + \frac{c}{m+3} - \beta_j\right);$$

(iv) if all  $\mu_k = 0$  and  $c$  is equal to  $\beta_{j_0}$  (for a certain  $j_0$ ), then

$$p_m(t) = \left(t + \frac{c}{m+3} - \gamma\right)^2 \prod_{j=1}^s \left(t + \frac{c}{m+3} - \beta_j\right),$$

except when  $\alpha = 0$  (and  $\lambda \neq 0$ ), when

$$p_m(t) = \left(t + \frac{c}{m+3} - \gamma\right) \prod_{j=1}^s \left(t + \frac{c}{m+3} - \beta_j\right).$$

*Proof.* Let  $t$  be a non-constant root of the polynomial  $P_1$ . Then

$$\begin{aligned} \left(t - \frac{(m+2)c}{m+3}\right) \hat{p}_r \left(t + \frac{c}{m+3}\right) &= -\frac{f}{r^2} \hat{q}_1 \left(t + \frac{c}{m+3}\right), \\ \left(t - \frac{(m+2)c}{m+3}\right)^2 \hat{p}'_r \left(t + \frac{c}{m+3}\right) &= \left(t - \frac{(m+2)c}{m+3}\right) P'_1(t) + \frac{f}{r^2} \hat{q}_1 \left(t + \frac{c}{m+3}\right) \\ &\quad - \frac{f}{r^2} \left(t - \frac{(m+2)c}{m+3}\right) \hat{q}'_1 \left(t + \frac{c}{m+3}\right). \end{aligned}$$

Using these relations and Lemma 18, we can calculate the square norm of the covector  $(d^H P_1)(t)$  as follows:

$$\begin{aligned} g((d^H P_1)(t), (d^H P_1)(t)) &= -4\hat{q} \left(t + \frac{c}{m+3}\right) \left(t - \frac{(m+2)c}{m+3}\right) P'_1(t) \\ &\quad - \frac{4f}{r^2} \left(t - \frac{(m+2)c}{m+3}\right) \hat{q}' \left(t + \frac{c}{m+3}\right) \hat{q}_1 \left(t + \frac{c}{m+3}\right) \\ &\quad - \frac{4f}{r^2} \hat{q} \left(t + \frac{c}{m+3}\right) \hat{q}_1 \left(t + \frac{c}{m+3}\right) \\ &\quad + \frac{4f}{r^2} \left(t - \frac{(m+2)c}{m+3}\right) \hat{q} \left(t + \frac{c}{m+3}\right) \hat{q}'_1 \left(t + \frac{c}{m+3}\right) \\ &\quad + \frac{8f}{r^2} \left(t - \frac{(m+2)c}{m+3}\right) \left(t + \frac{c}{m+3} - \gamma\right) \hat{q}_1^2 \left(t + \frac{c}{m+3}\right). \end{aligned}$$

Since  $\hat{q}'(t)\hat{q}_1(t) - \hat{q}(t)\hat{q}'_1(t) = 2(t - \gamma)\hat{q}_1^2(t)$ , the above expression reduces to

$$\begin{aligned} g((d^H P_1)(t), (d^H P_1)(t)) &= -4\hat{q} \left(t + \frac{c}{m+3}\right) \left(t - \frac{(m+2)c}{m+3}\right) P'_1(t) \\ &\quad - \frac{4f}{r^2} \hat{q}_1^2 \left(t + \frac{c}{m+3}\right) \left(t - \gamma + \frac{c}{m+3}\right)^2. \end{aligned}$$

On the other hand, Lemma 16 together with the definition of  $P_1$  imply that

$$\left(\frac{d}{dr} P_1\right)(t) = -\frac{2f}{r} \hat{q}_1 \left(t + \frac{c}{m+3}\right) \left(t - \frac{(m+2)c}{(m+3)} - \lambda\right).$$

Since  $g(dr, dr) = 1/f$ , we obtain

$$\begin{aligned} g((dP_1)(t), (dP_1)(t)) &= g((d^H P_1)(t), (d^H P_1)(t)) + \left(\frac{d}{dr} P_1(t)\right)^2 g(dr, dr) \\ &= -4\hat{q} \left(t + \frac{c}{m+3}\right) \left(t - \frac{(m+2)c}{m+3}\right) P'_1(t). \end{aligned}$$

On the other hand, since  $P'_1(t) dt + d(P_1(t)) = 0$ , we get

$$g(dt, dt) = \frac{g((dP_1)(t), (dP_1)(t))}{P'_1(t)^2}. \tag{47}$$



We distinguish three cases: (i)  $P_1$  has no constant roots; (ii)  $P_1$  has a unique constant root, which is simple and equal to  $t_1 = \lambda + (m + 2)c/(m + 3)$ ; and (iii)  $P_1$  has two constant roots,  $t_1$  and  $t_2 = (m + 2)c/(m + 3)$ , which are simple and distinct (see Proposition 17). In all cases, the Bryant characteristic polynomial of  $(M, \omega, J)$  is

$$p_c(t) = \left(t - \frac{(m + 2)c}{m + 3}\right) \hat{Q}\left(t + \frac{c}{m + 3}\right).$$

In case (i), the Bryant minimal polynomial of  $(M, g, J)$  is

$$p_m(t) = \hat{q}\left(t + \frac{c}{m + 3}\right) \left(t - \frac{(m + 2)c}{m + 3}\right).$$

In case (ii) it is

$$p_m(t) = (t - t_1)^{-1} \left(t - \frac{(m + 2)c}{m + 3}\right) \hat{q}\left(t + \frac{c}{m + 3}\right)$$

and in case (iii) it is

$$p_m(t) = (t - t_1)^{-1} (t - t_2)^{-1} \left(t - \frac{(m + 2)c}{m + 3}\right) \hat{q}\left(t + \frac{c}{m + 3}\right).$$

The conclusion follows. □

### 5. Examples

We end the paper by considering some important classes of Bochner-flat Kähler manifolds and by showing how they can be realised locally as generalised Kähler cones.

(i) *Bryant Bochner-flat Kähler structures.* Let  $N = S^{2m+1} \subset \mathbb{C}^{m+1}$  with its standard CR structure and  $(k_1, \dots, k_{m+1})$  a system of non-negative real numbers. Define, for every  $r > 0$ , the vector field

$$T_r(z) := \sum_{j=1}^{m+1} (1 + k_j r^2) \left(x_j \frac{\partial}{\partial y_j} - y_j \frac{\partial}{\partial x_j}\right),$$

which is the Reeb vector field of a Sasaki structure on  $S^{2m+1} \subset \mathbb{C}^{m+1}$ . Here  $z = (z_1, \dots, z_{m+1})$  belongs to  $S^{2m+1}$ ,  $z_j = x_j + iy_j$  for any  $j \in \{1, \dots, m + 1\}$  and  $r^2 = |z_1|^2 + \dots + |z_{m+1}|^2$ . The family of Sasaki Reeb vector fields  $\{T_r, r > 0\}$  defines a Bochner-flat generalised Kähler cone on  $\mathbb{C}^{m+1} \setminus \{0\}$ , which belongs to the first case of Theorem 11; the solution of equation (19) is  $\mu(t) = -2/t$  and the hermitian operator  $B$  is semi-simple, with eigenvalues

$$k'_j = k_j - \frac{1}{m + 2} \sum_{i=1}^{m+1} k_i, \quad \text{for } j \in \{1, \dots, m + 1\}.$$

This Bochner-flat Kähler structure was discovered in [Bry01] and further studied in [DG06]; it can be extended as a complete Bochner-flat Kähler structure on  $\mathbb{C}^{m+1}$ .

(ii) *Bochner-flat Kähler–Einstein structures.* Let  $(M, g, J)$  be a Bochner-flat generalised Kähler cone. With the notation of Theorem 11, suppose that  $B = eA$ , for  $e \in \mathbb{R}$ . If  $(M, g, J)$  belongs to cases (i) and (ii) of Theorem 11, then it is Kähler–Einstein if and only if  $e^2 + 2d = 0$ ; moreover, the Bryant modified Ricci operator of  $(M, \omega, J)$  is

$$\Theta = \frac{e}{m + 3} \text{Id}$$

in case (i) and

$$\Theta = -\frac{e}{m+3}\text{Id}$$

in case (ii). If  $(M, g, J)$  belongs to the case (iii) of Theorem 11 and  $B = eA$  then it is never Einstein; if it belongs to case (iv) of Theorem 11 then it is Kähler–Einstein if and only if  $e = 0$  (and  $\lambda < 0$ ); the Bryant modified Ricci operator is

$$\Theta = \frac{\lambda}{m+3}\text{Id}$$

in this case.

(iii) *Bochner-flat generalised Kähler cones of order one.* If  $B = eA$  but  $(M, g, J)$  is not Einstein, then it must have order one. The Bryant minimal and characteristic polynomials have the following expressions. If  $(M, g, J)$  belongs to case (i), respectively case (ii) of Theorem 11, then

$$p_c(t) = \left(t \mp \frac{e}{m+3}\right)^{m+1} \left( \left(t \mp \frac{e}{m+3}\right)^2 \pm e \left(t \mp \frac{e}{m+3}\right) + \frac{e^2 + 2d}{4} \right),$$

$$p_m(t) = \left(t \mp \frac{e}{m+3}\right) \left( \left(t \mp \frac{e}{m+3}\right)^2 \pm e \left(t \mp \frac{e}{m+3}\right) + \frac{e^2 + 2d}{4} \right).$$

Moreover,  $(M, g, J)$  is of hyperbolic type when  $d > 0$ , of one-step parabolic type when  $d = 0$  and of elliptic type when  $d < 0$ . If  $(M, J, g)$  belongs to case (iii) and case (iv) of Theorem 11, then

$$p_c(t) = \left(t + \frac{(m+2)\lambda}{m+3}\right) \left(t - \frac{\lambda}{m+3}\right)^{m+2},$$

$$p_m(t) = \left(t + \frac{(m+2)\lambda}{m+3}\right) \left(t - \frac{\lambda}{m+3}\right)^2.$$

(Recall that  $\lambda = 0$  in case (iii) and  $\lambda \neq 0$  in case (iv).) When  $(M, \omega, J)$  belongs to case (iii), it is one-step parabolic; when it belongs to case (iv), it is two-step parabolic. Bochner-flat Kähler structures of order one have been studied in [ACG06]. As shown in [ACG06, DG06], they fibre over a Kähler manifold with constant holomorphic sectional curvature (in our formalism, the fibration is  $M \rightarrow N_r/\tilde{T}_r$ ).

(iv) *Weighted projective spaces as generalised Kähler cones.* Let  $\mathbb{P}_a^{m+1}$  be a weighted projective space, of weights  $(a_1, \dots, a_{m+1})$ , where  $a_j$  are positive integers. As shown in [Bry01, DG06],  $\mathbb{P}_a^{m+1}$  has a canonical Bochner-flat Kähler structure, of semi-simple type, isomorphic with  $M_C$ , where  $C$  is a hermitian semi-simple operator of  $\mathbb{C}^{m+2,1}$ , with eigenvalues  $-\sum_{j=1}^{m+2} \lambda_j, \lambda_1, \dots, \lambda_{m+2}$ , where  $\lambda_j$  are related to the weights  $a_j$  by

$$\lambda_j = a_j - \frac{1}{m+3} \sum_{i=1}^{m+2} a_i,$$

for any  $j \in \{1, \dots, m+2\}$ . As a Bochner-flat generalised Kähler cone,  $\mathbb{P}_a^{m+1}$  belongs to case (i) of Theorem 11;  $\mu$  is any solution of equation (19), with  $d = 2a_{m+2}^2/(m+3)^2$ , and the hermitian operator  $B$  is semi-simple, with eigenvalues

$$-\frac{\sum_{j=1}^{m+1} a_j}{m+2}, \quad a_1 - \frac{\sum_{j=1}^{m+1} a_j}{m+2}, \quad \dots, \quad a_{m+1} - \frac{\sum_{j=1}^{m+1} a_j}{m+2}.$$

(v) *Tachibana and Liu Bochner-flat generalised Kähler cones.* Consider a Bochner-flat generalised Kähler cone structure  $(g, J)$  which belongs to case (i) of Theorem 11. With the notation of § 4, assume that  $k_1 = \dots = k_{m+1} := \bar{k}$ . On the set  $D \subset \mathbb{C}^{m+1}$  defined by (24), the Kähler structure

$(g, J)$  has a Kähler potential  $x$  which depends only on  $r^2 := |z_1|^2 + \cdots + |z_{m+1}|^2$ . As a function of  $r^2 = t$ ,  $x$  satisfies the implicit equation

$$e^{ax(t)} \dot{\mu}(x(t))^{-1/2} = t, \tag{48}$$

where  $a := (m + 2)\bar{k}$ . In general, a Kähler structure which is defined on an open subset of the standard  $\mathbb{C}^m$  and has a global Kähler potential, say  $h$ , which depends only on  $r^2$  is Bochner-flat if and only if  $h$ , as a function of  $r^2 = t$ , satisfies a differential equation of the form [TL70]

$$\ddot{h}(t) = \lambda_1 t \dot{h}^3(t) + \lambda_2 \dot{h}^2(t), \tag{49}$$

where  $\lambda_1, \lambda_2 \in \mathbb{R}$ . It can be easily verified that, if  $x$  satisfies (48), then it also satisfies (49), with  $\lambda_1 := a^2 + d/2$  and  $\lambda_2 := -2a$ .

ACKNOWLEDGEMENTS

I warmly thank Paul Gauduchon and David Calderbank for encouraging me to study Bochner-flat generalised Kähler cones and for many useful comments on a first version of this paper. I thank the anonymous referee for his detailed report and his deep observations and suggestions.

REFERENCES

ACG04 V. Apostolov, D. Calderbank and P. Gauduchon, *Hamiltonian 2-forms in Kähler geometry two, Global classification*, J. Differential Geom. **68** (2004), 277–345.

ACG06 V. Apostolov, D. Calderbank and P. Gauduchon, *Hamiltonian 2-forms in Kähler geometry one*, J. Differential Geom. **73** (2006), 359–412.

Bry01 R. L. Bryant, *Bochner–Kähler manifolds*, J. Amer. Math. Soc. **14** (2001), 623–715.

BS76 D. Burns and S. Shnider, *Spherical hypersurfaces in complex manifolds*, Invent. Math. **33** (1976), 223–246.

Cal02 D. M. J. Calderbank, Featured review of [Bry01], *MathSciNet* MR1824987(2002i:53096).

CM74 S. S. Chern and J. Moser, *Real hyper-surfaces in complex manifolds*, Acta Math. **133** (1974), 219–271.

Dav04 L. David, *Weyl connections and curvature properties of CR manifolds*, Ann. Global Anal. Geom. **26** (2004), 59–72.

DG06 L. David and P. Gauduchon, *The Bochner-flat geometry of weighted projective spaces*, CRM Proceedings and Lecture Notes, vol. 40 (American Mathematical Society, Providence, RI, 2006), 109–156.

Gau01 P. Gauduchon, *The Bochner tensor of a weakly Bochner-flat Kähler manifold*, Anal. Univ. Timisoara **39** (2001), 189–137.

GO98 P. Gauduchon and L. Ornea, *Locally conformally Kähler metrics on Hopf surfaces*, Ann. Inst. Fourier (Grenoble) **48** (1998), 1107–1127.

TL70 S. Tachibana and R. Liu, *Notes on Kählerian metrics with vanishing Bochner curvature tensor*, Kodai Math. Sem. Rep. **22** (1970), 3131–321.

Tan75 N. Tanaka, *A differential geometric study on strongly pseudo-convex manifolds*, Lectures in Mathematics, vol. 9 (Kyoto Univ., Kinokuniya, Tokyo, 1975).

Web77 S. M. Webster, *On the pseudo-conformal geometry of a Kähler manifold*, Math. Z. **157** (1977), 265–270.

Liana David liana.david@imar.ro; l.david@sns.it

Institute of Mathematics of the Romanian Academy ‘Simion Stoilow’, Calea Grivitei nr. 21, Sector 1, 010702 Bucharest, Romania

*Current address:* Centro di Ricerca Matematica ‘Ennio de Giorgi’, Scuola Normale Superiore, Collegio Puteano Piazza dei Cavalieri nr. 3, I-56100 Pisa, Italy