

symptoms were similar in SARS-CoV-2 versus influenza patients. Vigilance should be undertaken in treatment of children presenting with all respiratory illnesses.

241

### Utilizing real-world evidence to increase efficiency of randomized controlled trials with application to repurposed therapeutics for COVID-19

Lillian M F Haine<sup>1</sup>, Thomas A Murray<sup>1</sup> and Joseph S Koopmeiners<sup>1</sup>  
<sup>1</sup>University of Minnesota

**OBJECTIVES/GOALS:** We aim to extend a novel statistical method called the Semi-Supervised Mixture Multisource Exchangeability Model (SS-MIX-MEM) and to implement the SS-MIX-MEM approach to supplement ALPS-COVID data with N3C data to achieve analyses with greater precision and actionable conclusions. **METHODS/STUDY POPULATION:** We will apply the SS-MIX-MEM to supplement the Angiotensin receptor blocker-based Lung Protective Strategy for COVID-19 (ALPS-COVID) RCTs with the National COVID Cohort Collaborative (N3C) database. ALPS-COVID includes both an inpatient and outpatient trial, which investigate losartan as a treatment for COVID-19. The outpatient trial sought to randomize 580 individuals but only enrolled 117, whereas the inpatient trial met its enrollment target and randomized 205 individuals. The N3C database has 3,237,344 COVID-19 cases alongside demographics, lab values, and more. **RESULTS/ANTICIPATED RESULTS:** In simulation studies, the proposed SS-MIX-MEM approach effectively leveraged a subgroup of supplemental real world data for RCT analyses, improving trial efficiency by increasing precision of treatment effect estimates, decreasing necessary sample size, and introducing minimal bias. In an influenza trial real world data application, the SS-MIX-MEM approach was able to effectively provide insight into treatment effect heterogeneity found in an RCT analogous to incorporating around 80 individuals into a subgroup analysis. We anticipate that leveraging external real world data in a re-analysis of the ALPS-COVID RCTs could provide new insights into losartan, a readily available, potentially beneficial therapeutic for COVID-19. **DISCUSSION/SIGNIFICANCE:** The high blood pressure drug, losartan, is readily available, has an established safety profile, and might be effective as a treatment for COVID-19. Given that we have very few effective treatment options and are still in the midst of a global pandemic, patients with COVID-19 would greatly benefit from a repurposed, readily available treatment.

242

### Characterization of Maternal Stress During Pregnancy

Satvinder K. Dhaliwal<sup>1</sup>, Greta Wilkening<sup>2</sup>, Angela Lee-Winn<sup>1</sup>, Deborah Glueck<sup>1</sup>, Dana Dabelea<sup>1</sup> and Wei Perng<sup>1</sup>

<sup>1</sup>University of Colorado, Denver and <sup>2</sup>Colorado Children's Hospital

**OBJECTIVES/GOALS:** 1. To characterize domains of maternal psychosocial stress from the Edinburgh Postnatal Depression Scale (EPDS) and Cohens Perceived Stress Scale (PSS) administered during pregnancy using principal components analysis (PCA). 2. To identify sociodemographic, perinatal, and lifestyle correlates of maternal psychosocial stress domains. **METHODS/STUDY POPULATION:** Using data from 1,079 pregnant women in the Healthy Start Study who completed both the EPDS and PSS in early pregnancy, we ran PCA and retained factors representative of

uncorrelated domains of maternal psychosocial stress based on the Scree plot and Eigenvalues >1. We then used linear regression to identify sociodemographic, perinatal, and lifestyle correlates of each maternal stress domain, followed by multivariable models that mutually adjusted for all characteristics that were statistically significant at alpha = 0.10. **RESULTS/ANTICIPATED RESULTS:** We identified three domains of maternal psychosocial stress based on PCA results: Feeling Overwhelmed (Domain 1), Anhedonia (Domain 2), and Lack of Control (Domain 3). In unadjusted analyses, lower household income and poor diet quality were associated with higher scores for all three domains. In adjusted analyses, lower household income, being multiparous, inadequate or excessive GWG, and poor diet quality were associated with Feeling Overwhelmed. Older age, Hispanic ethnicity, and poor diet quality were associated with Anhedonia. Non-Hispanic, Black race/ethnicity, lower educational attainment, having a partner born outside the US, larger household size, receiving public assistance, and smoking during pregnancy were associated with Lack of Control. **DISCUSSION/SIGNIFICANCE:** We identify three unique domains of maternal psychosocial stress that are differentially related to socio-demographic, perinatal, and lifestyle characteristics. Correlates of stress domains shed light on upstream determinants and biological and psychosocial mechanisms through which experiences of stress manifest.

244

### A Systematic Review of Epileptiform Changes During Sevoflurane Anesthesia In Infants and Children

Luciana Gizzo<sup>1</sup>, Daria Ivenitsky<sup>1</sup>, Alexander Ferrera<sup>2</sup>, Matthew Tam<sup>2</sup>, Alan D. Legatt<sup>2,5</sup>, Elissa G. Yozawitz<sup>2,3,6</sup>, Yungtai Lo<sup>2,7</sup>, Guohua Li<sup>8</sup>, Shlomo Shinnar<sup>2,3,6,7</sup> and Jerry Chao<sup>2,9</sup>

<sup>1</sup>The University of New England College of Osteopathic Medicine, <sup>2</sup>Albert Einstein College of Medicine, Montefiore Medical Center, <sup>3</sup>The Saul R. Korey Department of Neurology, <sup>4</sup>Dominick P. Purpura Department of Neuroscience, <sup>5</sup>Department of Medicine (Critical Care), <sup>6</sup>Department of Pediatrics, <sup>7</sup>Department of Epidemiology and Population Health, <sup>8</sup>Departments of Anesthesiology and Epidemiology, Columbia University Mailman School of Public Health and <sup>9</sup>Department of Anesthesiology

**OBJECTIVES/GOALS:** Early clinical case reports have described incidental epileptiform changes during electrophysiological monitoring. The objective of this study was to perform a systematic review of all existing investigations of epileptiform activity during sevoflurane use in pediatric anesthesia. The heterogenous EEG data will be analyzed in a meta analysis **METHODS/STUDY POPULATION:** A targeted, PICO-based clinical question was crafted and registered a priori on PROSPERO on 3/19/21. Under the guidance of a librarian from the Albert Einstein College of Medicine, a boolean search string was generated to search articles and gray literature for terms such as pediatric, sevoflurane and electroencephalogram in PubMed, OVID, Cochrane, Google Scholar, etc. We utilized the software platform tool COVIDENCE to manage our review. 495 references were imported for initial screening. 56 English-language, full-text studies were included for further review. The final 13 references were included in data extraction and Newcastle-Ottawa bias assessment. The characteristics of the studies and their primary outcomes were collected in tabular form. Strategies for data synthesis were discussed weekly. **RESULTS/ANTICIPATED RESULTS:** Epileptiform changes reported in the literature during pediatric sevoflurane

anesthesia ranged from 0 - 95%. EEG data were acquired using a variety of recording systems with variable number of leads and heterogeneous outcomes reported. The periods of anesthesia monitoring were also heterogeneous. Characteristics of the studies are presented in Table 1. 495 references were imported for screening with 13 final references for data extraction. EEG abnormalities were reported in 204/649 (31.4%) subjects ranging in age from neonate to 18 years; the majority of studies utilized less than 16 channels of (10/13, 76.9%) (Table 1). There was variability in sevoflurane dosing, premedication (e.g., midazolam, hydroxyzine), and periods of anesthesia monitored. DISCUSSION/SIGNIFICANCE: There was heterogeneity noted across reviewed literature including study design, phases of anesthesia, ventilation methods, number of EEG leads recorded and adjuvant anesthetics administered. Nevertheless, this review rigorously classified epileptiform activity during Sevoflurane thereby influencing modern anesthesia.

245

### Likelihood of live birth following fertility preserving treatment among reproductive-age women diagnosed with gynecologic malignancies or pre-malignancies\*

Ruoxi Yu<sup>1</sup>, Mindy S. Christianson<sup>1</sup>, Anna L. Beavis<sup>1</sup>, Rebecca L. Stone<sup>1</sup> and Johns Hopkins<sup>2</sup>

<sup>1</sup>Johns Hopkins Medicine, Department of GYN/OB and <sup>2</sup>University School of Medicine

OBJECTIVES/GOALS: To determine the impact of fertility preserving treatment (FPT) on likelihood of live birth in a cohort of reproductive-age women (18-45 y) after diagnosis of gynecologic malignancy or pre-malignancy METHODS/STUDY POPULATION: We performed a retrospective cohort study of women ages 18-45 seen by gynecologic oncologists for newly diagnosed cervical cancer (CC), endometrial intraepithelial neoplasia (EIN) or endometrial cancer (EC), and borderline ovarian tumor (BOT) or invasive ovarian cancer (OC) at an academic center from 2015-2019, excluding women who completed childbearing. Our primary outcome was live birth after diagnosis and our exposure was FPT defined as services received by reproductive endocrinology and infertility specialists. We performed Pearsons Chi-squared and log binomial regression to assess association between live birth and FPT with adjustment for patient demographic and disease factors. RESULTS/ANTICIPATED RESULTS: Out of 220 women (median age 36 y), most were White (54% vs. 25% Black) and 37% percent were diagnosed with BOT/OC (vs. 35% EIN/EC; 28% CC). After diagnosis of disease, 19% of women (n=41) had documented FPT and 8% of women (n= 17) had a live birth. By the end of follow-up, 6% of women who did not receive FPT had a live birth (n=11/178) compared to 15% of those who did (n=6/40, p=0.12). In univariate regression, women who received FPT were 2.4 times more likely to have a live birth after disease diagnosis than those who did not receive FPT (p-value = 0.06). However, after adjusting for age at diagnosis, relationship status, disease stage and disease type, the association between FPT and live birth was less robust (RR = 1.4, p-value = 0.6). DISCUSSION/SIGNIFICANCE: In this study, a minority of women had FPT or live births. Our data suggest that FPT benefit should be considered in context of age, relationship status, and disease characteristics for reproductive-age women diagnosed with gynecologic malignancies. Given the complexity, women should be offered referral for consultation with a fertility specialist.

248

### Use of a Propensity Score to Examine Association between Rates of In-Hospital Decongestion and Mortality and Cardiovascular Outcomes Among Patients admitted for Acute Heart Failure

Wendy McCallum<sup>1</sup>, Hocine Tighiouart<sup>2</sup>, Jeffrey M. Testani<sup>3</sup>, Matthew Griffin<sup>3</sup>, Marvin A. Konstam<sup>4</sup>, James E. Udelson<sup>4</sup> and Mark J. Sarnak<sup>5</sup>

<sup>1</sup>Tufts Medical Center, <sup>2</sup>Institute for Clinical Research and Health Policy Studies, Tufts Medical Center, Boston, Massachusetts; Tufts Clinical and Translational Science Institute, Tufts University, <sup>3</sup>Division of Cardiovascular Medicine, Yale School of Medicine, New Haven, Connecticut, <sup>4</sup>Division of Cardiology and the CardioVascular Center, Tufts Medical Center, Boston, Massachusetts and <sup>5</sup>Division of Nephrology, Tufts Medical Center, Boston, Massachusetts

OBJECTIVES/GOALS: Decongestion, or fluid removal, is an important goal in the management of acute heart failure (AHF) among patients with heart failure with reduced ejection fraction (HFrEF). We sought to examine whether the rate of decongestion is associated with mortality and cardiovascular (CV) outcomes. METHODS/STUDY POPULATION: Using data from the Efficacy of Vasopressin Antagonism in Heart Failure Outcome Study With Tolvaptan (EVEREST) trial (n=4133), we evaluated the rate of decongestion by using linear mixed models to derive the in-hospital slope of b-type natriuretic peptide (BNP) and hematocrit as proxies of volume overload and hemoconcentration, respectively. A propensity score was developed to match patients from the quartile with most rapid rates of decongestion to the three quartiles with slower rates. Cox proportional hazards regression models were fitted to assess the association between rate of decongestion with risk of all-cause mortality and a composite of CV mortality or AHF hospitalization. RESULTS/ANTICIPATED RESULTS: Slower rates of in-hospital decongestion were associated with increased risk of both outcomes over a median 10-month follow-up. Those with slower rates of BNP decline, in comparison to the propensity-score matched patients with the most rapid rates of BNP decline, had higher hazards of mortality (HR=1.73 [1.23, 2.42]) and the composite outcome (HR=1.48 [1.18, 1.86]). Those with slower rates of hematocrit increase, in comparison to the propensity-score matched patients with the most rapid rates of hematocrit increase, showed a trend toward higher hazard of mortality (HR=1.17 [0.95, 1.43]) and an increased risk of the composite outcome (HR=1.26 [1.08, 1.47]). DISCUSSION/SIGNIFICANCE: Among patients with HFrEF admitted for AHF, slower rates of decongestion are associated with increased risk of mortality, CV mortality and AHF hospitalization. It remains unknown whether more rapid decongestion provides cardiovascular benefit or if it serves as a proxy for less treatment resistant heart failure.

250

### Defining Developmentally High-Risk Full Term and Late Preterm Infants in the Neonatal Intensive Care Unit

Katherine Carlton<sup>1</sup>, Erwin Cabacungan<sup>1</sup> and Susan Cohen<sup>1</sup>

<sup>1</sup>Medical College of Wisconsin

OBJECTIVES/GOALS: We aim to describe the preschool age developmental outcomes of children born full term or late preterm requiring care in the Childrens Wisconsin (CW) neonatal intensive care unit (NICU). Our objective is to develop a model to predict which NICU infants are at high risk for abnormal preschool age