

which proves our claim. In particular, this shows that for the sequence defined in (3),  $\lim_{n \rightarrow \infty} w_n = 1$ , and then for the sequence defined in (2),  $\lim_{n \rightarrow \infty} v_n = -1$ .

Finally, let us observe that the study of the families of recursive sequences

$$\tilde{u}_{n+1}^{\beta,\gamma} = \min \{ \tilde{u}_n^{\beta,\gamma}, \cos(\gamma n + \beta) \}, \quad n \geq 1,$$

$$z_{n+1}^{\beta,\gamma} = \max \{ z_n^{\beta,\gamma}, \sin(\gamma n + \beta) \}, \quad n \geq 1,$$

and

$$z_{n+1}^{\beta,\gamma} = \min \{ z_n^{\beta,\gamma}, \sin(\gamma n + \beta) \}, \quad n \geq 1,$$

reduces to the obvious equalities:

$$\tilde{u}_n^{\beta,\gamma} = -u_n^{\beta + \pi,\gamma}, \quad z_n^{\beta,\gamma} = u_n^{\beta - \frac{1}{2}\pi,\gamma} \quad \text{and} \quad z_n^{\beta,\gamma} = -u_n^{\beta + \frac{1}{2}\pi,\gamma}.$$

*References*

1. J. Haas, M. D. Weir, G. B. Thomas, Jr., *University Calculus: Alternate Edition*, Pearson Education (2008).
2. K.-L. Kueh, A note on Kronecker's approximation theorem, *Amer. Math. Monthly* **93**, no. 7, (1986) pp. 555-556.
3. W. Schmidt, Diophantine Approximations and Diophantine Equations, *Lecture Notes in Mathematics*, Vol. **1467**, Springer (1991).

10.1017/mag.2024.125 © The Authors, 2024

ÁRPÁD BÉNYI

Published by Cambridge University Press

*Department of Mathematics,*

on behalf of

*516 High St, Western Washington University,*

The Mathematical Association

*Bellingham, WA 98225, USA*

*e-mail: benyia@wwu.edu*

## 108.42 On matrices whose elements are integers with given determinant

*Introduction*

For matrices with large positive integer elements with a small determinant is an interesting question in a linear algebra course. In this paper, we investigate matrices of order  $n$  with large positive integer elements and having a small determinant. In [1], the author explains the method for finding an infinite family of square matrices of order 2 with large positive integer entries and small positive integer determinant. Motivated by this fact, we generalise it for the case of square matrices of any arbitrary order  $n \geq 2$ . More precisely, we prove the following result.

*Theorem 1:* Given positive integers  $d$  and  $M$ , there exist infinitely many matrices  $A = [a_{ij}]_{1 \leq i,j \leq n}$  with integer elements satisfying  $a_{ij} \geq M$  and  $\det A = d$ .



For the proof, first we consider matrices of order 3. For this purpose, we employ a linear Diophantine equation. A linear Diophantine equation (in three variables) is an equation of the general form  $ax + by + cz = d$  where  $a$ ,  $b$  and  $c$  are given integers and  $x$ ,  $y$  and  $z$  are unknown integers. It is a classical fact that (see [2] for more detail) this Diophantine equation has integer solution in  $x$ ,  $y$  and  $z$  if, and only if,  $d$  is a multiple of the greatest common divisor of  $a$ ,  $b$  and  $c$ . In order to deal with matrices of higher order, instead of considering Diophantine equation on more variables, we try to build it by using the notion of the Kronecker product, running over an inductive procedure. We recall that for the matrices  $A = [a_{ij}]$  and  $B$  of order  $m \times n$  and  $p \times q$ , respectively, the Kronecker product of the matrices  $A$  and  $B$ , denoted by  $A \otimes B$ , is the  $mp \times nq$  matrix defined by

$$A \otimes B = \begin{bmatrix} a_{11}B & \dots & a_{1n}B \\ \vdots & & \vdots \\ a_{m1}B & \dots & a_{mn}B \end{bmatrix},$$

where the block  $a_{ij}B$  is formed by multiplying each  $a_{ij}$  element by the entire matrix  $B$ .

*Theorem 1:* (for the case  $n = 3$ )

In this section, we prove Theorem 1 for the case  $n = 3$ . We give a numerical example to ensure that the method of proof is constructive.

*Lemma:*

Given positive integers  $d$  and  $M$ , there exist infinitely many matrices  $A = [a_{ij}]_{1 \leq i,j \leq 3}$  with integer elements satisfying  $a_{ij} \geq M$  and  $\det A = d$ .

*Proof:* For the given integer  $M > 0$  we consider the following  $3 \times 3$  matrix with integer entries

$$A = \begin{bmatrix} M & M + 1 & x \\ M + 1 & M + 2 & y \\ 2M + 1 & 2M + 2 & z \end{bmatrix}.$$

Note that  $\det A = -Mx + (M + 1)y - z$ . Thus,  $\det A = d$  reads as the following linear Diophantine equation in three variables

$$-Mx + (M + 1)y - z = d. \quad (1)$$

Since  $\gcd(M, M + 1, 1) = 1$ , (1) has infinitely many solutions. We can solve (1) by reducing it into a two variable equation. Let  $(M + 1)y - z = W$ . By (1) we have the following linear Diophantine equations in two variables

$$-Mx + W = d, \quad (2)$$

$$(M + 1)y - z = W. \quad (3)$$

We observe that  $x_0 = 0$  and  $W_0 = d$  is a particular integer solution of (2). Thus, all integer solutions of (2), are

$$x = t, \quad W = d + Mt, \quad (t \in \mathbb{Z}). \tag{4}$$

From (3) and (4) we have  $(M + 1)y - z = d + Mt$ . Since  $\gcd(M + 1, 1) = 1$ , this equation has infinitely many solutions for all integers  $M$ . It is easy to check that  $y_0 = t$  and  $z_0 = t - d$  is a particular integer solution. So, all of its integer solutions are

$$y = t - k, \quad z = t - d - (M + 1)k, \quad (t, k \in \mathbb{Z}).$$

Attaching  $x = t$  to this, we obtain the general solution of (1) as well. To obtain conditional matrices of the Lemma, it is sufficient to take  $t > M$  and  $k < \lfloor \frac{t-d-M}{M+1} \rfloor$ , simultaneously. This completes the proof.

*Remark:* We can investigate the Lemma above with several Diophantine equations. Let  $p, q$  and  $r$  be different prime numbers greater than  $M$ , and

$$A = \begin{bmatrix} p & p & x \\ q & q + 1 & y \\ r & r & z \end{bmatrix}.$$

Then, the equality  $\det A = d$  reads as the Diophantine equation  $pz - rx = d$ . The condition  $\gcd(p, r) = 1$  ensures the existence of infinitely many solutions for this equation. Therefore, the Lemma can be considered with at least 5 of 9 prime entries.

*Example*

Consider three prime numbers 15485863, 32452843 and 49979687 greater than 15000000. Let

$$\begin{aligned} a_{11} &= a_{12} = 15485863, \\ a_{21} &= a_{22} - 1 = 32452843, \\ a_{31} &= a_{32} = 49979687. \end{aligned}$$

By the above Remark, we obtain

$$15485863z - 49979687x = 2 \quad \text{with} \quad M = 15000000 \quad \text{and} \quad d = 2.$$

It is easy to check that

$$x_0 = 13228138 \quad \text{and} \quad z_0 = 42693016$$

is a particular integer solution and so all integer solutions to the equation are of the form

$$x = 13228138 + 15485863t, \quad z = 42693016 + 49979687t, \quad t \in \mathbb{Z}.$$

We take  $y = 160481183$ , which is a prime number, and  $t = 1$ . Thus

$$A = \begin{bmatrix} 15485863 & 15485863 & 28714001 \\ 32452843 & 32452844 & 160481183 \\ 49979687 & 49979687 & 92672703 \end{bmatrix},$$

with  $\det A = 2$ .

*Theorem 1* (for the general case)

We give some notations to simplify our description of the proof of Theorem 1 for the general case  $n \geq 3$ . We let  $J_n$  and  $I_n$  denote the  $n \times n$  all-one matrix and the identity matrix, respectively and  $\mathbf{0}_{m \times n}$  be the  $m \times n$  zero-matrix. For square matrices  $A_1, \dots, A_n$  we define  $\text{diag}(A_1, \dots, A_n)$  as the block matrix where the blocks along the diagonal are  $A_1, \dots, A_n$  and all other blocks are  $\mathbf{0}$ . Now we give the proof of Theorem 1.

*Proof of Theorem 1:* First, we assume that  $n$  is even, and we let  $n = 2k$  for some integer  $k$ . By the result of [1], for  $l \in \mathbb{N}$ , there are infinitely many sets  $\mathcal{A}_l = \{A_{l1}, \dots, A_{lk}\}$  of  $2 \times 2$  matrices with integer elements greater than  $M$  that satisfy the condition  $\prod_{i=1}^k \det A_{li} = d$ . We consider the following block matrix

$$B_l = J_{2k} \otimes A_{l1} + \text{diag}(\mathbf{0}_{2 \times 2}, A_{l2}, \dots, A_{lk}).$$

By the definition of  $B_l = [b_{ij}]_{1 \leq i, j \leq 2k}$ , we observe that  $b_{ij} \geq M$ . It is easy to see that  $B_l$  is equivalent to  $\text{diag}(A_{l1}, \dots, A_{lk})$  by using row operations. Since  $\det(\text{diag}(A_{l1}, \dots, A_{lk})) = \prod_{i=1}^k \det A_{li}$ , we deduce that  $\det(B_l) = d$ .

For odd values of  $n$  we follow an inductive procedure. The case  $n = 3$  has been considered in the Lemma. For  $n = 2k + 1$  with  $k \geq 2$  we let  $A_{2k+1}$  be the  $(2k + 1) \times (2k + 1)$  block matrix defined recurrently by

$$A_{2k+1} = \begin{bmatrix} A_{2k-1} & B \\ C & A_2 + D \end{bmatrix}$$

where  $A_2$  is the  $2 \times 2$  matrix by methods of [1],  $B$ ,  $C$  and  $D$  obtained from  $A_{2k-1}$  as follows:

- $B$  is the first two columns of matrix  $A_{2k-1}$ ,
- $C$  is the first two rows of matrix  $A_{2k-1}$ ,
- $D$  is the upper left block of size  $2 \times 2$  of the matrix  $A_{2k-1}$ .

Accordingly,  $A_{2k+1}$  and  $\text{diag}(A_{2k-1}, A_2)$  are equivalent, and consequently

$$\det A_{2k+1} = \det A_{2k-1} \det A_2 = d.$$

Note that we can take infinitely many  $2 \times 2$  and  $(2k - 1) \times (2k - 1)$  matrices same as  $A_2$  and  $A_{2k-1}$  with integer elements larger than  $M$  and  $\det A_{2k-1} \det A_2 = d$ . Therefore, there exist infinitely many matrices  $A_{2k+1}$  satisfying the conditions of the theorem.

References

1. M. Hassani, Conditional  $2 \times 2$  matrices with three prime elements and given determinant, *Math. Gaz.* **105** (July 2021) pp. 305-306.
2. D. M. Burton, *Elementary number theory*, McGraw-Hill, New York, 2007.
3. D. C. Lay, *Linear algebra and its applications*, 3rd edn., Addison-Wesley Publishing Co., Reading, MA, 2003.

10.1017/mag.2024.126 © The Authors, 2024

HOSSEIN MOSHTAGH

Published by Cambridge

Department of Computer Science,

University Press on behalf of

University of Garmsar,

The Mathematical Association

Garmsar, Iran

e-mail: [h.moshtagh@fmgarmsar.ac.ir](mailto:h.moshtagh@fmgarmsar.ac.ir)

### 108.43 An alternating recursion: proof of a conjecture by Erik Vigen

The following construction was considered by Erik Vigen in [1]. With positive numbers  $a_0, a_1$  given,  $a_n$  is defined for  $n \geq 2$  by an alternating recursion:

$$a_{2n} = \sqrt{a_{2n-2}a_{2n-1}},$$

the geometric mean of the previous two terms, while

$$a_{2n+1} = \frac{2a_{2n-1}a_{2n}}{a_{2n-1} + a_{2n}},$$

the harmonic mean of the previous two terms (which we denote by  $H(a_{2n-1}, a_{2n})$ ).

It was conjectured in [1], with support from numerical calculations, that  $a_n$  converges to  $\gamma(a_0, a_1)$ , where for  $x < y$ ,

$$\gamma(x, y) = \frac{y}{\sqrt{\frac{y}{x} - 1}} \tan^{-1} \sqrt{\frac{y}{x} - 1} \tag{1}$$

while for  $x > y$ ,

$$\gamma(x, y) = \frac{y}{\sqrt{1 - \frac{y}{x}}} \tanh^{-1} \sqrt{1 - \frac{y}{x}}. \tag{2}$$

Here we give a proof for the case where  $a_0 < a_1$ , so that (1) applies. The case  $a_0 > a_1$  can then be proved similarly, or derived from (1) using  $\tanh^{-1} z = \frac{1}{2} \ln \frac{1+z}{1-z}$  and  $\ln(1 + ic) - \ln(1 - ic) = 2i \tan^{-1} c$ .

*Lemma 1:* ( $a_n$ ) tends to a limit.

*Proof:* First, since either type of mean of  $x$  and  $y$  lies between  $x$  and  $y$ , an easy induction shows that  $a_0 < a_n < a_1$  for all  $n \geq 2$ . Now  $a_2 = \sqrt{a_0 a_1}$ , so