

Menopause and Risk-Taking Behaviours: A Cross-Sectional, Online Survey

Dr Dan Reisel*, Ms Aini Kamal, Dr Sarah Glynn and Dr Louise Newson

Newson Health, Stratford-upon-Avon, United Kingdom

*Presenting author.

doi: 10.1192/bjo.2024.232

Aims. Limited data suggest that negative mood symptoms in the menopause transition may be associated with a higher prevalence of alcohol misuse and other risk-taking behaviours in menopausal women. Excessive alcohol consumption can exacerbate menopausal symptoms, reduce quality of life and is associated with chronic morbidity that overlaps with the consequences of long-term oestrogen deficiency (such as osteoporosis and cardiovascular disease). The aim of this survey was to explore the impact of mental ill-health on alcohol consumption and gambling habits in menopausal women.

Methods. We constructed an anonymous survey consisting of multiple-choice and free-text questions. The survey was distributed online via social media channels on the 22 August 2023 and was open for 6 weeks. All perimenopausal and menopausal women were invited to participate. Responses were collected using the Qualtrics survey platform and analysed in Excel for descriptive statistics.

Results. 1,178 responses were submitted. One in three women reported drinking more alcohol during the perimenopause/menopause; 15% of women drink more than the recommended maximum of 14 units per week, and 24% (286) are spending up to £50 per week on alcohol. 70% (332) cited anxiety, stress, and/or depression as the reason for their increased alcohol consumption, whilst 29% (135) said they drank to alleviate menopause symptoms. Further, 5% (54) of respondents admitted gambling more since the onset of perimenopause/menopause; 43% (27) said it was due to anxiety, stress, and/or depression, whilst 13% (9) said they do so to help manage their menopause symptoms.

Conclusion. This anonymous, cross-sectional survey found evidence of an association between menopause and addiction. Increased awareness of this association should facilitate earlier recognition and more timely access to support and effective treatment for addiction, including hormone replacement therapy to treat menopausal symptoms that may underlie and/or exacerbate unhealthy lifestyle behaviours.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

Neurodevelopmental Disorders and Their Association With Neurodegenerative Disorders: A Systematic Narrative Review

Dr Rebecca Restorick^{1,2*}, Dr Huw Dunstall³ and Dr Katja Umla-Runge²

¹NHS Borders, Selkirk, United Kingdom; ²Cardiff University, Cardiff, United Kingdom and ³Swansea Bay UHB, Swansea, United Kingdom

*Presenting author.

doi: 10.1192/bjo.2024.233

Aims. Neurodevelopmental disorders (NDDs), such as dyslexia, dyspraxia, and dyscalculia affect cognitive function and therefore share symptomology with neurodegenerative disorders, such as

Alzheimer's disease, vascular dementia, and frontotemporal lobe dementia. The primary aim of this narrative systematic review is to ascertain if there is an association between NDDs and neurodegenerative disorders. Secondary aims are what the prevalence of NDDs within a dementia population is and what effect these early life learning disorders have on patients as they get older. It was hypothesised that NDDs would overestimate the severity of cognitive impairment, thereby increasing the severity of dementia staging, and impacting patient care.

Methods. Using a Population, Exposure, Comparator, Outcome, Setting, and Study design (PECOS) framework, keywords of "dementia", "dyslexia", "Dyspraxia/clumsy child syndrome/developmental apraxia/motor learning difficulty/disorder of attention and motor perception" and "dyscalculia/mathematical learning disability" were searched for on 4 databases (SCOPUS, OVID, Cochrane Central Register of Controlled Trials and Web of Science) from January 1, 1960 – June 10, 2022. Studies were included if they discussed both neurodegenerative and neurodevelopmental disorders or compared an intervention typically used in one disorder on the other (e.g., dementia intervention being used on neurodevelopmental disorder). Studies were excluded from grey literature articles, or if they only discussed a neurodevelopmental or neurodegenerative disorder without reference to the other, or if it included acquired, rather than neurodevelopmental dyslexia, dyscalculia, or dyspraxia.

Results. A total of 8 studies were included for narrative synthesis. The main finding was an association between dyslexia and both Alzheimer's disease and frontotemporal dementia. Many studies suggested this was due to a genetic phenotype that caused a vulnerability in the language regions of patients' cortices. There was also evidence of structural changes associated with NDDs and increased levels of grey and white matter atrophy in dementia subtypes, particularly in the language areas of the brain.

Conclusion. Due to screening and consequently formal diagnosis of neurodevelopmental disorders only recently coming into education systems, many adults currently attending memory clinics did not have a formal diagnosis. As there was limited research on dyspraxia and dementia, partly due to limited research into dyspraxia itself and without a standardized diagnostic tool for adolescents and adults, further research is needed in this area. The hypothesis of NDDs increasing the severity of dementia staging was also not supported by the literature results, and on the contrary, some studies suggested greater global preservation of cognitive function in patients with NDDs and dementia.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

Decrypting the Thalamic Subnuclei and Functional Composites in Adolescents With Psychotic Experiences

Dr Sahar Riaz*, Michael O'Connor, Anurag Nasa, Dr Mary Cannon and Dr Darren Roddy

Royal College of Surgeons in Ireland, Dublin, Ireland

*Presenting author.

doi: 10.1192/bjo.2024.234

Aims. The thalamus, a dual grey matter formation within the diencephalon is thought to be involved in psychosis. It consists of distinct nuclei with specific functions. To date no study has investigated the volumes of the thalamic nuclei in young adolescents with Psychotic Experiences (PEs).

Methods. This study used T1 imaging with Freesurfer analysis to investigate the differences in thalamic nuclei in 98 young people (53 with PEs) over three time points, from ages 11 to 18. A linear mixed effects (LME) model was used to examine the longitudinal nature of the data.

Results. The findings were entirely left sided – specifically a smaller left whole thalamus ($p = 0.04$), significant reduction in the size of the left pulvinar ($p = 0.008$) and a slight increase in the size of left ventral nucleus ($p = 0.005$).

Conclusion. This study found significant volumetric differences in thalamic functional composite nuclei between adolescents with a history of PE compared with healthy controls. Two such nuclear groups survived post-hoc DTR testing, the left ventral and left pulvinar nuclei. The pulvinar nucleus demonstrated a reduced volume over time in PE groups compared with healthy controls whilst the left ventral nucleus demonstrated an increased volume over time in PE groups compared with controls. The thalamus has been shown to be actively involved in the modulation of cortico-cortical communication via cortico-thalamo-cortical pathways, thus synchronizing the activity of the cortex during tasks that require attention. One of the core deficits believed to be a part of psychotic illnesses is the inappropriate modulation of attention through various cortical networks. This disrupted modulation results in a lack of control of goal-directed behaviour and can be attributed to the changes seen in pulvinar in psychotic illnesses, thus resulting in impairment in the integrity of sensory information and context processing. The affiliation of the ventral thalamic nucleus to the dopaminergic system, particularly the substantia nigra, may aid in explaining why this nucleus demonstrates larger volumes in adolescents with PEs compared with healthy controls over time.

More research needs to be done on following this cohort up, specifically investigating changes in thalamic nuclei in those who develop a diagnosable psychotic disorder.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

Linguistic Creativity and Formal Thought Disorder in Schizophrenia

Dr Louise Robinson^{4*}, Professor Dawn Archer¹, Dr Alex Bartha³, Dr Gerasimos Chatzidamianos² and Mr Oliver Delgaram-Nejad¹

¹Manchester Metropolitan University, Manchester, United Kingdom;

²Americian College of Greece, Greece; ³East London NHS Foundation Trust, East London, United Kingdom and ⁴Lancashire and South Cumbria NHS Foundation Trust, Lancashire, United Kingdom

*Presenting author.

doi: 10.1192/bjo.2024.137

Aims. The present investigation was interested in whether formal thought disorder (FTD) in schizophrenia was in any way related to linguistic creativity. The project's main aims and research questions were the development of operational definitions of linguistic creativity and FTD in schizophrenia, an investigation of creative language processing in schizophrenia, and an investigation of creative language output in schizophrenia.

Methods. We designed a psycholinguistic experiment and collected natural language data to build a specialised schizophrenia corpus. Recruitment for the psycholinguistic experiment was challenged by the COVID pandemic and the technical abilities of

clinical participants. Those data are thus underpowered and not reported in the results. We collected sufficient data for the construction of the specialised corpus.

Results. We tested an operational definition of FTD in schizophrenia (the '4TD Framework') against our natural language dataset. There was good support for the framework, with grammatical and discourse tracking features reliably distinguishing clinical and comparison speakers ($p < 0.05$). We also examined concordance lines and grouped random concordances into error types. Error types were consistently similar across groups, suggesting that speech disturbances in schizophrenia are on a continuum with those of nonclinical speakers. We also conducted a keyness analysis to examine the key terms and semantic categories present in the corpus and noted significant differences in the clinical cohort. Clinical participants found discussion of the topic of linguistic creativity more challenging, deviating from topic more often. They also involved topics of emotional and personal concern at rates of up to 16 to 32 times more often than comparison participants in some cases.

Conclusion. Our results provide support for the dysexecutive and dyssemantic hypotheses of FTD, as well as work on the Thought Language Index (TLI) that also suggests that language disturbances in schizophrenia and FTD are on a continuum with nonclinical speech. Further research is needed to understand how these phenomena are positioned in relation to FTD as a transdiagnostic entity.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

Patterns of Psychiatric Help Seeking Behaviour Among Patients With Psychotic Illness, Presenting at Mental Health Institute in Pakistan

Dr Rajesh Kumar^{1,2}, Dr Rahul Roy^{1,3*} and Dr Priyanka Devi⁴

¹LUMHS, Jamshoro, Pakistan; ²SIR CJIP, Hyderabad, Pakistan; ³West London NHS Foundation Trust, Crowthorne, United Kingdom and ⁴PUMHSw, Nawabshah, Pakistan

*Presenting author.

doi: 10.1192/bjo.2024.235

Aims. To identify the patterns of psychiatric help seeking behaviour among patients with psychotic illness presenting to mental health institute in Pakistan.

Methods. This Descriptive, Cross-Sectional Design comprised of a sample of 103 patients with psychotic diseases, chosen via non-probability – consecutive sampling at Outpatient Department of Sir Cowasjee Jehangir Institute of Psychiatry, Hyderabad. The relatives of psychotic patients who were between 18 and 65 years of age with either gender and living with at least one family member were interviewed after taking informed written consent. An anonymous self-structured questionnaire containing inquiries pertaining to basic biodata, sociodemographic details, psychiatric diagnosis and disease particulars, pattern of help seeking and time and reasons for delay.

Results. The mean age of the sample stood at 32 years (± 9.5 SD). 1/3 of the sample comprised of male patients while only 24% were comprised of female population. The mean time elapsed after first episode psychosis till interview was 82 ± 32 months (7 years) while mean delay in help seeking to any helper was reported to be 41 ± 17 months (3.5 years). Majority of the patients approached first to faith healers (Aamil Baba, Witch Doctor,