

## Research Article

**Cite this article:** Squair C, Proudfoot K, Montelpare W, Doucette T and Overall KL (2024). Effects of changing veterinary handling techniques on canine behaviour and physiology Part 2: Behavioural measurements. *Animal Welfare*, **33**, e43, 1–16  
<https://doi.org/10.1017/awf.2024.37>

Received: 21 July 2023  
 Revised: 12 March 2024  
 Accepted: 11 June 2024

**Keywords:**

adaptive care; animal welfare; collaborative care; low stress handling; veterinary examination; WDQ-Pet

**Corresponding author:**

Karen L Overall;  
 Email: [koverall@upeu.ca](mailto:koverall@upeu.ca)

**Author contributions:**





Conceptualisation: KLO, CS; Formal analysis: KLO, TD, WM; Funding acquisition: KLO; Investigation: CS, KLO; Methodology: CS, KLO, KP, WM; Project administration: CS, KLO; Supervision: KLO; Writing – original draft: CS, KLO; Writing – review & editing: KP, WM, TD.

© The Author(s), 2024. Published by Cambridge University Press on behalf of The Universities Federation for Animal Welfare. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided that no alterations are made and the original article is properly cited. The written permission of Cambridge University Press must be obtained prior to any commercial use and/or adaptation of the article.



Twitter: @UFAW\_1926  
 webpage: <https://www.ufaw.org.uk/>

# Effects of changing veterinary handling techniques on canine behaviour and physiology Part 2: Behavioural measurements

Camille Squair<sup>1</sup> , Kathryn Proudfoot<sup>1</sup> , William Montelpare<sup>2</sup> ,  
 Tracy Doucette<sup>3</sup> and Karen L Overall<sup>1</sup> 

<sup>1</sup>Department of Health Management, Atlantic Veterinary College, University of Prince Edward Island, Charlottetown, PE, Canada, C1A 4P3; <sup>2</sup>Department of Applied Human Sciences, University of Prince Edward Island, Charlottetown, PE, Canada, C1A 4P3 and <sup>3</sup>Department of Psychology, University of Prince Edward Island, Charlottetown, PE, Canada, C1A 4P3

**Abstract**

Signs of distress in dogs during veterinary visits are often normalised rather than viewed as welfare concerns. Interventions designed to reduce fear during veterinary visits were evaluated to see if they affected dogs' behaviours compared to dogs without interventions. Twenty-eight dogs were examined at four visits across eight weeks. Dogs were randomised into intervention (distress reduction/adaptive care) and control groups (standard care) and evaluated via the Working Dog Questionnaire – Pet Dog Version (WDQ-Pet). At visit 1 (baseline) all dogs received the control protocol. Homework was assigned following visit 1 to practice collaborative examination (intervention) or to pet the dog (control) for the same allotted time. At each visit, behaviours were scored (clinical stress score) via video and in-person observations when dogs entered the hospital, stepped onto a scale to be weighed, entered the exam room, at the beginning and end of examination, and after venipuncture. There were no differences between groups at visit 1, or across visits entering the hospital or exam room. At visit 4, intervention scores either decreased or remained low when weighed, and at the beginning and end of the physical exam. Control scores were significantly higher than the intervention scores during these periods. Reduced clinical stress scores indicate intervention dogs had improved care experience compared to the control. The study results highlight the value of applying simple and adaptable interventions, ultimately leading to improved animal care and welfare.

**Introduction**

The welfare of dogs undergoing veterinary care has primarily focused on achieving a basic standard of care that includes redress of physical pain. Through this lens, signs of fear, anxiety and general distress have often been normalised — and even expected — as part of routine care. Instead, such signs should be viewed as animal welfare concerns, affecting both behavioural preferences regarding how to be handled and mental well-being (Broom 2016; Mellor 2016).

Fear is defined as an emotional and motivational state provoked by specific stimuli, resulting in withdrawal, active avoidance, and/or defensive behaviours (Steimer 2002; Overall 2013a). Furthermore, fear can result in physiological and behavioural indicators of response to stressors and/or distress. Fearful responses may begin as soon as dogs walk into a clinic. In one study, conducted at a veterinary hospital in Germany, fewer than half of the dogs entered the clinic calmly ( $n = 62/135$ ) and 13.3% ( $n = 18/135$ ) had to be dragged or carried into the building (Döring *et al.* 2009). Another study found that when walking into a veterinary clinic, 60% ( $n = 279/462$ ) of dogs showed apprehensive postures and 18% ( $n = 81/462$ ) showed signs of fear-related aggression (Stanford 1981). Two-thirds of dogs in a veterinary waiting room ( $n = 30/45$ ) spent more than 20% of the time exhibiting at least one sign of stress, and 53% ( $n = 24/45$ ) exhibited four or more signs of stress; the most common signs being nose licking, panting, lowered ears, and crying/vocalisation (Mariti *et al.* 2015). Elevated physiological stress markers in dogs, including serum cortisol and heart rate, have been found in dogs constrained to sit in some waiting-room environments (Perego *et al.* 2014). Other aspects of veterinary visits, including walking onto a scale, physical examination, and being placed onto an examination table, have shown to increase signs of distress — including trembling, tail-tucking, and avoidance/escape behaviours — in canine patients (Hernander 2008; Döring *et al.* 2009; Mariti *et al.* 2017; Stellato *et al.* 2019a).

The pervasive level of fear currently experienced by patients during veterinary visits has both immediate welfare concerns, long-term implications and consequences. Each negative event an animal experiences at the veterinary clinic conditions them for future negative responses to

similar events, causing subsequent visits to become more difficult and time-consuming for patient, owner, and staff members alike (Döring *et al.* 2009; Stellato *et al.* 2021). Dogs < 2 years old that visited the practice frequently were more fearful than older dogs that visited infrequently (Döring *et al.* 2009), suggesting that exposure to experiences they perceive as fearful are salient and that repeated exposure to veterinary practices and associated procedures may sensitise the patient (Domjan 2015). Such outcomes are behaviourally and emotionally problematic for our patients, and costly to the veterinary team in terms of time, staff costs, stress and burn-out (Steffey *et al.* 2023).

In many species, including dogs, fear can impair immune function and reproductive abilities, increase the risk of contracting infectious diseases, delay healing, and have a negative effect on lifespan (Moberg 2000; Dhabhar 2009; Dreschel 2010; Gouin & Kiecolt-Glaser 2011; Edwards *et al.* 2019). Stress responses experienced during veterinary visits negatively affect the animal, pet owner and veterinary staff in a variety of manifestations. Outcomes may include decreased frequency of veterinary care, negative effects of stress on the patient's long-term health, impaired ability to assess, accurately diagnose, and treat health concerns (e.g. high heart rate or tense abdomen, which could be due to disease processes or a stress response) (Beerda *et al.* 1997; Schubert *et al.* 2009). Furthermore, the physical and logistic struggle involved in providing basic care carries an increased risk of injury to the veterinary team when the patient experiences stress, and stress experienced by the pet — and by extension the owner — has been cited as a contributing factor for delaying veterinary care (Volk *et al.* 2011; Overall 2013b; Edwards *et al.* 2019; Stellato *et al.* 2021). Overall, this negatively affects treatment outcomes and the standard of care for both interventional and preventative veterinary care.

There is now an increased acknowledgment and awareness of the persistence of stress and fear in veterinary patients (e.g. [www.fearfreepets.com](http://www.fearfreepets.com)). Recognition has grown in the veterinary community for the importance of low stress handling in the veterinary hospital and clinic (Overall 2013b, 2019; Lloyd 2017; Mandese *et al.* 2021; Riemer *et al.* 2021). Some research has shown that dogs with positive experiences while at the veterinary clinic have been found to show fewer behavioural indicators of fear, such as trembling, tail tucking, and avoidance behaviours, than those with negative experiences (Döring *et al.* 2009), and researchers have begun to investigate the influence of collaborative care on dog fear levels during veterinary examination (Stellato *et al.* 2019b). Early veterinary visits set the foundation for subsequent interactions and can have lasting effects on the persistence of fearful behaviours — including panting, yawning, lip-licking — in patients into adulthood (Godbout & Frank 2011). There have been many recommendations regarding low stress handling, such as reducing restraint during procedures (blood draw, exam) (Overall 2013b; Lloyd 2017; Riemer *et al.* 2021) or using a blue bathmat and treats when weighing (Overall 2013b; Lloyd 2017; Edwards *et al.* 2019) and applying EMLA (2.5% lidocaine/2.5% prilocaine) cream prior to venipuncture (Overall 2013b; van Oostrom & Knowles 2018; Oliveira *et al.* 2019; Crisi *et al.* 2021). However, assessment of the value of implementing these techniques within a veterinary setting and the effectiveness of these techniques in reducing canine fear and distress is limited. Hence, the aim of our study was to determine whether simple, easy, and inexpensive interventions positively affected distress in canine patients at the veterinary clinic. Rather than singling out one

intervention, an examination protocol that altered most of the patterns of the standard physical exam was utilised for the intervention group, since what the dogs experience is the entire approach. Accordingly, responses were assayed for components of an overall pattern of changes, rather than to any single individual intervention.

This paper is part of a larger project where dogs were evaluated behaviourally and physiologically for their responses to the veterinary visit and intervention. The physiological differences between the control and intervention group at week 8 (visit 4) are discussed elsewhere (Squair *et al.* 2023). Here, we report on the behavioural data. Our objectives were to: (1) report on the demographic and owner-reported behavioural responses to the Working Dog Questionnaire – Pet Dog Version (WDQ-Pet) for each group of dogs to learn if dogs differ in behavioural history or exposure that could affect outcomes during veterinary exams; (2) compare the behavioural responses of dogs to veterinary clinic visits between in-person and video assessments; and (3) determine the effect of a series of interventions during the veterinary exam on behaviours that indicate fear and/or distress in dogs.

## Materials and methods

### Ethical approval

The animal study protocol was approved by the Animal Care Committee (ACC) and the Research Ethics Board (REB) of UPEI (Protocol 21-02; 19 May 2021). Written informed consent was obtained from all dog owners involved in the study, and they could withdraw from the study at any time.

### Enrolment and inclusion criteria

Dogs whose guardians expressed an interest in their dogs' behaviours during veterinary visits and who were interested in making such visits as happy as possible were solicited for the study via posters (Appendix 1; Supplementary material). Posters were placed in local businesses and veterinary offices within Charlottetown, Prince Edward Island, Canada and within the Atlantic Veterinary College (AVC) Veterinary Teaching Hospital (VTH) waiting room, and in the hallways of the hospital. A recruitment message was sent to AVC staff and veterinary students via the AVC dean's office. A total of 30 dogs were screened to enrol a target of 28 participants (Table 1) based on initial power calculations of 90% with a one-tailed probability of 0.1 and  $z_{\beta} = 1.28$ .

Inclusion criteria specified that participating dogs had to be at least six months of age and be in good health. Requiring the dogs be six months of age guaranteed some prior veterinary experience since dogs would have their initial veterinary administered core vaccines by then and proof of rabies vaccination was required. This is also the minimum age used for all pharmacological studies that have licensed behavioural medications, including those used prophylactically for veterinary evaluation. Exclusion criteria included females that were pregnant or lactating, animals that were receiving behaviour-altering medications, and those with a history of overt aggression during veterinary examinations. The pre-enrolment WDQ-Pet screened for overt aggression, and dogs that snarled, lifted their lip, growled, snapped, or bit when handled or approached by strangers or in veterinary situations were excluded for safety reasons.

**Table 1.** Signalment, treatment group (control or intervention), and study completion status of dogs (n = 30) that were screened and enrolled for participation

ID	Age (months)	Sex	Breed	Weight	Group	Status
01	76	MC	Dachshund	7.2 kg	Intervention	Completed
02	35	FS	Chihuahua mix	5.4 kg	Control	Withdrawn
03	102	MC	German Shepherd/Husky mix	30.2 kg	Control	Completed
04	46	MC	Springer Spaniel mix	19.2 kg	Intervention	Completed
05	61	FS	Beagle	13.2 kg	Control	Completed
06	73	FS	Shih Tzu mix	8.2 kg	Control	Completed
07	31	MC	Pitbull mix	30.4 kg	Intervention	Completed
08	120	MC	Dalmatian	27.0 kg	Intervention	Completed
09	31	MC	Bernese Mountain Dog	38.8 kg	Intervention	Completed
10	15	MC	Maltese/Lhasa Apso mix	4.6 kg	Intervention	Completed
11	38	FS	Springer Spaniel	21.2 kg	Control	Completed
12	48	MC	Dalmatian	29.4 kg	Intervention	Completed
13	84	FS	American Staffordshire Terrier Mix	20.8 kg	Control	Completed
14	24	FS	Belgian Groenendael Sheepdog	21.2 kg	Control	Completed
15	31	MC	Mastiff mix	34.0 kg	Intervention	Completed
16	50	FS	Chihuahua mix	3.0 kg	Intervention	Completed
17	80	FS	German Shepherd mix	27.6 kg	Control	Completed
18	29	FS	Labradoodle	28.8 kg	Control	Completed
19	160	MC	Golden Retriever	38.8 kg	Intervention	Completed
20	68	MC	Poodle Dachshund mix	11.4 kg	Intervention	Completed
21	102	MC	Labrador Retriever	64.0 kg	Control	Completed
22	127	FS	Chihuahua mix	4.2 kg	Control	Completed
23	7	FI	Golden Retriever	23.6 kg	Control	Completed
24	42	MC	Labrador mix	20.6 kg	Intervention	Completed
25	19	FS	Terrier mix	6.6 kg	Intervention	Completed
26	114	FS	Newfoundland dog	56.2 kg	Control	Completed
27	62	FS	Labrador Retriever	25.0 kg	Intervention	Completed
28	24	FS	English Bulldog	25.8 kg	Control	Withdrawn
29	52	MC	Toy Goldendoodle	6.5 kg	Control	Completed
30	9	MI	Australian Cattle Dog	21.5 kg	Control	Completed

Study ID number reflects enrolment order (MC = male, castrated; MI = male, intact; FS = female, spayed; FI = female, intact).

### Demographics and historical behaviour

All owners completed the WDQ-Pet, a survey tool used in previous studies (Overall *et al.* 2006, 2019). The WDQ-Pet collects information on demographics (age, sex, number of homes, source, etc) and responses to a series of environmental and behavioural stimuli as noted by the owner. All categories evaluated in the WDQ-Pet are included in Table 2. The WDQ-Pet, with and without the scoring rubric, is found in Appendix 2 (see Supplementary material). The WDQ-Pet was used for two reasons. First, we used it to ensure inclusion and exclusion criteria were met. Second, the WDQ-Pet was used to assess whether, upon enrolment, the control and intervention groups differed with respect to demographic, historical, or behavioural response parameters including global (e.g. hesitancy to walk in some environments or on some substrates) and specific (e.g. separation anxiety, noise reactivity) fears and anxieties.

### Experimental design and exam protocol

Dogs were assigned using 20 iterations of a random number generator (Randomizer.org). Dogs were numbered from 1 to 28 as they were enrolled, and using the random number generator, were assigned by one investigator to group A or B. Assignment of control vs intervention group to coded group A or B was done by another investigator by a coin toss. Both groups were treated the same at the first visit, undergoing the control protocol. After the first visit, 14 dogs received the intervention protocol and 14 dogs continued to receive the control protocol for visits 2–4. The protocols differed when weighing the dog, during the physical exam and blood draw, and with respect to the type of homework assigned for the dog owners between visits 2, 3, and 4. For the intervention group homework consisted of practicing a collaborative physical exam for 5 min, three times per week, whereas owners in the control group

**Table 2.** WDAQ-Pet (Working Dog Questionnaire-Pet version) completed by owners' prior enrolment of dogs (n = 28). Questionnaire consists of 58 categories of interest (both demographic and behavioural/environmental information)

Demographic information
<ul style="list-style-type: none"> <li>Sex, weight, source of dog, age adopted, age at neuter</li> <li>More than 1 home after natal home (Y/N)/Previous home</li> <li>Training history - type of training (group, private, advanced – agility, herding etc)</li> <li>Method of maintaining at home (kennel, crate, loose in house, in yard, etc)</li> </ul>
Reward/Reinforcement-based questions
<ul style="list-style-type: none"> <li>Type of rewards used to train (food, toy, clicker, praise, other)</li> <li>Type of punishment/correction used with dog</li> <li>Use of ball, and/or pull toy/towel, and/or Kong/ball on a rope as a reward (ie. hold it, throw it, bounce it, other)</li> <li>How the dog ends playing fetch (activity measure)</li> <li>Fetch in the dark (Y/N) (activity measure)</li> </ul>
Dog's response to:
<ul style="list-style-type: none"> <li>Owner standing quietly with reward and squatting with reward</li> <li>Receiving a verbal praise and dog can see owner's face, and when they cannot see owner's face</li> <li>Receiving a verbal request to respond (e.g. sit) <ul style="list-style-type: none"> <li>Owner looking at dog and gives verbal praise</li> <li>Owner NOT looking at dog and gives verbal praise</li> <li>Owner looking at dog and gives a treat or toy</li> <li>Owner NOT looking at dog and gives a treat or toy</li> </ul> </li> <li>Receiving an object reward and throwing a toy (or other non-food reward object) (activity measure)</li> </ul>
Questions About reacting to the environment
<ul style="list-style-type: none"> <li>General events or activities of dog when outside walking</li> <li>Sniffing behaviour of dog on walks</li> </ul>
Dog's response to:
<ul style="list-style-type: none"> <li>Going out in rain – with and without thunder and wind, encountering puddles, new types of surfaces on walks (e.g. board placed over a hole, subway grate)</li> <li>Continuous loud noise, intermittent loud noise, and sharp burst of noise</li> <li>Human, known dog, and unknown dog approaching within 3 feet/1 m <ul style="list-style-type: none"> <li>Note if/type of accompanying vocalisation during this event</li> </ul> </li> <li>Moving objects (bicycles/motorcycles)</li> <li>Being asked to walk across 22 standard surfaces</li> </ul>
General behavioural patterns shown by the dog
<ul style="list-style-type: none"> <li>Type of seeking interaction/pattern displayed by the dog</li> <li>What calms the dog when they barks, etc</li> </ul>
Dog's response to:
<ul style="list-style-type: none"> <li>Food found on the street</li> <li>Approaches (by unfamiliar dog, or human, or vehicle) when confined in vehicles/crates</li> <li>Amount of time (15, 30, 60 min) spent in vehicle/crate</li> </ul>
Husbandry information
<ul style="list-style-type: none"> <li>Maintenance style of dog when alone (e.g. free inside, outside, kennel etc)</li> <li>General behaviours displayed by dog when left alone</li> <li>Types of situations where the dog may have diarrhoea/vomiting</li> </ul>
General behavioural and medical history
<ul style="list-style-type: none"> <li>History of illness (hip dysplasia, cardiac disease, other disease)</li> <li>Dog's compliance with requests for sit, down/lie down, stay, wait, heel, fetch, leave it/drop it, take it</li> <li>SAIR (Separation Anxiety Intensity Rank) score to being separated/left – during real absence, virtual absence, and virtual and real absence combined</li> <li>AIR (Anxiety Intensity Rank) score to noises</li> <li>Aggression screen score</li> <li>Fear screen score</li> </ul>

**Table 3.** Treatment protocol summaries used for dogs (n = 28) within control (n = 14) and intervention (n = 14) groups at each of the four visits, originally published in Squair *et al.* (2023)

	Control treatment	Intervention treatment
Scale	<ul style="list-style-type: none"> <li>Walk-on stainless steel scale</li> <li>Weighed before examination</li> </ul>	<ul style="list-style-type: none"> <li>Walk-on stainless steel scale covered with blue, non-slip yoga mat, moved away from wall, with dog lured on with treats (Overall 2013b; Lloyd 2017; Edwards <i>et al.</i> 2019)</li> <li>Weighed after examination (Overall 2013b)</li> </ul>
Physical exam	<ul style="list-style-type: none"> <li>White coat</li> <li>Small dogs on table</li> </ul>	<ul style="list-style-type: none"> <li>No white coat (Marino <i>et al.</i> 2011; Cobos <i>et al.</i> 2015; Fanucchi 2022)</li> <li>All dogs on the floor or owner's lap if that was the dog's preference (Döring <i>et al.</i> 2009)</li> <li>Lickimat* (Innovative Pet Products PTY, Australia) and blue mat for non-slip examination (Overall 2013b; Westlund 2015; Lloyd 2017; Riemer <i>et al.</i> 2021)</li> </ul>
Blood draw	<ul style="list-style-type: none"> <li>Fake lidocaine application to three legs</li> <li>Standard needle and syringe</li> <li>Standard restraint</li> </ul>	<ul style="list-style-type: none"> <li>Application of lidocaine cream to three legs (Overall 2013b; van Oostrom &amp; Knowles 2018; Crisi <i>et al.</i> 2021)</li> <li>Closed double ended butterfly catheter system (Overall 2013b; Hefler <i>et al.</i> 2004; World Health Organization (WHO) 2010; Riddick 2023)</li> <li>Reduced to no restraint primarily using guidance and positioning (Overall 2013b; Lloyd 2017; Riemer <i>et al.</i> 2021)</li> </ul>
Homework	<ul style="list-style-type: none"> <li>Petting dog for 5 min, three times a week</li> </ul>	<ul style="list-style-type: none"> <li>Practice the steps of a collaborative physical exam for 5 minutes 3 times a week (Overall 2013b; Lloyd 2017; Stellato <i>et al.</i> 2019b, Riemer <i>et al.</i> 2021)</li> </ul>

were instructed to simply pet their dogs for the same duration and frequency. Table 3 summarises the key differences between each treatment group. The differences in the organisation and management of the visits and handling, including photograph illustration, for dogs in the control vs the intervention groups have been discussed in great detail in part one of this study series (Squair *et al.* 2023).

All dogs were weighed using the control group protocol at visit 1. The scale at visit 1 was a plain metal scale, with no interventions (step 1 of 4). For dogs unable to get onto the bare scale, that failure was noted, and interventions were sequentially offered starting with the addition of a blue yoga mat (step 2 of 4), moving the scale with the mat from the wall (step 3 of 4), and finally adding treats (step 4 of 4). After visit 1, all dogs in the control group continued with this weighing protocol, while the intervention group had all the interventions present from the beginning. This rubric helped to assess the level of intervention needed at visit 1 for all dogs, and the level of intervention across all visits required for control dogs. The time to get all four feet on the scale so that a weight could be accurately obtained was also measured across all visits using video data.

The same room was used for every visit, and physical examination and blood draws were conducted by the same clinician (CS) at every visit. Table 4 describes the order, timing, and frequency of the standardised physical exam used for both the intervention and



**Table 4.** Exam structure — including order, and timing or frequency — used for dogs (n = 28) in the intervention (n = 14) and control (n = 14) groups at each of four visits (Korpivaara *et al.* 2021). Numbers (1 through 15) are later referred to as different steps in the exam

Physical examination protocol
1. Dog stroked gently from head to base of tail three times
2. Hand placed over the thigh pulse point for 30 s
3. Lidocaine (2.5% lidocaine/2.5% prilocaine) put on legs (two saphenous and one cephalic) for intervention dogs (control dogs are just touched in these areas)
4. Auscultation of heart and lungs 15 s from each side of the chest
5. Manual manipulation of lymph nodes (in order submandibular, prescapular, popliteal)
6. Gentle abdominal palpation undertaken for 15 s
7. Each paw lifted for 5 s for testing placement; first hind limbs and then forelimbs
8. Lifting of upper lips (control of the oral mucous membranes)
9. Observation of external ear canals for 5 s each (without an otoscope)
10. Ear thermometer placed in position until reading
11. Eyes examined directly (observation of the conjunctiva, checking of the cornea) for 5 s each
12. Venipuncture
13. Gently place a hand on the dog's back and tell them they are good
14. Remove from table and give treat, or if the dog is on the floor, just give the treat (note whether the dog takes the treat on record)
15. Walk owner to parking lot and give treat mid-way to car (note response on record)

control groups, which has been used in other clinical studies (Godbout *et al.* 2007; Hauser *et al.* 2020; Korpivaara *et al.* 2021). During the exam in the intervention group, a blue mat was provided for the dog's use and Lickimat® with treats (cream cheese or Kong® cheese spray) was provided during the exam and venipuncture. Other treats offered throughout included string cheese, dehydrated liver treats, or hypoallergenic treats, depending on individual preference. The blue mat and Lickimat® were sent home with the dogs in the intervention group, so the owners could practice their physical exam procedure homework.

To obtain laboratory samples, dogs in the control group underwent routine venipuncture (using either saphenous or cephalic veins) involving physically holding off a vein and whole body-restraint (i.e. holding when standing or when lying in lateral recumbence). Blood was taken with a needle and Luer-lock syringe without the application of lidocaine. Since it was difficult to find a cream that felt or smelled to us like the 2.5% lidocaine/2.5% prilocaine cream (EMLA® cream) used for the dogs in the intervention group, the fur was rubbed over the area of both saphenous and cephalic veins as exam step 3 for the control dogs to mimic the effects of applying the lidocaine cream. For dogs in the intervention group, a researcher wearing gloves applied 2.5% lidocaine/2.5% prilocaine cream by rubbing it over the area of both saphenous and cephalic veins as in exam step 3. Additionally, dogs in the intervention group had blood taken with low-to-no restraint, using a butterfly vacuette, double-ended closed system catheter that did not require that the vein be held off, and in an adaptive manner that allowed them to choose their posture while they were licking a

Lickimat®. Owners were present for both the physical exam and blood draw in both groups and sat in one of the two chairs available. In the intervention group, owners were given the option to hold the Lickimat® for their pet or leave it on the mat on the floor. Owners were allowed to assist with gentle restraint (e.g. holding a collar/leash or petting the dog's head) in both groups if they felt comfortable/wanted to assist. Photographs of physical examination set-up and restraint methods have been published elsewhere (Squair *et al.* 2023).

Following each visit the investigators offered dogs a treat (dehydrated liver, string cheese, or a hypoallergenic treat) once they were in the parking lot to determine whether the groups responded differently after their veterinary visit experience was completed (yes or no). Each of the 14 dogs in each group had four opportunities for parking lot treats. This step was added based on the findings of Lind *et al.* (2017) that dogs that refused treats when in the hospital, willingly took them when outside the clinical environment.

### Clinical stress scores

To estimate if dogs experienced fear or distress during the visit, a series of in-person and video measures were used to assess the dogs' behaviour at each visit. All dogs were evaluated for their behavioural response using a six-point ordinal scale: 1 = calm; 5 = profound avoidance and distress, with 0 denoting excitement without distress (see Table 5 for the scale used at different stages of the veterinary exam). Dogs were scored as they entered the veterinary hospital, walked onto the scale, crossed the threshold into the

**Table 5.** Four clinic stress scales used to score behaviour of dogs (n = 28) within control (n = 14) and intervention (n = 14) groups for four specific events (entering the clinic, stepping onto the scale, entering the exam room, and at the beginning and end of the physical exam) at each of four visits (adapted from Overall 2013b)

Score	Definition of dog's behaviour and demeanour
<i>Scale 1: Entry to the clinic</i>	
0	Extremely friendly, outgoing, enthusiastically solicitous of attention
1	Calm, relaxed, seemingly unmoved
2	Alert, but calm and co-operative
3	Tense, stiff, but co-operative, panting slowly, not very relaxed but can still be easily led on lead; tail may be down but is not clamped hard – may vary in position; neck may be slightly lowered
4	Very tense, anxious, may be shaking or whining, with lowered or hunched body posture; will not sit or lie down; if exposed may do so behind owners' legs, panting, difficult to manoeuvre on lead; tail may be tucked
5	Extremely stressed, barking/howling, tries to hide, needs to be lifted up or forced to move; tail clamped hard; slinking, very low as resisting forward direction
<i>Scale 2: Weighing the dog</i>	
0	Extremely friendly, outgoing, enthusiastically solicitous of attention, eagerly gets onto scale
1	Calm, relaxed, seemingly unmoved, and walks easily onto scale and sits
2	Alert, but calm and co-operative, can get onto scale but may not sit on it

(Continued)

Table 5. (Continued)

Score	Definition of dog's behaviour and demeanour
3	Tense, but co-operative, panting slowly, not very relaxed but can still be easily led on lead, gets onto scale only with encouragement; tail may be down but not clamped and may vary in position as moves
4	Very tense, anxious, may be shaking or whining, may have hunched or lowered body posture, will not sit or lie down if exposed (may do so behind owners' legs), panting, difficult to maneuver on lead, must be helped/encouraged to get on or stay on scale for 10 s to get reading
5	Extremely stressed, barking/howling, tries to hide, needs to be lifted up or forced to get onto or stay on scale for 10 s to get reading; tail clamped and dog slinking, hiding or trying to escape
<i>Scale 3: Entering the exam room</i>	
0	Extremely friendly, outgoing, enthusiastic solicitous of attention
1	Calm, relaxed, seemingly unmoved and walks into room easily
2	Alert, but calm and co-operative; walks into room but is neither interactive or tense
3	Tense, but cooperative, panting slowly, not very relaxed but can still be easily led on lead; tail may lower as enters but variable in movement
4	Very tense, anxious, may be shaking or whining, will not sit or lie down if exposed (may do so behind owners' legs), panting, difficult to maneuver on lead, avoids room; may have hunched or lowered body posture and lowered or hidden tail
5	Extremely stressed, barking/howling, often has tail clamped, body lowered tries to hide or escape, needs to be lifted up or forced to move into room and moved forward manually
<i>Scale 4: Beginning and end of the physical exam</i>	
0	Extremely friendly, outgoing, enthusiastic, solicitous of attention
1	Calm, relaxed, seemingly unmoved
2	Alert, but calm and co-operative; neither interactive nor tense
3	Tense, but co-operative, panting slowly, not very relaxed but can still be easily manipulated for exam and co-operates with procedures; tail variable in movement and posture but may be low
4	Very tense, anxious, may be shaking, whining or frozen, difficult to manoeuvre, tries to avoid exam, hunched or lowered body posture possible, tail may be low, hidden, may hold onto table but not co-operate – more endures the process without active escape
5	Extremely stressed, barking/howling, tries to hide, body and tail lowered, escape risk, needs to be lifted up or held to be examined or would bolt from table or room, difficult to control and may be tempted to bite, may not be able to complete exam.

examination room, and at the beginning and the end of the examination (Table 5). The 'entry to clinic' score was taken as the dog crossed the threshold of entry doors with both front legs. The 'weighing the dog' score was taken when all four feet were on the scale and the dog stayed on long enough for a weight to be obtained. Weighing occurred on the way into the exam room for the control group, whereas for the intervention it occurred following the exam and blood draw as the dog was exiting the exam room. The 'entering

the exam room score' was taken when the dogs' full weight on their first paw crossed the threshold into the exam room. The 'beginning and the end of the exam' score was taken as the experimenter first touched the dog (beginning) and as they told the dog he or she was good (end).


Whether dogs took a treat in the parking lot at the end of the visit (yes or no) was evaluated. In-person measurements and camera management were performed by KLO and/or 1–3 trained observers. Three video cameras (one Sony 4K FDR-AX43 and two Sony Handicams HDR-CX405) — two on tripods and one hand-held — were used for video analysis. One tripod camera was used to record dogs entering the hospital and when being weighed, and all three cameras were used to record the dog entering the exam room and during examination and blood draw. Figure 1 illustrates a schematic of the exam room and camera set-up. Videos were analysed by CS and a trained veterinary student.

### Descriptive behaviour

In addition to the clinical stress scales, a descriptive ethogram was used to subjectively assess the behaviour of dogs at each step of the physical examination (see Table 4) during visits 1 and 4. A score ranging from 1–5 was assigned for (i) body posture, (ii) tail posture, (iii) ear posture, (iv) gaze, (v) mouth posture, (vi) activity and (vii) vocalisation for each step of the exam for visits 1 and 4 (Table 6). Two categories from the ethogram were omitted from the final analysis: pupillary changes and respiration. Pupillary changes could not be adequately visualised in most dogs to score them and, as noted in Squair *et al.* (2023), the ambient temperature in the exam room, given that this study was done in summer and early autumn, did not allow for the use of respiration as a measure in any context. Mean stress scores were derived by summing individual scores for regions and behaviours commonly noted to be indicative of fear/stress responses during each step of the physical examination. Table 6 shows the ethogram scale used, an evaluation of a series of body parts commonly assessed for signaling anxious or fearful responses in dogs (adapted from Overall 2013b; Korpivaara *et al.* 2021; Jokela *et al.* 2023). The ethogram scores for each of the eleven steps of the physical exam were only determined by video assessment. Two trained observers evaluated the ethogram scores of the dogs across the eleven steps of the physical examination and all scoring of all body regions: one was the investigator (CS) and another was a student, both masked to the visit number and group assignment. An overlap of 10% of the data were scored by both raters. Since Spearman's rho, an indication of concordance in scoring, was extremely high,  $r_s(254) = 0.94$ ;  $P < 0.001$ , the student completed the remaining scaled ethogram scores.

Time to get onto the scale to be weighed was measured from when the cue from the camera operator (if a verbal cue was not heard on video, time was calculated from when the dog entered the frame) to when all four paws were on the scale (when a successful weight could be obtained).

Behaviour during venipuncture was assessed using clinical stress scale 4 and time for the procedure in seconds. Both assessments were carried out using video analysis. Time for venipuncture was measured from when the needle first entered the skin to the needle exiting the skin following successful blood collection. Due to constraints of visibility on video, there were usable data for scores for only nine dogs in the control group and nine in the intervention group, and usable data for time measurement for ten dogs in the control group and nine in the intervention group. *A priori* it was specified that there would be a maximum of three sticks in two legs



**Figure 1.** Schematic of examination room set-up used to assess the dogs (n = 28) at each of four visits (m: meters). Cameras (cam) 1 and 2 were stationed on tripods. Camera 3 was hand-held to be able to fully assess the close up view of the dogs’ facial features.

**Table 6.** Ethogram scale for evaluating body regions and behaviours commonly noted to be indicative of fear/stress responses in dogs. This scale was used for video scoring behaviour of dogs (n = 28) undergoing each step of the exam in Table 4

Score	Body region/ activity	Behavioural responses
1	Body posture	Relaxed, moves on own Easily manipulated
	Tail posture	Relaxed, normal position for the breed or held high
	Ear posture	Relaxed, normal position for breed or high and softly forward
	Gaze/eyes	Relaxed Looks steadily and easily veterinarian and can meet gaze, or calmly looking elsewhere in the room
	Mouth posture	Normal – jaw relaxed Lips relaxed May be panting, but mouth is relaxed
	Activity	Flexible Sits or stands relaxed where it is placed or moves slightly to sit more comfortably.
	Vocalisation	None
2	Body posture	Slightly tense Can be manipulated
	Tail posture	Slightly lower than normal for the breed (but not completely down)
	Ear posture	Slightly moved backwards or tense Can change position back and forth.

(Continued)

**Table 6.** (Continued)

Score	Body region/ activity	Behavioural responses
	Gaze/eyes	Eyes slightly tense, blinking Looks only intermittently at veterinarian
	Mouth posture	Jaw tense Lips firm Corner of mouth may be slightly moved back May be panting
	Activity	Inactive but alert May pace a bit or stands/sits quietly
	Vocalisation	Occasional whine, cry
3	Body posture	Stiff, possibly rigid Body slightly lower Can still be manipulated but less fluid (more rigid)
	Tail posture	Completely down and low for the breed (but not tucked)
	Ear posture	Fully back Permanently moved backwards
	Gaze/eyes	Tense Scans room and may not look at veterinarian (or do so only seldom)
	Mouth posture	Lips are tighter and may be slightly back May lick lip occasionally Panting with increased tension – dry
Activity	Some rigidity and/or slight trembling May have increased/decreased locomotor activity May lean/step away from veterinarian	

(Continued)

Table 6. (Continued)

Score	Body region/activity	Behavioural responses
	Vocalization	Whines or whimpers half of the time May growl
4	Body posture	Hunched (crouched) Low posture Difficult to manoeuvre
	Tail posture	Tucked between legs
	Ear posture	Ears fixed back and down Tense
	Gaze/eyes	Tense Not scanning – looks steadily at distance or owner
	Mouth posture	Lips are tight and pulled back May be yawning May lick lips Panting with increased tension – dripping
	Activity	Tense and rigid Periodic trembling May try to gain distance from veterinarian/can freeze in this attempt May lie down – leg may be up and show parts of belly
	Vocalization	Constant whine or whimper May snarl, snap – without apparent intent to make contact
5	Body posture	Curled Completely withdrawn – belly maximally tucked
	Tail posture	Clamped hard up to the belly
	Ear posture	Ears fixed as low and back as it possible Extremely tense
	Gaze/eyes	Tense, eyes might be wide open (may see whites of the eyes) Cannot meet veterinarian's gaze Stares fixedly and steadily at immediate fore-distance
	Mouth posture	Lips and mouth are extremely tense and pulled back. Profound panting, salivating, gasping
	Activity	Uncontrollable trembling Tries to jump from table or completely freezes Needs to be held or will bolt Tries to hide May not be able to complete exam procedure
	Vocalisation	Yelp, howl Bite or bite attempt (intend to make contact)

(three received lidocaine cream) so this time estimate is a very rough gauge of the time required for all the machinations (including giving the dog a break or getting a new syringe or needle) needed to obtain the sample.

### Statistical analysis

Interval data were analysed using two-way (visit × treatment) repeated measures ANOVAs and independent samples *t*-tests. Ordinal data were analysed using Mann-Whitney *U* and Spearman rho, and nominal data were analysed using Chi-squared goodness

of fit test and the Fisher exact test. The specific tests used are described for each measurement as follows.

All categories for the WDQ-Pet were compared for dogs in the control and intervention groups using parametric or non-parametric tests as appropriate. Clinical stress scores obtained from in-person observation and video assessments were compared using Spearman's rho, a measure of concordance. In-person scoring was used to assess behaviour when dogs entered the clinic. Since there was some overlapping of the benchmarks for the rating scales, it is likely that the scoring was not truly discrete, so the ordinal data were usually treated as continuous (Norman 2010; Sullivan & Artino 2013). Scoring via video analysis was used to assess behaviour when weighed and during the physical exam and venipuncture. This was to ensure that multiple camera views could be used to see the entire dog, if needed. Video analysis was also used to measure duration when weighed, examined, and during venipuncture. Video analysis was used to determine clinic stress score for entry into the exam room, at the start and end of the physical examination, for ethogram physical exam scores, and during venipuncture. In-person scores were used when assessing dogs' entry into the clinic.

In addition to clinic stress scores, video analysis was also used to assess how long it took for the dog to successfully be weighed. To determine whether dogs were less stressed on their fourth visit compared with visit 1 (baseline) for any procedure, we converted raw data to nominal (Yes/No: did the score for behaviour stay the same, increase, or decrease). Each dog was assigned to either one of two categories (e.g. at visit 4 was their score lower or same/higher when compared to their score at visit 1). Intervention clinical stress scores when entering the clinic (Scale 1), being weighed (Scale 2), crossing the threshold into the examination room (Scale 3), and behavioural scores at the beginning and the end of the exam (Scale 4) were compared to control scores using a Chi-squared goodness-of-fit test.

A total examination ethogram stress score was obtained by summing the scores across all categories and steps of the exam. With the ethologically benchmarked scoring system in Table 6, Mann-Whitney *U* tests (two-tailed) compared control vs intervention groups for each part of the physical exam during visit 1 and visit 4.

The time (s) required to complete each step (1–11) of the physical examination, (recorded and summed to reflect the total time required to examine the dog), and the total time (s) elapsed from the start to the end of the physical examination were analysed using a two-way repeated measures ANOVA (visit × treatment). This total elapsed time included the pauses and breaks between steps of the physical examination.

Control and intervention venipuncture scores were compared (between groups and across time) using independent samples *t*-tests. Mann-Whitney *U* tests (two-tailed) compared the time to obtain a blood sample across groups and visits.

## Results

### Demographics and historical behaviour

Of the 58 categories, only two (one demographic and one behavioural response) showed a significant difference between the groups. Despite the randomisation protocol, the control group had more females (FI + FS) than males (MI + MC) and the intervention group having more males (MI + MC) than females (FI + FS) (Fisher exact test statistic 0.0213; *P* < 0.05).



The time spent to react when humans were squatted quietly in front of the dogs while holding a toy or treat was the only situation where dogs from the control and intervention groups differed in responses ( $t[26] = -3.33; P = 0.0026$ ). The dogs in the intervention group watched the human for at least 30 s and those in the control group watched for at least 60 s. No differences between the groups were found when the human was standing instead of squatting quietly in front of the dogs while holding a toy or treat ( $t[26] = -2.020; P = 0.054$ ).

**Comparison of in-person and video scores**

One in-person score was missing for dog 20 at visit 1. Concordance between the in-person and video clinical stress scores when dogs entered the exam room was  $r_s(109) = 0.406; P < 0.001$ . For the start of the physical exam (one in-person score was missing for dog 12 at visit 3 and dog 9 at visit 4),  $r_s$  was  $(108) = 0.586; P < 0.001$ . For the end of the physical examination (one in-person score was missing for dog 5 at visit 3, and two scores – dogs 9 and 13 – were missing for the in-person assessment at visit 4),  $r_s$  was  $(107) = 0.443; P < 0.001$ .

**Clinical stress scores**

**Entry to the clinic**

In-person clinical stress scores for entering the building (Table 5; Clinic stress scale 1) across visits decreased for 21.4% (3), did not change for 57.1% (8) and increased for 21.4% (3) of the dogs, respectively. Scores between visits 1 and 4 in the intervention group decreased for 28.6% (4), remained unchanged for 42.9% (6), and increased for 28.6% (4) of the dogs, respectively. A Chi-squared test showed no significant difference between the behaviour of the control vs intervention group across visits for entering the building ( $\chi^2(1, n = 28) = 0.190; P > 0.05$ ) (Figure 2).


**Weighing the dog**

The distribution of dogs needing various steps for weighing, by group, is shown in Table 7. There was no difference in distribution of steps needed between groups at the first visit ( $\chi^2[1,54] = 2.571; P = 0.46$ ).

There was also no significant difference between the control and intervention group for clinical stress scores at visit 1 when weighing

**Table 7.** Level of scale intervention that allowed intervention (n = 14) and control (n = 14) dogs to be weighed. No difference in distribution of scale interventions (steps) between groups at visit 1. Step 1 = bare, metal scale, flush against the wall without a lure; Step 2 = sequential addition of a blue yoga mat; Step 3 = then also moving the scale with the mat from the wall; Step 4 = adding a treat


Group	Step 1	Step 2	Step 3	Step 4
Intervention	9	5	0	0
Control	7	5	1	1




**Figure 3(a).** Video clinical stress scores once dogs (n = 28) were weighed on the scale at visits 1 and 4. No significant difference in scores between groups at visit 1; however, at visit 4 intervention (n = 14) scores were significantly lower compared to control (n = 14). The box and whisker plots show the means (x), the medians (lines), and the values for 75% of index scores (whiskers) for each group.

the dog ( $t$ -test;  $t[26] = -0.16; P = 0.87$ ). Both groups decreased their clinical stress scores when being weighed between visits 1 and 4 (Control group:  $t[26] = -2.22; P = 0.022$ , one-tailed; Intervention group:  $t$ -test;  $t[26] = -4.58$ , one-tailed;  $P = 0.00026$ ). However, at visit 4, dogs in the intervention group had a considerably lower score compared to those in the control ( $t[26] = 2.087; P = 0.023$ , one-tailed). The effect size for this finding was considerable (Cohen’s  $d = 0.79$ ) (Figure 3[a]).

Time for all four feet to rest on the scale for an accurate weight was obtained via video footage. These times across visits did not



**Figure 2.** Difference in visits 1 and 4 clinical stress score when dogs (n = 28) entered the building. No significant difference appreciated between intervention (n = 14) and control (n = 14) groups. Numbers above the bars are dogs in each group that fell within each category.



**Figure 3(b).** Time (min) to get all four feet onto the scale to be weighed across visits and groups. Dogs ( $n = 28$ ) within intervention ( $n = 14$ ) and control ( $n = 14$ ) groups both decreased the amount of time it took to get an accurate weight, with no significant difference between groups for overall time, and reduction in time across visits. The box and whisker plots show the means (x), the medians (lines), and the values for 75% of index scores (whiskers) for each group.

differ between groups ( $t[28] = -0.60$ ;  $P = 0.55$ ). Time to successfully obtain a weight decreased in both the control and intervention groups across visits (Control group: one-tailed  $t$ -test;  $t[28] = -2.60$ ;  $P = 0.011$ ; Intervention group: one-tailed  $t$ -test;  $t[28] = -3.57$ ;  $P = 0.0008$ ). There was no difference between groups in the reduction of time it took to successfully walk onto the scale by visit 4 ( $t[28] = -1.74$ ;  $P = 0.09$ ) (Figure 3[b]).

#### Entering the exam room

A Chi-squared test showed no significant difference between the video clinical stress scores of the control vs intervention group across visits 1 and 4 with respect to entering the exam room ( $\chi^2[1, n = 28] = 3.59$ ;  $P > 0.05$ ).

#### Beginning of the physical exam

A Chi-squared test showed a significant difference between the behaviour of the control vs intervention group between visits 1 and 4 with respect to the start of the physical exam ( $\chi^2[1, n = 28] = 5.60$ ;  $P = 0.018$ ) (Figure 4). At the start of the physical examination, control clinical stress scores decreased across visits for 14.3%


( $n = 2$ ), did not change for 64.3% ( $n = 9$ ) and increased for 21.4% ( $n = 3$ ) of the dogs, respectively. In the intervention group, scores between visits 1 and 4 decreased for 57.1% ( $n = 8$ ), remained unchanged for 35.7% ( $n = 5$ ), and increased for 7.1% ( $n = 1$ ), of the dogs, respectively. In the control group, of the nine scores that were unchanged, only two were for scores of 1 – the lowest score possible – at visit 1. The other seven unchanged scores in the control group represented a lack of improvement with time. One dog in the control group worsened with time with respect to the scoring of behaviour at the start of the physical exam. In the intervention group, of the five scores that were unchanged, all had scores of 1 (the lowest score) at visit 1.

#### End of the physical exam


A Chi-squared test showed a significant difference between the behaviour of the control vs intervention group between visits 1 and 4 with respect to the end of the physical exam ( $\chi^2[1, n = 28] = 7.036$ ;  $P = 0.008$ ) (Figure 5). At the end of the physical examination, the control group clinical stress scores decreased for 28.6% ( $n = 4$ ), did not change for 42.9% ( $n = 6$ ) and increased for 28.6% ( $n = 4$ ) of the dogs, between visits 1 and 4, respectively. In the intervention group, scores between visits 1 and 4 decreased for 79.0% ( $n = 11$ ), remained unchanged for 21.0% ( $n = 3$ ), and increased for 0% ( $n = 0$ ) of the dogs.

#### Descriptive behaviour

No differences were found between the behaviour of dogs from each group for any step of the physical exam during visit 1. However, at visit 4, dogs in the intervention group had significantly lower ethogram behavioural scores at Step 1 of the physical exam when the dog was stroked gently from head to base of tail ( $P = 0.048$ ), when manually manipulating lymph nodes (Step 5; submandibular, prescapular, popliteal) ( $P = 0.020$ ), during gentle abdominal palpation (Step 6) ( $P = 0.024$ ), while lifting each paw for testing placement (Step 7) ( $P = 0.002$ ), when lifting the upper lips (Step 8) ( $P = 0.024$ ), and during an examination of the eyes (Step 11; conjunctiva; cornea) ( $P = 0.018$ ) when compared to dogs in the control group. Starting at Step 5, the majority of the physical examination, except for manipulation of the ears and ear temperature, had lower stress scores at visit 4 in the intervention group. The



**Figure 4.** Clinical stress score changes across visits at the start of the exam for dogs ( $n = 28$ ) within intervention ( $n = 14$ ) and control ( $n = 14$ ) groups. Change in behaviour from visits 1 to 4 at the start of the exam is significantly different between intervention and control. Numbers above the bars represent the number of dogs in each group. \* Represents the significant effect of clinical stress scores that did not change/decrease in the control group ( $P = 0.018$ ).



**Figure 5.** Clinical stress score changes across visits at the end of the exam for dogs ( $n = 28$ ) within intervention ( $n = 14$ ) and control ( $n = 14$ ) groups. Change in behaviour from visits 1 to 4 at the end of the exam is significantly different between intervention and control. Numbers above the bars represent the number of dogs in each group. \* Represents the significant effect of decreased clinical stress in the intervention group ( $P = 0.008$ ).

total ethogram stress score during the examination for dogs in the intervention group was significantly lower than that for dogs in the control group on visit 4 ( $P = 0.030$ ) (see Table 8).

#### Duration of time to examine the dog

For the total time to examine the dog, no significant main effects were found for visit ( $F[1,15] = 1.01$ ;  $P = 0.33$ ) or for treatment ( $F[1,15] = 3.40$ ;  $P = 0.085$ ), nor was an interaction between variables present ( $F[1,15] = 1.63$ ;  $P = 0.22$ ).

#### Total duration of time to complete physical examination

For the total time to complete the physical examination, significant main effects were found for visit ( $F[1,20] = 5.15$ ;  $P = 0.034$ ) and for treatment ( $F[1, 20] = 4.78$ ;  $P = 0.041$ ). A significant visit  $\times$  treatment interaction was also present ( $F[1,20] = 11.53$ ;  $P = 0.003$ ) (Figure 6). Time to complete the exam was less for visit 4 than for visit 1 for dogs in the control group, but greater than visit 1 for dogs in the intervention group. When time to complete the physical examination at visit 4 was compared, dogs in the intervention group had longer exams compared to those in the control group ( $t[20] = -4.130$ ,  $p_{\text{tukey}} = 0.003$ ). The difference between visits 1 and 4 was an average decrement of 30 s for control group dogs (average time of exam at visit 4 = 321 s, or roughly 5 min), and an average increase of 139.4 s (2.5 min) for intervention group dogs (average time of exam at visit 4 = 468 s, or roughly 8 min), which included pauses to adapt to the dog's behaviour.

#### Venipuncture

Scores for venipuncture are graphed in Figure 7(a). As the scorer had to be able to see the needle both enter and exit the vein after the sample was obtained to determine time elapsed, some dogs were lost from our sample. The control group did not differ from the intervention group for scores at visit 1 ( $t[18] = 0.3133$ ;  $P = 0.76$ ). The intervention group score was not significantly different for visits 1 and 4 ( $t[18] = -0.16$ ;  $P = 0.44$ , one-tailed), but the control group score significantly increased between visits 1 and 4 ( $t[18] = 2.29$ ;  $P = 0.025$ , one-tailed). Accordingly, when comparing the scores for visit 4, the intervention group showed significantly lower scores than the control group ( $t[18] = 1.912$ ;  $P = 0.037$ , one-tailed).

There were no significant pair-wise comparisons across group or visit for time to obtain blood sample (Mann-Whitney  $U$  test: control vs intervention visit 1 ( $P = 0.81$ ); control visit 1 vs 4 ( $P = 0.41$ ); intervention visit 1 vs 4 ( $P = 0.154$ ); control vs intervention visit 4 ( $P = 0.052$ ). However, when assessing how many dogs in each group (control vs intervention) experienced a decrease in the amount of time necessary for venipuncture from visit 1 to 4, only 2/10 control dogs experienced decreased time for venipuncture, but 6/9 dogs in the intervention group did so ( $z = -2.057$ ;  $P = 0.02$ ) (Figure 7[b]).

#### Taking a treat in the parking lot at the end of the visit


None of the dogs in the control group declined a treat on any visit (56/56 treat opportunities accepted). One dog in the intervention group declined a parking lot treat on all four visits, and one declined a treat at visit 3, only (52/56 treat opportunities accepted). There was no difference between the groups in treat acceptance (Fisher exact test;  $P = 0.12$ ), despite the provisioning of treats to dogs in the intervention group at most of the major stages of the experimental process.

#### Discussion

Our study objectives were to compare the control and intervention group for differences in outcomes using two different handling protocols. To do this, we used both WDQ-Pet scores and the in-person and video scoring of behavioural responses, with the intent to ultimately determine the effect of targeted interventions on behaviours indicating fear and/or distress in canine patients during veterinary procedures. Numerous studies have evaluated the effect veterinary procedures and exams have on canine patients (Hernander 2008; Döring *et al.* 2009; Mariti *et al.* 2017; Stellato *et al.* 2019a). The literature assessing the application of low-stress interventions during veterinary visits and how these may affect canine patients' responses to veterinary care is not as robust and only subsets of collaborative care have previously been investigated in dogs. Van Oostrom and Knowles (2018) found that applying topical analgesia prior to venipuncture decreased stress-related behaviours including withdrawal and/or defensive movements. Stellato *et al.* (2019b) and Wess *et al.* (2022) found only mild effects


**Table 8.** Behavioural scores for body regions and behaviours of dogs (n = 28) indicative of fear/stress responses during each step of the physical examination. At visit 4, intervention (n = 14) behavioural scores were significantly lower than control (n = 14) at Steps 1, 5, 6, 7, 8, and 11, in addition to a significantly lower total comprehensive stress score. Exam Stages = Sequential steps of the physical examination performed on each dog (see Table 3 for details of each step); N = number; IQR = Interquartile Range; r = effect size (Rank biserial correlation). \* P < 0.05 compared to control, with exam step highlighted. P-values reported only for significant findings

Exam	Treatment	Visit 1		Visit 4		r	P
		N	Median (IQR)	N	Median (IQR)		
1. Stroke	<b>Control</b>	<b>14</b>	<b>15.0 (10.3)</b>	<b>13</b>	<b>13.0 (4.00)</b>		
	<b>Intervention</b>	<b>14</b>	<b>11.0 (8.50)</b>	<b>13</b>	<b>8.0 (2.00)*</b>	<b>0.4556</b>	<b>0.048</b>
2. Pulse	Control	14	16.0 (5.25)	13	15.0 (5.00)		
	Intervention	14	15.5 (6.00)	12	10.0 (9.00)		
3. Lidocaine/ Touch	Control	13	18.0 (5.00)	12	15.0 (4.25)		
	Intervention	14	14.5 (3.00)	13	16.0 (8.00)		
4. Auscultation Chest	Control	14	18.5 (3.00)	14	17.0 (2.75)		
	Intervention	14	17.5 (3.00)	13	15.0 (7.00)		
5. Lymph Nodes	<b>Control</b>	<b>13</b>	<b>18.0 (8.00)</b>	<b>13</b>	<b>18.0 (3.00)</b>		
	<b>Intervention</b>	<b>14</b>	<b>16.0 (7.75)</b>	<b>12</b>	<b>12.5 (4.50)*</b>	<b>0.5513</b>	<b>0.02</b>
6. Abdominal Palpation	<b>Control</b>	<b>14</b>	<b>18.5 (5.25)</b>	<b>14</b>	<b>17.0 (4.75)</b>		
	<b>Intervention</b>	<b>14</b>	<b>16.5 (4.50)</b>	<b>12</b>	<b>13.9 (7.00)*</b>	<b>0.5238</b>	<b>0.024</b>
7. Paws	<b>Control</b>	<b>14</b>	<b>20.0 (5.00)</b>	<b>13</b>	<b>18.0 (4.00)</b>		
	<b>Intervention</b>	<b>14</b>	<b>18.0 (5.25)</b>	<b>12</b>	<b>14.0 (3.00)*</b>	<b>0.7179</b>	<b>0.002</b>
8. Lift Lip	<b>Control</b>	<b>14</b>	<b>19.0 (4.00)</b>	<b>11</b>	<b>18.0 (3.50)</b>		
	<b>Intervention</b>	<b>14</b>	<b>16.5 (5.50)</b>	<b>12</b>	<b>14.0 (1.75)*</b>	<b>0.5606</b>	<b>0.024</b>
9. Ear Exam	Control	14	19.5 (3.00)	13	17.0 (6.00)		
	Intervention	14	17.5 (5.00)	12	13.5 (1.25)		
10. Ear Temperature	Control	14	18.5 (4.75)	13	17.0 (5.00)		
	Intervention	14	21.0 (2.50)	13	16.0 (3.00)		
11. Eyes	<b>Control</b>	<b>13</b>	<b>17.0 (6.00)</b>	<b>13</b>	<b>16.0 (3.00)</b>		
	<b>Intervention</b>	<b>13</b>	<b>19.0 (5.00)</b>	<b>10</b>	<b>11.5 (4.75)*</b>	<b>0.5923</b>	<b>0.018</b>
Total Comprehensive Stress Score	<b>Control</b>	<b>12</b>	<b>202 (50.80)</b>	<b>9</b>	<b>190 (24.00)</b>		
	<b>Intervention</b>	<b>13</b>	<b>188 (42.00)</b>	<b>8</b>	<b>126 (37.00)*</b>	<b>0.6390</b>	<b>0.03</b>




**Figure 6.** Time (s) to complete the physical exam. Time to complete the physical exam at visit 4 was significantly longer for dogs in the intervention group (n = 14) compared to the control (n = 14). Values represent mean (± SEM). \* Denotes significant difference (P = 0.003; see text).





**Figure 7(a).** Venipuncture video behaviour scores across visits for dogs ( $n = 18$ ) within intervention ( $n = 9$ ) and control ( $n = 9$ ) groups. No significant differences were observed between intervention and control scores at visit 1. Across visits, no significant difference in intervention scores, but control scores significantly increased. Visit 4 intervention scores were significantly lower than the control. The box and whisker plots show the means (x), medians (lines), and the values for 75% of index scores (whiskers) for each group.



**Figure 7(b).** Time (s) for blood to be drawn across visits for dogs ( $n = 19$ ) within intervention ( $n = 9$ ) and control ( $n = 10$ ) groups. No significant difference across visits or group for time to obtain a blood sample; however, significantly more intervention dogs (6/9) had a reduction in time from visits 1 to 4 compared to control dogs (2/10). The box and whisker plots show the means (x), medians (lines), and the values for 75% of index scores (whiskers) for each group.

of practising physical exam steps at home, but these studies used only two visits, a baseline and intervention visit. Changing behaviour takes time and these designs did not test what could happen over time with enhanced exposure and practice which is, after all, what we expose dogs to in fearful veterinary circumstances and one reason why they worsen with frequency of visits (Döring *et al.* 2009; Stellato *et al.* 2021). Finally, dogs were more willing to approach and take treats from a veterinarian not wearing a white laboratory coat than one not doing so (Fanucchi 2022). Dogs do not experience the components of a physical exam as individual stressors. All components interact to shape the dog's final response to veterinary care. To the authors' knowledge, no other study has investigated the combined effect of multiple low-stress handling techniques in canine patients over time.

The WDQ-Pet questionnaire was used to assess whether dogs were randomly assigned to each group with respect to their behavioural profiles entering the study. Of the 58 demographic and behavioural categories assessed by the WDQ-Pet (the aggression and fear screens each have 53 categories of assessment that sum to a final score), 56 did not differ between the control and intervention groups. Despite an adequate randomisation protocol, dogs in the intervention group were overwhelmingly male ( $n = 11/14$ ) while dogs in the control group were overwhelmingly female ( $n = 9/14$ ). Rare statistical findings are, by their very nature, rare. This rare

outcome prohibited the investigation of any putative effects of sex on outcomes. However, the WDQ-Pet is an extremely detailed and robust assessment tool for general behaviours of dogs across broad categories and specific behaviours in provocative categories (e.g. walking in the rain, experiencing an unpredictable loud noise), and these groups of dogs – despite the sex distribution – did not differ in their scores for aggression, fear, environmental reactivity, reactivity during training, or SAIR (Separation Anxiety Intensity Rank) or AIR (Anxiety Intensity Rank) scores, suggesting that there were no effects of sex on these historical behaviours.

For no behavioural category was there any difference between the dogs in the control and intervention group except when a human crouched in front of the dog with a treat or toy. In an assessment of 58 broad categories, it is unlikely that a difference in focus when crouched (but not standing) of 30 s (over a range of times of 0 s/does not focus, a few seconds, 10 s, 30 s, 60 s, more than 60 s) in one setting would be clinically meaningful. While few studies examine behavioural histories and patterns of dogs entering into experiments with this degree of rigor, it is safe to say that with the exception of the statistically significant but rare finding of differences in each sex by group, the two groups were well randomised and evenly distributed in terms of demographics, training history, and historical behavioural and environmental responses when entering the study.

In-person assessments of the various clinic stress level scores correlated with the scores taken from the videos with at least moderate concordance. If the veterinary team pays attention, standardises their assessment (e.g. one front paw fully in the room across the threshold) and sets the criteria amongst themselves, they will be able to gain valuable information about dogs' responses to specific clinical stressors just using a tick-sheet. This is valuable and practical knowledge for those wishing to integrate these findings into their daily practice.

There were no differences between the control and intervention groups entering the hospital and the exam room between visits 1 and 4, and these scores were relatively low for most dogs. However, there were significant, and likely behaviourally important, differences across time for the groups for stress assessments made both at the beginning and end of the exam. Intervention dogs had lower exam clinical stress scores across visits and were more likely to decrease their scores over time at both the start and end of the exam compared to control dogs. This finding shows a clear effect of the altered physical and behavioural approach that an adaptive, collaborative examination entails. That the score at the beginning of the exam was significantly lower at visit 4 suggests that dogs in the intervention group were not anticipating an unpleasant experience. That intervention group scores were lower at the end of the exam at visit 4 suggests that dogs did not have an unpleasant experience.

Obtaining a weight may be viewed as a benign procedure; however, this human viewpoint may differ from what the dog experiences. Instead, we should consider that weighing the dog may be the first manipulation where dogs begin to exhibit behaviours associated with responses to stressors (Hernander 2008). Stressors are additive and the longer we can delay their appearance and the more we can lower their effects, the better for our patients (Main 2022).

Measurements of how long it took for a veterinarian to perform a task are often used as indicators of how easy it was to ostensibly accomplish the task. Stress scores, on the other hand, indicate how easy it was for the dog to experience to experience the task. Here, both groups had a reduction in the time it took to obtain an accurate

weight from visits 1–4, with no difference in the time decrement by visit 4. The lack of time difference at visit 4 suggests that the adaptive, step-wise design we used for weighing dogs in the control group helped to ensure that they successfully went onto the scale. Interestingly, dogs in the control group may have needed additional interventions to be weighed across all four visits but the intervention was always at an equal or lower step level (1–4) than it was at visit 1. When assessing the initial stress scores for dogs in the intervention and control groups, no difference was found between groups when being weighed or in the number of step-wise interventions needed to get them to the point of being weighed at visit 1. When being weighed, dogs in both groups experienced a decrease in their recorded stress scores from visits 1 to 4; however, dogs in the intervention group had a high probability of a significantly lower clinical stress score at visit 4 than dogs in the control group. Furthermore, the effect size of this result is substantial. That the control group complied with being weighed and in the same time as the intervention group by visit 4, but had higher stress scores at visit 4 than those in the intervention group (Figure 4), is a cautionary warning to the veterinary staff. While time may be related to money and how rushed staff feel, it is critical to assess not just that the patient is complying, but also how the patient views the cost of that compliance. Clearly, the dogs in the intervention group experienced less stress being weighed at the last visit, although time costs to staff of weighing dogs in both groups was the same here. If dogs become more fearful across visits, the cost of procedures to the veterinary staff in the currencies of time and job satisfaction may increase. These data show that this pattern need not be the case. In fact, one dog in the intervention group who had been reluctant to be weighed at the first visit, bounded onto the scale at the last visit and when told he was done, did it again (see supplementary video in [Supplementary materials](#)).

We cannot know whether the relatively high stress scores shown by dogs in the control group at visit 4 are due to the placement of the weighing procedure prior to the physical examination or due to the overall, unrelieved stress level of the examination associated with standard handling. The combination of outcomes suggests that when the veterinary staff change their behaviours to respond to the dog's signals and needs, everyone benefits.

Both venipuncture and physical exam assessments had a lack of continuous, interpretable video for some dogs. This was a field study and, as such, showed inherent limitations.

The assessment of time and benchmarked, ethogram stress scores during the step-wise physical examination were revealing. While dogs in the control group did not differ from those in the intervention group at visit 1, in terms of time to complete the physical examination, changes were seen by visit 4 that depended on both treatment and visit. Dogs in the control group experienced less change when comparing the total elapsed time for physical exam between visits 1 and 4, with an average decrease of 30 s in the time for the exam. In contrast, dogs in the intervention group experienced an average increase of 139 s (2.5 min) in the time for the exam across visits. Since pauses between exam steps were included in the measurement of time for the complete physical exam, it is likely this increase in time for the dogs in the intervention group reflected the adaptive nature of a truly collaborative exam.

Pairing these data with those from the benchmarked, ethogram stress assessment of the step-wise and total stress scores for the exam reveals an important pattern. While the exam may have taken longer for dogs in the intervention group, the total ethogram stress score for the combined steps of the exam for dogs in the intervention group was significantly lower than that for dogs in the control

group on visit 4. Furthermore, when the individual steps of exam were analysed with respect to the ethogram stress scores, dogs in the intervention group at visit 4 had significantly lower stress scores when stroked gently from head to base of tail, when palpating lymph nodes, during gentle abdominal palpation, while lifting each paw for testing placement, when lifting the upper lips, and during an examination of the eyes (conjunctiva; cornea) compared to dogs in the control group. Starting at step 5, the majority of the steps in the physical examination had lower stress scores at visit 4 in the intervention vs the control group. This finding is somewhat surprising with respect to lifting the paws, given the results of another study (Jokela *et al.* 2023) that found picking up of feet to be provocative for a large number of dogs.

Venipuncture is a potentially aversive and stressful procedure for dogs (Chebroux *et al.* 2015; van Oostrom & Knowles 2018) and, by extension, may be upsetting for their owners and the veterinary team. Generally, more than one veterinary team member is required, dogs' veins are held off physically with a tourniquet or by hand, and the dog is physically restrained so that they cannot move. No topical anaesthetic is generally used. This description matches the methodology used here for the control group. When scores at the end of the venipuncture procedure were compared, the intervention group had significantly lower scores than the control group at visit 4, but due to a significant increase in scores for the control group between visits 1 and 4. These data may indicate that dogs in the control group became increasingly sensitised to the entire examination and venipuncture procedure, a finding that our physiology data, and specifically the stress response index evaluating heart rate, serum cortisol and neutrophil lymphocyte ratio, also suggested (Squair *et al.* 2023).

The amount of time from inserting the needle to removing it after the sample is obtained is a rough, but informative estimate of how many adjustments and steps were needed by the veterinary team to obtain the sample from the dog. While the control group experienced standard hospital procedures, there was also an effort to use no force beyond mild restraint, and instead readjust the behaviour of the researchers conducting the visit. None of these dogs experienced extreme force when handling (such as multiple people forcibly holding them down). It is interesting that, regarding time, there were no significant pair-wise comparisons across group or visit, but when number of dogs experiencing decreased venipuncture times at visit 4 was compared with visit 1, the intervention group dogs were strongly favoured. Given the small sample size, caution in interpreting this significant result is urged, but it may be a good representation of how the dogs perceived the combined exam and venipuncture strategies. Neither initial scores nor times for venipuncture differed between groups, suggesting that this result was due to the differential handling and not prior experience. Additionally, the results suggest that using the intervention approach for drawing blood is viewed more favourably by the dog, while not taking more time for the veterinary staff compared to standard methods. It is also important to remember that venipuncture was the last procedure that the dogs experienced prior to being petted, told they were good, and offered a treat. As such, it may represent a cumulative effect of the entire experience for the dog and may have taken a borderline concerned dog and pushed him or her into the realm of a greater stress and distress response.

### Study limitations

There are limitations to this type of study, which include the inability to be completely blinded. While the human participants

did not witness other owners' tests, there were physical differences in the study protocol (e.g. white coats, blue mats, Lickimats®, techniques for drawing blood) which indicated assignment group and could have been informative to participants. As far as was possible great care was taken to blind researchers and limit bias. The laboratory technicians who processed the blood samples were blinded to the group status of the samples. The student who helped analyse videos was blinded to treatment and the order of the videos and had not been involved in the study. Videos were coded so that the order of the visits was not known. Additionally, the owners did not know what group they were in until the end of the study and many of them in each group were surprised when we unblinded their dog's group for them.

This was a 'real-world' field study, and these can be difficult to manage since owners and patients are involved and the situation within which behaviours are being evaluated remains dynamic. The intensive nature of the evaluations means sample sizes are small, and they became smaller videos were being relied upon because, despite three cameras being available, continuous observations were not possible. This is especially true for the data on venipuncture, a very intimate procedure. While we found a significant difference between groups, the sample size is tiny and subject to errors of false attribution (Type I error). Such results would require to be replicated in larger studies to have any stand-alone meaning. However, as one part of a study that evaluated numerous aspects of the dogs' responses to stressors, it is helpful to note that even the venipuncture measures follow the same direction of significance as the other measures. While one should worry were this not the case, it is reassuring that even the messiest measures converged into the same pattern.

### Animal welfare implications and conclusion

Within veterinary medicine, signs of fear and distress have been normalised, which negatively affects the patients' care, health, and welfare. The behaviour scores and ethogram score analysis described reveal a strong pattern of improvement in canine patients' well-being when participating in collaborative care procedures. This data, combined with the physiological data previously published (Squair *et al.* 2023), strongly suggest that patients receiving low-stress interventions during veterinary visits have a greater reduction in stress responses over time compared to patients not experiencing those interventions. Overall, by meeting the patient's needs, and reducing extraneous duress, there can be improvement in the welfare of dogs within the veterinary clinic setting. Furthermore, our findings have widespread implications for members of the veterinary community, revealing that simple adjustments made in handling patients, can lead to significant improvements in animals' well-being during visits to the veterinarian. Ultimately, these changes result in providing an improved standard of care.

**Supplementary material.** The supplementary material for this article can be found at <http://doi.org/10.1017/awf.2024.37>.

**Data availability statement.** The data presented in this study are available on request from KLO. The data are not publicly available due to the ethical agreement with the participants that only anonymised, summary data would be publicly presented. Anyone wishing anonymised summary data can contact us.

**Acknowledgements.** The authors would like to thank Lickimat® for donating their product for this research study, and the student volunteers and summer research students for their assistance in the examinations and video recording.

Tsukasa Iwamoto, an AVC veterinary student, helped analyse the video data when she was a summer VetSRA student with KLO's group. We would also like to thank the dog owners and their dogs for their participation for this study.

This research was fully funded by a grant from the Sir James Dunn Animal Welfare Centre at the Atlantic Veterinary College to KLO (A615846). UPEI (including the G Murray and Hazel Scholarship and Zoetis Graduate Student Research Award), the American College of Veterinary Behaviorists (ACVB) Veterinary Behavior Symposium (VBS) (including the 2022 Fear Free Award and the 2023 RK Anderson Resident Award), and VIN provided additional support to CS through scholarships and grants.

**Competing interest.** None.

### References

- Beerda B, Schilder MBH, van Hooff JanARAM and de Vries HW 1997 Manifestations of chronic and acute stress in dogs. *Applied Animal Behaviour Science* **52**: 307–319. [https://doi.org/10.1016/S0168-1591\(96\)01131-8](https://doi.org/10.1016/S0168-1591(96)01131-8)
- Broom DM 2016 Considering animals' feelings: Précis of Sentience and animal welfare (Broom 2014). *Animal Sentience* **1**. <https://doi.org/10.51291/2377-7478.1015>
- Chebrox A, Leece EA and Brearley JC 2015 Ease of intravenous catheterisation in dogs and cats: a comparative study of two peripheral catheters. *Journal of Small Animal Practice* **56**: 242–246. <https://doi.org/10.1111/jsap.12318>
- Cobos B, Haskard-Zolnieriek K and Howard K 2015 White coat hypertension: improving the patient-health care practitioner relationship. *Psychology Research and Behavior Management*: **133**. <https://doi.org/10.2147/PRBM.S61192>
- Crisi PE, De Santis F, Giordano MV, Cerasoli I, Colucci F, Di Tommaso M and Luciani A 2021 Evaluation of eutectic lidocaine/prilocaine cream for jugular blood sampling in cats. *Journal of Feline Medicine and Surgery* **23**: 185–189. <https://doi.org/10.1177/1098612X20917309>
- Dhabhar FS 2009 Enhancing versus suppressive Effects of Stress on Immune Function: Implications for Immunoprotection and Immunopathology. *Neuroimmunomodulation* **16**: 300–317. <https://doi.org/10.1159/000216188>
- Domjan M 2015 Elicited behavior, habituation, and sensitization. In: Domjan M (ed) *The Principles of Learning and Behavior, Seventh Edition* pp 29–58. Cengage Learning: Stamford, CT, USA.
- Döring D, Roscher A, Scheipl F, Küchenhoff H and Erhard MH 2009 Fear-related behaviour of dogs in veterinary practice. *The Veterinary Journal* **182**: 38–43. <https://doi.org/10.1016/j.tvjl.2008.05.006>
- Dreschel NA 2010 The effects of fear and anxiety on health and lifespan in pet dogs. *Applied Animal Behaviour Science* **125**: 157–162. <https://doi.org/10.1016/j.applanim.2010.04.003>
- Edwards PT, Smith BP, McArthur ML and Hazel SJ 2019 Fearful Fido: Investigating dog experience in the veterinary context in an effort to reduce distress. *Applied Animal Behaviour Science* **213**: 14–25. <https://doi.org/10.1016/j.applanim.2019.02.009>
- Fanucchi L 2022 Dogs' preference for white coat versus no white coat when offered a food reward in the exam room. *Open Access Journal of Veterinary Science & Research* **7**: 1–5. <https://doi.org/10.23880/oajvsr-16000219>
- Godbout M and Frank D 2011 Persistence of puppy behaviors and signs of anxiety during adulthood. *Journal of Veterinary Behavior* **6**: 92. <https://doi.org/10.1016/j.jveb.2010.08.023>
- Godbout M, Palestrini C, Beauchamp G and Frank D 2007 Puppy behavior at the veterinary clinic: A pilot study. *Journal of Veterinary Behavior* **2**: 126–135. <https://doi.org/10.1016/j.jveb.2007.06.002>
- Gouin J-P and Kiecolt-Glaser JK 2011 The Impact of psychological stress on wound healing: Methods and mechanisms. *Immunology and Allergy Clinics of North America* **31**: 81–93. <https://doi.org/10.1016/j.jiac.2010.09.010>
- Hauser H, Campbell S, Korpivaara M, Stefanovski D, Quinlan M and Siracusa C 2020 In-hospital administration of dexmedetomidine oromucosal gel for stress reduction in dogs during veterinary visits: A randomized, double-blinded, placebo-controlled study. *Journal of Veterinary Behavior* **39**: 77–85. <https://doi.org/10.1016/j.jveb.2020.05.002>
- Hefler L, Grimm C, Leodolter S and Tempfer C 2004 To butterfly or to needle: The pilot phase. *Annals of Internal Medicine* **140**: 935–936. <https://doi.org/10.7326/0003-4819-140-11-200406010-00027>



- Hernander L** 2008 *Factors influencing dogs' stress level in the waiting room at a veterinary clinic*. Student Report, Swedish University of Agricultural Sciences, Department of Animal Environment and Health, Skara, Sweden.
- Jokela F, Mercier P, Honeckman L, Dunham AE and Overall KL** 2023 Sensitivity of benchmarked behavioral assays for distress – should we attend to certain behaviors during exams? *Journal of Veterinary Behavior*. <https://doi.org/10.1016/j.jveb.2023.06.012>
- Korpivaara M, Huhtinen M, Aspegren J and Overall K** 2021 Dexmedetomidine oromucosal gel reduces fear and anxiety in dogs during veterinary visits: A randomised, double-blind, placebo-controlled clinical pilot study. *Veterinary Record* **189**. <https://doi.org/10.1002/vetr.832>
- Lind AK, Hydrbring-Sandberg E, Forkman B and Keeling LJ** 2017 Assessing stress in dogs during a visit to the veterinary clinic: Correlations between dog behavior in standardized tests and assessments by veterinary staff and owners. *Journal of Veterinary Behavior: Clinical Applications and Research* **17**: 24–31. <https://doi.org/10.1016/j.jveb.2016.10.003>
- Lloyd J** 2017 Minimising stress for patients in the veterinary hospital: Why it is important and what can be done about it. *Veterinary Sciences* **4**: 22. <https://doi.org/10.3390/vetsci4020022>
- Main K** 2022 Canine stress in the veterinary environment. *Companion Animal* **27**: 2–6. <https://doi.org/10.12968/coan.2022.0020>
- Mandese WW, Griffin FC, Reynolds PS, Blew AC, Deriberprey AS and Estrada AH** 2021 Stress in client-owned dogs related to clinical exam location: a randomised crossover trial. *Journal of Small Animal Practice* **62**: 82–88. <https://doi.org/10.1111/jsap.13248>
- Marino CL, Cober RE, Iazbik MC and Couto CG** 2011 White-coat effect on systemic blood pressure in retired racing greyhounds: Blood pressure in retired racing greyhounds. *Journal of Veterinary Internal Medicine* **25**: 861–865. <https://doi.org/10.1111/j.1939-1676.2011.00735.x>
- Mariti C, Pierantoni L, Sighieri C and Gazzano A** 2017 Guardians' perceptions of dogs' welfare and behaviors related to visiting the veterinary clinic. *Journal of Applied Animal Welfare Science* **20**: 24–33. <https://doi.org/10.1080/10888705.2016.1216432>
- Mariti C, Raspanti E, Zilocchi M, Carlone B and Gazzano A** 2015 The assessment of dog welfare in the waiting room of a veterinary clinic. *Animal Welfare* **24**: 299–305. <https://doi.org/10.7120/09627286.24.3.299>
- Mellor DJ** 2016 Updating animal welfare thinking: Moving beyond the 'Five Freedoms' towards 'A Life Worth Living'. *Animals* **6**: 21. <https://doi.org/10.3390/ani6030021>
- Moberg GP** 2000 Biological response to stress: implications for animal welfare. In: Mench JA and Moberg GP (eds) *The Biology of Animal Stress: Basic Principles and Implications for Animal Welfare* pp 1–21. CABI Books: Wallingford, UK.
- Norman G** 2010 Likert scales, levels of measurement and the "laws" of statistics. *Advances in Health Sciences Education* **15**: 625–632. <https://doi.org/10.1007/s10459-010-9222-y>
- Oliveira RL, Soares JH, Moreira CM, Silva CP, Carrasco LP and Souza HJ** 2019 The effects of lidocaine–prilocaine cream on responses to intravenous catheter placement in cats sedated with dexmedetomidine and either methadone or nalbuphine. *Veterinary Anaesthesia and Analgesia* **46**: 492–495. <https://doi.org/10.1016/j.vaa.2019.03.005>
- Overall K, Hamilton S and Chang M** 2006 Understanding the genetic basis of canine anxiety: Phenotyping dogs for behavioral, neurochemical, and genetic assessment. *Journal of Veterinary Behavior-Clinical Applications and Research* **1**: 124–141. <https://doi.org/10.1016/j.jveb.2006.09.004>
- Overall KL** 2013a Abnormal canine behaviors and behavioral pathologies not primarily involving pathological aggression. *Manual of Clinical Behavioral Medicine for Dogs and Cats, First Edition* pp 231–309. Elsevier: St Louis, MO, USA.
- Overall KL** 2013b Embracing behavior as a core discipline: Creating the behavior-centered practice. *Manual of Clinical Behavioral Medicine for Dogs and Cats, First Edition* pp 2–44. Elsevier: St Louis, MO, USA.
- Overall KL** 2019 Evidence-based paradigm shifts in veterinary behavioral medicine. *Journal of the American Veterinary Medical Association* **254**: 798–807. <https://doi.org/10.2460/javma.254.7.798>
- Overall KL, Dunham AE, Scheifele P and Sonstrom Malowski K** 2019 Fear of noises affects canine problem solving behavior and locomotion in standardized cognitive tests. *Applied Animal Behaviour Science* **221**: 104863. <https://doi.org/10.1016/j.applanim.2019.104863>
- Perego R, Proverbio D and Spada E** 2014 Increases in heart rate and serum cortisol concentrations in healthy dogs are positively correlated with an indoor waiting-room environment. *Veterinary Clinical Pathology* **43**: 67–71. <https://doi.org/10.1111/vcp.12118>
- Riddick L** 2023 Paediatric blood sampling: how to improve your chances of getting it right. *Paediatrics and Child Health* **33**: 114–117. <https://doi.org/10.1016/j.paed.2023.01.006>
- Riemer S, Heritier C, Windschnurer I, Pratsch L, Arhant C and Affenzeller N** 2021 A review on mitigating fear and aggression in dogs and cats in a veterinary setting. *Animals* **11**: 158. <https://doi.org/10.3390/ani11010158>
- Schubert C, Lambertz M, Nelesen RA, Bardwell W, Choi J-B and Dimsdale JE** 2009 Effects of stress on heart rate complexity—A comparison between short-term and chronic stress. *Biological psychology* **80**: 325–332. <https://doi.org/10.1016/j.biopsycho.2008.11.005>
- Squair C, Proudfoot K, Montelpare W and Overall KL** 2023 Effects of changing veterinary handling techniques on canine behaviour and physiology part 1: Physiological measurements. *Animals* **13**: 1253. <https://doi.org/10.3390/ani13071253>
- Stanford TL** 1981 Behavior of dogs entering a veterinary clinic. *Applied Animal Ethology* **7**: 271–279. [https://doi.org/10.1016/0304-3762\(81\)90083-3](https://doi.org/10.1016/0304-3762(81)90083-3)
- Steffey MA, Griffon DJ, Risselada M, Scharf VF, Buote NJ, Zamprogno H and Winter AL** 2023 Veterinarian burnout demographics and organizational impacts: a narrative review. *Frontiers in Veterinary Science* **10**: 1184526. <https://doi.org/10.3389/fvets.2023.1184526>
- Steimer T** 2002 The biology of fear- and anxiety-related behaviors. *Dialogues in Clinical Neuroscience* **4**: 231–249.
- Stellato AC, Flint HE, Dewey CE, Widowski TM and Niel L** 2021 Risk-factors associated with veterinary-related fear and aggression in owned domestic dogs. *Applied Animal Behaviour Science* **241**: 105374. <https://doi.org/10.1016/j.applanim.2021.105374>
- Stellato AC, Hoffman H, Gowland S, Dewey CE, Widowski TM and Niel L** 2019a Effect of high levels of background noise on dog responses to a routine physical examination in a veterinary setting. *Applied Animal Behaviour Science* **214**: 64–71. <https://doi.org/10.1016/j.applanim.2019.03.009>
- Stellato AC, Jajou S, Dewey CE, Widowski TM and Niel L** 2019b Effect of a standardized four-week desensitization and counter-conditioning training program on pre-existing veterinary fear in companion dogs. *Animals* **9**: 767. <https://doi.org/10.3390/ani9100767>
- Sullivan GM and Artino AR** 2013 Analyzing and interpreting data from Likert-type scales. *Journal of Graduate Medical Education* **5**: 541–542. <https://doi.org/10.4300/JGME-5-4-18>
- van Oostrom H and Knowles TG** 2018 The clinical efficacy of EMLA cream for intravenous catheter placement in client-owned dogs. *Veterinary Anaesthesia and Analgesia* **45**: 604–608. <https://doi.org/10.1016/j.vaa.2018.03.009>
- Volk JO, Felsted KE, Thomas JG and Siren CW** 2011 Executive summary of the Bayer veterinary care usage study. *Journal of the American Veterinary Medical Association* **238**: 1275–1282. <https://doi.org/10.2460/javma.238.10.1275>
- Wess L, Böhm A, Schützinger M, Riemer S, Yee JR, Affenzeller N, and Arhant C** 2022 Effect of cooperative care training on physiological parameters and compliance in dogs undergoing a veterinary examination—A pilot study. *Applied Animal Behaviour Science* **250**: 105615. <https://doi.org/10.1016/j.applanim.2022.105615>
- Westlund K** 2015 To feed or not to feed: Counterconditioning in the veterinary clinic. *Journal of Veterinary Behavior* **10**: 433–437. <https://doi.org/10.1016/j.jveb.2015.05.008>
- World Health Organisation (WHO)** 2010 *Paediatric and neonatal blood sampling: WHO guidelines on Drawing Blood. Best Practices in Phlebotomy*. World Health Organisation: Geneva, Switzerland.