

REMARK ON THE TRICOMI EQUATION

TADATO MATSUZAWA

§1. As an application of the Carleman-type estimation Hörmander [4], p. 221, has proved the following:

A solution (distribution) of the Tricomi equation

$$\frac{\partial^2 u}{\partial t^2} + t \frac{\partial^2 u}{\partial x^2} = 0$$

in an open set Ω in $R_{x,t}^2$ belongs to $C^\infty(\Omega)$ if it is in $C^\infty(\Omega_-)$ where $\Omega_- = \{(x, t); (x, t) \in \Omega, t < 0\}$.

In this note we shall consider the same problem for the inhomogeneous Tricomi equation

$$\frac{\partial^2 u}{\partial t^2} + t \frac{\partial^2 u}{\partial x^2} = f(x, t)$$

in a different manner. The existence of the solution in the generalized sense is well known. Furthermore we shall consider the propagation of analyticity. More precisely, the solution u is analytic in Ω if it is analytic in Ω_- and if $f(x, t)$ is analytic in Ω (Theorem 3.1). We shall use the results of [2] and [5] in the proof.

§2. The following theorem is obtained from the results of Berezin [2].

THEOREM 2.1. *Consider the following (backward) Cauchy problem:*

$$(2.1) \quad u_{tt} + tu_{xx} = f(x, t) \quad \text{in } D,$$

$$(2.2) \quad u(x, 0) = \varphi(x), \quad u_t(x, 0) = \psi(x) \quad \text{in } a \leq x \leq b$$

where D denotes a domain in the region $t < 0$ bounded by characteristics passing through $(a, 0)$ and $(b, 0)$, $(a < b)$. Assume $f(x, t)$ and $f_x(x, t)$ are continuous in \bar{D} and the initial data $\varphi(x), \psi(x)$ are thrice continuously differentiable in $[a, b]$. Then there exists one and only one solution

Received April 21, 1972.

$u(x, t)$ of the problem (2.1), (2.2) having continuous second derivatives in \bar{D} . Furthermore, if $f(x, t)$ and $\varphi(x), \psi(x)$ are infinitely differentiable in \bar{D} and in $[a, b]$ respectively, then the solution $u(x, t)$ is an infinitely differentiable function in \bar{D} .

By virtue of Theorem 2.1 it is shown that there exists a fundamental solution $E(x, t)$ for the backward Cauchy problem for the equation $Lu = u_{tt} + tu_{xx} = 0$. That is, there exists a distribution $E(x, t)$ in the region $t \leq 0$ such that

$$(2.3) \quad LE = E_{tt} + tE_{xx} = 0 \quad \text{for } t < 0,$$

$$(2.4) \quad E(x, 0) = 0, \quad E_t(x, 0) = \delta_x.$$

In fact, take $f(x, t) = 0, \varphi(x) = 0$ and

$$\psi(x) = \begin{cases} 0 & x < 0 \\ x^4/4! & x \geq 0 \end{cases}$$

in Theorem 2.1. Then there exists a solution $v(x, t)$ for the problem (2.1), (2.2) with these data having second continuous derivatives in the region $t \leq 0$. The desired fundamental solution is given by

$$(2.5) \quad E(x, t) = \frac{\partial^5}{\partial x^5} v(x, t) \quad t \leq 0,$$

where differentiation in x is interpreted in the sense of distributions. By Theorem 2.1 and (2.5) we have

$$(2.6) \quad \text{supp. } E(x, t) \subset \{(x, t); -\frac{2}{3}(-t)^{3/2} \leq x \leq \frac{2}{3}(-t)^{3/2}, t \leq 0\},$$

$$(2.7) \quad E(\cdot, t) \in C([-T, 0]; \mathcal{D}'(R_x)),$$

$$(2.8) \quad E_t(\cdot, t) \in C([-T, 0]; \mathcal{D}'(R_x))$$

for any $T > 0$, where $\mathcal{D}'(R_x)$ denotes the space of distributions in R_x .

Furthermore, by using the partial hypoellipticity of the Tricomi operator L in t (cf. [4], §§2.2, 4.3), we have the following.

COROLLARY 2.1. *Let Ω be an open set in $R_{x,t}^2$ such that $\{(x, 0); a < x < b\} \subset \Omega$. If $u \in \mathcal{D}'(\Omega)$ satisfies*

$$(2.9) \quad Lu = u_{tt} + tu_{xx} = 0 \quad \text{in } \Omega,$$

$$(2.10) \quad u = 0 \quad \text{in } \Omega_+ = \{(x, t); (x, t) \in \Omega, t > 0\}.$$

Then $u = 0$ in $\Omega_+ \cap (\bar{D} \cup \Omega)$ where D denotes a domain in the region $t < 0$ bounded by characteristics passing through $(a, 0)$ and $(b, 0)$.

For the proof we apply Theorem 2.1 by regularizing u with respect to x .

§ 3. Let Ω be an open set in $R^2_{x,t}$ which intersects x -axis.

THEOREM 3.1. Let $u = u(x, t) \in \mathcal{D}'(\Omega)$ be a solution of the equation

$$(3.1) \quad Lu = u_{tt} + tu_{xx} = f(x, t) \quad \text{in } \Omega$$

with $f \in C^\infty(\Omega)$. Then $u \in C^\infty(\Omega)$ if it is in $C^\infty(\Omega_-)$ where $\Omega_- = \{(x, t); (x, t) \in \Omega, t < 0\}$. Furthermore, u is an analytic function in Ω if it is analytic in Ω_- and if $f(x, t)$ is analytic in Ω .

We shall prove this theorem in several steps. First we shall show that $u(x, 0) \in C^\infty\{x; (x, 0) \in \Omega\}$.

Assume $\{(x, 0); 0 \leq x \leq b\} \subset \Omega, (0 < b)$. If we take $T > 0$ sufficiently small then the closed domain \bar{D} bounded by $\{(x, 0); 0 \leq x \leq b\}$, characteristics passing through $(0, 0)$ and $(b, 0)$ and $\{(x, -T); -\infty < x < +\infty\}$ is contained in $\Omega \cap \{(x, t); t \leq 0\}$. Let $u(x, t)$ and $f(x, t)$ be functions given in Theorem 3.1 and b, T be sufficiently small, then by the usual way (cf. [3]) we have

$$(3.2) \quad \begin{aligned} u(x, 0) = & \int E_t(x - y, -T)u(y, -T)dy - \int E(x - y, -T)u_t(y, -T)dy \\ & - \iint_{-T \leq \tau \leq 0} E(x - y, \tau)f(y, \tau)dyd\tau, \quad 0 < x < b, \end{aligned}$$

where the integral is taken in the sense of distributions. We note that there exists $u(x, 0) = \lim_{t \rightarrow 0} u(\cdot, t)$ in $\mathcal{D}'(0 < x < b)$ by the partial hypoellipticity of L in t (cf. [4], § 4). The formula (3.2) is justified because of the assumptions for u, f and the properties of $E(x, t)$: (2.6), (2.7), (2.8). Thus we have proved that $u(x, 0) \in C^\infty(0, b)$, and hence

$$u(x, 0) \in C^\infty\{x; (x, 0) \in \Omega\}.$$

Similarly, if u and f are analytic in Ω_- and Ω respectively, then we see that $u(x, 0)$ is analytic in $\{x; (x, 0) \in \Omega\}$. We omit the detail.

In the next section we shall show that

$$(3.3) \quad u \in C^\infty(\Omega \cap \{(x, t); t \geq 0\})$$

from which we see that $u(x, 0)$ and $u_t(x, 0)$ are in $C^\infty\{x; (x, 0) \in \Omega\}$. Then, applying Theorem 2.1 and Corollary 2.1, we have

$$(3.4) \quad u \in C^\infty(\Omega \cap \{(x, t); t \leq 0\}).$$

By (3.3), (3.4) and noting that the form of the equation is $u_{tt} + tu_{xx} = f$ in Ω we have $u \in C^\infty(\Omega)$ by the usual method of calculation (cf. § 4).

In the analytic case, from the assumption the $u(x, 0)$ is analytic in $\{x; (x, 0) \in \Omega\}$ we shall show, in the next section, $u = u(x, t)$ is analytic in $\Omega \cap \{(x, t); t \geq 0\}$ from where we have $u(x, 0), u_t(x, 0)$ are analytic in $\{x; (x, 0) \in \Omega\}$. Then by Cauchy-Kowalevski theorem and Corollary 2.1, u is analytic in a neighbourhood of the x -axis contained in Ω . On the other hand, u is analytic in $\Omega_+ = \{(x, t) \in \Omega, t > 0\}$ because it is a solution of an elliptic equation in Ω_+ . Thus u is analytic in Ω .

§ 4. It remains for us to prove the regularity property of the solution u in $\Omega \cap \{(x, t); t \geq 0\}$.

THEOREM 4.1. *Let $f \in C^\infty(\Omega)$ ($\in C^\omega(\Omega)$) and $u \in \mathcal{D}'(\Omega)$ such that*

$$(4.1) \quad Lu = u_{tt} + tu_{xx} = f(x, t) \quad \text{in } \Omega,$$

$$(4.2) \quad u(x, 0) = \psi(x) \in C^\infty\{x; (x, 0) \in \Omega\} \quad (\in C^\omega\{x; (x, 0) \in \Omega\}).$$

Then we have $u \in C^\infty(\Omega \cap \{(x, t); t \geq 0\})$ ($\in C^\omega(\Omega \cap \{(x, t); t \geq 0\})$). Here C^ω denotes the set of analytic functions.

To prove this theorem we use the method employed in [5], §§ 5, 6. We note that it is sufficient to prove the case $u(x, 0) = \psi(x) = 0$. First we prepare the following theorem which is derived by a direct computation. Take $G = (a < x < b) \times [0, T)$ such that $\bar{G} \subset \Omega$ and introduce the notation:

$$(4.3) \quad \|v\|_{\mathfrak{H}(G)}^2 = \sum_{j=0}^2 \|D_t^j v\|_{L^2(G)}^2 + \|t^{1/2} v_{xt}\|_{L^2(G)}^2 + \|t^{1/2} v_x\|_{L^2(G)}^2 + \|tv_{xx}\|_{L^2(G)}^2.$$

$\mathfrak{H}(G)$ is a Hilbert space with the norm $\|\cdot\|_{\mathfrak{H}(G)}$.

THEOREM 4.2 (cf. [5], Theorem 4.2). *There exists a constant $C > 0$ such that*

$$(4.4) \quad \|v\|_{\mathfrak{H}(G)} \leq C \|Lv\|_{L^2(G)}$$

for all $v \in \mathfrak{H}(G)$ with $\text{supp. } v \subset G$ and $v(x, 0) = 0$.

Suppose $f(x, t) \in C^\infty(\Omega)$, then by the partial hypoellipticity of L in t (cf. [4], § 4.3) we conclude that for any $r (\geq 2)$ there exists a number $\beta = \beta(u, r)$ such that

$$(4.5) \quad \zeta u \in H_{(r, \beta)}(G) = H_{(r, \beta)}(R^2)|_G$$

for any $\zeta = \zeta(x, t) \in C_0^\infty(G)$, For the notation $H_{(r, \beta)}(R^2)$, we refer to [4], § 2.5.

For a real number s we define an operator T_s :

$$\widehat{T_s v}(\xi, t) = (1 + |\xi|^2)^{s/2} \hat{v}(\xi, t),$$

where $v \in \mathcal{S}'(R_{x,t}^2 \cap \{t \geq 0\})$ and $\hat{v}(\xi, t)$ denotes the partial Fourier transformation of v with respect to x . (cf. [4], § 1.7.)

For any $x_0 \in (a, b)$ take $\zeta \in C_0^\infty(G)$ such that $\zeta(x_0, 0) \neq 0$ and

$$\frac{\partial \zeta}{\partial t}(x, t) = 0 \quad \text{if } (x, t) \in G, \quad 0 \leq t \leq \frac{T}{2}.$$

Then by (4.5) we have

$$(4.6) \quad \varphi T_\beta \zeta u \in \mathfrak{S}(G)$$

for any $\varphi \in C_0^\infty(G)$. Starting with (4.6), by using the estimate (4.4) we can easily show that $\varphi T_s \zeta u \in \mathfrak{S}(G)$ for any s and $\varphi \in C_0^\infty(G)$ from where we have $\varphi D_x^j u \in \mathfrak{S}(G), j = 0, 1, 2, \dots$. And rewriting the form of the equation $u_{tt} = -tu_{xx} + f$, we have $\varphi D_t^i D_x^j \in L^2(G), 0 \leq i, j < \infty$. Then we have $u \in C^\infty(G)$, from where we have $u \in C^\infty(G)$.

Next we consider the case where $f \in C^\omega(G)$ and $u(x, 0) = 0$. In this case we have $u \in C^\infty(G)$ by the above result. To obtain the analyticity of u in $\Omega \cap \{(x, t); t \geq 0\}$, we have to estimate precisely the successive derivatives of u . We can pursue the manner employed in [6], § 6 where the analyticity of the solutions of the equations $u_{tt} + t^{2k}u_{xx} = f, k = 0, 1, 2, \dots$, was proved. In the following we shall give an outline of the reasoning.

Introduce the notations:

$$G_\epsilon = (a + \epsilon < x < b - \epsilon) \times [0 \leq t < T] \quad 0 < \epsilon < \text{Min} \left(\frac{b-a}{2}, \frac{T}{2} \right),$$

$$G_\epsilon^* = G_\epsilon \setminus (a + \epsilon < x < b - \epsilon) \times \left[0 \leq t < \frac{T}{2} \right],$$

$$N_\epsilon(v) = \|v\|_{L^2(G_\epsilon)}, \quad N_\epsilon^*(v) = \|v\|_{L^2(G_\epsilon^*)}.$$

LEMMA 4.1 (cf. [4], ch. 1). *Let $\varepsilon, \varepsilon_1$ be positive numbers with $0 < \varepsilon + \varepsilon_1 < \text{Min}((b - a)/2, T/2)$. Then there exists functions $\psi = \psi_{\varepsilon, \varepsilon_1} \in C_0^\infty(G_{\varepsilon_1})$ such that $\psi = \psi_{\varepsilon, \varepsilon_1} \equiv 1$ on $G_{\varepsilon + \varepsilon_1}$ and*

$$(4.7) \quad \begin{aligned} \text{Max } |D_x^j D_t^r \psi| &\leq C_{j+r} \varepsilon^{-(j+r)} & 0 \leq j + r \leq 2 \\ D_t \psi &\equiv 0 & \text{on } (a + \varepsilon_1, b - \varepsilon_1) \times \left[0, \frac{T}{2}\right]. \end{aligned}$$

LEMMA 4.2 (cf. [6], Lemma 6.2). *There exists a constant $C > 0$ such that*

$$(4.8) \quad \begin{aligned} &\sum_{j=0}^2 \varepsilon^j N_{\varepsilon + \varepsilon_1}(D_t^j v) + \sum_{j=0}^2 \varepsilon^j N_{\varepsilon + \varepsilon_1}(t D_x^j v) + N_{\varepsilon + \varepsilon_1}^*(v) \\ &+ \varepsilon N_{\varepsilon + \varepsilon_1}^*(D_x v) + \varepsilon^2 N_{\varepsilon + \varepsilon_1}^*(D_t D_x v) \\ &\leq C\{\varepsilon^2 N_{\varepsilon_1}(Lv) + \sum_{j=0,1} \varepsilon^j N_{\varepsilon_1}(t D_x^j v) + N_{\varepsilon_1}^*(v) + \varepsilon N_{\varepsilon_1}^*(D_t v)\} \end{aligned}$$

for all $v \in C^\infty(G)$ and $v(x, 0) = 0$. The constant C does not depend on $\varepsilon, \varepsilon_1$ under the condition mentioned previously.

This lemma is obtained by substituting $\psi_{\varepsilon, \varepsilon_1} v$ in (4.4).

LEMMA 4.3 (cf. [4], ch. 7). *Let w be an analytic function in G . Then there exists a constant $C > 0$ such that*

$$(4.9) \quad \varepsilon^{j+r} N_{k\varepsilon}(D_x^j D_t^r w) \leq C^{j+r+1} \quad \text{if } j + r < k,$$

for all integer $k > 0$. Conversely, if $w \in C^\infty(G)$ satisfies (4.9), then w is analytic in G .

Proof of the analyticity of u in $\Omega \cap \{(x, t); t \geq 0\}$.

First we shall show that there exists a constant $B > 0$ such that, for any $\varepsilon > 0$ and for any integer $l > 0$,

$$(4.10) \quad \left. \begin{aligned} &\sum_{r=0}^2 \varepsilon^{r+j} N_{l\varepsilon}(D_t^r D_x^j u) \\ &\sum_{r=0}^2 \varepsilon^{r+j} N_{l\varepsilon}(t^{2k} D_x^{r+j} u) \\ &\sum_{r=0,1} \varepsilon^{r+j} N_{l\varepsilon}^*(D_x^{r+j} u) \\ &\varepsilon^{2+j} N_{l\varepsilon}^*(D_t D_x^{j+1} u) \end{aligned} \right\} \leq B^{l+1}$$

if $j < l$.

It we take B sufficiently large, we have (4.10) for $l = 1$ by Lemma 4.2. Next, since $f(x, t)$ is analytic in \bar{G} , there exists a constant $C_0 > 0$ such that

$$\varepsilon^{2+j}N_{j\varepsilon}(D_x^j f) \leq C_0^{j+1},$$

for $j = 1, 2, \dots$ and $0 < \varepsilon < (b - a)/2$.

Assuming that (4.10) have been proved for an $l > 0$, we shall prove (4.10) for $l + 1$. Replacing v by $\varepsilon^l D_x^l u$ and ε_1 by $l\varepsilon$ in (4.8), we see that the terms in the left hand side of (4.10) for the case $l + 1$ are smaller than $5C_0 B^{l+1}$ if $j < l + 1$. Hence we have (4.10) for $l + 1$ if $5C_0 B^{l+1} \leq B^{l+2}$. This condition is satisfied for all l if $B > \max(5C_0, 1)$.

From (4.10) (cf. Lemma 4.3) we obtain

$$(4.11) \quad \sum_{r=0}^2 \|D_t^r D_x^j u\|_{L^2(G_{\varepsilon_1})} \leq C_1^{j+1} j^j, \quad j = 0, 1, 2, \dots$$

for some constant $C_1 > 0$ where $G_{\varepsilon_1} = (a + \varepsilon_1, b - \varepsilon_1) \times [0, T/2]$ with $\varepsilon_1 > 0$ sufficiently small.

To obtain the successive estimates including the derivatives in both x and t , we rewrite the equation $Lu = f$ in the form $D_t^2 u = -tD_x^2 u + f$. And using (4.11) by the usual way (cf. [6] for example) we have

$$\|D_x^j D_t^r u\|_{L^2(G_{\varepsilon_1})} \leq C_2^{j+r+1} (j + r)^{j+r} \quad 0 \leq j, r < \infty$$

for some constant $C_2 > 0$, from which we have the analyticity of u in G_{ε_1} by the Sobolev lemma.

REFERENCES

- [1] Baouendi M. S., Sur une class d'opérateurs elliptiques dégénérés, Bull. Soc. Math. France, **95**, 45-87 (1967).
- [2] Berezin I. S., On Cauchy's problem for linear equations of the second order with initial conditions on a parabolic line (in Russian). Mat. Sbornik, N. S. t24 (66), 301-320 (1949).
- [3] Courant R. and Hilbert D.: Methods of mathematical physics. Vol. II. Interscience Pub. New York (1962).
- [4] Hörmander L.: Linear partial differential operators. Berlin, Springer-Verlag (1963).
- [5] Matsuzawa T., Sur les équations $u_{tt} + t^\alpha u_{xx} = f$ ($\alpha \geq 0$). Nagoya Math. J. Vol. **42**, (1971).
- [6] Matsuzawa T., Sur les équations quasi-elliptiques et les classes de Gevrey. Bull. Soc. Math. France, t96, 243-263 (1968).

Nagoya University