

RATIONAL REPRESENTATIONS AND RATIONAL GROUP ALGEBRA OF VZ p -GROUPS

RAM KARAN CHOUDHARY and SUNIL KUMAR PRAJAPATI

(Received 2 October 2023; accepted 14 June 2024; first published online 6 November 2024)

Communicated by Benjamin Martin

Abstract

In this article, we study rational matrix representations of VZ p -groups (p is any prime). Using our findings on VZ p -groups, we explicitly obtain all inequivalent irreducible rational matrix representations of all p -groups of order $\leq p^4$. Furthermore, we establish combinatorial formulae to determine the Wedderburn decompositions of rational group algebras for VZ p -groups and all p -groups of order $\leq p^4$, ensuring simplicity in the process.

2020 Mathematics subject classification: primary 20D15; secondary 20C15, 16S34.

Keywords and phrases: rational representations, Wedderburn decomposition, VZ-groups, p -groups.

1. Introduction

This paper consistently employs the following notation: G for a finite group, $\text{Irr}(G)$ for the set of all complex irreducible characters of G , \mathbb{F} for a field with characteristic 0 and p for a prime number. In representation theory, a challenging and crucial task is to compute all inequivalent irreducible matrix representations of G over \mathbb{F} , even for $\mathbb{F} = \mathbb{C}$. In this paper, we deal with $\mathbb{F} = \mathbb{Q}$. For a given character $\chi \in \text{Irr}(G)$, we define $\Omega(\chi)$ as follows:

$$\Omega(\chi) = m_{\mathbb{Q}}(\chi) \sum_{\sigma \in \text{Gal}(\mathbb{Q}(\chi)/\mathbb{Q})} \chi^{\sigma},$$

where $m_{\mathbb{Q}}(\chi)$ represents the Schur index of χ over \mathbb{Q} . Note that $\Omega(\chi)$ corresponds to the character of an irreducible \mathbb{Q} -representation ρ of G . Conversely, if ρ is an irreducible \mathbb{Q} -representation of G , then there exists $\chi \in \text{Irr}(G)$ such that $\Omega(\chi)$ is the character of ρ . Generally, obtaining an irreducible \mathbb{Q} -representation ρ of G that affords the

The first author acknowledges University Grants Commission, Government of India. The corresponding author acknowledges SERB, Government of India for financial support through grant (MTR/2019/000118).

© The Author(s), 2024. Published by Cambridge University Press on behalf of Australian Mathematical Publishing Association Inc.



character $\Omega(\chi)$ is a challenging task. A significant and interesting task is to determine all inequivalent irreducible matrix representations of G over \mathbb{Q} for several reasons. For example, one problem in rationality concerns the realizability of an \mathbb{F} -representation of G over its subfields, specially, the realizability of a \mathbb{C} -representation of G over \mathbb{R} or \mathbb{Q} .

In this article, we study irreducible rational matrix representations for some classes of p -groups. In Section 3, for any p -group G , we present Algorithms 15 and 20 for computing an irreducible rational matrix representation of G that affords the character $\Omega(\chi)$, where $\chi \in \text{Irr}(G)$. These algorithms rely heavily on results from [9, 31, 32]. To obtain an irreducible rational matrix representation of a finite p -group G affording the character $\Omega(\chi)$, where $\chi \in \text{Irr}(G)$ is equivalent to finding a pair (H, ψ) , where $H \leq G$ and $\psi \in \text{Irr}(H)$, with some suitable properties (see Algorithms 15 and 20). We refer to (H, ψ) as a required pair for an irreducible rational matrix representation of G that affords the character $\Omega(\chi)$. Importantly, using a required pair (H, ψ) , we can also compute an irreducible complex matrix representation of G that affords the character χ . In Section 4, we study required pairs for VZ p -groups. A group G is called a VZ-group if all its nonlinear irreducible characters vanish off the centre. VZ-groups have been extensively studied by various researchers in [8, 18, 19, 25, 26]. By using our results on VZ p -groups, we explicitly obtain all inequivalent irreducible rational matrix representations of all p -groups of order $\leq p^4$ (see Sections 5 and 6).

In parallel, this article delves into an investigation of the Wedderburn decomposition of $\mathbb{Q}G$ with a specific focus on VZ p -groups. For a semisimple group algebra $\mathbb{F}G$, the Wedderburn components are matrix algebras over finite extensions of \mathbb{F} in the case of positive characteristic, and Brauer-equivalent to cyclotomic algebras in the case of zero characteristic, as per the Brauer–Witt theorem (see [30]). Further, the Wedderburn decomposition of $\mathbb{F}G$ aids in describing the automorphisms group of $\mathbb{F}G$ (see [11, 22]) or studying the unit group of the integral group ring $\mathbb{Z}G$ when $\mathbb{F} = \mathbb{Q}$ (see [7, 14, 16, 28]). The Wedderburn decomposition of $\mathbb{Q}G$ has been extensively studied in [2–4, 17, 21, 23, 24]. They used various concepts such as computation of the field of character values, Shoda pairs, numerical representation of cyclotomic algebras and so forth, to compute simple components of $\mathbb{Q}G$. We prove Theorems 1 and 2, which provide a combinatorial description for the Wedderburn decomposition of rational group algebra of a VZ p -group. Our results formulate the computation of the Wedderburn decomposition of a VZ p -group G solely based on computing the number of cyclic subgroups of $Z(G)$ and $Z(G)/G'$, which is similar to the Perlis–Walker theorem for an abelian group (see Lemma 6). Indeed, we prove the following theorems.

THEOREM 1. *Let G be a finite VZ p -group (odd prime p). Let m_1, m_2 and m_3 denote the exponents of $G/G', Z(G)$ and $Z(G)/G'$, respectively. Then the Wedderburn decomposition of $\mathbb{Q}G$ is as follows:*

$$\mathbb{Q}G \cong \bigoplus_{d_1|m_1} a_{d_1} \mathbb{Q}(\zeta_{d_1}) \bigoplus_{d_2|m_2, d_2 \nmid m_3} a_{d_2} M_{|G/Z(G)|^{1/2}}(\mathbb{Q}(\zeta_{d_2}))$$

$$\bigoplus_{d_2|m_2, d_2|m_3} (a_{d_2} - a'_{d_2}) M_{|G/Z(G)|^{1/2}}(\mathbb{Q}(\zeta_{d_2})),$$

where a_{d_1} , a_{d_2} and a'_{d_2} are the number of cyclic subgroups of G/G' of order d_1 , the number of cyclic subgroups of $Z(G)$ of order d_2 and the number of cyclic subgroups of $Z(G)/G'$ of order d_2 , respectively.

THEOREM 2. *Let G be a VZ 2-group. Let m_1 , m_2 and m_3 denote the exponents of G/G' , $Z(G)$ and $Z(G)/G'$, respectively. Suppose $k = |\{\chi \in \text{nl}(G) : m_{\mathbb{Q}}(\chi) = 2\}|$ and $\mathbb{H}(\mathbb{Q})$ represents the standard quaternion algebra over \mathbb{Q} . Then the Wedderburn decomposition of $\mathbb{Q}G$ is as follows:*

$$\mathbb{Q}G \cong \bigoplus_{d_1|m_1} a_{d_1} \mathbb{Q}(\zeta_{d_1}) \bigoplus kM_{1/2|G/Z(G)|^{1/2}}(\mathbb{H}(\mathbb{Q})) \bigoplus (a_2 - a'_2 - k)M_{|G/Z(G)|^{1/2}}(\mathbb{Q}) \\ \bigoplus_{d_2|m_2, d_2 \nmid m_3} a_{d_2} M_{|G/Z(G)|^{1/2}}(\mathbb{Q}(\zeta_{d_2})) \bigoplus_{d_2|m_2, d_2|m_3} (a_{d_2} - a'_{d_2})M_{|G/Z(G)|^{1/2}}(\mathbb{Q}(\zeta_{d_2})),$$

where a_{d_1} , a_l and a'_l ($l \in \{2, d_2 \geq 4\}$) are the number of cyclic subgroups of G/G' of order d_1 , the number of cyclic subgroups of $Z(G)$ of order l and the number of cyclic subgroups of $Z(G)/G'$ of order l , respectively.

In Section 4.2, we derive several consequences from the above theorems. Further, if G is a nonabelian p -group of order p^4 of maximal class, then G has a unique abelian subgroup of index p (see Section 6.2). We prove Theorem 3, which formulates the computation of the Wedderburn decomposition of a non-VZ p -group G of order p^4 .

THEOREM 3. *Let G be a nonabelian p -group (odd prime p) of order p^4 of nilpotency class 3, and let H be its unique abelian subgroup of index p . Let m and m' denote the exponents of H and H/G' , respectively. Then the Wedderburn decomposition of $\mathbb{Q}G$ is as follows:*

$$\mathbb{Q}G \cong \bigoplus \mathbb{Q}(G/G') \bigoplus_{d|m, d \nmid m'} \frac{a_d}{p} M_p(\mathbb{Q}(\zeta_d)) \bigoplus_{d|m, d|m'} \frac{a_d - a'_d}{p} M_p(\mathbb{Q}(\zeta_d)),$$

where a_d and a'_d are the number of cyclic subgroups of order d of H and H/G' , respectively.

In this article, we also provide a brief analysis of primitive central idempotents and their corresponding simple components in the Wedderburn decomposition of the rational group ring of a VZ p -group (see Section 4.3).

2. Notation and some basic results

2.1. Notation. For a finite group G , the following notation is used consistently throughout this article.

- G' the commutator subgroup of G
- $|S|$ the cardinality of a set S

$\text{Core}_G(H)$	the normal core of H in G for $H \leq G$
$\text{Irr}(G)$	the set of irreducible complex characters of G
$\text{lin}(G)$	$\{\chi \in \text{Irr}(G) : \chi(1) = 1\}$
$\text{nl}(G)$	$\{\chi \in \text{Irr}(G) : \chi(1) \neq 1\}$
$\text{FIrr}(G)$	the set of faithful irreducible complex characters of G
$\text{Irr}^{(m)}(G)$	$\{\chi \in \text{Irr}(G) : \chi(1) = m\}$
$\text{cd}(G)$	$\{\chi(1) : \chi \in \text{Irr}(G)\}$
$\text{Irr}_{\mathbb{Q}}(G)$	the set of irreducible rational characters of G
$\text{Irr}_{\mathbb{Q}}^{(m)}(G)$	$\{\chi \in \text{Irr}_{\mathbb{Q}}(G) : \chi(1) = m\}$
$\mathbb{F}(\chi)$	the field obtained by adjoining the values $\{\chi(g) : g \in G\}$ to the field \mathbb{F} for some $\chi \in \text{Irr}(G)$
$m_{\mathbb{Q}}(\chi)$	the Schur index of $\chi \in \text{Irr}(G)$ over \mathbb{Q}
$\Omega(\chi)$	$m_{\mathbb{Q}}(\chi) \sum_{\sigma \in \text{Gal}(\mathbb{Q}(\chi)/\mathbb{Q})} \chi^{\sigma}$ for $\chi \in \text{Irr}(G)$
$\ker(\chi)$	$\{g \in G : \chi(g) = \chi(1)\}$ for $\chi \in \text{Irr}(G)$
$\text{Irr}(G N)$	$\{\chi \in \text{Irr}(G) : N \not\subseteq \ker(\chi)\}$, where $N \trianglelefteq G$
ψ^G	the induced character of ψ to G , where ψ is a character of H for some $H \leq G$
Ψ^G	the induced representation of Ψ to G , where Ψ is a representation of H for some $H \leq G$
$\chi \downarrow_H$	the restriction of a character χ of G on H , where $H \leq G$
$\mathbb{F}G$	the group ring (algebra) of G with coefficients in \mathbb{F}
$M_n(D)$	a full matrix ring of order n over the skewfield D
$Z(B)$	the centre of an algebraic structure B
$\phi(n)$	the Euler phi function
ζ_m	an m th primitive root of unity

2.2. Basic results. In this subsection, we discuss some basic concepts and results, which we use frequently throughout the article.

Let G be a finite group and $n = |G|$. Consider $\mathbb{Q}(\zeta_n)$, the n th cyclotomic field obtained by adjoining a primitive n th root of unity to \mathbb{Q} . Let $\chi \in \text{Irr}(G)$. If $\sigma \in \text{Gal}(\mathbb{Q}(\zeta_n)/\mathbb{Q})$, then define the function $\chi^{\sigma} : G \rightarrow \mathbb{C}$ as $\chi^{\sigma}(g) = \sigma(\chi(g))$ for $g \in G$. It is easy to observe that $\chi^{\sigma} \in \text{Irr}(G)$ and hence, $\text{Gal}(\mathbb{Q}(\zeta_n)/\mathbb{Q})$ acts on $\text{Irr}(G)$. Note that $\mathbb{Q}(\chi)$ is a finite degree Galois extension of \mathbb{Q} and the Galois group $\text{Gal}(\mathbb{Q}(\chi)/\mathbb{Q})$ is abelian. It is easy to see that $\text{Gal}(\mathbb{Q}(\chi)/\mathbb{Q})$ also acts on $\text{Irr}(G)$, with action given by $\sigma \cdot \chi := \chi^{\sigma}$. Under the above set-up, we have the following lemma.

LEMMA 4 [12, Lemma 9.17]. *Let $E(\chi)$ denote the Galois conjugacy class of $\chi \in \text{Irr}(G)$ under the action of $\text{Gal}(\mathbb{Q}(\zeta_n)/\mathbb{Q})$. Then,*

$$|E(\chi)| = [\mathbb{Q}(\chi) : \mathbb{Q}].$$

DEFINITION 5. Let $\chi, \psi \in \text{Irr}(G)$. We say that χ and ψ are Galois conjugates over \mathbb{Q} if $\mathbb{Q}(\chi) = \mathbb{Q}(\psi)$, and there exists $\sigma \in \text{Gal}(\mathbb{Q}(\chi)/\mathbb{Q})$ such that $\chi^{\sigma} = \psi$.

In [24], Perlis and Walker studied the group ring of a finite abelian group G over the field of rational numbers and proved the following result.

LEMMA 6 (Perlis–Walker theorem). *Let G be a finite abelian group of exponent m . Then the Wedderburn decomposition of $\mathbb{Q}G$ is as follows:*

$$\mathbb{Q}G \cong \bigoplus_{d|m} a_d \mathbb{Q}(\zeta_d),$$

where a_d is equal to the number of cyclic subgroups of G of order d .

REMARK 7. Let G be a finite abelian group of exponent m . Then by Lemma 6:

- (1) the number of rational irreducible representations of G of degree $\phi(d)$ is equal to a_d ; and
- (2) the total number of rational irreducible representations of G is equal to $\sum_{d|m} a_d$.

Let \mathbb{K} be an arbitrary field with characteristic zero and \mathbb{K}^* be the algebraic closure of \mathbb{K} . Let U be an irreducible \mathbb{K}^* -representation of G with character χ . The *Schur index* of U with respect to \mathbb{K} is defined as

$$m_{\mathbb{K}}(U) = \text{Min}[\mathbb{L} : \mathbb{K}(\chi)],$$

the minimum being taken over all fields \mathbb{L} in which U is realizable. Note that $m_{\mathbb{K}}(\chi) = m_{\mathbb{K}}(U)$.

Reiner [27] characterized the simple component of the Wedderburn decomposition of $\mathbb{K}G$ and proved the following result.

LEMMA 8 [27, Theorem 3]. *Let T be an irreducible \mathbb{K} -representation of G , and extend T (by linearity) to a \mathbb{K} -representation of $\mathbb{K}G$. Set*

$$A = \{T(x) : x \in \mathbb{K}G\}.$$

Then, A is simple algebra over \mathbb{K} , and we may write $A = M_n(D)$, where D is a division ring. Further,

$$Z(D) \cong \mathbb{K}(\chi) \quad \text{and} \quad [D : Z(D)] = (m_{\mathbb{K}}(U_i))^2 \quad (1 \leq i \leq k),$$

where U_i are irreducible \mathbb{K}^* -representations of G such that $T = m_{\mathbb{K}}(U_i) \bigoplus_{i=1}^k U_i$ as a \mathbb{K}^* -representation.

In this article, we use the classification of p -groups of order $\leq p^4$ (odd prime p) provided in [13], which is based on the isoclinism concept.

DEFINITION 9. Two finite groups G and H are said to be isoclinic if there exist isomorphisms $\theta : G/Z(G) \rightarrow H/Z(H)$ and $\phi : G' \rightarrow H'$ such that the following

diagram is commutative:

$$\begin{array}{ccc}
 G/Z(G) \times G/Z(G) & \xrightarrow{a_G} & G' \\
 \downarrow \theta \times \theta & & \downarrow \phi \\
 H/Z(H) \times H/Z(H) & \xrightarrow{a_H} & H'
 \end{array}$$

where $a_G(g_1Z(G), g_2Z(G)) = [g_1, g_2]$ for $g_1, g_2 \in G$, and $a_H(h_1Z(H), h_2Z(H)) = [h_1, h_2]$ for $h_1, h_2 \in H$.

The resulting pair (θ, ϕ) is called an *isoclinism* of G onto H . Isoclinism was first introduced by Hall [10] for the classification of p -groups. It is a generalization of the concept of isomorphism between two groups. It is well known that two isoclinic nilpotent groups have the same nilpotency class. Now, we end this subsection by quoting the following lemma.

LEMMA 10 [29, Theorem 3.2]. *Let G and H be isoclinic groups. Then, $|H||\text{Irr}^{(k)}(G)| = |G||\text{Irr}^{(k)}(H)|$.*

3. Algorithm

This section outlines the algorithm for computing irreducible rational matrix representations of p -groups. For $\chi \in \text{Irr}(G)$, there exists a unique irreducible \mathbb{Q} -representation ρ of G such that χ occurs as an irreducible constituent of $\rho \otimes_{\mathbb{Q}} \mathbb{F}$ with multiplicity $m_{\mathbb{Q}}(\chi)$, where \mathbb{F} is a splitting field of G . Therefore, the distinct Galois conjugacy classes give the distinct irreducible rational representations of G .

LEMMA 11 [32, Proposition 1]. *Let $\psi \in \text{lin}(G)$ and $N = \ker(\psi)$ with $n = [G : N]$. Suppose $G = \bigcup_{i=0}^{n-1} Ny^i$. Then,*

$$\psi(xy^i) = \zeta_n^i, \quad (0 \leq i < n; x \in N).$$

Now, let $f(X) = X^s - a_{s-1}X^{s-1} - \dots - a_1X - a_0$ be the irreducible polynomial over \mathbb{Q} such that $f(\zeta_n) = 0$, where $s = \phi(n)$ and

$$\Psi(xy^i) = \begin{pmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \\ 0 & 0 & \dots & 0 & 1 \\ a_0 & a_1 & \dots & \dots & a_{s-1} \end{pmatrix}^i, \quad (0 \leq i < n; x \in N).$$

Then Ψ is an irreducible \mathbb{Q} -representation of G , whose character is $\Omega(\psi)$.

LEMMA 12 [32, Proposition 3]. *Let H be a subgroup of G and $\psi \in \text{Irr}(H)$ be such that $\psi^G \in \text{Irr}(G)$. Then $m_{\mathbb{Q}}(\psi^G)$ divides $m_{\mathbb{Q}}(\psi)[\mathbb{Q}(\psi) : \mathbb{Q}(\psi^G)]$. Furthermore, the induced character $\Omega(\psi)^G$ of G is a character of an irreducible \mathbb{Q} -representation of G if and only if*

$$m_{\mathbb{Q}}(\psi^G) = m_{\mathbb{Q}}(\psi)[\mathbb{Q}(\psi) : \mathbb{Q}(\psi^G)].$$

In this case, $\Omega(\psi)^G = \Omega(\psi^G)$.

LEMMA 13 [9, Theorem 1]. *Let G be a p -group and $\chi \in \text{Irr}(G)$. Then, one of the following holds.*

- (1) *There exists a subgroup H of G and $\psi \in \text{lin}(H)$ such that $\psi^G = \chi$ and $\mathbb{Q}(\psi) = \mathbb{Q}(\chi)$.*
- (2) *$p = 2$ and there exist subgroups $H < K$ in G with $[K : H] = 2$ and $\lambda \in \text{lin}(H)$ such that $\lambda^K = \phi$, $[\mathbb{Q}(\lambda) : \mathbb{Q}(\phi)] = 2$, $\phi^G = \chi$ and $\mathbb{Q}(\phi) = \mathbb{Q}(\chi)$.*

LEMMA 14 [12, Corollary 10.14]. *Let G be a p -group and $\chi \in \text{Irr}(G)$. If p is an odd prime, then $m_{\mathbb{Q}}(\chi) = 1$; otherwise $m_{\mathbb{Q}}(\chi) \in \{1, 2\}$.*

Let G be a p -group (odd prime p) and $\chi \in \text{Irr}(G)$. According to Lemma 14, $m_{\mathbb{Q}}(\chi) = 1$. By Lemma 13, there exists a subgroup H of G with $\psi \in \text{lin}(H)$ such that $\psi^G = \chi$ and $\mathbb{Q}(\psi) = \mathbb{Q}(\chi)$. Therefore, from Lemma 12, we have $\Omega(\chi) = \Omega(\psi)^G$. Now by using Lemma 11, compute an irreducible matrix representation Ψ of H over \mathbb{Q} that affords the character $\Omega(\psi)$. Then Ψ^G is an irreducible \mathbb{Q} -representation of G that affords the character $\Omega(\psi)^G = \Omega(\psi^G) = \Omega(\chi)$. In summary, an algorithm to find an irreducible rational matrix representation of a p -group G (odd prime p) can be outlined as follows.

ALGORITHM 15. Input: An irreducible complex character χ of a finite p -group G (odd prime p).

- (1) Find a pair (H, ψ) , where $H \leq G$ and $\psi \in \text{lin}(H)$ is such that $\psi^G = \chi$ and $\mathbb{Q}(\psi) = \mathbb{Q}(\chi)$.
- (2) Find an irreducible \mathbb{Q} -representation Ψ of H that affords the character $\Omega(\psi)$.
- (3) Induce Ψ to G .

Output: Ψ^G , an irreducible \mathbb{Q} -representation of G whose character is $\Omega(\chi)$.

REMARK 16. Obtaining an irreducible rational matrix representation of a finite p -group G (odd prime p) affording the character $\Omega(\chi)$, where $\chi \in \text{Irr}(G)$ is equivalent to finding a pair (H, ψ) , where H is a subgroup of G and $\psi \in \text{lin}(H)$, satisfying $\psi^G = \chi$ and $\mathbb{Q}(\psi) = \mathbb{Q}(\chi)$. We refer to this pair (H, ψ) as a required pair for an irreducible rational matrix representation of G that affords the character $\Omega(\chi)$.

Now, we describe the algorithm for computing irreducible rational matrix representations of 2-groups.

LEMMA 17 [31, Theorem 2.12]. *Let G be a 2-group and $\chi \in \text{Irr}(G)$. Then there exists a pair (H, ψ) such that $H \leq G$, $\psi \in \text{Irr}(H)$, $\psi^G = \chi$, $\mathbb{Q}(\chi) = \mathbb{Q}(\psi)$, and one of the following holds:*

- (1) $H/\ker(\psi) \cong Q_n (n \geq 2)$, $m_{\mathbb{Q}}(\chi) = 2$, $\mathbb{Q}(\chi) = \mathbb{Q}(\zeta_{2^n} + \zeta_{2^n}^{-1})$;
- (2) $H/\ker(\psi) \cong D_n (n \geq 3)$, $m_{\mathbb{Q}}(\chi) = 1$, $\mathbb{Q}(\chi) = \mathbb{Q}(\zeta_{2^n} + \zeta_{2^n}^{-1})$;

- (3) $H/\ker(\psi) \cong SD_n (n \geq 3), m_{\mathbb{Q}}(\chi) = 1, \mathbb{Q}(\chi) = \mathbb{Q}(\zeta_{2^n} - \zeta_{2^n}^{-1});$
- (4) $H/\ker(\psi) \cong C_n (n \geq 0), m_{\mathbb{Q}}(\chi) = 1, \mathbb{Q}(\chi) = \mathbb{Q}(\zeta_{2^n}),$

where Q_n, D_n and SD_n are respectively the generalized quaternion, dihedral and semidihedral group of order 2^{n+1} , and C_n is the cyclic group of order 2^n .

LEMMA 18 [32, Example 7 and Proposition 8].

- (1) Let $G = Q_n = \langle a, b : a^{2^n} = 1, b^2 = a^{2^{n-1}}, bab^{-1} = a^{-1} \rangle$ be the generalized quaternion group of order 2^{n+1} , and let $\chi \in \text{FIrr}(G)$. Then, $\chi = \psi^G$, where $H = \langle a \rangle$ and $\psi \in \text{lin}(H)$ is such that $\psi(a) = \zeta_{2^n}, m_{\mathbb{Q}}(\chi) = 2$ and $\Omega(\chi) = \Omega(\psi)^G$.
- (2) Let $G = D_n = \langle a, b : a^{2^n} = b^2 = 1, bab^{-1} = a^{-1} \rangle$ be the dihedral group of order 2^{n+1} or $G = SD_n = \langle a, b : a^{2^n} = b^2 = 1, bab^{-1} = a^{2^{n-1}-1} \rangle$ be the semi-dihedral group of order 2^{n+1} , and let $\chi \in \text{FIrr}(G)$. Then, $\Omega(\chi) = \Omega(\psi)^G$, where $H = \langle a^{2^{n-1}}, b \rangle$ and $\psi \in \text{lin}(H)$ is such that $\psi(a^{2^{n-1}}) = -1, \psi(b) = 1$.

REMARK 19. Let G be a 2-group and let $\chi \in \text{nl}(G)$. By Lemma 17, there exists a pair (H, ψ) , with $H \leq G$ and $\psi \in \text{Irr}(H)$, satisfying the following properties: $\psi^G = \chi, \mathbb{Q}(\chi) = \mathbb{Q}(\psi)$ and $H/\ker(\psi)$ is isomorphic to one of the following groups: cyclic group, generalized quaternion group, dihedral group or semi-dihedral group. We define $\bar{\psi} \in \text{FIrr}(H/\ker(\psi))$ such that $\bar{\psi}(h \ker(\psi)) = \psi(h)$ for all $h \in H$. Now we have two cases.

Case 1 ($H/\ker(\psi) \cong C_n$). In this case, $\psi \in \text{lin}(H)$ and hence by using Lemma 11, we get an irreducible rational matrix representation Ψ of H that affords the character $\Omega(\psi)$. Then by Lemma 12, Ψ^G is an irreducible rational matrix representation of G that affords the character $\Omega(\chi)$.

Case 2 ($H/\ker(\psi) \cong Q_n, \text{ or } D_n, \text{ or } SD_n \text{ for some } n \in \mathbb{N}$). In this case, since $\bar{\psi} \in \text{FIrr}(H/\ker(\psi))$, by Lemmas 18 and 11, there exists an irreducible rational matrix representation of $H/\ker(\psi)$ that affords the character $\Omega(\bar{\psi})$. Indeed, we get an irreducible rational matrix representation Ψ of H that affords the character $\Omega(\psi)$. Then again by Lemma 12, Ψ^G is an irreducible rational matrix representation of G that affords the character $\Omega(\chi)$.

In view of Remark 19, an algorithm to find an irreducible rational matrix representation of a 2-group G can be outlined as follows.

ALGORITHM 20. Input: An irreducible complex character χ of a finite 2-group G .

- (1) Find a pair (H, ψ) , where $H \leq G$ and $\psi \in \text{Irr}(H)$ such that $\psi^G = \chi, \mathbb{Q}(\chi) = \mathbb{Q}(\psi)$ and $H/\ker(\psi)$ is a cyclic, generalized quaternion, dihedral or semi-dihedral group.
- (2) Find an irreducible \mathbb{Q} -representation Ψ of H that affords the character $\Omega(\psi)$.
- (3) Induce Ψ to G .

Output: Ψ^G , an irreducible \mathbb{Q} -representation of G whose character is $\Omega(\chi)$.

Here, we call such a pair (H, ψ) a *required pair* for an irreducible rational matrix representation of G that affords the character $\Omega(\chi)$.

4. VZ p -group

A group G is called a VZ-group if all its nonlinear complex irreducible characters vanish off centre (see [19]). In this case, $G' \subseteq Z(G)$ and hence the nilpotency class of G is 2. The character degree set is given by $\text{cd}(G) = \{1, |G/Z(G)|^{1/2}\}$. Furthermore, G has $|Z(G)| - |Z(G)/G'|$ many inequivalent nonlinear irreducible complex characters, and there is a one-to-one correspondence between the sets $\text{nl}(G)$ and $\text{Irr}(Z(G)|G')$ (see [26, Section 3.1]). For any $\mu \in \text{Irr}(Z(G)|G')$, the corresponding $\chi_\mu \in \text{nl}(G)$ is defined as follows:

$$\chi_\mu(g) = \begin{cases} |G/Z(G)|^{1/2} \mu(g) & \text{if } g \in Z(G), \\ 0 & \text{otherwise.} \end{cases} \quad (1)$$

Observe that, being a nilpotency class 2 group, G can be written as a direct product of its Sylow subgroups. Since $\text{cd}(G) = \{1, |G/Z(G)|^{1/2}\}$, all the Sylow subgroups of G , except one, are abelian.

4.1. Rational representations of VZ p -groups. Let G be a VZ-group and let $\chi_\mu \in \text{nl}(G)$ (as defined in (1)).

Since G is monomial, there exists a subgroup H of G with index $|G/Z(G)|^{1/2}$, and $\psi \in \text{lin}(H)$, satisfying $\psi^G = \chi_\mu$. We now present the following results that provide a description of H and ψ .

PROPOSITION 21. *Let G be a VZ-group. Suppose H is a subgroup of G with index $|G/Z(G)|^{1/2}$ and $\psi \in \text{lin}(H)$. Then, $\psi^G = \chi_\mu \in \text{nl}(G)$ (as defined in (1)) if and only if H is normal in G such that $Z(G) \subset H$ and $\psi \downarrow_{Z(G)} = \mu$ with $\mu \in \text{Irr}(Z(G)|G')$.*

PROOF. Let $\psi \in \text{lin}(H)$ be such that $\psi^G = \chi_\mu$. Let T be a set of right coset representatives of H in G . Then, for $g \in G$, we have $\psi^G(g) = \sum_{g_i \in T} \psi^\circ(g_i g g_i^{-1})$, where ψ° is defined by $\psi^\circ(x) = \psi(x)$ if $x \in H$ and $\psi^\circ(x) = 0$ if $x \notin H$. Now, for $z \in Z(G)$, we get $\psi^G(z) = |G/Z(G)|^{1/2} \psi^\circ(z)$ and since $\psi^G = \chi_\mu$, we obtain $\psi^\circ(z) = \mu(z) = \psi(z)$. This implies that $Z(G) \subseteq H$ and $\psi \downarrow_{Z(G)} = \mu$. Since $G' \subset Z(G)$, H is normal in G .

Conversely, assume H is a subgroup of G with index $|G/Z(G)|^{1/2}$, $\psi \in \text{lin}(H)$ and $Z(G) \subseteq H$ with $\psi \downarrow_{Z(G)} = \mu$, where $\mu \in \text{Irr}(Z(G)|G')$. Claim: $\psi^G \in \text{nl}(G)$. In contrast, suppose that $\psi^G \notin \text{nl}(G)$, then ψ^G must be a sum of some linear characters of G . Hence, $G' \subseteq \ker(\psi^G)$, which is a contradiction as $\psi \downarrow_{Z(G)} = \mu$, where $\mu \in \text{Irr}(Z(G)|G')$. This proves the claim. \square

We prove Proposition 22 which describes a required pair of a VZ 2-group.

PROPOSITION 22. *Let G be a VZ 2-group and $\chi \in \text{nl}(G)$. Consider (H, ψ) as a required pair for an irreducible rational matrix representation of G that affords the character $\Omega(\chi)$. Then $H/\ker(\psi)$ is isomorphic to one of the following groups: cyclic group, quaternion group of order 8 denoted as Q_8 or dihedral group of order 8 denoted as D_8 .*

PROOF. Since H is a monomial group, there exists a subgroup K of H such that $\lambda^H = \psi$, where $\lambda \in \text{lin}(K)$. Consequently, we have $\lambda^G = \chi$. From Proposition 21, it follows that $Z(G) \leq K$, which implies that $G' \leq Z(G) \leq H$. It is worth noting that $H' \leq Z(H)$ and therefore, $(H/\ker(\psi))' \leq Z(H/\ker(\psi))$. Hence, $H/\ker(\psi)$ must be isomorphic to one of the groups: cyclic, Q_8 or D_8 . \square

Let G be a VZ 2-group and $\chi \in \text{Irr}(G)$. Consider a required pair (H, ψ) for a rational representation of G that affords the character $\Omega(\chi)$. Then by Proposition 22, $H/\ker(\psi)$ is isomorphic to one of the following: cyclic, Q_8 or D_8 .

Case 1 ($H/\ker(\psi)$ is cyclic). See Case 1 of Remark 19, to get an irreducible rational matrix representation of G that affords the character $\Omega(\chi)$.

Case 2 ($H/\ker(\psi) \cong D_8$). Suppose $H/\ker(\psi) = \langle a, b : a^4 = b^2 = 1, bab^{-1} = a^{-1} \rangle$ (that is $\psi \in \text{nl}(H)$). Then, $\bar{\psi} \in \text{FIrr}(H/\ker(\psi))$ and $\bar{\lambda} \in \text{lin}(K/\ker(\psi))$ are such that $\bar{\lambda}^{H/\ker(\psi)} = \bar{\psi}$ and $\mathbb{Q}(\bar{\lambda}) = \mathbb{Q}(\bar{\psi}) = \mathbb{Q}$, where $K/\ker(\psi) = \langle a^2, b \rangle$ and $\bar{\lambda}$ is defined as $\bar{\lambda}(a^2) = -1, \bar{\lambda}(b) = 1$. This shows that $(K/\ker(\psi), \bar{\lambda})$ is a required pair for the rational representation of $H/\ker(\psi)$ that affords the character $\Omega(\bar{\psi})$. This implies that (K, λ) is also a required pair for the rational representation of G that affords the character $\Omega(\chi)$. Note that $\mathbb{Q}(\lambda) = \mathbb{Q}(\chi) = \mathbb{Q}, [G : H] = \frac{1}{2}|G/Z(G)|^{1/2}$ and $[G : K] = |G/Z(G)|^{1/2}$.

Case 3 ($H/\ker(\psi) \cong Q_8$). Suppose $H/\ker(\psi) = \langle a, b : a^4 = b^4 = 1, bab^{-1} = a^{-1} \rangle$ (that is, $\psi \in \text{nl}(H)$). Then $\bar{\psi} \in \text{FIrr}(H/\ker(\psi))$ and $\bar{\lambda} \in \text{lin}(K/\ker(\psi))$ are such that $\bar{\lambda}^{H/\ker(\psi)} = \bar{\psi}$ and $[\mathbb{Q}(\bar{\lambda}) : \mathbb{Q}(\bar{\psi})] = 2$, where $K/\ker(\psi) = \langle a \rangle$ and $\bar{\lambda}$ is defined as $\bar{\lambda}(a) = \zeta_4$. Since $m_{\mathbb{Q}}(\bar{\psi}) = 2$, from Lemma 12, we get $\Omega(\bar{\lambda})^{H/\ker(\psi)} = \Omega(\bar{\psi})$. This implies that there exists a subgroup K of G and $\lambda \in \text{lin}(K)$ such that $\Omega(\lambda)^G = \Omega(\chi)$. Note that $[\mathbb{Q}(\lambda) : \mathbb{Q}(\chi)] = 2, [G : H] = \frac{1}{2}|G/Z(G)|^{1/2}$ and $[G : K] = |G/Z(G)|^{1/2}$.

REMARK 23. In view of the above discussion and Algorithm 15, to obtain an irreducible rational matrix representation of a VZ p -group G (p is any prime) that affords the character $\Omega(\chi)$, where $\chi \in \text{Irr}(G)$, we need to do the following.

- (1) If $m_{\mathbb{Q}}(\chi) = 1$, then find $H \leq G$ and $\psi \in \text{lin}(H)$ such that $\psi^G = \chi, \mathbb{Q}(\psi) = \mathbb{Q}(\chi)$.
- (2) If $m_{\mathbb{Q}}(\chi) = 2$, then find $H \leq G$ and $\psi \in \text{lin}(H)$ so that $\psi^G = \chi, [\mathbb{Q}(\psi) : \mathbb{Q}(\chi)] = 2$.

We call such a pair (H, ψ) a special required pair for an irreducible rational matrix representation of a VZ p -group G that affords the character $\Omega(\chi)$, where $\chi \in \text{Irr}(G)$.

Now, we prove Lemma 24 which provides a description of the character fields, which is useful to obtain a special required pair of a VZ p -group.

LEMMA 24. Let G be a VZ-group and let $\chi_{\mu} \in \text{nl}(G)$ (as defined in (1)). Consider a subgroup H of G with index $|G/Z(G)|^{1/2}$ and $\psi_{\mu} \in \text{lin}(H)$ such that $\psi_{\mu}^G = \chi_{\mu}$. Then, $\mathbb{Q}(\psi_{\mu}) = \mathbb{Q}(\chi_{\mu})$ if and only if $|\ker(\psi_{\mu})/\ker(\mu)| = |G/Z(G)|^{1/2}$, and $[\mathbb{Q}(\psi_{\mu}) : \mathbb{Q}(\chi_{\mu})] = 2$ if and only if $|\ker(\psi_{\mu})/\ker(\mu)| = \frac{1}{2}|G/Z(G)|^{1/2}$.

PROOF. By Proposition 21, $\psi_\mu \downarrow_{Z(G)} = \mu$ with $\mu \in \text{Irr}(Z(G)|G')$ and $\mathbb{Q}(\chi_\mu) = \mathbb{Q}(\mu)$. Observe that

$$\begin{aligned} \mathbb{Q}(\psi_\mu) = \mathbb{Q}(\chi_\mu) = \mathbb{Q}(\mu) &\iff \mathbb{Q}(\zeta_{|H/\ker(\psi_\mu)|}) = \mathbb{Q}(\zeta_{|Z(G)/\ker(\mu)|}) \\ &\iff |H/\ker(\psi_\mu)| = |Z(G)/\ker(\mu)| \\ &\iff |\ker(\psi_\mu)| = |H/Z(G)||\ker(\mu)| \\ &\iff |\ker(\psi_\mu)| = |G/Z(G)|^{1/2}|\ker(\mu)|. \end{aligned}$$

Again,

$$\begin{aligned} [\mathbb{Q}(\psi_\mu) : \mathbb{Q}(\chi_\mu)] = [\mathbb{Q}(\psi_\mu) : \mathbb{Q}(\mu)] = 2 &\iff |\mathbb{Q}(\zeta_{|H/\ker(\psi_\mu)|}) : \mathbb{Q}(\zeta_{|Z(G)/\ker(\mu)|})| = 2 \\ &\iff |H/\ker(\psi_\mu)| = 2|Z(G)/\ker(\mu)| \\ &\iff |\ker(\psi_\mu)| = \frac{1}{2}|H/Z(G)||\ker(\mu)| \\ &\iff |\ker(\psi_\mu)| = \frac{1}{2}|G/Z(G)|^{1/2}|\ker(\mu)|. \end{aligned}$$

This completes the proof. □

COROLLARY 25. Let G be a VZ-group. Suppose H is a subgroup of G with index $|G/Z(G)|^{1/2}$ and $\psi \in \text{lin}(H)$ is such that $\psi^G \in \text{nl}(G)$. If one of the following is satisfied

- (a) $\text{cd}(G) = \{1, p\}$;
- (b) $|G'| = p$,

then H is abelian.

PROOF. Let H be a subgroup of G of index $|G/Z(G)|^{1/2}$ and let $\psi \in \text{lin}(G)$ be such that $\psi^G \in \text{nl}(G)$. Then by Proposition 21, $Z(G) \subseteq H$ and $\psi \downarrow_{Z(G)} = \mu$.

Case (a): Suppose $\text{cd}(G) = \{1, p\}$. This implies $|G/Z(G)| = p^2$ and $|H/Z(G)| = p$. This shows that H is abelian.

Case (b): Suppose $|G'| = p$. In this case, $|H'| \in \{1, p\}$. Since $H' \subseteq \ker(\psi^G) = \text{Core}_G(\ker(\psi))$ and $\psi^G \in \text{nl}(G)$, we get $|H'| = 1$. Thus, H is abelian. □

As we know, for a VZ-group G , G' is an elementary abelian subgroup, then by Corollary 25, we get the following result.

COROLLARY 26. Let G be a VZ-group. Suppose H is a subgroup of G with index $|G/Z(G)|^{1/2}$ and $\psi \in \text{lin}(H)$ is such that $\psi^G \in \text{nl}(G)$. If $Z(G)$ is cyclic, then H is abelian.

COROLLARY 27. Suppose G is a VZ p -group of order $\leq p^5$ (p is any prime). Let H be a subgroup G of index $|G/Z(G)|^{1/2}$ with $\psi \in \text{lin}(H)$ such that $\psi^G \in \text{nl}(G)$. Then, H is abelian.

PROOF. Suppose G is a VZ p -group of order $\leq p^5$. In this case, $\text{cd}(G) \in \{\{1, p\}, \{1, p^2\}\}$. If $\text{cd}(G) = \{1, p\}$, then by Corollary 25, H is abelian. If $\text{cd}(G) = \{1, p^2\}$, then $\sqrt{|G/Z(G)|} = p^2$ and hence, $|G'| = |Z(G)| = p$. Again, by Corollary 25, it follows that H is abelian. □

REMARK 28. The conclusion of Corollary 27 need not hold for higher order VZ p -groups. For instance, consider the group

$$G = G_{(15,1)} = \langle \alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6 : [\alpha_3, \alpha_5] = \alpha_1, [\alpha_4, \alpha_5] = \alpha_2, [\alpha_3, \alpha_6] = \alpha_2, [\alpha_4, \alpha_6] = \alpha_1^\nu, \alpha_1^p = \alpha_2^p = \alpha_3^p = \alpha_4^p = \alpha_5^p = \alpha_6^p = 1 \rangle,$$

of order p^6 ($p \geq 7$), where ν denotes the smallest positive integer that is a quadratic nonresidue (mod p) (see [20]). Here, $Z(G) = G' = \langle \alpha_1, \alpha_2 \rangle$. It is easy to observe that G is a VZ p -group and $\text{cd}(G) = \{1, p^2\}$. Set $H = \langle \alpha_1, \alpha_2, \alpha_3, \alpha_5 \rangle$. Then $H' = \langle \alpha_1 \rangle$ and $H/H' = \langle \alpha_2H', \alpha_3H', \alpha_5H' \rangle \cong C_p \times C_p \times C_p$. Define $\bar{\psi} \in \text{Irr}(H/H')$ such that $\bar{\psi}(\alpha_2H') = \zeta_p, \bar{\psi}(\alpha_3H') = 1$ and $\bar{\psi}(\alpha_5H') = 1$. Now, define a character ψ of H by taking the lift of $\bar{\psi} \in \text{Irr}(H/H')$. Then, $\psi \in \text{lin}(H)$ and $\psi \downarrow_{Z(G)} \in \text{Irr}(Z(G)|G') = \text{Irr}(Z(G)) \setminus 1_{Z(G)}$, where $1_{Z(G)}$ is the trivial character of $Z(G)$. Then by Proposition 21, $\psi^G \in \text{nl}(G)$; however, H is nonabelian.

4.2. Rational group algebra of a VZ p -group. Let G be a VZ-group. Then the Wedderburn decomposition of $\mathbb{C}G$ is as follows:

$$\mathbb{C}G \cong |G/G'| \mathbb{C} \bigoplus (|Z(G)| - |Z(G)/G'|) M_{|G/Z(G)|^{1/2}}(\mathbb{C}),$$

where $M_{|G/Z(G)|^{1/2}}(\mathbb{C})$ denotes the ring of matrices of order $|G/Z(G)|^{1/2}$ over \mathbb{C} . In this subsection, we compute the Wedderburn decomposition of the rational group algebra of a finite VZ p -group.

Let G be a finite group and let $\chi, \psi \in \text{Irr}(G)$. It is well known that if χ and ψ are Galois conjugates over \mathbb{Q} , then $\ker(\chi) = \ker(\psi)$. Now, we begin with the following easy observations.

LEMMA 29. Let G be a finite group and let $\chi, \psi \in \text{lin}(G)$ be such that $\ker(\chi) = \ker(\psi)$. Then, χ and ψ are Galois conjugates over \mathbb{Q} .

In general, if $\chi, \psi \in \text{nl}(G)$ are such that $\ker(\chi) = \ker(\psi)$, then χ may not be Galois conjugate to ψ over \mathbb{Q} . However, in the case of VZ groups, this is true.

LEMMA 30. Let G be a VZ-group and let $\chi, \psi \in \text{Irr}(G)$. Then χ and ψ are Galois conjugates over \mathbb{Q} if and only if $\ker(\chi) = \ker(\psi)$.

PROOF. If $\chi, \psi \in \text{lin}(G)$, then the result follows from Lemma 29. Now let $\chi, \psi \in \text{nl}(G)$ be such that $\ker(\chi) = \ker(\psi)$. Then, in view of (1), there exist $\mu, \nu \in \text{Irr}(Z(G) | G')$ such that $\chi \downarrow_{Z(G)} = \mu$ and $\psi \downarrow_{Z(G)} = \nu$. Observe that $\ker(\chi) = \ker(\mu)$ and $\ker(\psi) = \ker(\nu)$. Thus, μ and ν are Galois conjugates and hence, χ and ψ are Galois conjugates over \mathbb{Q} . This completes the proof. □

LEMMA 31. Consider a finite abelian group G , where d divides the exponent of G . Let a_d denote the number of cyclic subgroups of G with order d . Then the number of non-Galois conjugate characters χ satisfying $\mathbb{Q}(\chi) = \mathbb{Q}(\zeta_d)$ is precisely a_d .

PROOF. See [1, Lemma 1]. □

Analogously to Lemma 31, we have the following lemma for VZ-groups.

LEMMA 32. Consider a VZ-group G . Let $\chi_\mu \in \text{nl}(G)$ (defined in (1)). Assume that a_d and a'_d represent the number of cyclic subgroups of order d of $Z(G)$ and $Z(G)/G'$, respectively. Then the following statements hold.

- (1) The number of non-Galois conjugate nonlinear characters χ_μ of G satisfying $\mathbb{Q}(\chi_\mu) = \mathbb{Q}(\zeta_d)$, where $d \mid \exp(Z(G))$ but $d \nmid \exp(Z(G)/G')$, is equal to a_d .
- (2) The number of non-Galois conjugate nonlinear characters χ_μ of G satisfying $\mathbb{Q}(\chi_\mu) = \mathbb{Q}(\zeta_d)$, where $d \mid \exp(Z(G))$ and $d \mid \exp(Z(G)/G')$, is equal to $a_d - a'_d$.

PROOF. By the one-to-one correspondence between $\text{nl}(G)$ and $\text{Irr}(Z(G)|G')$, we get that $\ker(\chi_\mu) = \ker(\mu)$ and $\mathbb{Q}(\chi_\mu) = \mathbb{Q}(\mu)$. Observe that $\text{Irr}(Z(G)) = \text{Irr}(Z(G)|G') \sqcup \text{Irr}(Z(G)/G')$. Hence, if $d \mid \exp(Z(G))$ but $d \nmid \exp(Z(G)/G')$, then by Lemma 31, the number of non-Galois conjugate characters $\mu \in \text{Irr}(Z(G)|G')$ such that $\mathbb{Q}(\mu) = \mathbb{Q}(\zeta_d)$ is equal to a_d . This proves Lemma 32(1).

Similarly, if $d \mid \exp(Z(G))$ and $d \mid \exp(Z(G)/G')$, then the number of non-Galois conjugate characters $\mu \in \text{Irr}(Z(G)|G')$ such that $\mathbb{Q}(\mu) = \mathbb{Q}(\zeta_d)$ is the difference between the number of non-Galois conjugate characters in $\text{Irr}(Z(G))$ whose character field is $\mathbb{Q}(\zeta_d)$ and the number of non-Galois characters in $\text{Irr}(Z(G)/G')$ whose character field is $\mathbb{Q}(\zeta_d)$. Hence, by using Lemma 31, we get Lemma 32(2). □

Now, we prove Theorem 1, which provides the Wedderburn decomposition of a VZ p -group, where p is an odd prime.

PROOF OF THEOREM 1. Let G be a finite VZ p -group (odd prime p) and $\chi \in \text{Irr}(G)$. Suppose ρ is an irreducible \mathbb{Q} -representation of G that affords the character $\Omega(\chi)$. Let $A_{\mathbb{Q}}(\chi)$ be the simple component of the Wedderburn decomposition of $\mathbb{Q}G$ corresponding to ρ , that is isomorphic to $M_n(D)$ for some $n \in \mathbb{N}$ and a division ring D . From Lemma 14, $m_{\mathbb{Q}}(\chi) = 1$ and from Lemma 8, we have $[D : Z(D)] = m_{\mathbb{Q}}(\chi)^2$ and $Z(D) = \mathbb{Q}(\chi)$. Therefore, $D = Z(D) = \mathbb{Q}(\chi)$. Now consider $\rho = \rho_1 \oplus \rho_2 \oplus \dots \oplus \rho_k$, where for $1 \leq i \leq k$, ρ_i is a complex irreducible representation of G affording χ^{σ_i} for some $\sigma_i \in \text{Gal}(\mathbb{Q}(\chi) : \mathbb{Q})$. Here, $k = [\mathbb{Q}(\chi) : \mathbb{Q}]$. Since $m_{\mathbb{Q}}(\chi) = 1$, we observe that $n = \chi(1)$.

Let $\chi \in \text{lin}(G)$ and suppose ρ is the irreducible \mathbb{Q} -representation of G affording $\Omega(\chi)$. Let $\bar{\chi} \in \text{Irr}(G/G')$ be such that $\bar{\chi}(gG') = \chi(g)$. Hence, $A_{\mathbb{Q}}(\bar{\chi}) \cong \mathbb{Q}(\bar{\chi})$. Since G/G' is abelian, according to Lemma 6, the simple components of the Wedderburn decomposition of $\mathbb{Q}G$ corresponding to all irreducible \mathbb{Q} -representations of G whose kernels contain G' contribute

$$\bigoplus_{d_1 \mid m_1} a_{d_1} \mathbb{Q}(\zeta_{d_1})$$

in $\mathbb{Q}G$, where m_1 is the exponent of G/G' and a_{d_1} is the number of cyclic subgroups of G/G' of order d_1 .

Now, let ρ be an irreducible \mathbb{Q} -representation of G that affords the character $\Omega(\chi_\mu)$, where $\chi_\mu \in \text{nl}(G)$ as defined in (1). Here, $\chi_\mu(1) = |G/Z(G)|^{1/2}$ and $\mathbb{Q}(\chi_\mu) = \mathbb{Q}(\mu)$.

Therefore, by the above discussion, $A_{\mathbb{Q}}(\chi_{\mu}) \cong M_{|G/Z(G)|^{1/2}}(\mathbb{Q}(\mu))$. Observe that $\mathbb{Q}(\chi_{\mu}) = \mathbb{Q}(\mu) = \mathbb{Q}(\zeta_d)$ for some $d \mid \exp(Z(G))$. Now, we have two cases.

Case 1 ($d \mid \exp(Z(G))$ but $d \nmid \exp(Z(G)/G')$). In this case, from Lemma 32(1), the number of irreducible \mathbb{Q} representations of G that afford the character $\Omega(\chi_{\mu})$ is equal to the number of cyclic subgroups of $Z(G)$ of order d , where $\mathbb{Q}(\chi_{\mu}) = \mathbb{Q}(\mu) = \mathbb{Q}(\zeta_d)$.

Case 2 ($d \mid \exp(Z(G))$ and $d \mid \exp(Z(G)/G')$). In this case, from Lemma 32(2), the number of irreducible \mathbb{Q} representations of G that afford the character $\Omega(\chi_{\mu})$ is equal to the difference of the number of cyclic subgroups of $Z(G)$ of order d and the number of cyclic subgroups of $Z(G)/G'$ of order d , where $\mathbb{Q}(\chi_{\mu}) = \mathbb{Q}(\mu) = \mathbb{Q}(\zeta_d)$.

Let m_2 and m_3 be the exponents of $Z(G)$ and $Z(G)/G'$, respectively. Then by the above discussion, the simple components of the Wedderburn decomposition of $\mathbb{Q}G$ corresponding to all irreducible \mathbb{Q} -representations of G whose kernels do not contain G' contribute

$$\bigoplus_{d_2 \mid m_2, d_2 \nmid m_3} a_{d_2} M_{|G/Z(G)|^{1/2}}(\mathbb{Q}(\zeta_{d_2})) \quad \bigoplus_{d_2 \mid m_2, d_2 \mid m_3} (a_{d_2} - a'_{d_2}) M_{|G/Z(G)|^{1/2}}(\mathbb{Q}(\zeta_{d_2}))$$

in $\mathbb{Q}G$, where a_{d_2} and a'_{d_2} are the number of cyclic subgroups of $Z(G)$ of order d_2 and the number of cyclic subgroups of $Z(G)/G'$ of order d_2 , respectively. Therefore, the result follows. \square

COROLLARY 33. *Let G be a finite VZ p -group (odd prime p) with cyclic centre $Z(G)$. Then the Wedderburn decomposition of the group algebra $\mathbb{Q}G$ is given by*

$$\mathbb{Q}G \cong \mathbb{Q}(G/G') \bigoplus M_{|G/Z(G)|^{1/2}}(\mathbb{Q}(\zeta_{|Z(G)|})).$$

PROOF. Let G be a VZ p -group (odd prime p). It is known that $G' \subseteq Z(G)$ and G' is an elementary abelian p -group. Since $Z(G)$ is cyclic, $|G'| = p$. Let $\mu \in \text{Irr}(Z(G)|G')$. It follows that μ is faithful and $\mathbb{Q}(\mu) = \mathbb{Q}(\zeta_{|Z(G)|})$. Therefore, all nonlinear complex irreducible characters G are faithful and Galois conjugate to each other. This completes the proof. \square

COROLLARY 34. *Let G be an extra special p -group (odd prime p) of order p^{1+2n} . Then,*

$$\mathbb{Q}G \cong \mathbb{Q} \bigoplus (p^{2n-1} + p^{2n-2} + \dots + p + 1) \mathbb{Q}(\zeta_p) \bigoplus M_{p^n}(\mathbb{Q}(\zeta_p)).$$

PROOF. One can easily observe that G is a VZ p -group and $|G/Z(G)|^{1/2} = p^n = \chi(1)$, where $\chi \in \text{nl}(G)$. Therefore, the result follows from Lemma 6 and Corollary 33. \square

REMARK 35

- (1) Corollary 34 shows that the rational group algebras of two nonisomorphic groups may be isomorphic.

- (2) Suppose G is a nonabelian p -group of order p^3 (odd prime p). Then by Corollary 34,

$$\mathbb{Q}G \cong \mathbb{Q} \bigoplus (p + 1)\mathbb{Q}(\zeta_p) \bigoplus M_p(\mathbb{Q}(\zeta_p)).$$

This is also computed in [4, Theorems 3 and 4]. The authors computed Wedderburn components of $\mathbb{Q}G$ by using the Shoda pair concept.

COROLLARY 36. *Let G and H be two isoclinic VZ p -groups (odd prime p) of the same order. Then the Wedderburn decompositions of $\mathbb{Q}G$ and $\mathbb{Q}H$ are isomorphic if and only if $G/G' \cong H/H'$ and $Z(G) \cong Z(H)$.*

PROOF. The result follows from Theorem 1. □

Now, we discuss the rational group algebra of a VZ 2-group.

LEMMA 37. *Let G be a VZ 2-group. Suppose $\chi \in \text{nl}(G)$ is such that $m_{\mathbb{Q}}(\chi) = 2$. Then, $\mathbb{Q}(\chi) = \mathbb{Q}$.*

PROOF. The result follows from Lemma 17 and Proposition 22. □

Now we prove Theorem 2, which provides a combinatorial description for the Wedderburn decomposition of the rational group algebra of a VZ 2-group.

PROOF OF THEOREM 2. Let $\chi_{\mu} \in \text{nl}(G)$ as defined in (1). We have two cases for nonlinear irreducible complex characters of G .

Case 1 ($m_{\mathbb{Q}}(\chi_{\mu}) = 2, \chi_{\mu} \in \text{nl}(G)$). By Lemma 37, there are k rational irreducible representations of G , which correspond to k simple components of $\mathbb{Q}G$, denoted as $A_{\mathbb{Q}}(\chi_{\mu})$. We know that $A_{\mathbb{Q}}(\chi_{\mu}) \cong M_n(D)$ for some $n \in \mathbb{N}$ and a division ring D . By [31, Theorem 2.4], we have $n = \frac{1}{2}|G/Z(G)|^{1/2}$. Now by using Lemmas 8 and 37, we get $Z(D) = \mathbb{Q}(\chi) = \mathbb{Q}$ and $[D : \mathbb{Q}] = 4$.

Claim. $A_{\mathbb{Q}}(\chi_{\mu}) = M_{1/2|G/Z(G)|^{1/2}}(\mathbb{H}(\mathbb{Q}))$.

To prove our claim, let (H, ψ) be a required pair to compute an irreducible rational matrix representation of G that affords the character χ_{μ} . Then, $H/\ker(\psi) \cong Q_8$ (by Algorithm 20 and Proposition 22). Therefore, $\bar{\psi} \in \text{FIrr}(H/\ker(\psi))$ and $\bar{\psi} = (\bar{\lambda})^{H/\ker(\psi)}$, where $(\bar{\lambda}) \in \text{FIrr}(N/\ker(\psi))$ and $N/\ker(\psi) \cong C_4$. Hence, by Proposition 21, $N \trianglelefteq G$ with $\lambda \in \text{lin}(N)$ such that $\lambda^G = \chi_{\mu}$. Now, assume that $K = \ker(\psi)$ and observe that $N_G(K) = H$. Further, N/K is cyclic and a maximal abelian subgroup of H/K . Hence, (N, K) is an extremely strong Shoda pair (see [4]). Furthermore, from [15, Theorem 3.5.5],

$$A_{\mathbb{Q}}(\chi_{\mu}) = A_{\mathbb{Q}}(G, N, K) \cong M_{1/2|G/Z(G)|^{1/2}}(\mathbb{Q}(\zeta_4) * H/N) \cong M_{1/2|G/Z(G)|^{1/2}}(\mathbb{H}(\mathbb{Q})).$$

This completes the proof of the above claim and Case 1.

Case 2 ($m_{\mathbb{Q}}(\chi_{\mu}) = 1, \chi_{\mu} \in \text{nl}(G)$). Here, $A_{\mathbb{Q}}(\chi_{\mu}) \cong M_n(D)$ for some $n \in \mathbb{N}$ and a division ring D . By [31, Theorem 2.4], we have $n = |G/Z(G)|^{1/2}$. Now again, by using Lemma 8, we get $D = \mathbb{Q}(\chi_{\mu})$. Now we have two sub-cases:

Sub-case 2(1)($\mathbb{Q}(\chi_\mu) = \mathbb{Q}$). From Case 1 and Lemma 37, there are $a_2 - a'_2 - k$ Galois conjugacy classes of complex irreducible characters χ_μ such that $m_{\mathbb{Q}}(\chi_\mu) = 1$ and $\mathbb{Q}(\chi_\mu) = \mathbb{Q}$. Therefore, $\mathbb{Q}G$ contains $(a_2 - a'_2 - k)M_{|G/Z(G)|^{1/2}}(\mathbb{Q})$.

Sub-case 2(2)($\mathbb{Q}(\chi_\mu) \neq \mathbb{Q}$). In this sub-case, $\mathbb{Q}(\chi_\mu) = \mathbb{Q}(\mu) = \mathbb{Q}(\zeta_d)$, where $d \geq 4$. Observe that either $d \mid \exp(Z(G))$ but $d \nmid \exp(Z(G)/G')$, or $d \mid \exp(Z(G))$ and $d \mid \exp(Z(G)/G')$. Now by using a similar argument to that mentioned in the proof of Theorem 1, $\mathbb{Q}G$ contains

$$\bigoplus_{d_2|m_2, d_2 \nmid m_3} a_{d_2} M_{|G/Z(G)|^{1/2}}(\mathbb{Q}(\zeta_{d_2})) \oplus \bigoplus_{d_2|m_2, d_2|m_3} (a_{d_2} - a'_{d_2}) M_{|G/Z(G)|^{1/2}}(\mathbb{Q}(\zeta_{d_2})).$$

This completes the discussion of Case 2.

Now by using Lemma 6, Case 1 and Case 2, we get the result. □

COROLLARY 38. *Let G be a VZ 2-group G such that $Z(G)$ is cyclic and $|Z(G)| \geq 4$. Then the following hold:*

- (1) $m_{\mathbb{Q}}(\chi) = 1$ for each $\chi \in \text{Irr}(G)$;
- (2) $\mathbb{Q}G \cong \mathbb{Q}(G/G') \oplus M_{|G/Z(G)|^{1/2}}(\mathbb{Q}(\zeta_{|Z(G)|}))$.

PROOF. It follows from Theorem 2 and Lemma 37. □

COROLLARY 39. *Suppose G is a VZ 2-group with elementary abelian centre. Then,*

$$\mathbb{Q}G \cong \mathbb{Q}(G/G') \oplus kM_{1/2|G/Z(G)|^{1/2}}(\mathbb{H}(\mathbb{Q})) \oplus k'M_{|G/Z(G)|^{1/2}}(\mathbb{Q}),$$

where k and k' denote the number of nonlinear complex irreducible characters of G with Schur index 2 and the number of nonlinear complex irreducible characters of G with Schur index of 1, respectively.

The counting of rational irreducible representations of an abelian group can be determined using Lemma 6 and Remark 7. Corollary 40 provides a characterization for counting irreducible rational representations of VZ-groups.

COROLLARY 40. *For a VZ-group G , let n, x, y and z represent the total numbers of irreducible rational representations of $G, G/G', Z(G)$ and $Z(G)/G'$, respectively. It follows that $n = x + y - z$.*

PROOF. Observe that $|\{\eta \in \text{Irr}_{\mathbb{Q}}(G) : G' \subseteq \ker(\eta)\}| = x$. Further,

$$\begin{aligned} &|\{\eta \in \text{Irr}_{\mathbb{Q}}(G) : G' \not\subseteq \ker(\eta)\}| \\ &= \text{the number of Galois conjugacy classes of } \text{Irr}(Z(G)) \text{ over } \mathbb{Q} \\ &\quad - \text{the number of Galois conjugacy classes of } \text{Irr}(Z(G)/G') \text{ over } \mathbb{Q}. \end{aligned}$$

This completes the proof. □

In the case of a cyclic centre, we have the following corollary.

COROLLARY 41. *If G is a VZ p -group and $Z(G)$ is cyclic, then G has only one rational irreducible representation whose kernel does not contain G' .*

4.3. Primitive central idempotents in rational group algebras of VZ-groups. Let G be a finite group. An element e in $\mathbb{Q}G$ is an idempotent if $e^2 = e$. A primitive central idempotent e in $\mathbb{Q}G$ is one that belongs to the centre of $\mathbb{Q}G$ and cannot be expressed as $e = e' + e''$, where e' and e'' are nonzero idempotents such that $e'e'' = 0$. It is well known that a complete set of primitive central idempotents of $\mathbb{Q}G$ determines the decomposition of $\mathbb{Q}G$ into a direct sum of simple sub-algebras. Specifically, if e is a primitive central idempotent of $\mathbb{Q}G$, then the corresponding simple component of $\mathbb{Q}G$ is $\mathbb{Q}Ge$. For $\chi \in \text{Irr}(G)$, the expression

$$e(\chi) = \frac{\chi(1)}{|G|} \sum_{g \in G} \chi(g)g^{-1}$$

defines a primitive central idempotent of $\mathbb{C}G$. In fact, the set $\{e(\chi) : \chi \in \text{Irr}(G)\}$ forms a complete set of primitive central idempotents of $\mathbb{C}G$. Moreover, for $\chi \in \text{Irr}(G)$, we define

$$e_{\mathbb{Q}}(\chi) := \sum_{\sigma \in \text{Gal}(\mathbb{Q}(\chi)/\mathbb{Q})} e(\chi^\sigma).$$

Then $e_{\mathbb{Q}}(\chi)$ is a primitive central idempotent in $\mathbb{Q}G$. For a subset X of G , define

$$\widehat{X} = \frac{1}{|X|} \sum_{x \in X} x \in \mathbb{Q}G,$$

and for a normal subgroup N of G , define

$$\epsilon(G, N) = \begin{cases} \widehat{G} & \text{if } G = N; \\ \prod_{D/N \in M(G/N)} (\widehat{N} - \widehat{D}) & \text{otherwise,} \end{cases}$$

where $M(G/N)$ represents the set of minimal nontrivial normal subgroups D/N of G/N , with D being a subgroup of G that contains N . In this subsection, we compute a complete set of primitive central idempotents of the rational group algebra of a VZ-group. Let us start with a general result.

LEMMA 42 [15, Lemma 3.3.2]. *Let G be a finite group. If $\chi \in \text{lin}(G)$ and $N = \ker(\chi)$, then the following hold:*

- (1) $e_{\mathbb{Q}}(\chi) = \epsilon(G, N)$;
- (2) $\mathbb{Q}G\epsilon(G, N) \cong \mathbb{Q}(\zeta_{|G/N|})$.

Theorem 43 provides a characterization of primitive central idempotents and their corresponding simple components in $\mathbb{Q}G$ for a VZ p -group.

THEOREM 43. *Let G be a VZ-group and let $\chi_\mu \in \text{nl}(G)$ (as defined in (1)) with $N = \ker(\chi_\mu) = \ker(\mu)$. Then the following statements hold:*

- (1) $e_{\mathbb{Q}}(\chi_\mu) = \epsilon(Z(G), N)$;
- (2) if G is a VZ p -group (odd prime p), then $\mathbb{Q}G\epsilon(G, N) \cong M_{|G/Z(G)|^{1/2}}(\mathbb{Q}(\zeta_{|Z(G)/N|}))$;

- (3) if G is a VZ 2-group, then $\mathbb{Q}G\epsilon(G, N) \cong M_{|G/Z(G)|^{1/2}}(\mathbb{Q}(\zeta_{|Z(G)/N|}))$ when $m_{\mathbb{Q}}(\chi_{\mu}) = 1$, and $\mathbb{Q}G\epsilon(G, N) \cong M_{1/2|G/Z(G)|^{1/2}}(\mathbb{H}(\mathbb{Q}))$ when $m_{\mathbb{Q}}(\chi_{\mu}) = 2$.

PROOF. From (1), it is easy to observe that $e(\chi_{\mu}) = e(\mu)$.

Furthermore, we can observe:

$$\begin{aligned} e_{\mathbb{Q}}(\chi_{\mu}) &= \sum_{\sigma \in \text{Gal}(\mathbb{Q}(\chi_{\mu})/\mathbb{Q})} e(\chi_{\mu}^{\sigma}) \\ &= \sum_{\sigma \in \text{Gal}(\mathbb{Q}(\mu)/\mathbb{Q})} e(\chi_{\mu}^{\sigma}) \\ &= \sum_{\sigma \in \text{Gal}(\mathbb{Q}(\mu)/\mathbb{Q})} e(\mu^{\sigma}) \\ &= e_{\mathbb{Q}}(\mu) \\ &= \epsilon(Z(G), N) \quad (\text{from Lemma 42}), \end{aligned}$$

where $N = \ker(\mu)$. Moreover, if G is a VZ p -group (odd prime p), from Theorem 1, the simple component of $\mathbb{Q}G$ corresponding to $\chi_{\mu} \in \text{nl}(G)$ is given by

$$A_{\mathbb{Q}}(\chi_{\mu}) \cong M_{|G/Z(G)|^{1/2}}(\mathbb{Q}(\mu)) = M_{|G/Z(G)|^{1/2}}(\mathbb{Q}(\zeta_{|Z(G)/N|})).$$

Hence,

$$\mathbb{Q}Ge_{\mathbb{Q}}(\chi_{\mu}) = \mathbb{Q}G\epsilon(G, N) \cong M_{|G/Z(G)|^{1/2}}(\mathbb{Q}(\zeta_{|Z(G)/N|})).$$

Similarly, statement (3) follows from Theorem 2. □

REMARK 44. In [5, Corollary 2], primitive central idempotents of the rational group algebra of VZ-groups have been computed. However, our approach, which is based on character properties, offers a direct proof.

5. p -group of order p^3

Let G be a nonabelian p -group (odd prime p) of order p^3 . It is well known that G is isomorphic to one of the following two groups:

$$\begin{aligned} \Phi_2(21) &= \langle \alpha, \alpha_1, \alpha_2 : [\alpha_1, \alpha] = \alpha^p = \alpha_2, \alpha_1^p = \alpha_2^p = 1 \rangle, \quad \text{and} \\ \Phi_2(111) &= \langle \alpha, \alpha_1, \alpha_2 : [\alpha_1, \alpha] = \alpha_2, \alpha^p = \alpha_1^p = \alpha_2^p = 1 \rangle, \end{aligned}$$

(see [13, Section 4.3]). It is easy to check that both the groups are VZ p -groups. In both cases, we have $Z(G) = G' = \langle \alpha_2 \rangle \cong C_p$, $cd(G) = \{1, p\}$, $|\text{nl}(G)| = |Z(G)| - 1$ and $G/G' = \langle \alpha G', \alpha_1 G' \rangle \cong C_p \times C_p$. Since G is a VZ p -group, the nonlinear characters of G can be defined as follows:

$$\chi_{\mu}(g) = \begin{cases} p\mu(g) & \text{if } g \in Z(G), \\ 0 & \text{otherwise,} \end{cases} \tag{2}$$

where $\mu \in \text{Irr}(Z(G)|G')$. The rational representations of G are characterized in Proposition 45.

PROPOSITION 45. *Let G be a nonabelian group of order p^3 (odd prime p). Then the following statements hold.*

- (1) $|\text{Irr}_{\mathbb{Q}}^{(1)}(G)| = 1$, $|\text{Irr}_{\mathbb{Q}}^{(\phi(p))}(G)| = p + 1$ and $|\text{Irr}_{\mathbb{Q}}^{(\phi(p^2))}(G)| = 1$.
- (2) *A special required pair (H, ψ_{μ}) to determine a rational matrix representation of G whose character is $\Omega(\chi_{\mu})$ (χ_{μ} is defined in (2)) is given by $H = \langle \alpha_1, \alpha_2 \rangle$, and $\psi_{\mu} \in \text{lin}(H)$ defined by $\psi_{\mu}(\alpha_1) = 1$ and $\psi_{\mu}(\alpha_2) = \mu(\alpha_2)$.*

PROOF. The proof of Proposition 45(1) is obvious. As $Z(G) \subset H$ and $\psi_{\mu} \downarrow_{Z(G)} = \mu$, we get $\psi_{\mu}^G = \chi_{\mu}$ (from Proposition 21). Furthermore, since $\psi_{\mu}(\alpha_1) = 1$ and $\psi_{\mu}(\alpha_2) = \mu(\alpha_2)$, it follows that $\mathbb{Q}(\psi_{\mu}) = \mathbb{Q}(\mu) = \mathbb{Q}(\chi_{\mu})$. This completes the Proof of Proposition 45(2). \square

REMARK 46. In general, a required pair to find an irreducible rational matrix representation of G whose character is $\Omega(\chi)$ may not be unique. For example, consider $G = \Phi_2(111)$ and $\chi_{\mu} \in \text{nl}(G)$, as defined in (2). Take $H_1 = \langle \alpha, \alpha_2 \rangle$ and choose $\psi'_{\mu} \in \text{lin}(H_1)$ such that $\psi'_{\mu}(\alpha) = 1$ and $\psi'_{\mu}(\alpha_2) = \mu(\alpha_2)$. The pair (H_1, ψ'_{μ}) is also a special required pair to find an irreducible rational matrix representation of G that affords the character $\Omega(\chi_{\mu})$.

In Example 47, we show how to find an irreducible rational matrix representation associated with a required pair.

EXAMPLE 47. Consider

$$G = \Phi_2(21) = \langle \alpha, \alpha_1, \alpha_2 : [\alpha_1, \alpha] = \alpha^p = \alpha_2, \alpha_1^p = \alpha_2^p = 1 \rangle.$$

We have $Z(G) = G' = \langle \alpha_2 \rangle \cong C_p$. Let $\mu \in \text{Irr}(Z(G)|G')$ be such that $\mu(\alpha_2) = \zeta_p$. The character χ_{μ} defined in (2) is a nonlinear irreducible complex character of G . From Proposition 45, a special required pair to find an irreducible rational matrix representation of G affording the character $\Omega(\chi_{\mu})$ is (H, ψ_{μ}) , where $H = \langle \alpha_1, \alpha_2 \rangle$ and $\psi_{\mu} \in \text{lin}(H)$ is such that $\psi_{\mu}(\alpha_1) = 1$ and $\psi_{\mu}(\alpha_2) = \mu(\alpha_2) = \zeta_p$. Now, let Ψ_{μ} denote an irreducible rational matrix representation of degree $p - 1$ of H that affords the character $\Omega(\psi_{\mu})$. The explicit form of Ψ_{μ} is given by

$$\Psi_{\mu}(\alpha_1) = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \\ 0 & 0 & \cdots & 1 & 0 \\ 0 & 0 & \cdots & \cdots & 1 \end{pmatrix} = I, \quad \text{and} \quad \Psi_{\mu}(\alpha_2) = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \\ 0 & 0 & \cdots & 0 & 1 \\ -1 & -1 & \cdots & \cdots & -1 \end{pmatrix}$$

(see Lemma 11). Set $\Psi_{\mu}(\alpha_2) = P$, where P denotes a matrix of order $(p - 1)$, and let O denote the zero matrix of the same order. Then an irreducible rational matrix

representation Ψ^G of degree $p^2 - p$ of G affording the character $\Omega(\chi_\mu)$ is given by

$$\Psi_\mu^G(\alpha) = \begin{pmatrix} O & O & O & \cdots & P \\ I & O & O & \cdots & O \\ O & I & O & \cdots & O \\ \vdots & \vdots & \vdots & \ddots & \\ O & O & \cdots & I & 0 \end{pmatrix}, \quad \text{and} \quad \Psi_\mu^G(\alpha_1) = \begin{pmatrix} I & O & O & \cdots & O \\ O & P & O & \cdots & O \\ O & O & P^2 & \cdots & O \\ \vdots & \vdots & \vdots & \ddots & \\ O & O & \cdots & \cdots & P^{p-1} \end{pmatrix}.$$

Here, Ψ^G is a rational matrix representation of G whose kernel does not contain G' . Now, use Lemma 11 to compute all irreducible rational matrix representations of G whose kernel contains G' . This gives the complete description of all irreducible rational matrix representations of G .

REMARK 48. If G is a nonabelian group of order 8, then either $G \cong Q_8$ or $G \cong D_8$. A special required pair to obtain an irreducible rational matrix representation of G whose kernel does not contain G' is determined in Section 4.1.

6. p -Group of order p^4

In this section, we provide a comprehensive description of all inequivalent irreducible rational matrix representations and the Wedderburn decompositions of the rational group rings for all nonabelian groups of order p^4 . It is easy to observe that if G is a nonabelian group of order p^4 , then $|Z(G)| = p$ or p^2 , and $\text{cd}(G) = \{1, p\}$. Moreover, if $|Z(G)| = p$, then $G/Z(G)$ is nonabelian. As we know, a group G is VZ-group if and only if $\text{cd}(G) = \{1, |G/Z(G)|^{1/2}\}$. Hence, a p -group of order p^4 of nilpotency class 2 is a VZ p -group. In the subsequent subsections, we separately discuss the cases of groups of order p^4 of nilpotency class 2 and nilpotency class 3.

6.1. p -Groups of order p^4 of nilpotency class 2. Let G be a p -group of order p^4 of nilpotency class 2. Then $|G'| = p$, $|Z(G)| = p^2$ and $\text{cd}(G) = \{1, p\}$. Furthermore, $|\text{lin}(G)| = p^3$ and $|\text{nl}(G)| = p^2 - p$. Note that G is a VZ p -group. Then by (1), a nonlinear irreducible complex character is of the form χ_μ and is given by

$$\chi_\mu(g) = \begin{cases} p\mu(g) & \text{if } g \in Z(G), \\ 0 & \text{otherwise,} \end{cases} \tag{3}$$

where $\mu \in \text{Irr}(Z(G)|G')$.

For an odd prime p , we rely on James' classification of p -groups of order p^4 (see [13]). There exist two distinct isoclinic families of nonabelian groups of order p^4 , denoted as Φ_2 and Φ_3 (see [13, Section 4.4]). Prajapati *et al.* [25, Proposition 4.2] have proved that all the groups belonging to isoclinic family Φ_2 are VZ-groups. Note that all the groups belonging to isoclinic family Φ_3 are non-VZ-groups. Theorem 49 provides a description of all inequivalent irreducible rational matrix representations for all VZ p -groups of order p^4 , where p is an odd prime.

TABLE 1. Special required pair (H, ψ_μ) to obtain an irreducible rational matrix representation of $G \in \Phi_2$ that affords the character $\Omega(\chi_\mu)$, where $\chi_\mu \in \text{nl}(G)$ (defined in (3)).

Group G	$Z(G)$	G'	H	$\psi_\mu \in \text{Irr}(H)$ and $\mu \in \text{Irr}(Z(G) G')$
$\Phi_2(211)a = \langle \alpha, \alpha_1, \alpha_2, \alpha_3 : [\alpha_1, \alpha] = \alpha^p = \alpha_2, \alpha_1^p = \alpha_2^p = \alpha_3^p = 1 \rangle$	$\langle \alpha^p, \alpha_3 \rangle$	$\langle \alpha^p \rangle$	$\langle \alpha^p, \alpha_1, \alpha_3 \rangle$	$\psi_\mu(h) = \begin{cases} \mu(\alpha^p) & \text{if } h = \alpha^p, \\ 1 & \text{if } h = \alpha_1, \\ \mu(\alpha_3) & \text{if } h = \alpha_3 \end{cases}$
$\Phi_2(1^4) = \langle \alpha, \alpha_1, \alpha_2, \alpha_3 : [\alpha_1, \alpha] = \alpha_2, \alpha^p = \alpha_1^p = \alpha_2^p = \alpha_3^p = 1 \rangle$	$\langle \alpha_2, \alpha_3 \rangle$	$\langle \alpha_2 \rangle$	$\langle \alpha, \alpha_2, \alpha_3 \rangle$	$\psi_\mu(h) = \begin{cases} 1 & \text{if } h = \alpha, \\ \mu(\alpha_2) & \text{if } h = \alpha_2, \\ \mu(\alpha_3) & \text{if } h = \alpha_3, \end{cases}$
$\Phi_2(31) = \langle \alpha, \alpha_1, \alpha_2 : [\alpha_1, \alpha] = \alpha^{p^2} = \alpha_2, \alpha_1^p = \alpha_2^p = 1 \rangle$	$\langle \alpha^p \rangle$	$\langle \alpha^{p^2} \rangle$	$\langle \alpha^p, \alpha_1 \rangle$	$\psi_\mu(h) = \begin{cases} \mu(\alpha^p) & \text{if } h = \alpha^p, \\ 1 & \text{if } h = \alpha_1, \end{cases}$
$\Phi_2(22) = \langle \alpha, \alpha_1, \alpha_2 : [\alpha_1, \alpha] = \alpha^p = \alpha_2, \alpha_1^{p^2} = \alpha_2^p = 1 \rangle$	$\langle \alpha^p, \alpha_1^p \rangle$	$\langle \alpha^p \rangle$	$\langle \alpha^{-i}\alpha_1, \alpha^p \rangle$ ($0 \leq i \leq p - 1$)	$\psi_\mu(h) = \begin{cases} 1 & \text{if } h = \alpha^{-i}\alpha_1, \\ \mu(\alpha^p) & \text{if } h = \alpha^p, \end{cases}$
$\Phi_2(211)b = \langle \alpha, \alpha_1, \alpha_2, \gamma : [\alpha_1, \alpha] = \gamma^p = \alpha_2, \alpha^p = \alpha_1^p = \alpha_2^p = 1 \rangle$	$\langle \gamma \rangle$	$\langle \gamma^p \rangle$	$\langle \alpha_1, \gamma \rangle$	$\psi_\mu(h) = \begin{cases} 1 & \text{if } h = \alpha_1, \\ \mu(\gamma) & \text{if } h = \gamma, \end{cases}$
$\Phi_2(211)c = \langle \alpha, \alpha_1, \alpha_2 : [\alpha_1, \alpha] = \alpha_2, \alpha^{p^2} = \alpha_1^p = \alpha_2^p = 1 \rangle$	$\langle \alpha^p, \alpha_2 \rangle$	$\langle \alpha_2 \rangle$	$\langle \alpha^p, \alpha_1, \alpha_2 \rangle$	$\psi_\mu(h) = \begin{cases} \mu(\alpha^p) & \text{if } h = \alpha^p, \\ 1 & \text{if } h = \alpha_1, \\ \mu(\alpha_2) & \text{if } h = \alpha_2. \end{cases}$

THEOREM 49. Let G be a nonabelian p -group (odd prime p) of order p^4 in the isoclinic family Φ_2 . Then Table 1 determines all inequivalent irreducible rational matrix representations of G whose kernels do not contain G' .

PROOF. Let $G \in \Phi_2$ and let $\chi_\mu \in \text{nl}(G)$ (as defined in (3)). Suppose (H, ψ_μ) is a special required pair to obtain an irreducible rational matrix representation of G which affords the character $\Omega(\chi_\mu)$. By Proposition 21, it follows that $Z(G) \subset H$ and $\psi_\mu \downarrow_{Z(G)} = \mu$. Further, from Corollary 27, H is abelian. Since (H, ψ_μ) is a special required pair, $\mathbb{Q}(\psi_\mu) = \mathbb{Q}(\chi_\mu)$ and hence by Lemma 24, we must choose $\psi_\mu \in \text{lin}(H)$ such that $|\ker(\psi_\mu)| = |G/Z(G)|^{1/2}|\ker(\mu)| = p|\ker(\mu)|$. Consider $G = \Phi_2(22)$. Now, for each $0 \leq i \leq (p - 1)$, take $H = \langle \alpha^{-i}\alpha_1, \alpha^p \rangle$ and define $\mu \in \text{Irr}(Z(G)|G')$ as follows:

$$\mu(z) = \begin{cases} \zeta_p & \text{if } z = \alpha^p, \\ \zeta_p^i & \text{if } z = \alpha_1^p, \end{cases}$$

where $z \in Z(G)$. Observe that $(\alpha^{-i}\alpha_1)^p = \alpha^{-ip}\alpha_1^p$. Then, $\psi_\mu \in \text{lin}(H)$ (given in Table 1) satisfies $\psi_\mu(\alpha_1^p) = (\psi_\mu(\alpha^{-i}\alpha_1))^p(\psi_\mu(\alpha^p))^i = (\psi_\mu(\alpha^p))^i = \mu(\alpha_1^p)$. It is easy to check that a pair (H, ψ_μ) satisfies the criteria of a special required pair.

It is routine to check that all the pairs (H, ψ_μ) for the rest of the groups mentioned in Table 1 also satisfy the criteria to being special required pairs. This shows that Table 1 presents all of the special required pairs (H, ψ_μ) to find all inequivalent irreducible rational matrix representations of G whose kernels do not contain G' , where $G \in \Phi_2$. This completes the proof of Theorem 49. \square

Proposition 50 provides the counting of rational irreducible representations of different degrees for all groups of order p^4 in Φ_2 .

PROPOSITION 50. *Let G be a nonabelian group of order p^4 (odd prime p) in Φ_2 .*

- (1) *If $G = \Phi_2(211)a$ or $G = \Phi_2(1^4)$, then $|\text{Irr}_{\mathbb{Q}}^{(1)}(G)| = 1$, $|\text{Irr}_{\mathbb{Q}}^{(\phi(p))}(G)| = p^2 + p + 1$ and $|\text{Irr}_{\mathbb{Q}}^{(\phi(p^2))}(G)| = p$.*
- (2) *If $G = \Phi_2(31)$, then $|\text{Irr}_{\mathbb{Q}}^{(1)}(G)| = 1$, $|\text{Irr}_{\mathbb{Q}}^{(\phi(p))}(G)| = p + 1$, $|\text{Irr}_{\mathbb{Q}}^{(\phi(p^2))}(G)| = p$ and $|\text{Irr}_{\mathbb{Q}}^{(\phi(p^3))}(G)| = 1$.*
- (3) *If $G = \Phi_2(22)$, then $|\text{Irr}_{\mathbb{Q}}^{(1)}(G)| = 1$, $|\text{Irr}_{\mathbb{Q}}^{(\phi(p))}(G)| = p + 1$ and $|\text{Irr}_{\mathbb{Q}}^{(\phi(p^2))}(G)| = 2p$.*
- (4) *If $G = \Phi_2(211)b$, then $|\text{Irr}_{\mathbb{Q}}^{(1)}(G)| = 1$, $|\text{Irr}_{\mathbb{Q}}^{(\phi(p))}(G)| = p^2 + p + 1$, $|\text{Irr}_{\mathbb{Q}}^{(\phi(p^3))}(G)| = 1$.*
- (5) *If $G = \Phi_2(211)c$, then $|\text{Irr}_{\mathbb{Q}}^{(1)}(G)| = 1$, $|\text{Irr}_{\mathbb{Q}}^{(\phi(p))}(G)| = p + 1$ and $|\text{Irr}_{\mathbb{Q}}^{(\phi(p^2))}(G)| = p + 1$.*

PROOF. Suppose $\chi \in \text{Irr}(G)$ and $E(\chi)$ denotes the Galois conjugacy class of χ over \mathbb{Q} . Then the degree of the rational representation affording the character $\Omega(\chi)$ is $|E(\chi)|\chi(1)$.

- (1) For $G = \Phi_2(211)a$, we have $Z(G) = \langle \alpha^p, \alpha_3 \rangle \cong C_p \times C_p$, $G' = \langle \alpha^p \rangle \cong C_p$ and $G/G' = \langle \alpha G', \alpha_1 G', \alpha_2 G' \rangle \cong C_p \times C_p \times C_p$. Observe that in $\text{Irr}^{(1)}(G)$, there is a single Galois conjugacy class over \mathbb{Q} with size 1 and $(p^3 - 1)/\phi(p) = p^2 + p + 1$ distinct Galois conjugacy classes over \mathbb{Q} with size $\phi(p)$. If $\mu \in \text{Irr}(Z(G)|G')$, then $[\mathbb{Q}(\mu) : \mathbb{Q}] = \phi(p)$. Hence, there are $(p^2 - p)/\phi(p) = p$ distinct Galois conjugacy classes over \mathbb{Q} with size $\phi(p)$ in $\text{Irr}^{(p)}(G)$. Similar statements hold for $G = \Phi_2(1^4)$. This proves part (1).
- (2) For $G = \Phi_2(31)$, we have $Z(G) = \langle \alpha^p \rangle \cong C_{p^2}$, $G' = \langle \alpha^{p^2} \rangle \cong C_p$ and $G/G' = \langle \alpha G', \alpha_1 G' \rangle \cong C_{p^2} \times C_p$. In $\text{Irr}^{(1)}(G)$, there is one Galois conjugacy class over \mathbb{Q} with size 1, $(1 \times \phi(p) + \phi(p) \times 1 + \phi(p) \times \phi(p))/\phi(p) = p + 1$ distinct Galois conjugacy classes over \mathbb{Q} with size $\phi(p)$, and $(\phi(p^2) \times p)/\phi(p^2) = p$ distinct Galois conjugacy classes over \mathbb{Q} with size $\phi(p^2)$. If $\mu \in \text{Irr}(Z(G)|G')$, then $[\mathbb{Q}(\mu) : \mathbb{Q}] = \phi(p^2)$. Consequently, there is only one $((p^2 - p)/\phi(p^2) = 1)$ Galois conjugacy class over \mathbb{Q} with size $\phi(p^2)$ in $\text{Irr}^{(p)}(G)$. This completes the proof of part (2).

By using similar arguments, we get the proofs of the remaining parts of Proposition 50. \square

TABLE 2. Special required pair (H, ψ_μ) to obtain an irreducible rational matrix representation of a VZ 2-group G of order 16 that affords the character $\Omega(\chi_\mu)$, where $\chi_\mu \in \text{nl}(G)$ (defined in (3)).

Group G	$Z(G)$	G'	H	$\psi_\mu \in \text{Irr}(H)$ and $\mu \in \text{Irr}(Z(G) G')$
$G_1 = \langle x, y, z : x^4 = y^2 = z^2 = 1, [x, y] = [x, z] = 1, [y, z] = x^2 \rangle$	$\langle x \rangle$	$\langle x^2 \rangle$	$\langle x, y \rangle$	$\psi_\mu(h) = \begin{cases} \mu(x) & \text{if } h = x, \\ 1 & \text{if } h = y, \end{cases}$
$G_2 = \langle x, y : x^8 = y^2 = 1, [x, y] = x^4 \rangle$	$\langle x^2 \rangle$	$\langle x^4 \rangle$	$\langle x^2, y \rangle$	$\psi_\mu(h) = \begin{cases} \mu(x^2) & \text{if } h = x^2, \\ \mu(\alpha_3) & \text{if } h = y, \end{cases}$
$G_3 = \langle x, y, z : x^4 = y^2 = z^2 = 1, [x, z] = [y, z] = 1, [x, y] = x^2 \rangle$	$\langle x^2, z \rangle$	$\langle x^2 \rangle$	$\langle x^2, y, z \rangle$	$\psi_\mu(h) = \begin{cases} \mu(x^2) & \text{if } h = x^2, \\ 1 & \text{if } h = y, \\ \mu(z) & \text{if } h = z, \end{cases}$
$G_4 = \langle x, y, z : x^4 = y^2 = z^2 = 1, [x, z] = [y, z] = 1, [x, y] = z \rangle$	$\langle x^2, z \rangle$	$\langle z \rangle$	$\langle x^2, y, z \rangle$	$\psi_\mu(h) = \begin{cases} \mu(x^2) & \text{if } h = x^2, \\ 1 & \text{if } h = y, \\ \mu(z) & \text{if } h = z, \end{cases}$
$G_5 = \langle x, y : x^4 = y^4 = 1, [x, y] = x^2 \rangle$	$\langle x^2, y^2 \rangle$	$\langle x^2 \rangle$	$\langle x^2, y \rangle$	$\psi_\mu(h) = \begin{cases} \mu(x^2) & \text{if } h = x^2, \\ (\mu(y^2))^{1/2} & \text{if } h = y, \end{cases}$
$G_6 = \langle x, y, z : x^4 = y^4 = z^2 = 1, [x, z] = [y, z] = 1, [x, y] = x^2, x^2 = y^2 \rangle$	$\langle x^2, z \rangle$	$\langle x^2 \rangle$	$\langle x, z \rangle$	$\psi_\mu(h) = \begin{cases} (\mu(x^2))^{1/2} & \text{if } h = x, \\ \mu(z) & \text{if } h = z, \end{cases}$

REMARK 51. Let G and H be isoclinic groups with the same order. According to Lemma 10, we have $|\text{Irr}^{(k)}(G)| = |\text{Irr}^{(k)}(H)|$. However, it is important to note that $|\text{Irr}_{\mathbb{Q}}^{(k)}(G)|$ may not be equal to $|\text{Irr}^{(k)}(H)|$ (see Proposition 50).

Now, let G be a nonabelian 2-group of order 16 of nilpotency class 2. Then $|\text{nl}(G)| = 2$. We take presentations of 2-groups from Burnside’s book [6]. Theorem 52 provides a description of all inequivalent irreducible rational matrix representations of G whose kernels do not contain G' .

THEOREM 52. Let G be a nonabelian 2-group of order 16 of nilpotency class 2. Then Table 2 determines all inequivalent irreducible rational matrix representations of G whose kernels do not contain G' .

PROOF. Let G be a nonabelian group of order 16 of nilpotency class 2. Observe that G is a VZ 2-group. Let $\chi_\mu \in \text{nl}(G)$ as defined in (3). Suppose (H, ψ_μ) is a special required pair to obtain an irreducible rational matrix representation of G that affords the character $\Omega(\chi_\mu)$. By Proposition 21, it follows that $Z(G) \subset H$ and $\psi_\mu \downarrow_{Z(G)} = \mu$. Further, from Corollary 27, H is abelian. In view of Remark 23, we have the following: if $m_{\mathbb{Q}}(\chi_\mu) = 1$, then $\mathbb{Q}(\psi_\mu) = \mathbb{Q}(\chi_\mu)$ and if $m_{\mathbb{Q}}(\chi_\mu) = 2$, then $[\mathbb{Q}(\psi_\mu) : \mathbb{Q}(\chi_\mu)] = 2$. Therefore, from Lemma 24, we must choose $\psi_\mu \in \text{lin}(H)$ such that

$|\ker(\psi_\mu)| = 2|\ker(\mu)|$ whenever $m_{\mathbb{Q}}(\chi_\mu) = 1$, and $\psi_\mu \in \text{lin}(H)$ such that $|\ker(\psi_\mu)| = |\ker(\mu)|$ whenever $m_{\mathbb{Q}}(\chi_\mu) = 2$. Note that $m_{\mathbb{Q}}(\chi_\mu) = 2$ for $\chi_\mu \in \text{nl}(G_5)$, where $\mu \in \text{Irr}(Z(G_5)|G'_5)$ is given by $\mu(x^2) = -1, \mu(y^2) = -1$, and $m_{\mathbb{Q}}(\chi_\mu) = 2$ for all $\chi_\mu \in \text{nl}(G_6)$, where $\mu \in \text{Irr}(Z(G_6)|G'_6)$. It is routine to check that the pairs (H, ψ_μ) mentioned in Table 2 are special required pairs. This shows that Table 2 presents special required pairs (H, ψ_μ) to find irreducible rational matrix representations of G whose kernels do not contain G' , where $G \in \Phi_2$. This completes the proof of Theorem 52. \square

6.2. p -Groups of order p^4 of nilpotency class 3. Let G be a p -group of order p^4 of nilpotency class 3. Then $|Z(G)| = p, |G'| = p^2$ and $Z(G) \subset G'$. Further, $|\text{lin}(G)| = p^2, |\text{nl}(G)| = p^2 - 1$ and $\text{cd}(G) = \{1, p\}$.

We begin by presenting a few results that enable us to determine all the inequivalent irreducible rational matrix representations of G .

LEMMA 53. *Let G be a p -group (odd prime p) and let $1 \neq \chi \in \text{Irr}(G)$. Then, $\mathbb{Q}(\chi) \neq \mathbb{Q}$.*

PROOF. The proof is obvious. \square

LEMMA 54. *Let G be a nonabelian group with nilpotency class ≥ 3 . Suppose that there exists a maximal normal subgroup H of G such that both H and G/H are abelian. Then H is unique.*

PROOF. Since G/H is abelian, $G' \subseteq H$. As H is abelian, we get $C_G(G') \supseteq H$, where $C_G(G')$ denotes the centralizer subgroup of G' . Since nilpotency class of G is ≥ 3 , $G' \not\subseteq Z(G)$. This implies that $C_G(G') \neq G$. Thus, we conclude that $C_G(G') = H$, which implies the uniqueness of H . \square

COROLLARY 55. *If G is a nonabelian group of order p^4 of nilpotency class 3, then G has a unique abelian subgroup of index p , namely $C_G(G')$.*

PROOF. Let G be a nonabelian p -group of order p^4 of nilpotency class 3. Then $Z(G) \subset G'$ and there exists an abelian subgroup H of G with index p . Therefore, the proof follows from Lemma 54. \square

For an odd prime p , we follow James' classification of p -groups of order p^4 (see [13]). All the relevant groups in this subsection belong to Φ_3 (refer to [13, Section 4.4]).

REMARK 56. Let G be a nonabelian p -group ($p \geq 5$) of order p^4 in Φ_3 (see [13, Section 4.4]). Then the description of unique abelian subgroup H of G of index p is as follows.

- If $G = \Phi_3(1^4)$, then $H = \langle \alpha_1, \alpha_2, \alpha_3 \rangle \cong C_p \times C_p \times C_p$.
- If $G = \Phi_3(211)a$, then $H = \langle \alpha^p, \alpha_1, \alpha_2 \rangle \cong C_p \times C_p \times C_p$.
- If $G = \Phi_3(211)b_r$ ($r = 1, \nu$), then $H = \langle \alpha_1, \alpha_2 \rangle \cong C_{p^2} \times C_p$.

LEMMA 57. *Let G be a nonabelian p -group of order p^4 of nilpotency class 3. Let H be the unique abelian subgroup of G of index p . If $\psi \in \text{Irr}(H|G')$, then $\psi^G \in \text{nl}(G)$. Further, for $\chi \in \text{nl}(G)$, there exists some $\psi \in \text{Irr}(H|G')$ such that $\chi = \psi^G$.*

PROOF. Consider $\psi \in \text{Irr}(H|G')$. In contrast, suppose that $\psi^G \notin \text{nl}(G)$. This implies that ψ^G is a sum of some linear characters of G . This implies that $G' \subseteq \ker(\psi^G) \subseteq \ker(\psi)$, which is a contradiction.

Now, let $\psi \in \text{Irr}(H|G')$. Then $\psi^G \in \text{nl}(G)$, and hence the inertia group $I_G(\psi)$ of ψ in G is equal to H [12, Problem 6.1]. Furthermore, $\psi^G \downarrow_H = \sum_{i=1}^p \psi_i$, where the ψ_i terms are conjugates of ψ in G and $p = |G/I_G(\psi)|$. Hence, there are p conjugates of ψ and observe that $\psi^G = \psi_i^G \in \text{nl}(G)$ for each i . Thus, $|\text{nl}(G)| = |\text{Irr}(H|G')|/p = p^2 - 1$. This completes the proof of Lemma 57. \square

Theorem 58 provides the necessary information to determine all inequivalent irreducible rational matrix representations for all nonabelian p -groups of order p^4 of nilpotency class 3, where p is an odd prime.

THEOREM 58. *Let G be a nonabelian p -group (odd prime p) of order p^4 belonging to Φ_3 . Suppose H is the unique abelian subgroup of G with index p and $\chi \in \text{nl}(G)$ is such that $\chi = \psi^G$ for some $\psi \in \text{Irr}(H|G')$. Then (H, ψ) is a required pair to determine an irreducible rational matrix representation of G that affords the character $\Omega(\chi)$.*

PROOF. Suppose $p \geq 5$. If $G = \Phi_3(1^4)$ or $\Phi_3(211)a$, then $H \cong C_p \times C_p \times C_p$ (see Remark 56). Suppose $\chi \in \text{nl}(G)$ is such that $\chi = \psi^G$ for some $\psi \in \text{Irr}(H|G')$. Then, $\mathbb{Q}(\psi) = \mathbb{Q}(\zeta_p)$. Observe that $\mathbb{Q}(\chi) = \mathbb{Q}(\psi^G) \subseteq \mathbb{Q}(\psi)$. From Lemma 53, we get $\mathbb{Q}(\chi) = \mathbb{Q}(\psi) = \mathbb{Q}(\zeta_p)$. Next, if $G = \Phi_3(211)b_r$ ($r = 1, \nu$), then $H = \langle \alpha_1, \alpha_2 \rangle \cong C_{p^2} \times C_p$ (see Remark 56). Again, suppose that $\chi \in \text{nl}(G)$ is such that $\chi = \psi^G$ for some $\psi \in \text{Irr}(H|G')$. Then, $\mathbb{Q}(\psi) = \mathbb{Q}(\zeta_p)$ or $\mathbb{Q}(\zeta_{p^2})$. If $\mathbb{Q}(\psi) = \mathbb{Q}(\zeta_p)$, then from Lemma 53, $\mathbb{Q}(\psi) = \mathbb{Q}(\chi) = \mathbb{Q}(\zeta_p)$. Now, suppose $\mathbb{Q}(\psi) = \mathbb{Q}(\zeta_{p^2})$. This implies that $\psi(\alpha_1) = \zeta_{p^2}$. Assume that $G = \bigcup \alpha^i H$ ($0 \leq i \leq p - 1$). Then we have the following:

$$\begin{aligned} \psi^G(\alpha_1) &= \sum_{i=0}^{p-1} \psi^\circ(\alpha^{-i} \alpha_1 \alpha^i), \quad \text{where } \psi^\circ(g) = \begin{cases} \psi(g) & \text{if } g \in H, \\ 0 & \text{if } g \notin H, \end{cases} \\ &= \psi(\alpha_1) + \psi(\alpha_1 \alpha_2) + \psi(\alpha_1^{1+p} \alpha_2^2) + \psi(\alpha_1^{1+3p} \alpha_2^3) + \dots + \psi(\alpha_1^{1+(p-1)(p-2)/2p} \alpha_2^{p-1}) \\ &= \psi(\alpha_1)[1 + \psi(\alpha_2) + \psi(\alpha_1^p \alpha_2^2) + \psi(\alpha_1^{3p} \alpha_2^3) + \dots + \psi(\alpha_1^{(p-1)(p-2)/2p} \alpha_2^{p-1})] \\ &= \theta \zeta_{p^2}, \text{ for some } 0 \neq \theta \in \mathbb{Q}(\zeta_p). \end{aligned}$$

Therefore, $\mathbb{Q}(\psi) = \mathbb{Q}(\psi^G) = \mathbb{Q}(\zeta_{p^2})$. Hence, (H, ψ) is a required pair. Now, let $p = 3$. If

$$\begin{aligned} G = \Phi_3(211)b_1 &= \langle \alpha, \alpha_1, \alpha_2, \alpha_3 : [\alpha_1, \alpha] = \alpha_2, [\alpha_2, \alpha] = \alpha_1^3 \alpha_3 = \alpha_3, \\ &\alpha^3 = \alpha_2^3 = \alpha_3^3 = 1 \rangle, \end{aligned}$$

then $H = \langle \alpha_1, \alpha_2, \alpha_3 \rangle \cong C_3 \times C_3 \times C_3$ is the unique normal abelian subgroup of G of index 3 (see [13, Section 4.4]). Suppose $\chi \in \text{nl}(G)$ is such that $\chi = \psi^G$ for some $\psi \in \text{Irr}(H|G')$. From Lemma 53, we get $\mathbb{Q}(\chi) = \mathbb{Q}(\psi) = \mathbb{Q}(\zeta_3)$. Next, if G is one of the following groups:

$$\begin{aligned} \Phi_3(211)a &= \langle \alpha, \alpha_1, \alpha_2, \alpha_3 : [\alpha_1, \alpha] = \alpha_2, [\alpha_2, \alpha] = \alpha^3 = \alpha_3, \alpha_1^3 \alpha_3 = \alpha_2^3 = \alpha_3^3 = 1 \rangle \\ \Phi_3(1^4) &= \langle \alpha, \alpha_1, \alpha_2, \alpha_3 : [\alpha_1, \alpha] = \alpha_2, [\alpha_2, \alpha] = \alpha_3, \alpha^3 = \alpha_1^3 \alpha_3 = \alpha_2^3 = \alpha_3^3 = 1 \rangle \\ \Phi_3(211)b_2 &= \langle \alpha, \alpha_1, \alpha_2, \alpha_3 : [\alpha_1, \alpha] = \alpha_2, [\alpha_2, \alpha]^2 = \alpha_1^3 \alpha_3 = \alpha_2^3, \alpha^3 = \alpha_2^3 = \alpha_3^3 = 1 \rangle, \end{aligned}$$

then $H = \langle \alpha_1, \alpha_2 \rangle \cong C_9 \times C_3$ is the unique normal subgroup of G of index 3 (see [13, Section 4.4]). Again, suppose $\chi \in \text{nl}(G)$ is such that $\chi = \psi^G$ for some $\psi \in \text{Irr}(H|G')$. By a similar argument as in the case of $p \geq 5$, we can establish that $\mathbb{Q}(\chi) = \mathbb{Q}(\psi)$. Hence, (H, ψ) is a required pair. This completes the proof of Theorem 58. \square

In Proposition 59, the counting of rational irreducible representations of all p -groups (odd prime p) of order p^4 in Φ_3 is described.

PROPOSITION 59. *Let G be a nonabelian p -group ($p \geq 5$) of order p^4 in Φ_3 . Then we have the following.*

- (1) *If $G = \Phi_3(211)a$ or $\Phi_3(1^4)$, then $|\text{Irr}_{\mathbb{Q}}^{(1)}(G)| = 1$, $|\text{Irr}_{\mathbb{Q}}^{(\phi(p))}(G)| = p + 1$ and $|\text{Irr}_{\mathbb{Q}}^{(\phi(p^2))}(G)| = p + 1$.*
- (2) *If $G = \Phi_3(211)b_r$ ($r = 1, \nu$), then $|\text{Irr}_{\mathbb{Q}}^{(1)}(G)| = 1$, $|\text{Irr}_{\mathbb{Q}}^{(\phi(p))}(G)| = p + 1$, $|\text{Irr}_{\mathbb{Q}}^{(\phi(p^2))}(G)| = 1$ and $|\text{Irr}_{\mathbb{Q}}^{(\phi(p^3))}(G)| = 1$.*

PROOF. Suppose $\chi \in \text{Irr}(G)$ and $E(\chi)$ denotes the Galois conjugacy class of χ over \mathbb{Q} . Then the degree of the rational representation affording the character $\Omega(\chi)$ is $|E(\chi)|\chi(1)$.

- (1) Let

$$\begin{aligned} G = \Phi_3(211)a &= \langle \alpha, \alpha_1, \alpha_2, \alpha_3 : [\alpha_1, \alpha] = \alpha_2, [\alpha_2, \alpha] = \alpha^p = \alpha_3, \\ &\alpha_1^p = \alpha_2^p = \alpha_3^p = 1 \rangle. \end{aligned}$$

Then $Z(G) = \langle \alpha^p \rangle \cong C_p$, $G' = \langle \alpha^p, \alpha_2 \rangle \cong C_p \times C_p$ and $G/G' = \langle \alpha G', \alpha_1 G' \rangle \cong C_p \times C_p$. Thus, there are one Galois conjugacy class over \mathbb{Q} of size 1, and $(p^2 - 1)/\phi(p) = p + 1$ distinct Galois conjugacy classes over \mathbb{Q} of size $\phi(p)$ in $\text{Irr}^{(1)}(G)$. Further, $H = \langle \alpha^p, \alpha_1, \alpha_2 \rangle \cong C_p \times C_p \times C_p$ is the abelian subgroup of $\Phi_3(211)a$ of index p . Suppose $\chi \in \text{Irr}^{(p)}(G)$. Then from Theorem 58, $\chi = \psi^G$ for some $\psi \in \text{Irr}(H|G')$ and $\mathbb{Q}(\chi) = \mathbb{Q}(\psi)$. Moreover, from Lemma 53, $\mathbb{Q}(\chi) = \mathbb{Q}(\psi) = \mathbb{Q}(\zeta_p)$. Thus, there are $(p^2 - 1)/\phi(p) = p + 1$ distinct Galois conjugacy classes over \mathbb{Q} of size $\phi(p)$ in $\text{Irr}^{(p)}(G)$. We get similar results for $G = \Phi_3(1^4)$. This completes the proof of part (1) of Proposition 59.

- (2) Let

$$\begin{aligned} G = \Phi_3(211)b_1 &= \langle \alpha, \alpha_1, \alpha_2, \alpha_3 : [\alpha_1, \alpha] = \alpha_2, [\alpha_2, \alpha] = \alpha_1^p = \alpha_3, \\ &\alpha^p = \alpha_2^p = \alpha_3^p = 1 \rangle. \end{aligned}$$

Then $Z(G) = \langle \alpha_1^p \rangle = C_p$, $G' = \langle \alpha_1^p, \alpha_2 \rangle = C_p \times C_p$ and $G/G' = \langle \alpha G', \alpha_1 G' \rangle \cong C_p \times C_p$. Thus, there are one Galois conjugacy class over \mathbb{Q} of size 1, and

$(p^2 - 1)/\phi(p) = p + 1$ distinct Galois conjugacy classes over \mathbb{Q} of size $\phi(p)$ in $\text{Irr}^{(1)}(G)$. Further, $H = \langle \alpha_1, \alpha_2 \rangle \cong C_{p^2} \times C_p$ is the abelian subgroup of $\Phi_3(211)b_1$ of index p . Suppose $\chi \in \text{Irr}^{(p)}(G)$. Then from Theorem 58, $\chi = \psi^G$ for some $\psi \in \text{Irr}(H|G')$ and $\mathbb{Q}(\chi) = \mathbb{Q}(\psi)$. Observe that $|\{\psi \in \text{Irr}(H|G') : \mathbb{Q}(\psi) = \mathbb{Q}(\zeta_p)\}| = p^2 - p$ and $|\{\psi \in \text{Irr}(H|G') : \mathbb{Q}(\psi) = \mathbb{Q}(\zeta_{p^2})\}| = p^3 - p^2$. Since there are p conjugates of each $\psi \in \text{Irr}(H|G')$, the numbers of complex irreducible characters of degree p ($\chi \in \text{Irr}^{(p)}$) such that $\mathbb{Q}(\chi) = \mathbb{Q}(\zeta_p)$ and $\mathbb{Q}(\chi) = \mathbb{Q}(\zeta_{p^2})$ are $(p^2 - p)/p = p - 1$ and $(p^3 - p^2)/p = p^2 - p$, respectively. Thus, there are $(p - 1)/\phi(p) = 1$ Galois conjugacy classes over \mathbb{Q} of size $\phi(p)$, and $(p^2 - p)/\phi(p^2) = 1$ Galois conjugacy classes over \mathbb{Q} of size $\phi(p^2)$ in $\text{Irr}^{(p)}(G)$. We get similar results for $G = \Phi_3(211)b_v$. This completes the proof of part (2) of Proposition 59. \square

Now, we prove Theorem 3, which provides the Wedderburn decomposition of $\mathbb{Q}G$, where G is a p -group (odd prime p) of order p^4 of nilpotency class 3.

PROOF OF THEOREM 3. Observe that $G \in \Phi_3$. Suppose $\chi \in \text{nl}(G)$ and H is the unique abelian subgroup of G . Then from Lemma 57, there exists $\psi \in \text{Irr}(H|G')$ such that $\chi = \psi^G$, and if $\psi \in \text{Irr}(H|G')$, then $\psi^G \in \text{nl}(G)$. Now by Theorem 58, $\mathbb{Q}(\chi) = \mathbb{Q}(\psi)$. Observe that $\chi^\sigma = (\psi^\sigma)^G$, where $\sigma \in \text{Gal}(\mathbb{Q}(\chi)/\mathbb{Q})$ and there are exactly p distinct conjugates $\psi \in \text{Irr}(H|G')$ such that $\chi = \psi^G$ (see the proof of Lemma 57). Let X and Y be the representative sets of distinct Galois conjugacy classes of $\text{Irr}(G)$ and $\text{Irr}(H)$, respectively. Let d be a divisor of $\text{exp}(H)$ such that $\mathbb{Q}(\chi) = \mathbb{Q}(\psi) = \mathbb{Q}(\zeta_d)$. Set $m = \text{exp}(H)$ and $m' = \text{exp}(H/G')$. Then we have two cases.

Case 1 ($d \mid m$ but $d \nmid m'$). In this case,

$$\begin{aligned} |\{\chi \in X : \chi(1) = p, \mathbb{Q}(\chi) = \mathbb{Q}(\zeta_d)\}| &= \frac{1}{p} |\{\psi \in Y : \psi \in \text{Irr}(H|G'), \mathbb{Q}(\psi) = \mathbb{Q}(\zeta_d)\}| \\ &= \frac{a_d}{p}, \end{aligned}$$

where a_d denotes the number of cyclic subgroups of order d of H (see Lemma 31).

Case 2 ($d \mid m$ and $d \mid m'$). In this case,

$$\begin{aligned} |\{\chi \in X : \chi(1) = p, \mathbb{Q}(\chi) = \mathbb{Q}(\zeta_d)\}| &= \frac{1}{p} |\{\psi \in Y : \psi \in \text{Irr}(H|G'), \mathbb{Q}(\psi) = \mathbb{Q}(\zeta_d)\}| \\ &= \frac{a_d - a'_d}{p}, \end{aligned}$$

where a_d and a'_d denote the numbers of cyclic subgroups of order d of H and H/G' , respectively (see Lemma 31).

Now, let $A_{\mathbb{Q}}(\chi)$ be the simple component of the Wedderburn decomposition of $\mathbb{Q}G$ corresponding to the rational representation of G that affords the character $\Omega(\chi)$. Then $A_{\mathbb{Q}}(\chi) \cong M_n(D)$ for some $n \in \mathbb{N}$ and a division ring D . Observe that $n = p$ and $D = \mathbb{Q}(\chi)$ (see Lemma 8). Therefore, all the irreducible rational representations of G

whose kernels do not contain G' contribute

$$\bigoplus_{d|m, d|m'} \frac{a_d}{p} M_p(\mathbb{Q}(\zeta_d)) \bigoplus_{d|m, d|m'} \frac{a_d - a'_d}{p} M_p(\mathbb{Q}(\zeta_d))$$

in the Wedderburn decomposition of $\mathbb{Q}G$. This completes the proof of Theorem 3. \square

Corollary 60 immediately follows from Theorem 3.

COROLLARY 60. *Suppose G is a nonabelian p -group ($p \geq 5$) of order p^4 in Φ_3 .*

(1) *If $G = \Phi_3(211)a$ or $\Phi_3(1^4)$, then*

$$\mathbb{Q}G \cong \mathbb{Q} \bigoplus (p+1)\mathbb{Q}(\zeta_p) \bigoplus (p+1)M_p(\mathbb{Q}(\zeta_p)).$$

(2) *If $G = \Phi_3(211)b_r$ ($r = 1, \nu$), then*

$$\mathbb{Q}G \cong \mathbb{Q} \bigoplus (p+1)\mathbb{Q}(\zeta_p) \bigoplus M_p(\mathbb{Q}(\zeta_p)) \bigoplus M_p(\mathbb{Q}(\zeta_{p^2})).$$

REMARK 61. For $p = 3$, let $G \in \Phi_3$. For $G = \Phi_3(211)b_1$,

$$\mathbb{Q}G \cong \mathbb{Q} \bigoplus 4\mathbb{Q}(\zeta_3) \bigoplus 4M_3(\mathbb{Q}(\zeta_3)).$$

Additionally, for $G = \Phi_3(1^4)$ or $\Phi_3(211)a$ or $\Phi_3(211)b_2$,

$$\mathbb{Q}G \cong \mathbb{Q} \bigoplus 4\mathbb{Q}(\zeta_3) \bigoplus M_3(\mathbb{Q}(\zeta_3)) \bigoplus M_3(\mathbb{Q}(\zeta_9)).$$

We end this subsection with the following remark.

REMARK 62. It is well known that if G is a 2-group of maximal class, then G is isomorphic to one of the following: a dihedral group, a semi-dihedral group or a generalized quaternion group. Let G be a 2-group of order 16 of maximal class (that is, of nilpotency class 3). Then G has two inequivalent irreducible two-dimensional faithful complex representations and one irreducible two-dimensional nonfaithful complex representation. Observe that the irreducible two-dimensional nonfaithful complex representation of G is realizable over \mathbb{Q} . From Lemma 18, one can compute an irreducible rational matrix representation of G that affords the character $\Omega(\chi)$, where $\chi \in \text{Flrr}(G)$.

Acknowledgement

We thank the anonymous referee for their comments on the article.

References

- [1] R. G. Ayoub and C. Ayoub, 'On the group ring of a finite abelian group', *Bull. Aust. Math. Soc.* **1** (1969), 245–261.
- [2] G. K. Bakshi and G. Kaur, 'Character triples and Shoda pairs', *J. Algebra* **491** (2017), 447–473.
- [3] G. K. Bakshi and G. Kaur, 'A generalization of strongly monomial groups', *J. Algebra* **520** (2019), 419–439.

- [4] G. K. Bakshi and S. Maheshwary, 'The rational group algebra of a normally monomial group', *J. Pure Appl. Algebra* **218**(9) (2014), 1583–1593.
- [5] G. K. Bakshi and I. B. S. Passi, 'Primitive central idempotents in rational group algebras', *Comm. Algebra* **40**(4) (2012), 1413–1426.
- [6] W. Burnside, *Theory of Groups of Finite Order* (Cambridge University Press, Cambridge, 1911).
- [7] Á. del Río and M. Ruiz, 'Computing large direct products of free groups in integral group rings', *Comm. Algebra* **30**(4) (2002), 1751–1767.
- [8] G. A. Fernández-Alcober and A. Moretó, 'Groups with two extreme character degrees and their normal subgroups', *Trans. Amer. Math. Soc.* **353**(6) (2001), 2171–2192.
- [9] C. E. Ford, 'Characters of p -groups', *Proc. Amer. Math. Soc.* **101**(4) (1987), 595–601.
- [10] P. Hall, 'The classification of prime-power groups', *J. reine angew. Math.* **182** (1940), 130–141.
- [11] A. Herman, 'On the automorphism group of rational group algebras of metacyclic groups', *Comm. Algebra* **25**(7) (1997), 2085–2097.
- [12] I. M. Isaacs, *Character Theory of Finite Groups* (Dover Publications Inc., New York, 1994).
- [13] R. James, 'The groups of order p^6 (p an odd prime)', *Math. Comp.* **34**(150) (1980), 613–637.
- [14] E. Jespers and Á. del Río, 'A structure theorem for the unit group of the integral group ring of some finite groups', *J. reine angew. Math.* **521** (2000), 99–117.
- [15] E. Jespers and Á. del Río, *Group Ring Groups, Volume 1: Orders and Generic Constructions of Units* (De Gruyter, Berlin, 2016).
- [16] E. Jespers and G. Leal, 'Generators of large subgroups of the unit group of integral group rings', *Manuscripta Math.* **78**(3) (1993), 303–315.
- [17] E. Jespers, G. Leal and A. Paques, 'Central idempotents in rational group algebras of finite nilpotent groups', *J. Algebra Appl.* **2**(1) (2003), 57–62.
- [18] M. L. Lewis, 'Generalizing Camina groups and their character tables', *J. Group Theory* **12**(2) (2009), 209–218.
- [19] M. L. Lewis, 'The vanishing-off subgroup', *J. Algebra* **321**(4) (2009), 1313–1325.
- [20] M. F. Newman, E. A. O'Brien and M. R. Vaughan-Lee, 'Presentations for the groups of order p^6 for prime $p \geq 7$ ', Preprint, 2023, [arXiv:2302.02677](https://arxiv.org/abs/2302.02677) [math.GR].
- [21] A. Olivieri, Á. del Río and J. J. Simón, 'On monomial characters and central idempotents of rational group algebras', *Comm. Algebra* **32**(4) (2004), 1531–1550.
- [22] A. Olivieri, Á. del Río and J. J. Simón, 'The group of automorphisms of a rational group algebra of a finite metacyclic group', *Comm. Algebra* **34**(10) (2006), 3543–3567.
- [23] G. Olteanu, 'Computing the Wedderburn decomposition of group algebras by the Brauer–Witt theorem', *Math. Comp.* **76**(258) (2007), 1073–1087.
- [24] S. Perlis and G. L. Walker, 'Abelian group algebras of finite order', *Trans. Amer. Math. Soc.* **68**(3) (1950), 420–426.
- [25] S. K. Prajapati, M. R. Darafsheh and M. Ghorbani, 'Irreducible characters of p -group of order $\leq p^5$ ', *Algebr. Represent. Theory* **20**(5) (2017), 1289–1303.
- [26] S. K. Prajapati and R. Sharma, 'Total character of a group G with $(G, Z(G))$ as a generalized Camina pair', *Canad. Math. Bull.* **59**(2) (2016), 392–402.
- [27] I. Reiner, 'The Schur index in the theory of group representations', *Michigan Math. J.* **8** (1961), 39–47.
- [28] J. Ritter and S. K. Sehgal, 'Construction of units in integral group rings of finite nilpotent groups', *Trans. Amer. Math. Soc.* **324**(2) (1991), 603–621.
- [29] J. Tappe, 'On Isoclinic groups', *Math. Z.* **148** (1976), 147–153.
- [30] T. Yamada, *The Schur Subgroup of the Brauer Group*, Lecture Notes in Mathematics, 397 (Springer Verlag, Berlin, 1974).
- [31] T. Yamada, 'Types of faithful metacyclic 2-groups', *SUT J. Math.* **28**(1) (1992), 23–46.
- [32] T. Yamada, 'Remarks on rational representations of a finite group', *SUT J. Math.* **29**(1) (1993), 71–77.

RAM KARAN CHOUDHARY, Indian Institute of Technology, Bhubaneswar,
Arugul Campus, Jatni, Khurda 752050, India
e-mail: ramkchoudhary1997@gmail.com

SUNIL KUMAR PRAJAPATI, Indian Institute of Technology, Bhubaneswar,
Arugul Campus, Jatni, Khurda 752050, India
e-mail: skprajapati@iitbbs.ac.in