

SUBJECT INDEX

VOLUME 38, NOS. 2 AND 3, 1996¹

- Accelerator mass spectrometry (AMS), 347–353, 389–406, 415–423, 597–602
- Antarctic Ocean, 394
- Archaeological dates, 355–373
- Arctic soils, 241–245
- Arctic Circle, 597–602
- Arctic Ocean, 394
- Atlantic Ocean, 389–406; North, 407–414; tropical, 563–572
- Bering Sea, 583–595
- Bomb ¹⁴C, 389–406, 407–414
- Brazil, 563–572
- Calibration, ¹⁴C, 607–610
- Canada, 597–602
- Canary Islands, 389–406
- Cape Verde Islands, 563–572
- Carbon cycling: in soil, 181–190, 231–239; models, 209–217, 253–270
- Carbon dioxide (CO₂): fertilization, 187–188; storage in soil, 209–217
- Carbon pools, in soil, 231–239
- Cesium (¹³⁷Cs), 211–213
- Charcoal, in soil, 194–196
- Chernobyl event of 1986, 393
- Coral, 563–572
- δ¹³C, of soil organic matter, 191–201
- Data base, soils, 177–180, 271–275, 277–293, 295–345
- Deep-sea profiles, 396–403
- Dissolved inorganic carbon (DIC), 407–414, 563–572, 583–595
- Dissolved organic carbon (DOC), 247–251
- DNA adduction, 347–353
- El Niño – Southern Oscillation (ENSO) events, 597–602
- Factor of merit (FM), 375–384
- Fractionation, of soil organic matter, 181, 219–229, 231–239
- GEOSCEX expeditions: 1972, 389–406, 407–414; 1973 to 1974, 415–423
- Global Carbon Cycle Models (GCCM), 177–180
- Global warming, 241–245
- Greenland Sea, 389–406
- Gulf of Mexico, 407–414
- High Resolution Biosphere Model (HRBM), 177–180
- Holocene: climate, 191–201; vegetation changes, 191–201, 203–208
- Humic acids, 241–245
- Isopycnals, 407–414
- Laboratory reliability, 375–384
- Marine reservoir ages, 573–582
- Minnesota, 597–602
- Mollusk shells, 573–582
- Nicotine, 347–353
- Norwegian Sea, 389–406
- Pacific Northwest (USA), 597–602
- Pacific Ocean, 415–423, 415–423, 519–561, 573–582
- Paleovegetation, 198–199
- Particulate inorganic carbon (PIC), 583–595
- Regional ¹⁴C effects, 597–602, 607–610
- San Francisco Bay, 573–582
- San Miguel Island (California), 355–373
- Schwabe cycle, 597–602
- Sea of Okhotsk, 583–595
- Sediment trap samples, 583–595
- Shell, ¹⁴C dates on, 355–373
- Soil: bulk ¹⁴C measurements, 181–190; cultivated vs. uncultivated, 181–190, 209–217
- Soil locations: Arizona, 231–239; Brazil, 191–201, 203–208; Canada, 271–275; Germany, 247–251; Great Plains (USA), 231–239; Michigan, 231–239; Nebraska, 209–217; New Zealand, 184–185, 253–270; Norway, 241–245; Russia, 241–245; South Carolina, 186
- Soil organic matter (SOM): chronology, 203–208; global inventory, 186–187; stable isotope composition, 191–201, 203–208; turnover, 181–190, 209–217, 247–251, 253–270
- Surface waters, 389–406
- Thermocline, 407–414
- Tillage, 209–217
- Tree rings, 597–602, 607–610
- TTO expeditions: 1981, 389–406
- Varve chronology, 603–605
- Vegetation: C₃ vs. C₄, 191–201, 203–208
- Vegetation changes, 191–201, 203–208
- World Ocean Circulation Experiment (WOCE), 415–423, 425–518, 519–561

¹Vol. 38, No. 1, the AMS-7 Abstracts issue, is not included.