

later for patients who had not developed a psychosis. The MRI data from the 2 time points were compared within each group.

Results: (a) *Cross-sectional comparison:* relative to the group who did not become psychotic, those going on to develop psychosis had smaller grey matter volumes in right temporal temporal and inferior frontal cortex, and in the cingulate cortex bilaterally. (b) *Longitudinal comparison:* in the group who became psychotic there were reductions in grey matter volume in the medial temporal and anterior cingulate cortex bilaterally, the left fusiform and inferior frontal cortex and in the cerebellar cortex. There were no changes in the group who remained non-psychotic.

Discussion: There were marked differences in regional grey matter volume between high-risk subjects who later developed psychosis and those who did not, despite the absence of clinical differences at the time of scanning. The group who went on to develop psychosis showed longitudinal reductions in regional grey matter volume in association with the expression of frank psychotic symptoms. These data suggest that in psychotic disorders some abnormalities of grey matter volume predate the onset of frank symptoms while others appear in association with the first episode of psychosis.

S09.5

Disordered brain development and abnormal connectivity

E.T. Bullmore. *Department of Psychiatry, University of Cambridge, UK*

There is some evidence both for schizophrenia as a disorder of brain development and for schizophrenia as a disconnection syndrome. Imaging evidence for dysconnectivity in adult schizophrenia is reviewed and possible mechanisms by which abnormal early development might lead to adult dysconnectivity are rehearsed. One experimental approach to securing a more robust link in general between mechanisms of abnormal neurodevelopment and patterns of abnormal adult brain structure is to study rare neurogenetic syndromes where a specific genetic lesion is associated with well-characterised developmental abnormalities in animal models. An example of this approach is provided by a structural and functional MRI study of a human family with heterozygous mutation in PAX6, a highly-conserved neuro-developmental control gene which is important for inter-regional boundary demarcation and guided axonal growth in mice. The adult human phenotype is characterised by deficits in major white matter tracts and distributed functional deficits in fronto-striatal circuits. The implications for schizophrenia as a syndrome of abnormal development of neurocognitive networks are discussed.

S10. Eating disorders

Chairs: H. Wijbrand Hoek (NL), J. Treasure (GB)

S10.1

Evidence based treatments for anorexia nervosa

J. Treasure*. *GKT & IOP, Kings College London, UK*

Evidence based medicine (EBM) is the *integration* of best research evidence, together with clinical expertise and patient values (Sackett et al., 2000). All too often when the evidence based medicine approach has been considered there is a tendency to focus on the quality of the evidence for treatment. The second part of Sackett's definition, which discusses clinical expertise and patient's values,

tends to be overlooked. There is very little in the way of Level I and II evidence about the efficacy treatment in anorexia nervosa. However the fact that there is no evidence from RCT's should not be interpreted as if these treatments are of no value. It is not appropriate to dismiss treatment of the starvation state because of paucity of specific evidence in anorexia nervosa, as the natural history of starvation is known and effective treatment of starvation is also known. Thus there is an argument for not requiring evidence from RCTs to resuscitate and embark on treatment. There is a detailed, coherent body of research, which documents prognostic features and the factors that have to be considered in terms of the acute medical risk. Medical risk is critically important to guide the acute management of anorexia nervosa. The acute risk management involves a combination of the medical risk and psychological capacity set against the possible resources of motivation and psychosocial support.

Once we are out of these "fire fighting" stages there is evidence that specific psychotherapies are more effective than supportive counselling and dietary advice. It is useful to involve families in management but how and by how much is less certain. The early phase of research into pharmacotherapy produced little benefit, but new drugs and new paradigms such as using drugs to prevent relapse rather than to treat starvation are of interest.

S10.2

Evidence based treatment for bulimia nervosa

H. Wijbrand Hoek*. *Parnassia – The Hague Psychiatric Institute, The Hague, The Netherlands*

Eating disorders are mental disorders occurring mainly among young females. The prevalence of bulimia nervosa according to DSM-IV criteria among young females is 1%. Bulimia nervosa leads to serious physical, psychological and social consequences. Women suffering from bulimia nervosa are so ashamed of their disturbed eating behaviour that they hardly look for professional help. Only 6% of all women with bulimia nervosa in the population do come into mental health care.

Systematic reviews of large randomized controlled trials found that cognitive behavioural therapy compared with remaining on a waiting list reduced the symptoms of bulimia nervosa and improved non-specific symptoms such as depression. The NNT (Number Needed to Treat) of CBT is 3. The absolute remission rate of bulimia nervosa for CBT is around 40%. One 5-year follow-up study showed that the effect of CBT remained. Selfhelp based on CBT seems also to be effective.

Systematic reviews of RCT's with antidepressants compared to placebo have found a significant short-term reduction of bulimic symptoms.

S10.3

Osteopenia and bone mass increase in adolescent anorexia nervosa

J. Castro Fornieles. *University of Barcelona Hospital Clinic, Section of Child & Adolescent Psychiatry, Spain*

The percentage of patients with osteopenia, the variables related and the bone mass increase after recovery were studied. Bone mass was measured by dual-energy-x-ray absorptiometry in 180 female and 20 male adolescents with anorexia nervosa. The results were compared with normative values for bone mass in Spanish adolescents. In 108 females and 15 males a second examination was carried out after a follow-up of six to thirty four months. At lumbar spine 44.1% of girls and 35% of boys had osteopenia. The variables related to osteopenia were duration of illness and

amenorrhea ($p < .001$), physical activity ($p < .001$), calcium intake ($p < .001$) and body mass index ($p < .001$). At follow-up, subjects with good outcome had a higher percentage of bone mass increase than normal adolescents, whereas patients with poor outcome had a bone mass loss. Conclusions: 35–44 % of patients had osteopenia and related variables were body mass index, duration of illness, calcium intake and physical activity. Patients with good outcome had a high bone mass increase.

S10.4

Genes, environment, and eating disorders: twin study findings

A. Keski-Rahkonen¹*, A. Rissanen¹, J. Kaprio¹, R.J. Rose².

¹University of Helsinki, Finland

²Indiana University, Bloomington, USA

Eating disorders are highly heritable conditions. Body dissatisfaction and dieting-oriented behavior are common in young people and often predispose vulnerable individuals to eating disorders.

We assessed pairwise twin correlations and heritability estimates of dieting frequency and Eating Disorder Inventory subscales body dissatisfaction and drive for thinness in a population sample of 936 female and 811 male Finnish twins aged 24–25. Body dissatisfaction was much influenced by genes, with different sets of genes operating in females and males, and with possible genetic dominance in males. Drive for thinness and dieting frequency had a moderate heritability in females and a lower heritability in males. Individual-specific environmental factors were relatively important for body dissatisfaction, drive for thinness, and dieting frequency in both males and females, but environmental factors shared by the twin pair were of negligible importance.

The genetic factors influencing body dissatisfaction and dieting-oriented behavior may constitute a part of the genetic vulnerability to eating disorders. These influences are likely to be age-specific and sex-specific.

S10.5

Somatic consequences of eating disorders

T. Bruna*. Robert Fleury Stichting, Leidschendam, The Netherlands

More than other psychiatric disorders eating disorders are conditions in which a disturbed psyche directly contributes to a disturbed soma.

This study aims to assess the status at admission and course of the somatic consequences and laboratory findings in a sample of patients with serious and long lasting eating disorders.

Data were analysed per diagnostic group. Associations with ED-symptom severity were examined. Survival analyses were conducted to examine whether the physical status at admission could predict treatment outcome.

The sample consisted of 167 DSM-IV ED patients (mean age 28 years, mean duration of illness 11 years) referred to a tertiary care centre for inpatient treatment or day care. A physical examination and an extensive laboratory investigation were carried out.

Abnormal findings included anaemia, leucopenia and disturbed liver and renal function tests especially for the AN-group and electrolyte disturbances especially for the AN B/P type patients and BN patients. Most abnormal findings improved during treatment.

Empirically based guidelines for routine laboratory investigations are presented.

S11. Reforming psychiatry in Eastern Europe

Chairs: W. Rutz (WHO, Europe), L. Jacobsson (S)

S11.1

Swedish support to the restructuration of psychiatry in Eastern Europe

L. Jacobsson. Department of Psychiatry, Umeå, Sweden

The Swedish government has supported the restructuration of mental health services in North-western Russia, the Baltic countries and Bosnia Herzegovina since the early 1990'ies. This support has been channeled through the East Europe Committee of the Swedish Health Care Community, which is a joint body of all major actors in the Swedish health care community e.g. all professional organizations and the major care providers. Since 1995 more than 70 million Swedish crowns (6 million dollars) have been spent on hundreds of collaborative projects in which Swedish clinics and university institutions have been engaged in teaching and training of staff, study visits and supervision activities. Several thousands of professionals have been engaged in these activities. Some of these will be presented. Some evaluations of these projects have also been done and will be described.

S11.2

Reforming psychiatry in Eastern Europe – the WHO perspective

W. Rutz*. WHO Regional Office for Europe, Copenhagen, Denmark

Since 1999 the mental health program of the WHO Regional Office for Europe has been intensively involved in supporting necessary mental health reforms in the countries of transition in eastern, central and southeastern Europe. Focus of the work of the mental health program have been:

- Suicidality and other premature mortality, related to transitional stress and mental ill health.
- National mental health audits and planning.
- Restructuring, modernizing and humanizing services, leading to de-hospitalization, decentralization and community-based mental health services.
- And finally, counteracting taboo and discrimination.

This work is made possible with the help of WHO collaborating organizations, WHO task forces and WHO networks this work is made possible and it will be described. Outcomes as well as possible shortcomings will be analyzed.

S11.3

Reforming psychiatric services in Bosnia Hercegovina after the war

S. Loga*. University of Sarajevo, Bosnia and Hercegovina

Before the war in Bosnia (1992–1995) the organization of psychiatric service was on a relatively high level and not different from the other republics in former Yugoslavia. During the beginning of the war, most of the psychiatric institutions were closed, damaged or devastated. In spite of this psychiatric services continued to work, very often in improvised and poor conditions with the help of WHO and some NGO:s. After the war, in 1996, Federation of BiH government made an agreement with the World Bank on building