

THE DETERMINATION OF THE PRINCIPAL NUTATION TERMS FROM THE
OBSERVATIONS WITH ZTF-135 IN 1948.7-1989.0

L.A. GLEBOVA, L.D. KOSTINA, Z.M. MALKIN AND N.R. PERSIYANINOVA
Central Astronomical Observatory, USSR Academy of Sciences
Pulkovo
196140 Leningrad
USSR

ABSTRACT. A re-reduction of observations made with ZTF-135 in Pulkovo during 1948.7-1989.0 (all together about 87 000 observations) has been carried out with the aim of determining the amplitude of the principal terms of nutation in longitude and obliquity. The derived coefficients of nutation $M=6''.8445 \pm 0''.0033$ and $N=9''.2028 \pm 0''.0025$ are in good agreement with similar astrometric determinations.

The two observational programmes were used for ZTF-135 in 1948.7- 1989.0. During the first programme (Gordon, 1948) from 1948.7 to 1968.1 about 48 000 observations were made and during the second one (Kostina, 1970) from 1968.1 to 1989.0 about 39 000 observations were made. All individual latitudes were recalculated using the MERIT Standard algorithm and were corrected for polar motion. As the observations are not yet reduced to a uniform system, we estimated nutation coefficients for the two programmes independently. For further calculations we have formed the normal points for each Talcott pair as the mean value from observations made in one season, *i.e.* one normal point in a year for each pair.

Two methods were used in the present work. According to the first one, previously used for ZTF-135 observations (Kulikov, 1949; Romanskaya, 1961; Kostina et al., 1988), we solved by least squares the equation:

$$\varphi = \varphi_0 - \Delta\delta - \Delta\mu(t - t_0) - dN(\sin \alpha \cos \Omega - m \cos \alpha \sin \Omega)$$

where dN is the correction to the conventional coefficient of nutation in obliquity, $m = 0.7434$ is the relation between the axes of nutation ellipse and the other designations are obvious.

Thus we found for the two programmes
 $N = 9''.2028 \pm 0''.0022$ (epoch 1958.4),
 $N = 9''.2147 \pm 0''.0049$ (epoch 1978.5)
and the weighted mean
 $N = 9''.2049 \pm 0''.0020$ (epoch 1968.5).

However, as analysis of the algorithm shows, the coefficient of nutation obtained using such a method significantly depends on the conventional value m . For control we applied this method to

the latitudes reduced using the Woolard nutation coefficients. In this case we found $N=9''.1955$. It is clear that the method discussed is not satisfactory.

Therefore we estimated the coefficients of nutation solving by least square the well known equations:

$$\varphi = \varphi_0 - \Delta\delta - \Delta\mu(t - t_0) - dN \sin \alpha \cos \Omega + dM \cos \alpha \sin \Omega$$

where dM and dN are the corrections to the conventional coefficients of nutation in longitude and obliquity respectively.

Thus we found for the same epochs, given earlier, the values

$$M_1 = 6''.8414 \pm 0''.0035, \quad N_1 = 9''.2009 \pm 0''.0028,$$

$$M_2 = 6''.8645 \pm 0''.0094, \quad N_2 = 9''.2101 \pm 0''.0056$$

and the weighted mean values

$$M = 6''.8445 \pm 0''.0033, \quad N = 9''.2028 \pm 0''.0025,$$

which are in good agreement with other astrometric determinations (Capitaine *et al.*, 1988).

The authors wish to express their thanks to A.R. Valsky for his help in calculations.

REFERENCES

- Capitaine, N., Li, Z.X., and Nie, S.Z.: 1988, *Astron. Astrophys.*, **202**, 306–308.
 Gordon, Ya.E.: 1948, *Izvestiya GAO AN SSSR*, **141**, 130–139 (in Russian).
 Kostina, L.D.: 1970, *Izvestiya GAO AN SSSR*, **185**, 49–69 (in Russian).
 Kostina, L.D., Malkin Z.M., and Persiyaninova N.R.: 1988, In: *The study of the Earth as a planet by methods of geophysics, geodesy and astronomy*, Kiev, 8–10 (in Russian).
 Kulikov, K.A.: 1949, *Astron. Zhurnal*, **26**, 165–174.
 Romanskaya, C.V.: 1961, In: *Preliminary results of the investigations of the latitude oscillations and Polar motion*, Moscow, 81–85 (in Russian).