

Short Communication

Food insecurity among veterans of the US wars in Iraq and Afghanistan

Rachel Widome^{1,*}, Agnes Jensen², Ann Bangerter² and Steven S Fu^{2,3}

¹Division of Epidemiology and Community Health, University of Minnesota School of Public Health, 1300 South 2nd Street #300, Minneapolis, MN 55454, USA: ²VA HSR&D Center for Chronic Disease Outcomes Research (CCDOR), Minneapolis VA Health Care System, Minneapolis, MN, USA: ³Department of Medicine, University of Minnesota Medical School, Minneapolis, MN, USA

Submitted 16 December 2013: Final revision received 25 March 2014: Accepted 27 March 2014: First published online 8 May 2014

Abstract

Objective: Food insecurity, or lack of access to sufficient food for a healthful lifestyle, has been associated with many aspects of poor health. While the economic struggles among veterans of the wars in Iraq and Afghanistan have been documented, it is unknown how commonly this population struggles to afford food. Our purpose was to document the prevalence and correlates of food insecurity among US veterans of the wars in Iraq and Afghanistan.

Design: A cross-sectional survey.

Subjects: US military veterans who had served in the wars in Iraq and Afghanistan since October 2001.

Setting: Subjects responded to a survey mailed to them in summer 2012. Food security was measured by the US Household Food Security Module: Six Item Short Form. Demographic and behavioural health items were also included. Survey data were matched to medical record data from the Department of Veterans Affairs.

Results: Over one in four veterans reported past-year food insecurity with 12% reporting very low food security. Food-insecure veterans tended to be younger, not married/partnered, living in households with more children, earning lower incomes, had a lower final military pay grade, were more likely to use tobacco, reported more frequent binge drinking and slept less, compared with those who were food secure ($P < 0.05$ for all associations listed).

Conclusions: Previously undocumented, the problem of hunger among our newest veterans deserves attention.

Keywords
Food security
Veterans
Behavioural health

Since October 2001, 2.5 million American military members have served in the US wars in Iraq and Afghanistan⁽¹⁾. These veterans often face challenges upon their return to civilian life such as unemployment, difficulties in reintegrating with their family and into the community, mental health struggles and tobacco dependence⁽²⁾. While economic issues among returning veterans have been documented, less is known about how financial hardship is affecting veteran households.

Food security, or the consistent ability to access sufficient food for a healthful lifestyle, has been associated with many aspects of health including weight gain, diabetes and mental health issues^(3–9). In the USA food insecurity remains a problem; 14.5% of households were classified as food insecure in 2012⁽¹⁰⁾.

In the present paper, the prevalence of food insecurity is reported and the demographic and health-related

characteristics that may be associated with food insecurity are described among Iraq and Afghanistan war veterans.

Methods

Using the Department of Veteran Affairs' (VA) OEF/OIF/OND (Operation Enduring Freedom/Operation Iraqi Freedom/Operation New Dawn) Roster⁽¹¹⁾, which identifies all veterans who have served in the US wars in Iraq and Afghanistan since October 2001, 800 female and 1200 male veterans who had at least one out-patient health-care visit in the Minneapolis VA Healthcare System and had a telephone number listed in the record were randomly sampled. At the time of the survey (summer 2012), 70.8% of the current addresses on file were within the state of Minnesota and the rest were outside the state.

*Corresponding author: Email widome@umn.edu

Women were oversampled in order to have sufficient numbers to examine gender differences. Initially, potential participants were mailed a package that included an invitation to participate, informed consent material, a survey, a stamped return envelope and a \$US 20 incentive. We developed the Northstar survey to examine health behaviours related to chronic disease (such as tobacco use, physical activity, eating and sun exposure). A reminder postcard followed a week after the initial mailing; two weeks after the initial mailing non-respondents received a second survey packet that did not include an incentive⁽¹²⁾. The survey response rate was 52.3%, which exceeds the response rate of nearly all other population-based survey research in Iraq and Afghanistan war veterans^(13–17). Of the 922 respondents, there were fifty-seven individuals for whom food security status could not be calculated due to item non-response, service era misclassification or other reasons. Northstar survey data were supplemented with additional information on the veterans drawn from the VA's electronic medical record.

Food security was ascertained using the US Household Food Security Module: Six Item Short Form⁽¹⁸⁾, which measures food security over the prior 12 months and has been demonstrated to be a valid identifier of households that have low and very low food security⁽¹⁹⁾. This measure includes items such as: 'The food that (I/we) bought just didn't last, and (I/we) didn't have money to get more' and '(I/we) couldn't afford to eat balanced meals' (participants are instructed to indicate how often these statements were true)⁽¹⁸⁾. 'High/marginal food security' means that either there are no reported food access problems or that the household has some anxiety over food sufficiency but that there is minimal or no impact on diet. Households with 'low food security' report reduced diet quality but have little or no report of reducing intake. 'Very low food security' households report multiple past-year impacts on their eating which affected food intake⁽²⁰⁾.

On the Northstar survey, participants were asked to indicate income ('How much did you earn, before taxes and other deductions, during the past 12 months?') from a choice of nine brackets. We assigned the participant the median value of his/her selected bracket with the exception of the '\$US 100 000 or greater' bracket which we assigned a value of \$US 105 000. Employment was measured with the question, 'During the past 6 months, what were you doing on most days?' Past 30 d tobacco use was assessed with, 'In the past 30 d, did you use tobacco every day, some days or not at all?' Binge drinking was assessed with the item, 'Considering all types of alcoholic beverages, how many times during the past 30 d did you have five or more drinks on one occasion (four or more if you are female)?' Sleep was assessed with the question, 'How much sleep do you usually get at night on weekdays or workdays?' Physical activity was assessed using the International Physical Activity Questionnaire (IPAQ) Short Form⁽²¹⁾, which is a four-item measure of past 7 d physical activity.

For selected demographic and behavioural attributes, either the proportion of veterans in each of the three levels of food security (high/marginal, low and very low) or mean values for each level were computed. The χ^2 test or linear regression was used to test whether the demographic and behavioural factors had a bivariable association with level of food security. All variables that had a significant bivariable association with food security ($P < 0.05$) were entered into a multinomial logistic model. Variables in this initial mutually adjusted model that did not have a significant association ($P < 0.05$) with food security were removed to arrive at the final model presented. Analyses were conducted in 2013 using the SAS statistical software package version 9.2. All procedures were reviewed and approved by the Minneapolis VA institutional review board.

Results

Over one in four veterans (~27%) of the wars in Iraq and Afghanistan reported problems with food security. About 15% of veterans reported low food security and an additional 12% reported very low food security (Table 1). Veterans were more likely to be food insecure if they were younger, not married/partnered and not employed or on active duty. Food-insecure veterans had lower current income, reported lower final military pay grade and lived in households with more children. Those who were food insecure were more likely to use tobacco, report more frequent binge drinking and slept fewer hours at night. There was a gradient evident for self-reported general health status, with better health reported by those who were food secure.

When the characteristics that had significant bivariable associations with food security were combined into a multivariable model, marital status, general health status, tobacco use, income, children in the household and mean hours of sleep continued to have associations with food security at the $P < 0.05$ level (see Table 2). For instance, those who were married or partnered had 63% reduced odds of being at very low food security, compared with high/marginal level (adjusted OR = 0.37; 95% CI 0.19, 0.71). For each \$US 10 000 increase in reported income, the adjusted odds ratio of being at very low food security (compared with high/marginal food security) was 0.74 (95% CI 0.70, 0.79).

To estimate whether the survey responders differed systematically from non-responders, comparisons were made on several variables from the electronic medical record. Compared with non-responders, responders were more likely to be older (34.9 years *v.* 31.0 years, $P < 0.0001$), married or partnered (43.9% *v.* 32.5%, $P < 0.0001$), and less likely to have service-connected disability status (35.7% *v.* 43.2%, $P = 0.0006$) or to be male (55.1% *v.* 64.2%, $P < 0.0001$).

Table 1 Demographics and selected health behaviours by level of food security among veterans of the US wars in Iraq and Afghanistan, Northstar–2012

	Food security level						Total*	P
	High/marginal		Low		Very low			
	n	%	n	%	n	%		
Overall	633	73.2	127	14.7	105	12.1	865	
Gender†								
Male	355	74.1	66	13.8	58	12.1	479	0.700
Female	279	72.1	61	15.8	47	12.1	387	
Race/ethnicity								
Non-white	48	62.3	15	19.5	14	18.2	77	0.087
White, non-Hispanic	578	74.2	112	14.3	89	11.4	779	
Age (years)†, mean	35.5		32.8		33.0			0.007
Marital status								
Single	154	67.2	44	19.2	31	13.5	229	0.0001
Married or partnered	397	82.7	54	11.3	29	6.0	480	
Divorced, separated, widowed	79	53.4	29	19.6	40	27.0	148	
Income (individual) in past 12 months (\$US), mean	48 305		30 734		24 830			0.0001
Employment (individual)								
Employed for wages	352	76.2	66	14.3	44	9.5	462	0.0008
Self-employed	15	75.0	1	5.0	4	20.0	20	
On active duty	48	87.3	6	10.9	1	1.8	55	
Looking for work	36	60.0	15	25.0	9	15.0	60	
On disability/unable to work	12	48.0	7	28.0	6	24.0	25	
Caring for family	12	66.7	3	16.7	3	16.7	18	
Student	67	72.8	12	13.0	13	14.1	92	
Retired	14	87.5	0	0.0	2	12.5	16	
More than one selected	72	65.5	16	14.6	22	20.0	110	
No. of household members, mean	2.9		3.01		3.25			0.058
No. of children in household, mean	0.84		0.85		1.25			0.003
Highest military pay grade‡								
E1–E6	414	67.6	109	17.8	89	14.5	612	< 0.0001
E7–E9	92	79.3	14	12.1	10	8.6	116	
Any W or O grade	127	93.4	4	2.9	5	3.7	136	
Service-connected disability†,§								
No	231	74.8	48	15.5	30	9.7	309	0.244
Yes	402	72.3	79	14.2	75	13.5	556	
No. of deployments								
1	393	72.5	84	15.4	65	12.0	542	0.286
2	187	81.7	31	13.5	11	4.8	229	
3 or more	65	62.5	25	24.0	14	13.5	104	
Self-reported general health status								
Excellent	55	91.7	4	6.7	1	1.7	60	0.0001
Very good	249	73.9	31	9.2	57	16.9	337	
Good	254	70.2	57	15.7	51	14.1	362	
Fair	66	54.5	27	22.3	28	23.1	121	
Poor	7	30.4	8	34.8	8	34.8	23	
Past 30 d tobacco use								
None	433	79.4	70	12.8	42	7.7	545	0.0001
Some days	80	70.8	14	12.4	19	16.8	113	
Every day	120	58.0	43	20.8	44	21.3	207	
Binge drinking								
Never	309	77.1	49	12.2	43	10.7	401	0.019
1–3 times per month	210	70.7	55	18.5	32	10.8	297	
Once per week or more	108	68.4	21	13.3	29	18.4	158	
Hours of sleep, mean	6.7		6.2		5.9			0.0001
Physical activity								
High	358	74.1	62	12.8	63	13.0	483	0.349
Medium	141	73.8	32	16.8	18	9.4	127	
Low	134	70.2	33	17.3	24	12.6	105	

*Some demographic and behavioural category totals do not add up to 865 due to missing values.

†These data were obtained from the electronic medical record.

‡E1–E9 are enlisted ranks; W and O are officer ranks.

§'Service-connected disability' refers to veterans who are receiving compensation at any level for an injury, physical illness or mental illness that was incurred or exacerbated during active duty service.

Table 2 Adjusted odds ratios (AOR) depicting the associations between characteristics and food security among veterans of the US wars in Iraq and Afghanistan, Northstar – 2012. (All predictor variables in the table are mutually adjusted in the multinomial logistic model)

	Food security status						P
	Low			Very low			
	AOR	95 % CI	Ref.	AOR	95 % CI	Ref.	
Marital status							<0.0001
Single	Ref.	Ref.	Ref.	Ref.	Ref.	Ref.	
Married or partnered	0.59	0.35	1.00	0.37	0.19	0.71	
Divorced, separated, widowed	1.27	0.68	2.36	2.07	1.08	3.97	
Income (per \$US 10 000 difference)	0.82	0.78	0.86	0.74	0.70	0.79	<0.0001
No. of children in household	1.21	0.99	1.49	1.61	1.29	2.01	0.0002
General health status							0.0094
Excellent	Ref.	Ref.	Ref.	Ref.	Ref.	Ref.	
Very good	1.90	0.54	6.67	2.58	0.32	20.73	
Good	2.69	0.78	9.23	5.21	0.67	40.26	
Fair	4.34	1.19	15.87	7.49	0.92	60.89	
Poor	8.55	1.63	44.90	13.79	1.30	146.64	
Past 30 d tobacco use							0.0332
None	Ref.	Ref.	Ref.	Ref.	Ref.	Ref.	
Some days	0.98	0.50	1.91	1.81	0.91	3.61	
Every day	1.61	0.98	2.65	2.28	1.29	4.03	
Hours of sleep	0.86	0.73	1.00	0.79	0.66	0.94	0.0166

Ref., reference category.
For these analyses, n 771 due to missing values on items.

Discussion

At nearly 27 %, the prevalence of food insecurity in our sample of veterans who served in Iraq or Afghanistan was dramatically higher than the US prevalence of food insecurity (14.5 % in 2012⁽¹⁰⁾). Further, veterans reported very low food security at double the US rate (12.1 % *v.* 5.7 %).

To give context to this issue, we described food security by demographic and health characteristics. Factors generally associated with lower socio-economic status such as not being married/partnered and having a lower income were associated with food insecurity. However, neither the number of deployments nor having a service-connected disability was associated with reporting difficulty in accessing food, which would suggest that greater exposure to combat is not what links certain veterans to increased risk of being food insecure. In the bivariable models in Table 1, those who were food insecure were more likely to use tobacco, binge drink more frequently, sleep less and have poorer self-reported general health. The co-occurrence of food insecurity with these health behaviours suggests that food-insecure veterans face multiple serious threats to their well-being. Digging deeper, in the multivariable model, six variables retained their independent association with food security while mutually adjusted for each other, which may indicate that these factors play a role in setting veterans' food security trajectory. However, the study does not provide information on when various factors may have developed during each participant's timeline, and thus both whether there is causality between the various factors and food security and what direction it might go in are unclear. There are plausible pathways by which some of these

behavioural factors may be more than simply correlated with food insecurity. For instance, sleep issues may interfere with a veteran's ability to work and earn income, which in turn means there will be fewer resources for food. Alternatively the stress of worrying about obtaining food could be a reason why food-insecure veterans report less sleep. Spending on tobacco may deplete money that could be used to purchase food, thereby leading to food insecurity. To gain clarity on these issues, longitudinal research is needed. But what is known is that these issues cluster and can highlight who is at risk of hunger.

Several limitations of the current report deserve attention. First, while our response rate of 52 % was far higher than in the major survey studies of the new generation of veterans (for instance, the response rates of the baseline waves of the National Health Study for a New Generation of US Veterans and the Millennium Cohort Study were 34.3 %⁽¹⁷⁾ and 31 %⁽¹⁶⁾, respectively), there still may be important difference between survey responders and non-responders related to variables of interest. Indeed, we found differences in electronic medical record variables between responders and non-responders. However, these differences were in areas that were either not associated with food security (service connection status and gender) or these differences suggested that we might have underestimated the prevalence of food insecurity in the general Iraq and Afghanistan war veteran population as responders tended to be older and married/partnered, which are all factors associated with a lesser likelihood of food insecurity. Additional reasons why we might actually be under-reporting food insecurity are that Minnesota is a relatively economically prosperous⁽²²⁾ state and food

insecurity may be more common in non-white veterans whom our study under-represents. (In the general population of 7.9 million veterans in the VA health system, 80.8% are white⁽²³⁾, while 90.1% of the current sample was white.) Additionally, in 2011 the prevalence of food insecurity in Minnesota was reported to be 11.4% by one source⁽²⁴⁾ which was below the national average. Finally, while the US Household Food Security Module: Six Item Short Form has been demonstrated to be a valid tool for identifying households that have low or very low food security⁽¹⁹⁾, it has the disadvantage that unlike the eighteen-item US Household Food Security Survey Module, the six-item measure lumps households with 'marginal food security' into the food secure category. There is evidence that marginal food security may also be a risk factor for chronic disease^(6,10,25) and if so, it would have been advantageous for our report to be able to identify individuals in this category as well. An additional limitation is that food security is measured at the household level and many of our predictor variables, such as education, were asked just of the respondent and not of all household members. A strength of the study was use of the OEF/OIF/OND Roster, which completely enumerates those who served in Iraq and Afghanistan, as our sampling frame; this enhances the generalizability of our findings.

Future work should focus on connecting veterans with employment that can provide a liveable wage and food assistance for veterans in need. The USA is one of the wealthiest nations in the world⁽²⁶⁾ and was engaged in fighting two expensive wars for over a decade (with total costs estimated to be between \$US 4 and 6 trillion⁽²⁷⁾). In light of this, it is unacceptable that such a sizeable percentage of those who fought those wars struggle to afford food once they return home.

Acknowledgements

Acknowledgements: The authors would like to thank the many staff members at CCDOR who assisted with the implementation of the survey. *Financial support:* This material is the result of work supported with resources and the use of facilities at the Minneapolis VA Health Care System. Salary support for R.W. was provided by a VA Health Services Research and Development Career Development Award (CDA 09-012-2). The funder had no role in the design, analysis of writing of this article. The views expressed in this article are those of the authors and do not necessarily reflect the position or policy of the Department of Veterans Affairs or the US government. *Authorship:* R.W. took the lead in writing and conceptualized the study. A.J. worked on developing measures, editing drafts of the manuscript and recruitment. A.B. contributed to the study design and editing of the manuscript. S.S.F. assisted with study design and manuscript writing. *Ethics of human subject participation:* All procedures

were reviewed and approved by the Minneapolis VA institutional review board.

References

1. Iraq and Afghanistan Veterans of America (2011) IAVA – Model, Mission and History. <http://iava.org/about> (accessed October 2013).
2. Institute of Medicine (2013) *Returning Home from Iraq and Afghanistan: Assessment of Readjustment Needs of Veterans, Service Members, and Their Families*. Washington, DC: National Academies Press.
3. Parker ED, Widome R, Nettleton JA *et al.* (2010) Food security and metabolic syndrome in US adults and adolescents: findings from the National Health and Nutrition Examination Survey, 1999–2006. *Ann Epidemiol* **20**, 364–370.
4. Weiser SD, Young SL, Cohen CR *et al.* (2011) Conceptual framework for understanding the bidirectional links between food insecurity and HIV/AIDS. *Am J Clin Nutr* **94**, issue 6, 1729S–1739S.
5. Ivers LC & Cullen KA (2011) Food insecurity: special considerations for women. *Am J Clin Nutr* **94**, issue 6, 1740S–1744S.
6. Seligman HK, Bindman AB, Vittinghoff E *et al.* (2007) Food insecurity is associated with diabetes mellitus: results from the National Health Examination and Nutrition Examination Survey (NHANES) 1999–2002. *J Gen Intern Med* **22**, 1018–1023.
7. Martin KS & Ferris AM (2007) Food insecurity and gender are risk factors for obesity. *J Nutr Education Behav* **39**, 31–36.
8. Wilde PE & Peterman JN (2006) Individual weight change is associated with household food security status. *J Nutr* **136**, 1395–1400.
9. Laraia BA (2013) Food insecurity and chronic disease. *Adv Nutr* **4**, 203–212.
10. Coleman-Jensen A, Nord M & Singh A (2013) *Household Food Security in the United States in 2012. Economic Research Report* no. 155. Washington, DC: US Department of Agriculture, Economic Research Service.
11. VA Information Resource Center – VIREC (2014) OEF/OIF/OND Roster. <http://www.virec.research.va.gov/OEF-OIF-OND-Roster/Overview.htm> (accessed April 2014).
12. Dillman DA, Smyth JD & Christian LM (2009) *Internet, Mail, and Mixed-Mode Surveys: The Tailored Design Method*. Hoboken, NJ: John Wiley & Sons, Inc.
13. Elbogen EB, Sullivan CP, Wolfe J *et al.* (2013) Homelessness and money mismanagement in Iraq and Afghanistan veterans. *Am J Public Health* **103**, Suppl. 2, S248–S254.
14. Vogt D, Vaughn R, Glickman ME *et al.* (2011) Gender differences in combat-related stressors and their association with postdeployment mental health in a nationally representative sample of US OEF/OIF veterans. *J Abnorm Psychol* **120**, 797–806.
15. Beckham JC, Becker ME, Hamlett-Berry KW *et al.* (2008) Preliminary findings from a clinical demonstration project for veterans returning from Iraq or Afghanistan. *Mil Med* **173**, 448–451.
16. LeardMann CA, Powell TM, Smith TC *et al.* (2013) Risk factors associated with suicide in current and former US military personnel. *JAMA* **310**, 496–506.
17. Eber S, Barth S, Kang H *et al.* (2013) The National Health Study for a New Generation of United States Veterans: methods for a large-scale study on the health of recent veterans. *Mil Med* **178**, 966–969.
18. US Department of Agriculture (2008) *US Household Food Security Survey Module: Six-Item Short Form*. Washington, DC: US Department of Agriculture.

19. Blumberg SJ, Bialostosky K, Hamilton WL *et al.* (1999) The effectiveness of a short form of the Household Food Security Scale. *Am J Public Health* **89**, 1231–1234.
20. US Department of Agriculture, Economic Research Service (2013) Definitions of Food Security. <http://www.ers.usda.gov/topics/food-nutrition-assistance/food-security-in-the-us/definitions-of-food-security.aspx#.UwzMsyRRGr8> (accessed February 2014).
21. Craig CL, Marshall AL, Sjöström M *et al.* (2003) International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc* **35**, 1381–1395.
22. US Department of Commerce, Bureau of Economic Analysis (2013) Widespread economic growth in 2012. http://bea.gov/newsreleases/regional/gdp_state/gsp_newsrelease.htm (accessed October 2013).
23. US Department of Veterans Affairs (2012) *2011 Survey of Veteran Enrollees' Health and Reliance Upon VA*. Washington, DC: Department of Veterans Affairs, Veterans Health Administration, Office of the Assistant Deputy Under Secretary for Health for Policy and Planning.
24. Feeding America (2014) Map the Meal Gap. Food Insecurity in your County. <http://feedingamerica.org/hunger-in-america/hunger-studies/map-the-meal-gap.aspx> (accessed February 2014).
25. Cook JT, Black M, Chilton M *et al.* (2013) Are food insecurity's health impacts underestimated in the US population? Marginal food security also predicts adverse health outcomes in young US children and mothers. *Adv Nutr* **4**, 51–61.
26. Central Intelligence Agency (2013) *The World Factbook 2013–14*. Washington, DC: CIA; available at <https://www.cia.gov/library/publications/the-world-factbook/index.html>.
27. Bilmes LJ (2013) *The Financial Legacy of Iraq and Afghanistan: How Wartime Spending Decisions Will Constrain Future National Security Budgets*. Cambridge, MA: Harvard Kennedy School of Government.