

Consumption of fruits and vegetables among adolescents: a multi-national comparison of eleven countries in the Eastern Mediterranean Region

M. F. Al Ani¹, L. K. Al Subhi^{1*} and S. Bose²

¹Department of Food Science and Human Nutrition, College of Agricultural and Marine Sciences, Sultan Qaboos University, PO Box 34, Al Kboud, Muscat 123, Oman

²Department of Natural Resource Economics, College of Agricultural and Marine Sciences, Sultan Qaboos University, Oman, PO Box 34, Al Kboud, Muscat 123, Oman

(Submitted 17 March 2015 – Final revision received 31 July 2015 – Accepted 25 August 2015 – First published online 28 January 2016)

Abstract

Regional cross-country profile of fruit and vegetable (F&V) consumption is lacking in the Eastern Mediterranean Region (EMR). This study examines the prevalence and differences of consuming F&V ≥ 5 times/d among adolescents in eleven EMR countries, and also describes differences in the proportions of taking F&V ≥ 5 times/d by sex, age and BMI. The study included 26 328 school adolescents (13–15 years) with complete data on consumption of F&V, age, sex, weight and height taken from the Global School-based Student Health Survey conducted in the EMR between 2005 and 2009. Overall, only 19.4% of adolescents reported consuming F&V ≥ 5 times/d. The highest prevalence was reported in Djibouti (40.4%) and the lowest was reported in Pakistan (10.0%). Statistically significant differences in prevalence were observed across countries ($P < 0.05$). With the exception of Oman, Libya and Djibouti, significantly more males than females ate F&V ≥ 5 times/d. Proportion of students consuming F&V ≥ 5 times/d also varied significantly in all countries based on BMI ($P < 0.0001$), with students within normal BMI having the highest frequency. A negative trend was observed between age and the prevalence of taking F&V ≥ 5 times/d in most of the eleven EMR countries but Jordan, Djibouti and Morocco. The prevalence of adequate intake of F&V was low in the eleven EMR countries. There is a need for interventions to increase the prevalence of adolescents consuming F&V ≥ 5 times/d. Interventions should take into consideration psychosocial, environmental and socio-environmental factors influencing F&V intake within countries.

Key words: Adolescents: Cross-country comparison: Fruit and vegetable consumption: Eastern Mediterranean Region: Global School-based Student Health Survey

Fruits and vegetables (F&V) are essential components of a healthy diet in meeting requirements for growth, development and maintenance throughout the life cycle. Adequate daily intake of the F&V has been associated with reduced risks of CVD^(1–3), stroke⁽⁴⁾, type 2 diabetes⁽⁵⁾ and certain types of cancer^(6,7), all of which are major causes of mortality and morbidity in the Eastern Mediterranean Region (EMR)⁽⁸⁾. In addition, increased intake of F&V promotes satiety, decreases energy intakes and has been associated with better weight-related outcomes and successful weight-loss treatment for adolescents⁽⁹⁾.

The 2002 Joint FAO/WHO Expert Consultation on Diet, Nutrition and the Prevention of Chronic Diseases recommends a minimum of 400 g/d of F&V, an equivalent of ≥ 5 servings of F&V/d. The recommendation excludes potatoes and other starchy tubers⁽¹⁰⁾. As the protective role of F&V on non-communicable diseases (NCD) became well established, the World Health Organization⁽¹¹⁾ adopted an initiative in 2003 to promote their consumption and called to standardise their

definitions to facilitate multi-national comparison. In 2004, the World Health Assembly adopted the Global Strategy on Diet, Physical Activity and Health, which aims to promote F&V consumption around the world and assists member states in implementing national strategies emphasising in improving diet to reduce diseases and deaths associated with them⁽¹²⁾.

Although there is a trend for an increased global production of fifteen types of fruits and fifteen types of vegetables between 1998 and 2003⁽¹³⁾ and despite the evident benefits of adequate intake of F&V, multi-national Western dietary surveys indicate that most adolescents are not consuming the recommended amounts of F&V^(14–17). Longitudinal trends show that US adolescents decreased their daily intake of F&V along the spectrum of adolescence⁽¹⁸⁾. Only 9 and 14% of Chinese adolescents (n 2977) consumed the minimum recommended daily intakes of F&V⁽¹⁹⁾. Among Ghanaian adolescents, 56% (n 1195) rarely ate fruits and 48% rarely ate vegetables⁽²⁰⁾. Global School-based Student Health Survey (GSHS) data from seven developing African countries revealed that 77.5% of

Abbreviations: EMR, Eastern Mediterranean Region; F&V, fruits and vegetables; GSHS, Global School-based Student Health Survey; NCD, non-communicable diseases.

* **Corresponding author:** L. K. Al Subhi, fax +968 2441 3418, email lyraju@hotmail.com

adolescents (n 17 656) did not meet the daily recommended intake of F&V⁽²¹⁾.

To the best of the authors' knowledge, there is limited information available on the prevalence of F&V consumption among adolescents in EMR countries. EMR includes twenty-two diverse countries from two continents – Asia and Africa. Differences in risk behaviours such as inadequate intake of F&V have not been studied in the region. In addition, potential factors associated with F&V consumption are not well researched in the region.

The objectives of this study were as follows: (1) to estimate differences in the proportion of 13–15-year-old adolescents who reported consuming F&V ≥ 5 times/d across eleven EMR countries based on GSHS data, (2) to explore potential within-country differences in proportions of students taking F&V ≥ 5 times/d based on sex, age and BMI groups of the study sample.

Methods

Study data

The present study is based on a large cross-section of school students from eleven countries in EMR that participated in the GSHS between 2005 and 2009. The GSHS is a collaborative surveillance project coordinated by the WHO with the goal of helping countries measure and assess behavioural risk factors and protective factors among students aged 13–17 years^(22,23).

The GSHS is based on voluntary participation of nations. A two-staged cluster sampling standardised process is applied to produce representative country samples. First, schools are selected with probability proportional to sample size. Second, classes within schools are randomly selected. All students in the sampled classes are eligible to participate irrespective of age. Data are collected using a self-administered questionnaire. The questionnaire consists of ten sets of core questions that allow cross-country comparison, core-expanded questions and country-specific questions. Core sets address leading causes of morbidity and mortality among children and adults worldwide. One of the core sets is 'dietary behaviours', which is the source of data for the present study. Questionnaires were translated into appropriate languages and pilot tested for comprehension. Students in selected classes completed the survey during one regular period using computer-scannable answer sheets. Additional details on the GSHS methodology and study design are described elsewhere^(22,23).

By the end of January 2012, there were only eleven publically available country data sets (twenty-three) from the EMR countries that undertook the GSHS, and thus were included in this study. One country – Jordan – had data sets for 2 years, 2004 and 2007, and only the latest set was included to minimise the gaps of years among countries. The remaining countries had only one data set. The collective sample from the eleven countries was 47 256 school students. A stepwise exclusion process was applied. Observations with missing response to the core questions of fruit or vegetable consumption were excluded. Subsequently, students with missing age, sex, weight or height were excluded as they are all required to calculate the BMI. Finally, students aged <13 and >15 years were excluded, as the objective of the study was to describe the F&V consumption patterns of schoolchildren aged between 13 and

15 years. As a result, the valid study sample included 26 328 adolescents, 55.7% of the initial sample.

BMI (weight (kg)/height (m²)) was calculated using the self-reported weight and height information. Age- and sex-specific z scores for BMI of the study sample were calculated using the recommended 2007 least median of squares (LMS) method. BMI was classified based on the z score cut-off values, with $>+1$ SD being overweight, $+1$ to -2 SD being normal weight and <-2 SD being underweight^(24,25). Severe underweight and obesity were clustered with underweight and overweight, respectively.

Total country population and population of the adolescents aged 13–15 years, by GSHS year of survey, were obtained from the World Bank. Annual production, export and import of F&V for the respective years of the GSHS were obtained using the agriculture item groups 'fruit excl melons, total' and 'vegetables and melons, total' from FAOSTAT at (<http://faostat.fao.org/site/291/default.aspx>). The data were used to calculate the per capita availability of F&V of each country for the year of GSHS (Table 3).

Assessing consumption of fruits and vegetables

The dietary behaviours core module in version 2003–2008 of the GSHS has five questions and that of version 2009–2012 has seven questions. Both versions have two identical questions on F&V consumption^(22,23) where respondents were asked the following:

- During the past 30 d, how many times per day did you usually eat fruits?, and
- During the past 30 d, how many times per day did you usually eat vegetables?

Both questions were based on six response categories ranging from I did not eat, <1 time/d, 1 time/d, 2 times/d, 3 times/d, 4 times/d to ≥ 5 times/d. Country-specific examples of F&V were referenced in both questions⁽¹⁰⁾.

The recommended daily intake of F&V based on the Joint FAO/WHO Expert Consultation on Diet, Nutrition and Prevention of Chronic Diseases is a minimum of 400 g of F&V/d (excluding potatoes and other starchy tubers); an equivalent of ≥ 5 servings/d. However, the two questions in the GSHS noted above only provide frequency of intake rather than servings per day. The per day frequency of intake of F&V does not represent the actual servings, whereas the term serving/day implies both frequency and quantity of intake. For the purpose of this study, and because the GSHS data only provide frequency of daily intake of F&V with no reference to portion sizes, consumption of F&V ≥ 5 times/d was used as a general and proxy cut-off measure for adequate intake of F&V.

Statistical analyses

The χ^2 statistic was used to investigate whether the sample proportions with reference to 'adequate F&V consumption among adolescents' (≥ 5 servings/d) differ significantly across countries. A P value of <0.05 was considered statistically significant. Next, the Marascuilo procedure⁽²⁶⁾ was used to identify the proportions that differ significantly by comparing all possible pairs of proportions, and the results are presented in Table 4. Moreover, χ^2 statistic was used to investigate within-country differences in the proportion of adolescents consuming

F&V ≥ 5 times/d based on age, sex and BMI groups (Table 5). Statistical analyses were carried out using Microsoft Excel and SPSS 19.

Results

The overall initial response rate of the GSHS in the eleven EMR countries was 87.9%. The largest response rate was from Jordan (99.8%) and the lowest was from Pakistan (76%). The final valid study sample after all exclusions was 26 328 (55.7%) of the total initial sample (Table 1). The characteristics of the study population before and after exclusion are shown in Table 2. There were more male (55%) than female adolescents in the valid sample. The majority of students (66.9%) fall within the normal weight BMI. On the basis of the gross national incomes (GNI) of the year of data collection, seven of the countries were

Table 1. Global School-based Student Health Survey year of data collection, initial and valid sample sizes of eleven Eastern Mediterranean Region countries (Numbers and percentages)

Countries	Year of data collection	Initial sample	Response rate (%)	Valid sample	
				<i>n</i>	%
UAE	2005	15 790	89	9723	61.6
Oman	2005	2979	97	653	21.9
Lebanon	2005	5115	88	2423	47.4
Libya	2007	2242	98	1326	59.1
Jordan	2007	2197	99.8	1023	46.6
Djibouti	2007	1777	83	842	47.4
Egypt	2006	5249	87	3227	61.5
Morocco	2006	2670	84	833	31.2
Tunisia	2008	2870	83	974	33.9
Yemen	2008	1175	82	582	49.5
Pakistan	2009	5192	76	4722	91.0
Total		47 256	87.9	26 328	55.7

Table 2. Characteristics of the study sample before and after exclusion* (Numbers and percentages)

Characteristics	Before exclusion (<i>n</i>)	After exclusion (<i>n</i>)	%
Sex			
Male	24 648	14 489	55
Female	22 114	11 839	45
Missing	494		
Age (years)			
11	798		
12	5279		
13	11 287	8534	32.4
14	12 645	9664	36.7
15	10 478	8130	30.9
16+	6149		
Missing	620		
BMI (<i>z</i> scores)			
Underweight (< -2 sd)	–	1484	5.6
Normal weight (+1 to -2 sd)	–	17 600	66.9
Overweight (> +1 sd)	–	7244	27.5

* *n* before elimination for the BMI was not calculated as the calculation for the BMI according to the (*z* scores) method relies on critical points of sex and age. The exclusion of missed data on sex or age took place before BMI was calculated.

classified in either low- or lower-middle income category. Across the pooled sample, only 19.4% of adolescents reported consuming F&V ≥ 5 times/d. Djibouti as a lower middle-income country, with a low per capita availability of F&V (0.094 tonnes/year), had the highest reported prevalence of adolescents (40.4%, *n* 842) reported eating F&V ≥ 5 times/d. Pakistan had the lowest per capita availability of F&V (0.068) as well as the lowest prevalence of adolescents consuming F&V ≥ 5 times/d (10.0%, *n* 4722). It is interesting to note that Pakistan ranked second, after Egypt, in producing (1 192 330 tonnes) and third in exporting F&V (912 873 tonnes); nevertheless, it ranked first in F&V imports and had the lowest per capita availability of F&V. UAE, the only high-income country that ranked second after Pakistan in imports of F&V, had the highest per capita availability (0.409) (Table 3).

There were a number of cross-country statistically significant differences in the prevalence of adolescents reporting consuming F&V ≥ 5 times/d. For instance, the prevalence in the UAE was significantly lower ($P < 0.0001$) than that of Oman, Lebanon, Jordan, Djibouti, Morocco and Tunisia. On the other hand, it was significantly higher ($P < 0.0001$) than that of Libya and Pakistan. Pakistan had significantly ($P < 0.0001$) lower prevalence (10.0%) of adolescents consuming F&V ≥ 5 times/d compared with all countries except Libya (Table 4).

Comparisons of frequencies within age, sex and BMI categories are presented in Table 5. Overall, the proportion of students taking F&V ≥ 5 times/d tended to decrease as age increased; more 13-year-old (*n* 1806) and 14-year-old (*n* 1836) adolescents reported consuming F&V ≥ 5 times/d than 15-year-old students (*n* 1470). When countries were examined separately, a similar trend was found in seven of the countries. However, in the case of Jordan, Djibouti and Morocco, the proportions of teens consuming F&V ≥ 5 times/d increased with age, whereas it appeared to fluctuate in Pakistan with more 14-year-old adolescents (*n* 215) meeting the cut-off intake frequency for F&V. For most of the countries, there are significant differences ($P < 0.0001$) in the proportion of adolescents consuming F&V ≥ 5 times/d in the three age groups. Exceptions were seen in four countries – Oman, Libya, Morocco and Yemen – where the frequencies among the three age groups were not significant at $P < 5\%$. As for the sex factor, the total results show that more males (*n* 2972) than females (*n* 2140) reported intake of F&V ≥ 5 times/d; all countries showed similar trends except for Libya. The differences in the proportions between males and females were significant ($P < 0.04$ – $P < 0.0001$) within all counties but three – Oman, Libya and Djibouti. With respect to BMI, more adolescents with normal BMI (*n* 3346) than underweight (*n* 314) and overweight (*n* 1452) adolescents met the cut-off for adequate intake of F&V. The observed pattern was consistent and significant within all countries ($P < 0.0001$).

Discussion

It should be noted that the study only examined the prevalence of adequate consumption of F&V among adolescents using a proxy approach based on the frequency of intake per day. The GSHS data on F&V consumption is based on reported daily

Table 5. Within-country differences in proportions of fruit and vegetable consumption by age, sex and BMI

Countries	Adolescents consumed fruits and vegetables ≥ 5 times/d											
	Age (years)			Sex			BMI (kg/m ²)			χ^2 test	P	
	13	14	15	Male	Female	Underweight	Normal	Overweight				
	Count	Count	Count	Count	Count	Count	Count	Count				
UAE	661	591	480	961	771	51	988	693	795.13	<0.0001		
Oman	71	63	56	97	93	0.04	122	47	86.85	<0.0001		
Lebanon	227	227	162	355	261	14.04	430	158	409.88	<0.0001		
Libya	55	49	40	61	83	3.06	95	39	77.79	<0.0001		
Jordan	32	118	127	176	101	19.76	186	68	153.49	<0.0001		
Djibouti	40	119	181	186	154	2.82	239	64	212.23	<0.0001		
Egypt	390	180	40	338	272	6.92	374	204	287.62	<0.0001		
Morocco	102	106	121	184	145	4.38	230	66	203.02	<0.0001		
Tunisia	102	138	77	193	124	14.58	226	67	214.3	<0.0001		
Yemen	30	30	23	56	27	9.44	62	11	63.93	<0.0001		
Pakistan	96	215	163	365	109	137.18	394	35	529.08	<0.0001		
Total	1806	1836	1470	2972	2140	—	3346	1452	—	—		

frequency of intake with no reference to portion sizes. Thus, meeting the criteria of consuming F&V ≥ 5 times/d is not equivalent to consuming F&V ≥ 5 servings/d. Moreover, the GSHS data are obtained by means of a self-reported questionnaire – a method that is subject to recall errors and under/over-reporting⁽²⁷⁾. Nevertheless, the study is based on a large sample size from the GSHS that has a sound sampling process. Moreover, it included eleven EMR countries with diverse GNI and variable per capita availability of F&V.

The observed lack of trend in the relationship between adequate F&V consumption among adolescents and the GNI of the eleven EMR countries is not consistent with that reported by others among adults. On the basis of the Prospective Urban Rural Epidemiology study, a positive trend between healthy eating (including intake of higher amounts of F&V) and the gross domestic product (GDP) of countries has been noted; consumption of F&V increased with increased GDP⁽²⁸⁾. In addition, the per capita availability of F&V in the eleven countries was not seen to be in synchrony with the patterns of the prevalence of consumption. Such an observation could be affected by a lack of infrastructure of food supply chains of highly perishable commodities⁽²⁹⁾. It is strategically important to understand the inter-related aspects of accessibility and affordability and other contributing factors with consumption of F&V in these countries⁽³⁰⁾. EMR countries could explore available opportunities to expand availability of F&V presented by Aid for Trade that has a built-in venue to increase F&V supply⁽³¹⁾.

As noted, only 19.4% of adolescents in the eleven EMR countries met the cut-off for F&V consumption. These findings cause concern about the inadequacy of F&V intake among adolescents in light of the association between F&V consumption and NCD diseases (1–7). Overall, the prevalence of taking F&V ≥ 5 times/d in the eleven EMR countries is comparable with that reported in other nations. Using the same cut-off for adequate F&V, only 22.3% of American adolescents in the YRBS-2009 study consumed an adequate amount of F&V per day⁽³²⁾. In Brazil, only 23% of the teens consumed 5 F&V/d⁽³³⁾.

Intake of F&V in other countries within EMR is even lower. Although not included in the present study, Kuwaiti male and female adolescents were reported to consume vegetables (3.8 *v.* 3.5 times/week) and fruits (3.4 *v.* 2.8 times/week), respectively. The prevalence of F&V intake of ≥ 7 times/week was only reported in 26 *v.* 22.1% for vegetables and 17.5 *v.* 11.8% for fruits for males and females, respectively⁽³⁴⁾. The weekly mean frequency is extremely low, indicating a lower daily and therefore inadequate intake in this group. It is interesting to note that the intake frequency of vegetables is much higher than that of fruits in Kuwaiti adolescents. Such an observation could be attributed to not excluding potatoes in such types of studies. In Australian adolescents, the prevalence of adequate intake of both F&V decreased substantially when juices and potatoes were excluded⁽³⁵⁾. Variations in assessment approaches should be accounted for when comparing results.

The propensity for more males reporting intake of F&V ≥ 5 times/d seen in the present study is also similar to that of adolescents in the USA, where more males (23.9%) consumed adequate F&V compared with females (20.5%) in the Youth

Risk Behavior Survey (YRBS)-2009 study⁽³²⁾. Australian male adolescents had slightly higher intake than females (based on $\geq 1-3$ and $\geq 2-4$ servings/d of F&V, respectively)⁽³⁵⁾. Kuwaiti male teens were also reported to have significantly higher mean weekly intake frequency of vegetables ($P=0.02$) and fruits ($P=0.001$) than female teens⁽³⁴⁾. Nevertheless, female adolescents in other nations including Palestine⁽³⁶⁾, USA⁽¹⁴⁾ and Norway⁽³⁷⁾ were reported to be more prevalent in consuming F&V with different criteria for assessing adequate F&V intake.

Decreased intake with increased age seen in the eleven EMR countries is also reported among adolescents in other areas including the USA^(18,32) and Australia⁽³⁵⁾. The adverse relationship between F&V intake and age has been shown to start from early adolescence⁽³⁸⁾ and may persist into late adolescence and adulthood^(1,14,39). Several factors could be behind the decreased tendency of consuming adequate F&V in this cycle of life. As age increases, there is a lower preference towards F&V and a lower frequency of family dinners⁽⁴⁰⁾. The observed trend with BMI in this regional study is also in line with that seen among seventh-grade adolescents from the Teens Eating for Energy and Nutrition at School (TEENS) study and might be related to perceptions about weight. Boys and girls perceived as underweight significantly ate fewer servings of F&V per day than those perceived to be in the normal weight group. In addition, boys and girls perceived to be overweight consumed significantly fewer servings of F&V per day than those having normal weight⁽⁹⁾. Among 11-year-old Greek children, BMI did not associate with F&V intake significantly⁽⁴¹⁾. Similar to what is noted in the present study was reported in a large sample (n 246 995) of middle-aged Australian men and women. Compared with the overweight group, normal weight men were more likely to be in the highest vegetables, fruits and combined F&V intake quartiles and were less likely to have 5 servings of F&V/d⁽⁴²⁾. An increased intake of F&V may promote BMI reduction among adolescents⁽⁴³⁾ and may be an effective dietary strategy to control weight and reduce the risk of obesity. However, a study on the association between F&V intake and a change in BMI did not show a beneficial effect, and thus it was suggested that F&V recommendations should not be based on a beneficial effect on weight regulation⁽⁴⁴⁾.

Aside from age, sex and BMI, correlates of F&V intake among adolescent might be variable and inter-related. Less is known about the socio-economic, cultural, psychosocial and environmental determinants of the consumption of F&V among adolescents in the EMR. Findings from the project Eating Among Teens (EAT) interdicted that home availability explained 45% and test preference explained 28% of the variances in F&V intake among adolescents⁽⁴⁵⁾. Self-efficacy, parent and peer modelling, family and peer normative beliefs and social and health outcome expectations were found to be correlated with F&V intake among adolescents⁽⁴⁶⁾, whereas the socio-environmental factors related to school environment were availability and accessibility of F&V, nutrition education and food safety^(14,47). Such factors have not been evaluated, and further studies are warranted to explore the inter-association among them and intake of F&V in the EMR countries.

Interpretation of results from studies assessing adequate intake of F&V needs to be carried out in a careful manner. It is clear that consumption of French fries and juices are popular

among adolescents, and could inflate the estimated reported intake. The mean vegetable servings for American female adolescents decreased from 2 (SD 1.6) to 1.8 (SD 1.6) servings/d and that for male adolescents dropped from 1.8 (SD 1.6) to 1.6 (SD 1.5) servings/d when French fries were excluded⁽¹⁴⁾. The proportion of Australian children aged 9–16 years consuming $\geq 2-4$ servings/d of vegetables decreased from 19% to only 2% when potatoes were excluded. Likewise, the proportion of children reported consuming $\geq 1-3$ servings/d of fruits declined from 90 to 51% after exclusion of fruit juices among 9–13-year-old children and declined from 24 to 1% among 14–16-year-old adolescents⁽³⁵⁾. It is worth noting that the definition of fruit juice is very much distorted among the general public and might not mean 100% juice but fruit drinks, with mass processed products falling under juices. Thus, although a product might only contain no more than 30% juice concentrate by volume, the public perceives such products as fruit juices. Therefore, studies should carefully consider these aspects that might inflate the actual intake of F&V.

Adolescence is characterised by rapid growth and increased nutrient needs; with that, it is an important time to intervene with F&V intake to meet nutrient needs and to develop dietary patterns that may persist into adulthood. Efforts to increase F&V intake through promotion initiatives have been proved effective^(48,49). The WHO noted insufficient policy response and progress in curbing the burden of NCD; it has been and is calling countries to increase action against NCD risk factors, including unhealthy diet, to reduce the preventable and avoidable burden of morbidity, mortality and disability due to NCD. A multi-sectoral collaboration and cooperation at national, regional and global levels is called by WHO for populations to reach the highest attainable standards of health and productivity at every age⁽⁵⁰⁾.

Availability of policy documents for national NCD policies, programmes and action plans in low- and middle-income WHO member countries (n 140) is not clear. Only 47% of such countries were found to have policy documents, but only a small fraction of them proposed actions to promote healthier diets and physical activity. National policy actions to improve F&V consumption were particularly low in low- and middle-income EMR countries; only three countries were reported to have such documents that targeted the general public. Djibouti, which has the highest prevalence of adolescents (40.4%) taking F&V ≥ 5 times/d, has a document on a national policy action to promote family plantations and consumption of F&V. Jordan, ranked fifth among the eleven EMR countries in the prevalence (27.1%) of F&V intake, was found to have a national policy action to organise campaigns emphasising the importance of F&V. Policy action in Iran was to address poor dietary habits and promote consumption of F&V⁽³¹⁾. The impact of such action plans has to be studied to understand their effectiveness, to share information and to establish databases on best practices on national and regional action plans in promoting adequate intake of F&V. Other EMR countries should adapt policies aiming to increase F&V intake and prioritise action plans along such policies if a reduction of the low prevalence of F&V intake is desired to reduce the impact of such a modifiable risk factor associated with NCD.



The great majority of adolescents in the eleven EMR counties did not meet the adequate F&V consumption based on the proxy of eating F&V ≥ 5 times/d. Interventions in the EMR countries should be prioritised to increase the intake of F&V among adolescents. Adolescents are swamped with palatable low-dense nutrient foods. Understanding limitations would be strategically important if governments are to intervene in order to increase the prevalence of adequate F&V intake. Interventions should take into consideration the psychosocial, environmental and socio-environmental factors influencing F&V intake within countries. Further research is needed to identify effective interventions and policy approaches geared towards improving the availability and accessibility of F&V as well as increasing their consumption in EMR countries.

Acknowledgements

The authors thank all the reviewers who have contributed to shape this article.

This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

Authors of this study have contributed as follows – M. F. A. A. and L. K. A. S.: formulated the research questions. M. F. A. A.: conception of the idea, data extraction and coding and contributed to writing the initial draft. L. K. A. S.: guided data processing, and wrote the final version of the article. S. B.: data analyses, wrote the methods and critically reviewed the article. The authors declare that there are no conflicts of interest.

References

1. Mikkila V, Rasanen L, Raitakari OT, *et al.* (2005) Consistent dietary patterns identified from childhood to adulthood: the cardiovascular risk in young finns study. *Br J Nutr* **93**, 923–931.
2. Mirmiran P, Noori N, Zavareh MB, *et al.* (2009) Fruit and vegetable consumption and risk factors for cardiovascular disease. *Metabolism* **58**, 460–468.
3. Dauchet L, Amouyel P, Hercberg S, *et al.* (2006) Fruit and vegetable consumption and risk of coronary heart disease: a meta-analysis of cohort studies. *J Nutr* **136**, 2588–2593.
4. Bazzano LA, He J, Ogden LG, *et al.* (2002) Fruit and vegetable intake and risk of cardiovascular disease in US adults: the first national health and nutrition examination survey epidemiologic follow-up study. *Am J Clin Nutr* **76**, 93–99.
5. Carter P, Gray LJ, Troughton J, *et al.* (2010) Fruit and vegetable intake and incidence of type 2 diabetes mellitus: systematic review and meta-analysis. *BMJ* **341**, c4229.
6. Maynard M, Gunnell D, Emmett P, *et al.* (2003) Fruit, vegetables, and antioxidants in childhood and risk of adult cancer: the Boyd Orr cohort. *J Epidemiol Community Health* **57**, 218–225.
7. Papas MA, Giovannucci E & Platz EA (2004) Fiber from fruit and colorectal neoplasia. *Cancer Epidemiol Biomarkers Prev* **13**, 1267–1270.
8. Brug J, Tak NI, Te Velde SJ, *et al.* (2008) Taste preferences, liking and other factors related to fruit and vegetable intakes among schoolchildren: results from observational studies. *Br J Nutr* **29**, Suppl. 1, S7–S14.
9. Nystrom AA, Schmitz KH, Perry CL, *et al.* (2005) The relationship of weight-related perceptions, goals, and

- behaviors with fruit and vegetable consumption in young adolescents. *Prev Med* **40**, 203–208.
10. World Health Organization (2003) *Diet, Nutrition and the Prevention of Chronic Diseases*. WHO Technical Report Series. Geneva: WHO. <http://www.who.int/dietphysicalactivity/publications/trs916/en/> (accessed November 2014).
11. World Health Organization (2003) *Fruit and Vegetable Promotion Initiative/A Meeting Report/25-27/08/03*. Geneva: WHO. http://www.who.int/dietphysicalactivity/publications/f&v_promotion_initiative_report.pdf (accessed November 2014).
12. World Health Organization (2013) *Global Strategy on Diet, Physical Activity and Health: Diet and Physical Activity: A Public Health Priority*. Geneva: WHO. <http://www.who.int/dietphysicalactivity/en/> (accessed June 2013).
13. World Health Organization & Food and Agriculture Organization (2004) *Fruit and Vegetables for Health: Report of a Joint FAO/WHO Workshop, 1–3 September 2004, Kobe, Japan*. Geneva: WHO and FAO; www.fao.org/ag/magazine/FAO-WHO-FV.pdf (accessed November 2014).
14. Neumark-Sztainer D, Story M, Hannan PJ, *et al.* (2002) Overweight status and eating patterns among adolescents: where do youths stand in comparison with the healthy people 2010 objectives? *Am J Public Health* **92**, 844–851.
15. Centers for Disease Control and Prevention (2010) Youth risk behavior surveillance – United States, 2009. *MMWR Surveill Summ* **59**, 1–148.
16. Guenther PM, Dodd KW, Reedy J, *et al.* (2006) Most Americans eat much less than recommended amounts of fruits and vegetables. *J Am Diet Assoc* **106**, 1371–1379.
17. Vereecken CA, De Henauw S & Maes L (2005) Adolescents' food habits: results of the health behaviour in school-aged children survey. *Br J Nutr* **94**, 423–431.
18. Larson NI, Neumark-Sztainer D, Hannan PJ, *et al.* (2007) Trends in adolescent fruit and vegetable consumption, 1999–2004: project EAT. *Am J Prev Med* **32**, 147–150.
19. Zhang C-X, Chen Y-M, Chen W-Q, *et al.* (2012) Food group intake among adolescents in Guangzhou city compared with the Chinese dietary guidelines. *Asia Pac J Clin Nutr* **21**, 450–456.
20. Doku D, Koivusilta L, Raisamo S, *et al.* (2011) Socio-economic differences in adolescents' breakfast eating, fruit and vegetable consumption and physical activity in Ghana. *Public Health Nutr* **16**, 864–869.
21. Peltzer K & Pengpid S (2010) Fruits and vegetables consumption and associated factors among in-school adolescents in seven African countries. *Int J Public Health* **55**, 669–678.
22. Centers for Disease Control and Prevention (2012) Global School-based Student Health Survey (GSHS). Atlanta, GA: United States Centers for Disease Control. www.cdc.gov/gshs (accessed August 2012).
23. World Health Organization (2012) Global School-based Student Health Survey (GSHS). <http://www.who.int/chp/gshs/en> (accessed August 2012).
24. World Health Organization (2012) Computation of centiles and Z-scores for height-for-age, weight-for-age and BMI-for-age. <http://www.who.int/growthref/computation.pdf> (accessed August 2012).
25. World Health Organization (2012) Growth reference 5–19 years. http://www.who.int/growthref/who2007_bmi_for_age/en/index.html (accessed August 2012).
26. Marascuilo LA (1966) Large-sample multiple comparisons. *Psychol Bull* **65**, 280–290.
27. De Vriendt T, Huybrechts I, Ottevaere C, *et al.* (2009) Validity of self-reported weight and height of adolescents, its impact on classification into BMI-categories and the association with weighing behaviour. *Int J Environ Res Public Health* **6**, 2696–2711.

28. Teo K, Lear S, Islam S, *et al.* (2013) Prevalence of a healthy lifestyle among individuals with cardiovascular disease in high-, middle- and low-income countries: the Prospective Urban Rural Epidemiology (PURE) study. *JAMA* **309**, 1613–1621.
29. The Economist Intelligence Unit (2014) Special report: food loss and its intersection with food security. A report from The Economist Intelligence Unit. <http://foodsecurityindex.eiu.com/> (accessed November 2014).
30. Rahim HFA, Sibai A, Khader Y, *et al.* (2014) Non-communicable diseases in the Arab world. *Lancet* **383**, 356–367.
31. Lachat C, Otchere S, Roberfroid D, *et al.* (2013) Diet and physical activity for the prevention of noncommunicable diseases in low- and middle-income countries: a systematic policy review. *PLoS Med* **10**, e1001465.
32. Eaton DK, Kann L, Kinchen S, *et al.* (2010) Youth risk behavior surveillance—United States, 2009. *MMWR Surveill Summ* **59**, 1–142.
33. Rieth MA, Moreira MB, Fuchs FD, *et al.* (2012) Fruits and vegetables intake and characteristics associated among adolescents from Southern Brazil. *Nutr J* **11**, 1–7.
34. Allafi A, Al-Haifi AR, Al-Fayez MA, *et al.* (2014) Physical activity, sedentary behaviours and dietary habits among kuwaiti adolescents: gender differences. *Public Health Nutr*. **17**, 2045–2052.
35. Commonwealth Scientific Industrial Research Organisation, Preventative Health National Research Flagship & The University of South Australia (2007) *Australian National Children's Nutrition and Physical Activity Survey—Main Findings*. Canberra: Department of Health and Ageing, Astralina Food and Grocery Council, Department of Agriculture, Fisheries and Forestry. [http://www.health.gov.au/internet/main/publishing.nsf/Content/8F4516D5FAC0700ACA257BF0001E0109/\\$File/childrens-nut-phys-survey.pdf](http://www.health.gov.au/internet/main/publishing.nsf/Content/8F4516D5FAC0700ACA257BF0001E0109/$File/childrens-nut-phys-survey.pdf) (accessed February 2014).
36. Al Sabbah H, Vereecken C, Kolsteren P, *et al.* (2007) Food habits and physical activity patterns among Palestinian adolescents: findings from the national study of Palestinian schoolchildren (HBSC-WBG2004). *Public Health Nutr* **10**, 739–746.
37. Bere E, Brug J & Klepp K-I (2008) Why do boys eat less fruit and vegetables than girls? *Public Health Nutr* **11**, 321–325.
38. Di Noia J & Contento IR (2010) Fruit and vegetable availability enables adolescent consumption that exceeds national average. *Nutr Res* **30**, 396–402.
39. Hall JN, Moore S, Harper SB, *et al.* (2009) Global variability in fruit and vegetable consumption. *Am J Prev Med* **36**, 402–9 e5.
40. Granner ML, Sargent RG, Calderon KS, *et al.* (2004) Factors of fruit and vegetable intake by race, gender, and age among young adolescents. *J Nutr Educ Behav* **36**, 173–180.
41. Kouli E & Jago R (2008) Associations between self-reported fruit and vegetable consumption and home availability of fruit and vegetables among Greek primary-school children. *Public Health Nutr* **11**, 1142–1148.
42. Charlton K, Kowal P, Soriano MM, *et al.* (2014) Fruit and vegetable intake and body mass index in a large sample of middle-aged australian men and women. *Nutrients* **6**, 2305–2319.
43. Hart CN, Jelalian E, Raynor HA, *et al.* (2010) Early patterns of food intake in an adolescent weight loss trial as predictors of BMI change. *Eating Behav* **11**, 217–222.
44. Field AE, Gillman MW, Rosner B, *et al.* (2003) Association between fruit and vegetable intake and change in body mass index among a large sample of children and adolescents in the United States. *Int J Obes Relat Metab Disord* **27**, 821–826.
45. Neumark-Sztainer D, Wall M, Perry C, *et al.* (2003) Correlates of fruit and vegetable intake among adolescents: findings from project EAT. *Prev Med* **37**, 198–208.
46. Granner ML & Evans AE (2012) Measurement properties of psychosocial and environmental measures associated with fruit and vegetable intake among middle school adolescents. *J Nutr Educ Behav* **44**, 2–11.
47. Nago ES, Verstraeten R, Lachat CK, *et al.* (2011) Food safety is a key determinant of fruit and vegetable consumption in urban beninese adolescents. *J Nutr Educ Behav* **44**, 548–555.
48. Ammerman AS, Lindquist CH, Lohr KN, *et al.* (2002) The efficacy of behavioral interventions to modify dietary fat and fruit and vegetable intake: a review of the evidence. *Prev Med* **35**, 25–41.
49. Knai C, Pomerleau J, Lock K, *et al.* (2006) Getting children to eat more fruit and vegetables: a systematic review. *Prev Med* **42**, 85–95.
50. World Health Organization (2013) *Global Action Plan for the Prevention and Control of NCDs 2013–2020*. <http://www.who.int/nmh/publications/ncd-action-plan/en/>