

## Perspective

**Cite this article:** Blanco-Sacristán J, García FC, Johansen K, Maestre FT, Duarte CM and McCabe MF (2024). The Middle East as a natural laboratory to advance our understanding of global hyperarid drylands. *Cambridge Prisms: Drylands*, 1, e4, 1–9 <https://doi.org/10.1017/dry.2024.6>

Received: 28 March 2024

Revised: 20 October 2024

Accepted: 22 October 2024


**Keywords:**

drylands; hyper arid; conservation; restoration; biodiversity

**Corresponding author:**

Javier Blanco-Sacristán;  
Email: [javier.blancosacristan@kaust.edu.sa](mailto:javier.blancosacristan@kaust.edu.sa)

# The Middle East as a natural laboratory to advance our understanding of global hyperarid drylands

Javier Blanco-Sacristán<sup>1</sup> , Francisca C. García<sup>2</sup>, Kasper Johansen<sup>1</sup>, Fernando T. Maestre<sup>1</sup>, Carlos M. Duarte<sup>3</sup> and Matthew F. McCabe<sup>1</sup>

<sup>1</sup>Climate and Livability Initiative, Division of Biological and Environmental Sciences and Engineering, King Abdullah University of Science and Technology, Thuwal 23955-6900, Saudi Arabia; <sup>2</sup>Red Sea Research Center, Biological and Environmental Sciences and Engineering Division, King Abdullah University of Science and Technology, Thuwal 23955, Saudi Arabia and <sup>3</sup>Red Sea Research Center and Computational Bioscience Research Center, King Abdullah University of Science and Technology, Thuwal 23955-6900, Saudi Arabia

**Abstract**

Contrary to the common perception of hyperarid drylands as barren and lifeless, these regions are home to some of the planet's most unique biodiversity and support over 100 million people. Despite their ecological and human significance, hyperarid drylands remain among the least studied biomes in the world. In this article, we explore how improving our understanding of hyperarid ecosystems in the Middle East can yield valuable insights applicable to other hyperarid regions. We examine how ongoing greening initiatives in the Middle East offer a unique opportunity to deepen our knowledge of dryland ecology and advocate for the establishment of a comprehensive research program in the region. This program would focus on ecosystem functionality across spatial and temporal scales, setting the stage for a global monitoring network for hyperarid drylands. Such efforts would inform conservation strategies and climate change mitigation, while also shedding light on the resilience and adaptability of hyperarid ecosystems to environmental change. Ultimately, this monitoring would guide management practices to preserve biodiversity, enhance ecosystem services and promote sustainable development in hyperarid regions worldwide.

Hyperarid drylands, areas with an aridity index (precipitation/potential evapotranspiration) below 0.05, represent some of the most extreme environments on Earth. Despite the perception as being inhospitable to life, they host a diverse set of biota and ecosystems, including rangelands that provide grazing for nomadic tribes (Johnson, 1993), biocrusts that contribute to carbon sequestration (Kidron et al., 2015) or coastal mangroves and salt marshes that support fisheries and modulate nutrient cycling (El-Regal and Ibrahim, 2014). Encompassing an area of around 10 million km<sup>2</sup>, the extent of hyperarid regions is expected to grow by the end of the century due to increasing aridity driven by climate change. Current projections estimate the expansion of hyperarid land by 2050 to range from 6% under moderate scenarios to as much as 12% in the most pessimistic scenarios (Huang et al., 2016). While more than 100 million people currently live in hyperarid drylands (MEA, 2005), population growth rates as high as 65% by 2100 have been projected for developing countries in these regions (Huang et al., 2016), placing further strain on these ecosystems.


Hyperarid ecosystems remain poorly studied compared to other dryland and nondryland ecosystems (Brito et al., 2014; Šmíd et al., 2021). Research on their biodiversity, structure and function is limited, representing less than 3% of all dryland studies (Groner et al., 2023). These ecosystems are not only challenging to access (Ficetola et al., 2013) but also vastly under-protected, with just 6.7% of their total area designated for conservation (Lewin et al., 2024). The inaccessibility of hyperarid areas, coupled with the misconception that they are barren and devoid of life, has resulted in their neglect of conservation efforts (Durant et al., 2012). Consequently, there is a widespread but incorrect belief that these environments are either ecologically insignificant or incapable of further degradation (Martínez-Valderrama et al., 2020). Contrary to this view, hyperarid drylands are rich in biodiversity. For example, the Algerian Sahara alone is home to at least 1,200 plant species (Ozenda, 2004). Due to the unique adaptations of organisms in these extreme environments, hyperarid ecosystems offer valuable insights into how dryland systems might respond to future climate change. They serve as natural laboratories for studying the impacts of, and adaptations to, climatic change that could affect other dryland regions (Groner et al., 2023; Grünzweig et al., 2022). Furthermore, as nondryland areas face increasing water scarcity, mechanisms governing ecosystem functioning in drylands are expected to become relevant in these regions (Allan et al., 2020). Many of these changes are anticipated in densely populated regions, particularly in the subtropics and mid latitudes, with significant implications for food production and societal well-being (Grünzweig et al., 2022). Beyond ecological insights,

© The Author(s), 2024. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.



**Box 1.** Hyperarid drylands in the Middle East are much more than barren landscapes.

The Middle East is home to diverse ecosystems that, while also found in other regions under more favorable conditions, can thrive in some of the driest environments on Earth. Gaining a deeper understanding of these ecosystems offers valuable insights into their functioning, restoration potential and relevance for addressing climate change, land degradation and desertification. For instance, mangroves (1) along the coasts of the Red Sea and Arabian Sea (Almahasheer, 2018; Blanco-Sacristán *et al.*, 2022) are a key vegetation type in the Middle East. These ecosystems provide nursery grounds for marine life, support local communities through commercial species and protect coastlines from erosion. However, mangroves in this region endure extreme saline stress due to limited freshwater inputs and increasing groundwater extraction, on which they heavily rely (Adame *et al.*, 2021). Additionally, these mangroves face significant human pressures (Almahasheer *et al.*, 2016). Understanding how mangroves survive in such arid conditions offers a unique opportunity to predict how global mangrove ecosystems might respond to climate change, including the effects of human activities, sea-level rise and microclimatic shifts (Osland *et al.*, 2016). Similarly, the grasslands (2) of the Middle East, such as those in Iran's Taftan mountains (Burrascano *et al.*, 2018) and the southwestern Arabian Peninsula (Ghazanfar and Fisher, 1998), provide a valuable opportunity to study the interactions between abiotic and biotic factors across altitudinal and latitudinal gradients. As global aridity increases, understanding the dynamics of these grasslands – ranging from Mediterranean grasslands to semi-arid steppes – can offer crucial insights for improving grassland health in other dryland regions. This is particularly important given the extreme climatic conditions in which these grasslands exist, which mirror those in many other arid and semi-arid ecosystems globally, such as the grasslands in the Namib Desert (Evans *et al.*, 2020; Logan *et al.*, 2021) and Australia (Keast, 2013). By studying how these Middle Eastern grasslands thrive, researchers can gain a deeper understanding of the resilience and adaptive strategies of grassland ecosystems, crucial for managing the effects of climate change on grasslands worldwide. Shrublands (3), which dominate much of the Middle East – such as the eastern Arabian Peninsula, parts of Jordan (e.g., Jebel Ajloun) and northern Israel (Upper Galilee) – also play a critical ecological role. They provide habitats for insects and small rodents and host biocrusts – communities of photo- and heterotrophic organisms living on the soil surface in large, unvegetated drylands. Biocrusts are essential for maintaining dryland ecosystem health by influencing soil respiration, nutrient cycling and runoff dynamics. While biocrusts have been extensively studied in regions like the Negev Desert and the Arava Valley in Israel (e.g., Galun and Garty, 2003; Kidron and Tal, 2012), research across other Middle Eastern countries is limited. Studies from countries like Iran (Bashtian *et al.*, 2019), Iraq (Hamdi *et al.*, 1978), Jordan (El-Oqlah *et al.*, 1986), Oman (Abed *et al.*, 2013) and Saudi Arabia (Alotaibi *et al.*, 2020) suggest that biocrust composition is relatively uniform across the region (Galun and Garty, 2003), but more research is needed to fully understand their distribution and composition in the Middle East. With its long history of land use and anthropogenic impacts under climate change (Kaniewski *et al.*, 2012), the Middle East offers valuable insights into how human activities shape biocrust communities under extreme environmental conditions. Studying the interactions between biocrusts and human-induced changes – such as grazing, agriculture and urbanization – can inform strategies for managing and mitigating these impacts, both regionally and globally. Leveraging remote sensing technologies (e.g., satellites, drones and eddy-covariance towers) alongside in situ data collection could enhance ecosystem surveys, providing timely insights into their functioning. This data could also support the establishment of new monitoring networks, such as eddy covariance flux networks, which remain underrepresented in hyperarid drylands worldwide (Smith *et al.*, 2019).



studying the adaptations of organisms in hyperarid drylands holds great promise for biotechnological and biodiversity applications (Bull and Asenjo, 2013).

The Middle East accounts for over 30% of the world's hyperarid drylands. This region hosts diverse biomes that have developed unique ecoevolutionary adaptations over thousands of years of biotic and abiotic interactions. They support more than 8,000 unique species of vascular plants (Hegazy and Doust, 2016) and encompass a diverse range of ecosystems present in other regions, albeit under more favorable conditions. These ecosystems span from mangroves along the coastal fringes of the Red Sea to grasslands and shrublands extending across Turkey and Iraq (Box 1). Despite the geographic and historical interest the Middle East has generated, much of the research undertaken in this region has predominantly focused on the description of the flora in individual countries, such as Iran (e.g., Rechinger, 1963–2005), Israel and Palestine (e.g., Danin, 2004; Zohary, 1962), Lybia (e.g., Jafri and El-Gadi, 1977–1993), Oman (e.g., Ghazanfar, 1992; Ghazanfar and Fisher, 1998), Saudi Arabia (e.g., Mandaville, 2013; Migahid, 1978), Turkey (e.g., Davis et al., 1988, 1994) or Yemen (e.g., Brown and Mies, 2012; Kilian et al., 2002). Other studies described the vegetation of the Middle East from geobotanical and phytogeographical perspectives (e.g., Zohary, 1971, 1973). Similarly, the fauna of the Middle East has drawn significant interest due to the extreme environmental conditions these species endure, with several biodiversity hotspots in the region. For example, the Arabian Peninsula hosts a high number of endemic vertebrate species, 21.6% of which are unique to this region (Mallon, 2011). Additionally, the Middle East serves as an essential stopover for migratory bird species along major migratory routes that connect Africa, Asia and Europe (Schekler et al., 2022). Countries like Israel have been extensively studied for their key role in bird migration routes for decades already (e.g., Leshem and Yom-Tov, 1996). However, except for Hegazy and Doust (2016), the life stories of many Middle Eastern species have not been comprehensively investigated and described while concurrently considering this region's geography, plant evolution and ecology. Moreover, these studies have yet to integrate the complex interactions between human societies and ecosystems, particularly in the face of the additional pressures imposed by climate change.

Here, we elaborate on how research on the biodiversity and ecology of Middle East hyperarid drylands can advance our understanding of dryland ecosystems globally while also contributing to the success of ongoing Saudi and Middle East Green Initiatives (<https://www.greeninitiatives.gov.sa/>). With an initial investment of more than USD 180 billion, these green initiatives aim to restore degraded marine and terrestrial environments, enhance biodiversity and mitigate the impacts of climate change throughout the Middle East. We argue that if these initiatives are successfully developed and implemented, they might serve as the foundation for further experimental and theoretical studies on the impacts of extreme climates on dryland ecosystems globally. Furthermore, the Saudi and Middle East Green Initiatives could establish the base for applied solutions aimed at preserving and/or rehabilitating the biodiversity and ecosystem services of global drylands, mitigating climate change and addressing land degradation and desertification.

### Ongoing greening initiatives in the Middle East: An untapped potential to enhance our understanding of hyperarid ecosystems

To sustain its unique biodiversity into the future, it is crucial to promote the resilience and health of hyperarid ecosystems,

particularly given the compound pressures of anthropogenic influence and climate change. These are key objectives of the Saudi and Middle East Green Initiatives, which aim to protect up to 30% of Saudi Arabia's land and sea territories and plant up to 10 and 40 billion trees within the Kingdom and across the Middle East, respectively (<https://www.greeninitiatives.gov.sa/about-sgi/> and <https://www.greeninitiatives.gov.sa/about-mgi/>, respectively). Other actions supported by these initiatives include the increase of renewable energy capacity – which has already risen by 300% in Saudi Arabia, restoring degraded lands – 94,000 hectares have been rehabilitated across Saudi Arabia at the moment – and rewinding endangered species that play a key role in the ecological balance of these ecosystems. It is also expected that the Saudi Green Initiative will play a significant role in achieving the recent commitment of Saudi Arabia to reach net zero emissions by 2060, with the Middle East Green Initiative aiding broader regional objectives towards carbon neutrality. Moreover, the Saudi Green Initiative's ambition to protect at least 30% of Saudi Arabia's territories by 2030 is in harmony with the global “30x30” target adopted under the Kunming-Montreal Global Biodiversity Framework of the Convention on Biological Diversity (CBD, 2022). Although this is a challenging objective, 18.1% and 6.49% of Saudi Arabia's terrestrial and marine areas are already protected.

Both greening initiatives will protect some of the region's iconic terrestrial fauna, which are classified at varying levels of threat, ranging from vulnerable to critically endangered, according to the International Union for Conservation of Nature (IUCN) Red List of Threatened Species. Additionally, they will protect mangroves, coral reefs and salt marshes, which have coevolved in this region to create some of the most resilient marine ecosystems globally (McCabe et al., 2023). Some of the Saudi Green Initiative activities include the creation of national reserves, such as the King Salman bin Abdulaziz Royal Reserve, located in the north of the Arabian Peninsula. Covering approximately 130,000 km<sup>2</sup>, this reserve hosts vulnerable species of mammals (e.g., *Capra nubiana*, *Canis lupus arabs*) and birds (e.g., *Torgos tracheliotos*, *Falco cherrug*). Additionally, urban areas are targeted by these initiatives. Cities like Riyadh and Makkah in Saudi Arabia are seeing an increase in the number of trees planted and the creation of new green areas, enhancing human well-being and biodiversity (Cox et al., 2017; Gaston, 2010).

The Saudi and Middle East Green Initiatives should also learn from past actions and seek not only to ecologically transform broad landscapes but also to shape societies and economies. For example, the Great Green Wall for the Sahara and the Sahel Initiative (GGWSS), which emerged in 2007, involves over 20 countries bordering the Sahara to establish plantations on 100 million ha from Eritrea's Red Sea coast to Senegal's Atlantic coast (Sileshi et al., 2023). The GGWSS was built upon earlier initiatives aimed at combating desertification in the Sahel region's countries (Mbow, 2017). One such initiative was Algeria's Green Dam Initiative, started in 1972, which aimed to establish a three million ha band of plantations to halt the northward advance of the Sahara Desert (Benhizia et al., 2021). Other projects, such as the Acacia operation project and the Support for the rehabilitation and extension of the Nouakchott green belt in Mauritania, engaged local communities and national authorities in restoring inland and coastland ecosystems (Berte, 2010). Projects in the Sahel region have shown that where policies and incentives are favorable, farmers actively promote the natural regeneration of trees, resulting in vast areas now being covered by trees (e.g., Haglund et al., 2011). A participatory approach involves extensive community engagement and enhances accountability and stewardship in land-restoration efforts. Initially, a centralized approach, heavily reliant on forest department control



and substantial investment in equipment, marginalized local communities. Recognizing community ownership has enabled Sahelian countries to mitigate conflicts between development and environmental goals (Kumar, 2003). However, land privatization in the Sahel often fails due to diverse landscape uses and stakeholder needs (Schoneveld, 2017). These failures underscore the necessity for stakeholder-supported, site-specific solutions that enable ongoing improvement across countries and implementation sites. Learning from experiences in the Sahel region, local actions that can be scaled up with positive results include the zoning of grazing areas, ensuring water availability for livestock and promoting fodder trees (Mbow, 2017).

In Asia, the Great Green Wall of China (GGWC), initiated by the Chinese government in 1978, aims to combat desertification and reduce the eolian transport of dust from the Gobi Desert (Parungo *et al.*, 1994). Scheduled for completion in 2070 (Lu *et al.*, 2018), this project builds on China's experience with shelterbelt programs (Qi and Dauvergne, 2022). While the GGWC has yielded benefits, such as reduced dust movement and increased vegetation, during its first stages, many of the dryland areas targeted for afforestation were found to be better suited for grasslands and steppes than woodlands or forests (Cao *et al.*, 2010; Mátyás *et al.*, 2013), often leading to significant water pressures on water resources (Li *et al.*, 2021a). Not only tree survival rates were low but also irrigation was necessary in drier areas within many of these projects (e.g., Wang *et al.*, 2020). Nevertheless, subsequent research has demonstrated the benefits of shelterbelts in drylands for reducing net erosion (Su *et al.*, 2021) and improving crop productivity (Zheng *et al.*, 2016). Additionally, studies on biocrusts in China's drylands have shown that breeding them can effectively control land degradation (Li *et al.*, 2021b) by reducing dust emissions and increasing soil nutrient content (He *et al.*, 2019; Li *et al.*, 2010). Because of these experiences, new strategies in China now focus on science-based activities, encouraging natural regeneration, creating multispecies plantations, matching species to local conditions and emphasizing water conservation (Turner *et al.*, 2023).

Over the past four decades, Australia has also made significant advancements in restoring its drylands through sustained efforts and community involvement (Campbell *et al.*, 2017). Initiatives in Australia learned from small-scale efforts and led to a shift in policies towards large-scale activities, biodiversity conservation, water quality improvement and greenhouse gas mitigation. Successful restoration programs underscored community capacity and commitment, yet it was also recognized that community efforts alone were insufficient for sustainable resource management on a landscape or continental scale without technically and economically viable land use and farming systems. These lessons are particularly important in drylands, where synergistic interactions such as grazing intensification, drought, climate change, reduced fire frequency and changes in atmospheric chemistry or small animal populations can collectively overwhelm the effects of individual factors (Fu *et al.*, 2021a).

### Restoration in the Middle East cannot be based only on planting trees in the desert

Ambitious tree-planting objectives are not a new concept, even in drylands (Bond *et al.*, 2019). Unfortunately, many previous dryland afforestation efforts have often delivered tree monocultures, which risks reducing sustainable development by negatively affecting ecosystem functioning (Yao *et al.*, 2021). Apart from avoiding planting regimes that are incompatible with the landscape, the inherent

constraints of water availability in drylands and the increased pressures that large-scale tree planting places on these, are critical considerations when designing greening and restoration efforts (Schwärzel *et al.*, 2020). Although intrinsically appealing from a policy perspective (i.e., planting trees is a socially recognizable and acceptable climate action), excessive focus on afforestation using trees can miss opportunities for broader and longer-term benefits. For instance, mono-specific tree plantations may achieve a narrow accounting-based objective (in terms of trees planted or carbon captured) but they can reduce ecosystem diversity (e.g., Maestre and Cortina, 2004), jeopardize water resources for humans and ecosystems (e.g., Feng *et al.*, 2016) and amplify the risk of future carbon loss following any ecosystem disturbance (e.g., forest fires and pests; Anderegg *et al.*, 2020). In other parts of the world, regions deemed degraded have been mistakenly considered as potential areas for afforestation, simply by failing to carefully assess their suitability for tree planting (e.g., soil health, environmental gradients). Such areas have included grasslands and shrublands (Veldman *et al.*, 2019), which represent two of the more common environments found in the Middle East (Box 1; Hegazy and Doust, 2016).

Recognizing the limitations and unintended consequences of prior afforestation strategies underscores the importance of adopting a more nuanced approach to ecosystem restoration, particularly in hyperarid regions. Increased biodiversity is considered an indicator of healthier and more resilient ecosystems, allowing faster recovery from disturbance and providing ecosystem services that contribute to more sustainable and stable human development (Jactel *et al.*, 2017). Thus, restoration and conservation efforts should act in concert to increase biodiversity, thereby bolstering the resilience of all naturally occurring ecosystems. This holistic view is crucial if the goal is to restore the multifaceted ecosystems of hyperarid lands, considering the variety of services they provide (Box 1). For example, biocrusts are key players in dryland development and function that increase soil carbon and nutrient contents, impact multiple components of the hydrological cycle and reduce soil erosion and dust emissions (Eldridge *et al.*, 2020; Rodríguez-Caballero *et al.*, 2022), benefitting both the environment and human societies. Therefore, the development of a biocrust research program is urgently needed to understand their ecology, distribution and potential to restore degraded habitats and mitigate climate change in the Middle East.

Restoration and greening initiatives in the Middle East should focus not only on what is visible above ground but also on soils. Over 32% of the world's soil organic pool is stored in drylands worldwide (Plaza *et al.*, 2018a), with significant loss of carbon occurring in major cropland and grazing areas (Sanderman *et al.*, 2017). However, although soils' potential to mitigate climate change has been long recognized (Bossio *et al.*, 2020), their role in dryland restoration and mitigation efforts remains underexplored. Soil organic carbon can act as a stable carbon sink, showing resilience to land-use changes and disturbances, unlike above-ground biomass. Carbon-rich soils also enhance water and nutrient retention, enhancing ecosystem resilience to disturbances like droughts (e.g., Iizumi and Wagai, 2019). However, regional evaluations of soil organic carbon in drylands remain limited, with existing studies often producing inconsistent results (Fu *et al.*, 2021b). Furthermore, understanding the impact of land-use changes on regional soil carbon is hindered by insufficient data quality, poor representativeness and a lack of historical land-use information (Hendriks *et al.*, 2016). Comprehensive assessment of soil carbon stocks requires robust sampling methods that can scale

site-specific data to broader regional levels (Ciais et al., 2011), an ongoing challenge in terrestrial carbon studies (Zhang and Hartemink, 2017). As such, a regional dataset combining soil organic carbon, land-use and soil properties for the Middle East would enhance our understanding of how climate influences physical processes in global drylands. For instance, increasing aridity is known to reduce soil carbon and nitrogen levels (Delgado-Baquerizo et al., 2013) and disrupt the nutrient balance in dryland soils (Maestre et al., 2016). Carbon accumulation in soils is influenced by factors such as parent material, topography, microclimatic conditions and species diversity (Ramesh et al., 2019), while human activities can accelerate carbon emissions (Lal, 2004a; Schlesinger, 2000). Although improved management strategies (e.g., grazing regimes, organic amendments, cover crops, crop rotation and conservation tillage) can enhance carbon stocks in dryland soils (Lal, 2004b, 2018; Plaza et al., 2018b), they can be less effective in these environments due to their coarser texture and lower clay content, which protects organic matter from decomposition (Lehmann and Kleber, 2015; Six et al., 2002). It is, therefore, crucial to evaluate the interactions between biotic, abiotic and human factors to understand soil C dynamics in the Middle East.

### Concluding remarks

Hyperarid lands have been largely missing from existing large-scale global dryland field surveys (Maestre et al., 2012, 2022b). The Saudi and Middle East Green Initiatives provide a unique opportunity to gain insights into the processes that govern the structure, functioning and responses to climate change of hyperarid drylands. Knowledge gaps that need to be addressed include understanding: (i) the drivers for the unexpected high functional diversity in dryland plants (Gross et al., 2024); (ii) how plants will adapt to water scarcity and respond to increased inter-annual precipitation variability (Garcia-Pichel and Sala, 2022); (iii) developing a region-wide understanding of the distribution, characteristics and functioning of biocrusts (e.g., Abed et al., 2019) and (iv) the mechanisms, both physiological and genetic, behind the ability of soil microorganisms to endure extreme conditions (Makhalanyane et al., 2015). Further, many remote sensing-derived products ignore hyperarid drylands based on the assumption that vegetation is largely absent. As a result, hyperarid drylands are often excluded from remote sensing products typically used in global studies and vegetation estimates (e.g., Harris et al., 2021; Sabatini et al., 2022). This is problematic, as vegetation (and trees in particular) is more abundant in hyperarid areas than initially thought (Brandt et al., 2020; Reiner et al., 2023). More generally, international networks evaluating ecosystem carbon, water and energy fluxes, such as FLUXNET (Baldocchi et al., 2001), lack sites in hyperarid environments, despite these representing around 8% of the global land surface (Prävalie et al., 2019). Developing an augmented flux network that includes sites in the Middle East would provide invaluable information on hyperarid drylands and contribute to filling existing gaps in flux databases that preclude obtaining more precise carbon cycling and climate change impact estimates.

Many of the actions discussed here will occur in complex and unpredictable contexts, where human realities should be considered alongside ecological and biophysical factors. Drylands exhibit sensitivity to changes in structure–function relationships due to extreme climate conditions (D’Odorico and Bhattachan, 2012; Reynolds et al., 2007), and human interventions can alter the resilience and stability of these systems (Robinson et al., 2015).

Interactions between natural and human-induced processes affect dryland dynamics at specific scales (Fu et al., 2021a), varying by social, cultural and economic context (Stringer et al., 2017). Therefore, understanding the complex and adaptive nature of drylands in the Middle East should involve dynamic interactions between its ecosystems and human societies (Folke et al., 2016), requiring interdisciplinary efforts (Bautista et al., 2017).

Research in the Middle East should focus on the interplay between ecosystem services and human well-being to optimize services that enhance drylands’ health and human well-being in the long term (Fu et al., 2021b). Identifying local limiting factors and their impacts can improve knowledge of ecosystem functioning and livelihoods through sustainable development (Reed et al., 2015; Turner et al., 2003). An interdisciplinary approach, which evaluates ecological and social perspectives together, will allow for assessing ecological dynamics and their driving forces in the Middle East. It will not only enable an understanding of the macroscopic differences among various dryland systems in this region but also help identify management or policy responses likely to deliver successful outcomes in different types of drylands (Fu et al., 2021b).

Developing a comprehensive research program on ecosystem structure and functioning across multiple spatio-temporal scales in the Middle East is a critical step to provide the scientific underpinning needed for the success of ongoing green initiatives and climate change and desertification mitigation actions in this region. The creation of a Middle East collaborative network of researchers, practitioners and decision-makers, and the set-up of standardized regional surveys using standardized protocols following models successfully implemented in other large-scale and global surveys (e.g., Maestre et al., 2022a; Maestre and Eisenhauer, 2019), would be a fundamental step forward towards achieving this aim. Doing so would not only be key to creating the basis for long-term monitoring of ecosystem changes in the region but would provide us with invaluable insights to advance our understanding of hyperarid drylands and to better comprehend and react to the increasingly drier conditions being experienced and forecasted across the globe.

**Open peer review.** For open peer review materials, please visit <https://doi.org/10.1017/dry.2024.6>

**Author contribution.** J. B.-S. conceived the idea and wrote the initial draft; all authors contributed to the review and the final version of the manuscript.

**Financial support.** The work presented herein was funded by the King Abdullah University of Science and Technology’s Climate and Livability Initiative and Circular Carbon Initiative.

**Competing interest.** The authors declare that they have no conflict of interest.

### References

- Abed RM, Al-Sadi AM, Al-Shehi M, Al-Hinai S and Robinson MD (2013) Diversity of free-living and lichenized fungal communities in biological soil crusts of the Sultanate of Oman and their role in improving soil properties. *Soil Biology and Biochemistry* 57, 695–705. <https://doi.org/10.1016/j.soilbio.2012.07.023>.
- Abed RMM, Tamm A, Hassenrück C, Al-Rawahi AN, Rodríguez-Caballero E, Fiedler S, Maier S and Weber B (2019) Habitat-dependent composition of bacterial and fungal communities in biological soil crusts from Oman. *Scientific Reports* 9, 6468. <https://doi.org/10.1038/s41598-019-42911-6>
- Adame MF, Connolly RM, Turschwell MP, Lovelock CE, Fatoyinbo T, Lagomasino D, Goldberg LA, Holdorf J, Friess DA, Sasmito SD, Sanderman J, Sievers M, Buelow C, Kauffman JB, Bryan-Brown D and Brown CJ (2021) Future carbon emissions from global mangrove forest loss. *Global Change Biology* 27(12), 2856–2866. <https://doi.org/10.1111/gcb.15571>

- Allan RP, Barlow M, Byrne MP, Cherchi A, Douville H, Fowler HJ, Allan RA, Barlow M, Byrne MP, Cherchi A, Douville H, Fowler HJ, Gan TY, Pendergrass AG, Rosenfeld D, Swann ALS, Wilcox LJ and Zolina O (2020) *Advances in understanding large-scale responses of the water cycle to climate change. Annals of the New York Academy of Sciences* 1472(1), 49–75. <https://doi.org/10.1111/nyas.14337>
- Almahasheer H (2018) Spatial coverage of mangrove communities in the Arabian Gulf. *Environmental Monitoring and Assessment* 190, 85. <https://doi.org/10.1007/s10661-018-6472-2>.
- Almahasheer H, Aljowair A, Duarte CM and Irigoien X (2016) Decadal stability of Red Sea mangroves. *Estuarine, Coastal and Shelf Science* 169, 164–172. <https://doi.org/10.1016/j.ecss.2015.11.027>.
- Alotaibi MO, Sonbol HS, Alwakeel SS, Suliman RS, Fodah RA, Jaffal ASA, AlOthman NI, and Mohammed AE (2020) Microbial diversity of some sabkha and desert sites in Saudi Arabia. *Saudi Journal of Biological Sciences* 27, 2778–2789. <https://doi.org/10.1016/j.sjbs.2020.06.038>.
- Anderegg WR, Trugman AT, Badgley G, Anderson CM, Bartuska A, Ciais P, Cullenward D, Field CB, Freeman J, Goetz SJ, Hicke JF, Huntzinger D, Jackson RB, Nickerson J, Pacala S and Randerson JT (2020) Climate-driven risks to the climate mitigation potential of forests. *Science* 368, 1236570. <https://doi.org/10.1111/nph.15283>.
- Bashtian MH, Sepehr A, Farzam M and Bahreini M (2019) Biological soil crusts, plant functional groups, and soil parameters in arid areas of Iran. *Polish Journal of Ecology* 66, 337–351. <https://doi.org/10.3161/15052249PJE2018.66.4.003>.
- Baldocchi D, Falge E, Gu L, Olson R, Hollinger D, Running S, Anthoni P, Bernhofer C, Davis K, Evans R, Fuentes J, Goldstein A, Katul G, Law B, Lee X, Malhi Y, Meyers T, Munger W, Oechel W, Paw KT, Pilegaard K, Schmid HP, Valentini R, Verma S, Vesala T, Wilson K and Wofsy S (2001) FLUXNET: A new tool to study the temporal and spatial variability of ecosystem-scale carbon dioxide, water vapor, and energy flux densities. *Bulletin of the American Meteorological Society* 82(11), 2415–2434. [https://doi.org/10.1175/1520-0477\(2001\)082<2415:FANTTS&2.3.CO;2](https://doi.org/10.1175/1520-0477(2001)082<2415:FANTTS&2.3.CO;2).
- Bautista S, Llovet J, Ocampo-Melgar A, Vilagrosa A, Mayor ÁG, Murias C, Vallejo R and Orr BJ (2017) Integrating knowledge exchange and the assessment of dryland management alternatives – A learning-centered participatory approach. *Journal of Environmental Management* 195, 35–45.
- Benhizia R, Kouba Y, Szabó G, Négyesi G, Négyesi G and Ata B (2021). Monitoring the spatiotemporal evolution of the Green Dam in Djelfa Province, Algeria. *Sustainability* 13(14), 7953. <https://doi.org/10.3390/su13147953>
- Berte CJ (2010) *Fighting Sand Encroachment: Lessons from Mauritania*. Food and Agriculture Organization of the United Nations (FAO), Vol. 158.
- Blanco-Sacristán J, Johansen K, Duarte CM, Daffonchio D, Hoteit I and McCabe MF (2022). Mangrove distribution and afforestation potential in the Red Sea. *Science of the Total Environment* 843, 157098. <https://doi.org/10.1016/j.scitotenv.2022.157098>
- Bond WJ, Stevens N, Midgley GF and Lehmann CE (2019) The trouble with trees: afforestation plans for Africa. *Trends in Ecology & Evolution* 34, 963–965. <https://doi.org/10.1016/j.tree.2019.08.003>.
- Bossio DA, Cook-Patton SC, Ellis PW, Fargione J, Sanderman J, Smith P, Wood S, Zomer RJ, von Unger M, Emmer IM and Griscom BW (2020) The role of soils in mitigating global climate change. *Nature Sustainability* 3, 391–398. <https://doi.org/10.1038/s41893-020-0491-z>.
- Brandt M, Tucker CJ, Kariryaa A, Rasmussen K, Abel C, Small J, Chave J, Rasmussen LV, Hiernaux P, Diouf AA Kergoat L (2020) An unexpectedly large count of trees in the West African Sahara and Sahel. *Nature* 587, 78–82. <https://doi.org/10.1038/s41586-020-2824-5>.
- Brito JC, Godinho R, Martínez-Freiria F, Pleguezuelos JM, Rebelo H, Santos X, Vale CG, Velo-Antón G, Boratyński Z, Carvalho SB and Ferreira S. (2014) *Unravelling biodiversity, evolution and threats to conservation in the Sahara-Sahel. Biological Reviews. Biological Reviews* 89, 215–231. <https://doi.org/10.1111/brv.12049>.
- Brown G and Mies B (2012) *Vegetation Ecology of Socotra*. Springer Science & Business Media. Vol. 7.
- Bull AT and Asenjo JA (2013) Microbiology of hyper-arid environments: recent insights from the Atacama Desert, Chile. *Antonie Van Leeuwenhoek* 103, 1173–1179.
- Burrascano MVS, Naqinezhad A and Fernández MP (2018) Grasslands of the Mediterranean Basin and the Middle East and their Management. In *Grasslands of the World: Diversity, Management and Conservation*. CRC Press.
- Campbell A, Alexandra J and Curtis D (2017) Reflections on four decades of land restoration in Australia. *The Rangeland Journal* 39, 405–416. <https://doi.org/10.1071/RJ17056>.
- Cao S, Tian T, Chen L, Dong X, Yu X and Wang G (2010) Damage caused to the environment by reforestation policies in arid and semi-arid areas of China. *Ambio* 39, 279–283. <https://doi.org/10.1007/s13280-010-0038-z>.
- Convention on Biological Diversity (CBD) (2022) COP15: A new era for humanity - Earth's biodiversity will be restored. Available at <https://www.cbd.int/article/cop15-cbd-press-release-final-19dec2022> (accessed 12 March 2024).
- Ciais P, Bombelli A, Williams M, Piao SL, Chave J, Ryan CM, Henry M, Brender P and Valentini RJ (2011) The carbon balance of Africa: synthesis of recent research studies. *Philosophical Transactions of the Royal Society A* 369, 2038–2057. <https://doi.org/10.1098/rsta.2010.0328>.
- Cox DT, Shanahan DF, Hudson HL, Fuller RA, Anderson K, Hancock S and Gaston KJ (2017) Doses of nearby nature simultaneously associated with multiple health benefits. *International Journal of Environmental Research and Public Health* 14, 172. <https://doi.org/10.3390/ijerph14020172>.
- D'Odorico P and Bhattachan A (2012) Hydrologic variability in dryland regions: impacts on ecosystem dynamics and food security. *Philosophical Transactions of the Royal Society B: Biological Sciences* 367, 3145–3157. <https://doi.org/10.1098/rstb.2012.0016>.
- Danin A (2004) *Distribution Atlas of Plants in the Flora Palaestina Area*. Jerusalem, Israel: Israel Academy of Sciences and Humanities.
- Davis P, Mill R and Tan K (1988) *Flora of Turkey and the East Aegean Islands*. Edinburgh, U.K: Edinburgh University Press, Vol. 10.
- Davis S, Heywood V and Hamilton A (eds.) (1994) *Centres of Plant Diversity. Vol 1: Europe, Africa, Southwest Asia and Middle East*. Oxford, U.K: WWF and IUCN.
- Delgado-Baquerizo M, Maestre FT, Gallardo A, Bowker MA, Wallenstein MD, Quero JL, Ochoa V, Gozalo B, García-Gómez M, Soliveres S and García-Palacios P (2013) Decoupling of soil nutrient cycles as a function of aridity in global drylands. *Nature* 502, 672–676. <https://doi.org/10.1038/nature12670>.
- Durant SM, Pettorelli N, Bashir S, Woodroffe R, Wachter T, De Ornellas P, Ransom C, Abaigar T, Abdelgadir M, El Alqamy H and Beddief M (2012) Forgotten biodiversity in desert ecosystems. *Science* 336, 1379–1380. <https://doi.org/10.1126/science.336.6087.1379>.
- Eldridge DJ, Reed S, Travers SK, Bowker MA, Maestre FT, Ding J, Havrilla C, Rodriguez-Caballero E, Barger N, Weber B and Antoninka A (2020) *The pervasive and multifaceted influence of biocrusts on water in the world's drylands. Global Change Biology* 26, 6003–6014. <https://doi.org/10.1111/gcb.15232>.
- El-Oqlah AA, Hawksworth DL and Lahham JN (1986). Additions to the lichens of Jordan. *Candollea* 41, 69–73. <https://doi.org/10.5169/seals-879980>
- El-Regal MAA and Ibrahim NK (2014) Role of mangroves as a nursery ground for juvenile reef fishes in the southern Egyptian Red Sea. *Egyptian Journal of Aquatic Research* 40, 71–78. <https://doi.org/10.1016/j.ejar.2014.01.001>.
- Evans S, Todd-Brown KE, Jacobson K and Jacobson P (2020) Non-rainfall moisture: a key driver of microbial respiration from standing litter in arid, semiarid, and mesic grasslands. *Ecosystems* 23, 1154–1169. <https://doi.org/10.1007/s10021-019-00461-y>.
- Feng X, Fu B, Piao S, Wang S, Ciais P, Zeng Z, Lü Y, Zeng Y, Li Y, Jiang X and Wu B (2016) Revegetation in China's Loess Plateau is approaching sustainable water resource limits. *Nature Climate Change* 6, 1019–1022. <https://doi.org/10.1038/nclimate3092>.
- Ficetola GF, Bonardi A, Sindaco R and Padoa-Schioppa E (2013) Estimating patterns of reptile biodiversity in remote regions. *Journal of Biogeography* 40, 1202–1211. <https://doi.org/10.1111/jbi.12060>.
- Folke C, Biggs R, Norström AV, Reyers B and Rockström J (2016) Social-ecological resilience and biosphere-based sustainability science. *Ecology and Society* 21. <https://doi.org/10.5751/ES-08748-210341>.
- Fu C, Chen Z, Wang G, Yu X and Yu G (2021a) A comprehensive framework for evaluating the impact of land use change and management on soil organic



- carbon stocks in global drylands. *Current Opinion in Environmental Sustainability* **48**, 103–109. <https://doi.org/10.1016/j.cosust.2020.12.001>.
- Fu B, Stafford-Smith M, Wang Y, Wu B, Yu X, Lv N, Ojima DS, Lv Y, Fu C, Liu Y and Niu S (2021b) The Global-DEP conceptual framework—research on dryland ecosystems to promote sustainability. *Current Opinion in Environmental Sustainability* **48**, 17–28. <https://doi.org/10.1016/j.cosust.2020.12.002>.
- Galun M and Garty J (2003) Biological soil crusts of the Middle East. In *Biological Soil Crusts: Structure, Function, and Management*. Springer, pp. 95–106. [https://doi.org/10.1007/978-3-642-56475-8\\_8](https://doi.org/10.1007/978-3-642-56475-8_8).
- García-Pichel F and Sala O (2022) Expanding the pulse–reserve paradigm to microorganisms on the basis of differential reserve management strategies. *BioScience* **72**, 638–650. <https://doi.org/10.1093/biosci/biac036>.
- Gaston KJ (ed.) (2010) *Urban Ecology*. Oxford University Press. <https://doi.org/10.1017/CBO9780511778483>.
- Ghazanfar SA (1992) Quantitative and biogeographic analysis of the flora of the Sultanate of Oman. *Global Ecology and Biogeography Letters* **2**, 189–195. <https://doi.org/10.2307/2997660>.
- Ghazanfar SA and Fisher M (eds.) (1998) *Vegetation of the Arabian Peninsula*. Springer Science & Business Media, Vol. 25. <https://doi.org/10.1007/978-94-017-3637-4>.
- Grünzweig JM, De Boeck HJ, Rey A, Santos MJ, Adam O, Bahn M, Belnap J, Deckmyn G, Dekker SC, Flores O, Gliksmann D, Helman D, Hultine KR, Liu L, Meron E, Michael Y, Sheffer E, Throop HL, Tzok O and Yakir D (2022) Dryland mechanisms could widely control ecosystem functioning in a drier and warmer world. *Nature Ecology & Evolution* **6**(8), 1064–1076. <https://doi.org/10.1038/s41559-022-01779-y>.
- Groner E, Babad A, Berda Swiderski N and Shachak M (2023) Toward an extreme world: The hyper-arid ecosystem as a natural model. *Ecosphere* **14**. <https://doi.org/10.1002/ecs2.4586>.
- Gross N, Maestre FT, Liancourt P, Berdugo M, Martin R, Gozalo B, Ochoa V, Delgado-Baquerizo M, Maire V, Saiz H and Soliveres S (2024) Unforeseen plant phenotypic diversity in a dry and grazed world. *Nature* **632**, 808–814. <https://doi.org/10.1038/s41586-024-07731-3>.
- Haglund E, Ndjeunga J, Snook L and Pasternak D (2011) Dryland tree management for improved household livelihoods: farmer managed natural regeneration in Niger. *Journal of Environmental Management* **92**, 1696–1705. <https://doi.org/10.1016/j.jenvman.2011.01.027>.
- Hamdi YA, Yousef AN, Al-Azawi S, Al-Tai A and Al-Baquari MS (1978). Distribution of certain non-symbiotic nitrogen fixing organisms in Iraqi soils. *Ecological Bulletins* 110–115. <https://www.jstor.org/stable/20112669>
- Harris NL, Gibbs DA, Baccini A, Birdsey RA, De Bruin S, Farina M, Fatoyinbo L, Hansen MC, Herold M, Houghton RA and Potapov PV (2021) Global maps of twenty-first century forest carbon fluxes. *Nature Climate Change* **11**, 234–240. <https://doi.org/10.1038/s41558-020-00976-6>.
- He M, Hu R and Jia R (2019) Biological soil crusts enhance the recovery of nutrient levels of surface dune soil in arid desert regions. *Ecological Indicators* **106**, 105497. <https://doi.org/10.1016/j.ecolind.2019.105497>.
- Hegazy A and Doust JL (2016) *Plant Ecology in the Middle East*. Oxford University Press.
- Hendriks CMJ, Stoorvogel J and Claessens L (2016) Exploring the challenges with soil data in regional land use analysis. *Agricultural Systems* **144**, 9–21. <https://doi.org/10.1016/j.agsy.2016.01.007>.
- Huang J, Yu H, Guan X, Wang G and Guo R (2016) Accelerated dryland expansion under climate change. *Nature Climate Change* **6**, 166–171. <https://doi.org/10.1038/nclimate2837>.
- Iizumi T and Wagai R (2019) Leveraging drought risk reduction for sustainable food, soil and climate via soil organic carbon sequestration. *Scientific Reports* **9**, 19744. <https://doi.org/10.1038/s41598-019-55835-y>.
- Jactel H, Bauhus J, Boberg J, Bonal D, Castagnèrol B, Gardiner B, Gonzalez-Olabarria JR, Koricheva J, Meurisse N and Brockerhoff EG (2017) Tree diversity drives forest stand resistance to natural disturbances. *Current Forestry Reports* **3**, 223–243. <https://doi.org/10.1007/s40725-017-0064-1>.
- Jafri S and El-Gadi A (eds.) (1977–1993) *Flora of Libya*. Botany Dept., Tripoli University: Tripoli, Libya.
- Johnson DL (1993) Nomadism and desertification in Africa and the Middle East. *Geojournal* **31**, 51–66. <https://doi.org/10.1007/BF00815903>.
- Kaniewski D, Van Campo E and Weiss H (2012) Drought is a recurring challenge in the Middle East. *Proceedings of the National Academy of Sciences* **109**, 3862–3867. <https://doi.org/10.1073/pnas.1116304109>.
- Keast A (2013) *Biogeography and Ecology in Australia*. Springer. <https://doi.org/10.1007/978-94-017-6295-3>.
- Kidron GJ, Li XR, Jia RL, Gao YH and Zhang P (2015) Assessment of carbon gains from biocrusts inhabiting a dunefield in the Negev Desert. *Geoderma* **253**, 102–110. <https://doi.org/10.1016/j.geoderma.2015.04.015>.
- Kidron GJ and Tal SY (2012) The effect of biocrusts on evaporation from sand dunes in the Negev Desert. *Geoderma* **179**, 104–112. <https://doi.org/10.1016/j.geoderma.2012.02.021>.
- Kilian N, Hein P and Hubaishan MA (2002) New and noteworthy records for the flora of Yemen, chiefly of Hadhramout and Al-Mahra. *Willdenowia* **32**, 239–269. <https://doi.org/10.3372/wi.32.32207>.
- Kumar N (2003) *Community-Driven Development: Lessons from the Sahel: An Analytical Review*. World Bank, Operations Evaluation Department.
- Lal R (2004a) Soil carbon sequestration to mitigate climate change. *Geoderma* **123**, 1–22. <https://doi.org/10.1016/j.geoderma.2004.01.032>.
- Lal R (2004b) Soil carbon sequestration impacts on global climate change and food security. *Science* **304**, 1623–1627. <https://doi.org/10.1126/science.1097396>.
- Lal R (2018) Digging deeper: A holistic perspective of factors affecting soil organic carbon sequestration in agroecosystems. *Global Change Biology* **24**, 3285–3301. <https://doi.org/10.1111/gcb.14054>.
- Lehmann J and Kleber M (2015) The contentious nature of soil organic matter. *Nature* **528**, 60–68. <https://doi.org/10.1038/nature16069>.
- Leshem Y and Yom-Tov Y (1996) The magnitude and timing of migration by soaring raptors, pelicans and storks over Israel. *Ibis* **138**, 188–203. <https://doi.org/10.1111/j.1474-919X.1996.tb04328.x>.
- Lewin A, Murali G, Rachmilevitch S and Roll U (2024) Global evaluation of current and future threats to drylands and their vertebrate biodiversity. *Nature Ecology & Evolution* **8**, 1448–1458. <https://doi.org/10.1038/s41559-024-02450-4>.
- Li C, Fu B, Wang S, Stringer LC, Wang Y, Li Z, Liu Y and Zhou W (2021a) Drivers and impacts of changes in China's drylands. *Nature Reviews Earth & Environment* **2**, 858–873. <https://doi.org/10.1038/s43017-021-00226-z>.
- Li X, Hui R, Tan H, Zhao Y, Liu R and Song N (2021b) Biocrust research in China: recent progress and application in land degradation control. *Frontiers in Plant Science* **12**, 751521. <https://doi.org/10.3389/fpls.2021.751521>.
- Li XR, He MZ, Zerbe S, Li XJ and Liu LC (2010) Micro-geomorphology determines community structure of biological soil crusts at small scales. *Earth Surface Processes and Landforms* **35**, 932–940. <https://doi.org/10.1002/esp.1963>.
- Logan JR, Jacobson KM, Jacobson PJ and Evans SE (2021) Fungal communities on standing litter are structured by moisture type and constrain decomposition in a hyper-arid grassland. *Frontiers in Microbiology* **12**, 596517. <https://doi.org/10.3389/fmicb.2021.596517>.
- Lu F, Hu H, Sun W, Zhu J, Liu G, Zhou W, Zhang Q, Shi P, Liu X, Wu X and Zhang L (2018) Effects of national ecological restoration projects on carbon sequestration in China from 2001 to 2010. *Proceedings of the National Academy of Sciences* **115**, 4039–4044. <https://doi.org/10.1073/pnas.1700294115>.
- Maestre FT and Cortina J (2004) Are *Pinus halepensis* plantations useful as a restoration tool in semiarid Mediterranean areas? *Forest Ecology and Management* **198**, 303–317. <https://doi.org/10.1016/j.foreco.2004.05.040>.
- Maestre FT and Eisenhauer N (2019) Recommendations for establishing global collaborative networks in soil ecology. *Soil Organisms* **91**, 73–80. <https://doi.org/10.25674/so91iss3pp73>.
- Maestre FT, Eldridge DJ, Gross N, Bagousse-Pinguet L, Saiz H, Gozalo B, Ochoa V and Gaitán JJ (2022a) The BIODESERT survey: Assessing the impacts of grazing on the structure and functioning of global drylands. *Web Ecology* **22**, 75–96. <https://doi.org/10.5194/we-22-75-2022>.
- Maestre FT, Le Bagousse-Pinguet Y, Delgado-Baquerizo M, Eldridge DJ, Saiz H, Berdugo M, Gozalo B, Ochoa V, Guirado E, García-Gómez M and Valencia E (2022b) Grazing and ecosystem service delivery in global drylands. *Science* **378**, 915–920. <https://doi.org/10.1126/science.abq4062>.
- Maestre FT, Eldridge DJ, Soliveres S, Kéfi S, Delgado-Baquerizo M, Bowker MA, García-Palacios P, Gaitán J, Gallardo A, Lázaro R and Berdugo M (2016) Structure and functioning of dryland ecosystems in a changing world.

- Annual Review of Ecology, Evolution, and Systematics* 47, 215–237. <https://doi.org/10.1146/annurev-ecolsys-121415-032311>.
- Maestre FT, Quero JL, Gotelli NJ, Escudero A, Ochoa V, Delgado-Baquerizo M, García-Gómez M, Bowker MA, Soliveres S, Escolar C and García-Palacios P (2012) Plant species richness and ecosystem multifunctionality in global drylands. *Science* 335, 214–218. <https://doi.org/10.1126/science.1215442>.
- Makhalanyane TP, Valverde A, Gunnigle E, Frossard A, Ramond JB and Cowan DA (2015) Microbial ecology of hot desert edaphic systems. *FEMS Microbiology Reviews* 39, 203–221. <https://doi.org/10.1093/femsre/fuu011>.
- Mallon DP (2011) Global hotspots in the Arabian Peninsula. *Zoology in the Middle East* 54, 13–20. <https://doi.org/10.1080/09397140.2011.10648896>.
- Mandaville JP (2013) *Flora of Eastern Saudi Arabia*. Routledge.
- Martínez-Valderrama J, Guirado E and Maestre FT (2020) Desertifying deserts. *Nature Sustainability* 3, 572–575. <https://doi.org/10.1038/s41893-020-0561-2>.
- Mátyás C, Sun G and Zhang Y (2013) Afforestation and forests at the dryland edges: Lessons learned and future outlooks. In *Dryland East Asia: Land Dynamics Amid Social and Climate Change*. Berlin: Higher Education Press and De Gruyter, pp. 245–264. <https://doi.org/10.1038/10.13140/RG.2.1.43.25.4487>.
- Mbow C (2017) The Great Green Wall in the Sahel. *Oxford Research Encyclopedia*. <https://doi.org/10.1093/acrefore/9780190228620.013.559>.
- McCabe M, AlShalan M, Hejazi M, Beck H, Maestre FT, Guirado E, Wada Y, Al-Ghamdi SG, AlSaud N, Underwood M, Magistretti P, Gallouzi IE, KAUST, AEON Collective and KAPSARC (2023) *Climate futures report: Saudi Arabia in a 3 Degrees Warmer World*. Riyadh, Saudi Arabia: Riyadh University Publication. <https://doi.org/10.25781/KAUST-8XY63>.
- Migahid AM (1978) *Flora of Saudi Arabia*. Riyadh, Saudi Arabia: Riyadh University Publication.
- Millenium Ecosystem Assessment (MEA) (2005) *Ecosystems and Human Well-Being: Scenarios*. Island Press, pp. 623–662.
- Osland MJ, Enwright NM, Day RH, Gabler CA, Stagg CL and Grace JB (2016). Beyond just sea-level rise: Considering macroclimatic drivers within coastal wetland vulnerability assessments to climate change. *Global Change Biology* 22(1), 1–11. <https://doi.org/10.1111/gcb.13084>.
- Ozenda P (2004) *Flore et Végétation du Sahara*. Paris, France: Centre National de la Recherche Scientifique.
- Parungo F, Li Z, Li X, Yang D and Harris J (1994) Gobi dust storms and the Great Green Wall. *Geophysical Research Letters* 21, 999–1002. <https://doi.org/10.1029/94GL00879>.
- Plaza C, Zaccane C, Sawicka K, Méndez AM, Tarquis A, Gascó G, Heuvelink GB, Schuur EA and Maestre FT (2018a) Soil resources and element stocks in drylands to face global issues. *Scientific Reports* 8, 13788. <https://doi.org/10.1038/s41598-018-32229-0>.
- Plaza C, Gascó G, Méndez AM, Zaccane C and Maestre FT (2018b) Soil organic matter in dryland ecosystems. In García C, Nannipieri P and Hernández T (eds.), *The Future of Soil Carbon*. Academic Press, pp. 39–70. <https://doi.org/10.1016/B978-0-12-811687-6.00002-X>.
- Právilie R, Bandoc G, Patriche C and Sternberg T (2019) Recent changes in global drylands: Evidence from two major aridity databases. *Catena* 178, 209–231. <https://doi.org/10.1016/j.catena.2019.03.016>.
- Qi JJ and Dauvergne P (2022) China's rising influence on climate governance: Forging a path for the global South. *Global Environmental Change* 73, 102484. <https://doi.org/10.1016/j.gloenvcha.2022.102484>.
- Ramesh T, Bolan NS, Kirkham MB, Wijesekara H, Kanchikerimath M, Rao CS, Sandeep S, Rinklebe J, Ok YS, Choudhury BU and Wang H (2019) Soil organic carbon dynamics: Impact of land use changes and management practices: A review. *Advances in Agronomy* 156, 1–107. <https://doi.org/10.1016/bs.agron.2019.02.001>.
- Rechinger K (ed.) (1963–2005) *Flora Iranica* 1–175. Graz, Austria: Akademische Druck-u.-Verlagsanstalt.
- Reed MS, Stringer LC, Dougill AJ, Perkins JS, Athlipheng JR, Mulale K and Favretto N (2015) Reorienting land degradation towards sustainable land management: Linking sustainable livelihoods with ecosystem services in rangeland systems. *Journal of Environmental Management* 151, 472–485. <https://doi.org/10.1016/j.jenvman.2014.11.010>.
- Reiner F, Brandt M, Tong X, Skole D, Kariryaa A, Ciaia P, Davies A, Hiernaux P, Chave J, Mugabowindekwe M and Igel C (2023) More than one quarter of Africa's tree cover is found outside areas previously classified as forest. *Nature Communications* 14, 2258. <https://doi.org/10.1038/s41467-023-37880-4>.
- Reynolds JF, Smith DM, Lambin EF, Turner BL, Mortimore M, Batterbury SP, Downing TE, Dowlatabadi H, Fernández RJ, Herrick JE and Huber-Sannwald E (2007) Global desertification: building a science for dryland development. *Science* 316, 847–851. <https://doi.org/10.1126/science.1131634>.
- Robinson LW, Ericksen PJ, Chesterman S and Worden JS (2015) Sustainable intensification in drylands: what resilience and vulnerability can tell us. *Agricultural Systems* 135, 133–140. <https://doi.org/10.1016/j.agsy.2015.01.005>.
- Rodríguez-Caballero E, Stanelle T, Egerer S, Cheng Y, Su H, Canton Y, Belnap J, Andreae MO, Tegen I, Reick CH, Pöschl U and Weber B (2022). Global cycling and climate effects of aeolian dust controlled by biological soil crusts. *Nature Geoscience* 15, 458–463. <https://doi.org/10.1038/s41561-022-00942-1>.
- Sabatini FM, Jiménez-Alfaro B, Jandt U, Chytrý M, Field R, Kessler M, Lenoir J, Schrödt F, Wiser SK, Arfin Khan MA and Attorre F (2022) Global patterns of vascular plant alpha diversity. *Nature Communications* 13, 4683. <https://doi.org/10.1038/s41467-022-32063-z>.
- Sanderman J, Hengl T and Fiske GJ (2017) Soil carbon debt of 12,000 years of human land use. *Proceedings of the National Academy of Sciences, USA* 114, 9575–9580. <https://doi.org/10.1073/pnas.1706103114>.
- Schekler I, Smolinsky JA, Troupin D, Buler JJ and Sapir N (2022) Bird migration at the edge – geographic and anthropogenic factors but not habitat properties drive season-specific spatial topover distributions near wide ecological barriers. *Frontiers in Ecology and Evolution* 10, 822220. <https://doi.org/10.3389/fevo.2022.822220>.
- Schlesinger WH (2000) Soil respiration and the global carbon cycle. *Biogeochemistry* 48, 7–20. <https://doi.org/10.1023/A:1006247623877>.
- Schoneveld GC (2017) Host country governance and the African land rush: 7 reasons why large-scale farmland investments fail to contribute to sustainable development. *Geoforum* 83, 119–132. <https://doi.org/10.1016/j.geoforum.2016.12.007>.
- Schwärzel K, Zhang L, Montanarella L, Wang Y and Sun G (2020) How afforestation affects the water cycle in drylands: A process-based comparative analysis. *Global Change Biology* 26, 944–959. <https://doi.org/10.1111/gcb.14875>.
- Sileshi GW, Dagar JC, Kuyah S and Datta A (2023) The Great Green Wall Initiatives and Opportunities for Integration of Dryland Agroforestry to Mitigate Desertification. In *Agroforestry for Sustainable Intensification of Agriculture in Asia and Africa*. Singapore: Springer Nature Singapore, pp. 175–206. [https://doi.org/10.1007/978-981-19-4602-8\\_6](https://doi.org/10.1007/978-981-19-4602-8_6).
- Six J, Conant RT, Paul EA and Paustian K (2002) Stabilization mechanisms of soil organic matter: implications for C-saturation of soils. *Plant and Soil* 241, 155–176. <https://doi.org/10.1023/A:1016125726789>.
- Šmíd J, Sindaco R, Shobrak M, Busais S, Tamar K, Aghová T, Simó-Riudalbas M, Tarroso P, Geniez P, Crochet PA and Els J (2021) Diversity patterns and evolutionary history of Arabian squamates. *Journal of Biogeography* 48, 1183–1199. <https://doi.org/10.1111/jbi.14070>.
- Smith WK, Dannenberg MP, Yan D, Herrmann S, Barnes ML, Barron-Gafford GA, Biederman JA, Ferrenberg S, Fox AM, Hudson A and Knowles JF (2019) Remote sensing of dryland ecosystem structure and function: Progress, challenges, and opportunities. *Remote Sensing of Environment* 233, 111401. <https://doi.org/10.1016/j.rse.2019.111401>.
- Stringer LC, Reed MS, Flekens L, Thomas RJ, Le QB and Lala-Pritchard T (2017) A new dryland development paradigm grounded in empirical analysis of dryland systems science. *Land Degradation & Development* 28, 1952–1961. <https://doi.org/10.1002/ldr.2716>.
- Su ZA, Zhou T, Zhang XB, Wang XY, Wang JJ, Zhou MH, Zhang JH, He ZY and Zhang RC (2021) A preliminary study of the impacts of Shelter Forest on soil erosion in cultivated land: evidence from integrated 137 Cs and 210 Pbex measurements. *Soil & Tillage Research* 206, 104843. <https://doi.org/10.1016/j.still.2020.104843>.
- Turner II BL, Kasperson RE, Matson PA, McCarthy JJ, Corell RW, Christensen L, Eckley N, Kasperson JX, Luers A, Martello ML and Polsky C (2003) A framework for vulnerability analysis in sustainability science. *Proceedings of the National Academy of Sciences* 100, 8074–8079. <https://doi.org/10.1073/pnas.1231335100>.
- Turner MD, Davis DK, Yeh ET, Hiernaux P, Loizeaux ER, Fornof EM, Rice AM and Suiter AK (2023) Great green walls: hype, myth, and science.



- Annual Review of Environment and Resources* **48**, 263–287. <https://doi.org/10.1146/annurev-environ-112321-111102>.
- Veldman JW, Aleman JC, Alvarado ST, Anderson TM, Archibald S, Bond WJ, Boutton TW, Buchmann N, Buisson E, Canadell JG and Dechoum MD** (2019) Comment on “The global tree restoration potential”. *Science* **366** (6463), eaay7976. <https://doi.org/10.1126/science.aay7976>.
- Wang Z, Peng D, Xu D, Zhang X and Zhang Y** (2020) Assessing the water footprint of afforestation in Inner Mongolia, China. *Journal of Arid Environments* **182**, 104257. <https://doi.org/10.1016/j.jaridenv.2020.104257>.
- Yao Z, Xiao J and Ma X** (2021) The impact of large-scale afforestation on ecological environment in the Gobi region. *Scientific Reports* **11**. <https://doi.org/10.1038/s41598-021-93948-5>.
- Zhang YK and Hartemink AE** (2017) Sampling designs for soil organic carbon stock assessment of soil profiles. *Geoderma* **307**, 220–230. <https://doi.org/10.1016/j.geoderma.2017.08.013>.
- Zheng X, Zhu J and Xing Z** (2016) Assessment of the effects of shelterbelts on crop yields at the regional scale in Northeast China. *Agricultural Systems* **143**, 49–60. <https://doi.org/10.1016/j.agsy.2015.12.008>.
- Zohary M** (1962) *Plant Life of Palestine: Israel and Jordan*. Jerusalem: The Hebrew University Press.
- Zohary M** (1971) *The Phytogeographical Foundations of the Middle East*. Stuttgart: Gustav Fischer Verlag.
- Zohary M** (1973) *Geobotanical Foundations of the Middle East*. Stuttgart: Gustav Fischer Verlag.