

Research Article

Cite this article: Anettová L, Šípková A, Izquierdo-Rodríguez E, Velič V, Modrý D (2023). Rat lungworm survives winter: experimental overwintering of *Angiostrongylus cantonensis* larvae in European slugs. *Parasitology* **150**, 950–955. <https://doi.org/10.1017/S0031182023000781>

Received: 16 February 2023
Revised: 24 July 2023
Accepted: 7 August 2023
First published online: 24 August 2023




Keywords:

Angiostrongylus cantonensis; invasive nematode; *Limax maximus*; overwintering

Corresponding author:

David Modrý; Email: modrydav@sci.muni.cz

Rat lungworm survives winter: experimental overwintering of *Angiostrongylus cantonensis* larvae in European slugs

Lucia Anettová¹ , Anna Šípková¹ , Elena Izquierdo-Rodríguez^{2,3}, Vivienne Velič⁴ and David Modrý^{1,5,6} 

¹Department of Botany and Zoology, Faculty of Science, Masaryk University, Brno, Czech Republic; ²Instituto Universitario de Enfermedades Tropicales y Salud Pública de Canarias, Universidad de La Laguna, La Laguna, Canary Islands, Spain; ³Department of Obstetrics and Gynecology, Pediatrics, Preventive Medicine and Public Health, Toxicology, Legal and Forensic Medicine and Parasitology, Universidad de La Laguna, La Laguna, Canary Islands, Spain; ⁴University of Veterinary Sciences Brno, Palackého tř. 1946/1, 612 42 Brno, Czech Republic; ⁵Institute of Parasitology, Biology Center of Czech Academy of Sciences, Ceske Budejovice, Czech Republic and ⁶Department of Veterinary Sciences, Faculty of Agrobiological, Food and Natural Resources/CINeZ, Czech University of Life Sciences Prague, Prague, Czech Republic

Abstract

The rat lungworm *Angiostrongylus cantonensis* is a metastrongyloid nematode that causes neurological disorders in its accidental hosts, including humans. This invasive pathogen is native to Southeast Asia and adjacent regions and is gradually expanding its distribution to tropical and subtropical areas with new foci discovered near temperate regions. The parasite has a complex life cycle with a range of gastropods serving as intermediate hosts. A broad spectrum of poikilotherm vertebrates and invertebrates can serve as paratenic hosts. Since it has already been demonstrated that other, non-zoonotic metastrongyloids can survive in their intermediate hosts during the winter, the aim of our study was to evaluate the survival of *A. cantonensis* third-stage larvae in experimentally infected slugs (*Limax maximus*) kept at 4.5–7°C for 60 days. Third-stage larvae of *A. cantonensis* survived the period of low temperature and remained capable of infecting definitive hosts (laboratory rats) afterwards, even though their numbers dropped significantly. These results suggest that further spread to higher latitudes or altitudes is possible in areas with sufficient abundance of definitive hosts, since low winter temperatures are not necessarily an obstacle to the spread of the parasite.

Introduction

The rat lungworm, *Angiostrongylus cantonensis* (Chen, 1935) is a metastrongyloid nematode widely known for its zoonotic potential and unusually broad range of paratenic and accidental hosts. Besides humans and domestic animals as its well-described accidental hosts, the parasite has been reported as causing severe neurological disorders in wildlife. In all these accidental hosts, the parasite can lead to paralysis or death (Kuberski and Wallace, 1979; Lindo *et al.*, 2004; Monks *et al.*, 2005; Burns *et al.*, 2014; Cowie, 2017; Odani *et al.*, 2021). In most cases, humans get infected with *A. cantonensis* by eating raw or undercooked intermediate or paratenic hosts that contain infectious L3 larvae (Wang *et al.*, 2012). *Angiostrongylus cantonensis* originated in Southeast Asia and adjacent regions (Chen, 1935; Lv *et al.*, 2018), but is now well established in tropical and subtropical regions around the globe (Prociv and Carlisle, 2001; Caldeira *et al.*, 2007; Stockdale Walden *et al.*, 2017). In temperate regions, 6 species of the genus *Angiostrongylus* and 4 other species of closely related genera (*Rodentocaulus* spp., *Stefanskostrongylus* spp.) have been described (Cowie, 2019), and only one of them (*A. dujardini*) is known to harm animals other than its definitive hosts (Graille *et al.*, 2015; Eleni *et al.*, 2016). *Angiostrongylus cantonensis* cycles between definitive (several rat species mainly of the tribe Rattini) and intermediate (terrestrial and aquatic gastropods) hosts (Alicata, 1965; Yong and Eamsobhana, 2013). In addition, various poikilotherm vertebrates as well as invertebrates serve as paratenic hosts (Wallace and Rosen, 1967; Ash, 1968; Radomyos *et al.*, 1994; Wang *et al.*, 2018; Chaisiri *et al.*, 2019; Turck *et al.*, 2022).

Factors contributing to the emergence of *A. cantonensis* and its expanding distribution include climate change, global transport and the resulting facilitated movement of hosts (rodents and gastropods) and, finally, changes in population dynamics of hosts and cultural habits in endemic areas (York *et al.*, 2015; Cowie, 2017). Findings of new *A. cantonensis* foci in the last 2 decades show not only continuous extension of its distribution, but also an expansion into temperate regions of Europe and USA (Campbell and Little, 1988; Foronda *et al.*, 2010; Paredes-Esquivel *et al.*, 2019; Walden *et al.*, 2021; Delgado-Serra *et al.*, 2022; Galán-Puchades *et al.*, 2022). York *et al.* (2014) predicted an increase of suitable habitats for *A. cantonensis* within Europe.

There is an extensive range of metastrongyloid nematode species that thrive in the temperate realm. (González *et al.*, 2007; Ferdushy *et al.*, 2009; Majoros *et al.*, 2010; Traversa and Di

© The Author(s), 2023. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

Cesare, 2014; Čabanová *et al.*, 2018; Morgan *et al.*, 2021). The ability of their larvae to survive or even continue development in gastropods during the winter months in temperate climate has been experimentally demonstrated in *Troglostrongylus brevior* (Gerichter, 1948; Morelli *et al.*, 2020) and *Aelurostrongylus abstrusus* (Morelli *et al.*, 2021). To our knowledge, there have been no studies of *A. cantonensis* survival in overwintering intermediate hosts. We hypothesized and experimentally tested that *A. cantonensis* larvae can survive low winter temperatures in gastropods, which may be one of the key factors enabling their future expansion deeper into temperate regions.

Materials and methods

The experimental strain of *A. cantonensis* was obtained in 2017 in Fatu Hiva, French Polynesia and is maintained in laboratory conditions circulating between Wistar rats and experimental gastropods (*Subulina octona* and *Biomphalaria glabrata*). The identity of the isolate was confirmed by the morphology of adult nematodes obtained from infected rats as well as by *cox1* sequencing, with the haplotype identified as part of *A. cantonensis* clade 2 (Červená *et al.*, 2019).

Adult great grey slugs (*Limax maximus* – LM) were collected from a private garden near Brno, Czech Republic during Autumn 2021. Slugs were kept in plastic boxes containing 8–10 individuals at 20–24°C throughout the experiment except for adaptation to cool temperature and the overwintering period of the experimental group. Boxes containing slugs were equipped with shelters and moss, and coconut soil was used as a bedding. Slugs were checked daily and fed with fresh lettuce and fish flakes every other day. One experimental group (E, 34 individuals) and 2 control groups (K1, 22 individuals and K2; 30 individuals) with a total of 86 LM individuals were infected by offering approximately 1 gram of fresh rat feces (maximum 12 hours old) to slugs in all experimental and control groups for 2 weeks (Fig. 1A). The number of larvae per gram of feces was determined by modified Baermann larvoscopy, such that approximately 40 000 L1 larvae per experimental box were provided during the 2-week period.

Experimental group E contained 34 slugs. After the exposure to rat feces containing L1 larvae, all slugs were left for 4 weeks to allow larvae to develop to the third stage. After that period, slugs in group E were gradually adapted to the decreasing temperature by exposing them to 15°C for two 5-day periods with a 7-day break between them at room temperature. After this adaptation period, slugs from group E were deprived of food for 3 days to empty the digestive tract and placed in a laboratory cool room with temperature between 4.5 and 7°C for 60 days. These conditions simulated the natural conditions in which these slugs overwinter, hidden in a shelter or deep underground, where the temperature rarely dip below 0°C (South, 2012). During this period, the slugs were kept undisturbed and without food, except for weekly checks. All the slugs from group E exhibited typical signs of hibernation, i.e. clustered together with no apparent physical activity. After this period, all the slugs from groups E and K2 were weighed, sacrificed by decapitation and individually digested artificially (i.e. 0.3 g pepsin and 100 mL 0.7% HCl on a magnetic stirrer set to 200 RPM and 37°C) (Modrý *et al.*, 2021). Motile L3 larvae were isolated, counted and used for the confirmatory bioassay on 2 infection-naïve Wistar laboratory rats to assess their infectivity. Each rat was infected with 60 live motile L3 larvae *via* orogastric tube under general inhalation anaesthesia with diethyl ether. After 45 days post infection, Baermann larvoscopy of rat feces was performed to detect L1 larvae.

Slugs from group K1 (22) were used for confirmation of infection success. These slugs were sacrificed 4 weeks after the infection period, artificially digested as described and the number of



Figure 1. Hosts and developmental stages of *A. cantonensis* in described experiments. **A.** Slugs *Limax maximus* feeding on fresh rat feces; **B.** *Angiostrongylus cantonensis* L3 larva collected from artificially digested slugs, scale bar = 100 μ m.

larvae per slug was counted. Throughout the experiments, larvae were microscopically confirmed as *A. cantonensis* based on typical morphology, i.e. tail length, shape of the termination of the tail and oesophagus length (Ash, 1970); possible presence of other metastrongylids was excluded using the same criteria (Diakou *et al.*, 2016).

Slugs from group K2 (30) did not overwinter and remained at room temperature (20–24°C) throughout the experimental period, to check that *A. cantonensis* larvae survive in this slug species for this amount of time at room temperature. Slugs were checked daily and fed every other day as before the experiment. The remains of slugs from this group that died spontaneously less than 4 weeks before the end of the experiment were immediately examined for *A. cantonensis* larvae by artificial digestion. The slugs from K2 that died spontaneously more than 30 days before the end of experiment were excluded. Remaining slugs were sacrificed 4 weeks after the infection period, artificially digested as

described above and the number of larvae per individual slug was counted.

Statistical analysis was done using ANOVA and the Tukey HSD test. The average, *P* value and standard deviation were calculated in R Studio and Microsoft Excel was used to generate a boxplot graph.

Results

All 34 slugs from the experimental (i.e. overwintering) group E survived the entire experiment. In the control group K1 (22 individuals), there were no mortalities either since the slugs were sacrificed just after the larval development to L3. Of 30 slugs in control group K2, 21 survived the entire experiment, 9 of them died during the last month of the experiment and were excluded from the experiment. Six more slugs from K2 group died spontaneously in the last month of the experiment and were processed as those that were sacrificed. The average weight of slugs at the end of the experiment in groups K1, K2 and E was 2.0, 2.2 and 2.3 g, respectively.

In the overwintering group (E), 0–17 motile third-stage metatroglyoid larvae (Fig. 1B) were collected from individual slugs (an average of 4.97 larvae per slug). In control groups K1 and K2, average numbers of larvae retrieved were 17.6 and 18.4, respectively, with a maximum of 61 larvae per slug in K1 and 63 larvae per slug in K2. Detailed information on the number of larvae in each slug are given in Table 1. The difference of L3 survival between the controls and overwintering group is significant ($F_{2,74} = 20.05, P < 0.001$) according to the Tukey HSD test and the difference between the 2 control groups is negligible ($P = 0.95$). Figure 2 is a boxplot graph of numbers of the third-stage larvae collected in each group.

Both Wistar laboratory rats that served as positive controls for the infectivity of larvae retrieved from the overwintering group E had L1 in their feces on day 45 post infection.

Discussion

Our study confirmed the ability of third-stage larvae of *A. cantonensis* to survive winter conditions in gastropods native to

Table 1. Numbers of live motile L3 of *A. cantonensis* retrieved from each individual slug *Limax maximus* from control groups K1 and K2 and in overwintering group E. The first six slugs from K2 group died spontaneously during the last month of the experiment.

K1		K2		E	
Slug no.	Number of larvae	Slug no.	Number of larvae	Slug no.	Number of larvae
1	20	†1	20	1	5
2	15	†2	62	2	3
3	2	†3	15	3	14
4	12	†4	5	4	5
5	20	†5	15	5	17
6	25	†6	4	6	5
7	19	7	9	7	0
8	1	8	17	8	1
9	3	9	11	9	2
10	8	10	10	10	2
11	13	11	3	11	1
12	61	12	15	12	6
13	11	13	12	13	3
14	10	14	18	14	7
15	24	15	6	15	5
16	52	16	14	16	11
17	20	17	7	17	4
18	7	18	34	18	13
19	6	19	24	19	1
20	8	20	63	20	1
21	18	21	18	21	1
22	32			22	11
				23	9
		9 slugs from K2 died spontaneously before January 2022		24	2
				25	0
				26	7
				27	2
				28	2

(Continued)

Table 1. (Continued.)

K1		K2		E	
Slug no.	Number of larvae	Slug no.	Number of larvae	Slug no.	Number of larvae
				29	10
				30	1
				31	5
				32	6
				33	2
				34	5
AVERAGE	17.6	AVERAGE	18.19	AVERAGE	4.97

Boxplot graph of L3 larvae amounts per individual slugs




Figure 2. Boxplot graph of numbers of third-stage larvae collected from each slug. The X marks the average value. The median is indicated by a line across the box. The whiskers on box plots show the ranges of Q1 and Q4 up to the most extreme data points. K1 and K2 are control groups, E is the experimental overwintering group of slugs.

temperate Europe and remain infective for the definitive hosts. This result, together with the relatively long patent period in rats (Mackerras and Sandars, 1955), suggests that the range could further extend in Europe into areas with milder winters and sufficient abundance of definitive hosts.

Even though the collected larvae were infective for rats as definitive hosts, we observed a decrease in numbers of live motile larvae in slugs during the overwintering period (average 4.97 larvae per slug in group E compared to 18.19 in group K2). We chose *L. maximus* as a model species since it is a synanthropic slug with a lifespan of 2–3 years, naturally overwintering in shelters where the temperature almost never falls below 0°C (South, 2012). The highest mortality observed in group K2, which was kept at room temperature, may be attributable to their maladaptation to captive conditions, since overwintering is natural for these slugs. LM has been described as an intermediate host suitable for other metastrongyloid nematodes (Conboy *et al.*, 2017), although it usually carries relatively low numbers of L3 larvae (Ferdushy *et al.*, 2009). Other slug species in temperate Europe reported to host metastrongyloids also showed low abundance of larvae, barely reaching 100 larvae per individual (Lange

et al., 2018). For comparison, the numbers of *A. cantonensis* L3 recovered from tropical slugs may reach hundreds to thousands (or even millions) of larvae, depending on the species (Heyneman and Lim, 1967; Hollingsworth *et al.*, 2013; Kim *et al.*, 2014; Rollins *et al.*, 2021). Environmental temperature has an impact on activity of L1 larvae and their ability to develop in intermediate hosts (Ishii, 1984; Lv *et al.*, 2006). However, the massive load of *A. cantonensis* larvae in some tropical snails and slugs could also be related to differences in behaviour, food preferences and overall ecology of intermediate and definitive hosts.

Previous studies have shown that *A. cantonensis* larvae cannot develop to the third stage at less than 15°C (Ishii, 1984; Lv *et al.*, 2006), which is very similar to the case of *A. vasorum* (Ferdushy *et al.*, 2010). However, our results confirmed that already developed L3 larvae survive and maintain their infectivity like the latter species, which is widely distributed in carnivores in temperate Europe (Fuehrer *et al.*, 2021; Tieri *et al.*, 2021).

Our experiments indicate that once an infected mollusc as an intermediate host survives cold winter temperature, larvae of *A. cantonensis* most likely will stay infective within it for several

months. This result, together with the relatively long patent period in rats (Mackerras and Sandars, 1955) suggests further possible range expansion to higher latitudes or altitudes in areas with sufficient abundance of the definitive hosts as the low winter temperatures are not necessarily an obstacle to the spread of the parasite (until reaching certain very high latitude or altitude points).

Data availability. All data generated and analysed during this study are included in this published article.

Acknowledgements. We thank Kateřina Kintrová for her help with the statistical analysis. We also thank the Veterinary University, Brno for providing the facilities for the experiment.

Authors' contributions. Conceptualization was done by D.M. and L.A.; methodology by L.A. and D.M.; investigation was carried out by L.A, A.Š., E.I.R., V.V.; writing was by L.A, D.M.; visualization by L.A.; supervision by D.M. and project administration by D.M. All authors have read and agreed to the published version of the manuscript.

Financial support. The study was supported by SEAEUROPEJFS19IN-053 and Czech Science Foundation grant no. 22-26136S. Work of Lucia Anettová and Anna Šipkova was also supported by Specific research – support of student projects, n. MUNI/A/1488/2021. E. Izquierdo-Rodriguez was funded by the scholarship M-ULL (becas M-ULL, convocatoria 2019).

Competing interest. None.

Ethical standards. The authors declare that all ethical standards and requirements were fulfilled.

References

- Alicata JE (1965) Biology and distribution of the rat lungworm, *Angiostrongylus cantonensis*, and its relationship to eosinophilic meningoencephalitis and other neurological disorders of man and animals. *Advances in Parasitology* 3, 223–248.
- Ash LR (1968) The occurrence of *Angiostrongylus cantonensis* in frogs of New Caledonia with observations on paratenic hosts of metastrongyles. *Journal of Parasitology* 54, 432–436.
- Ash LR (1970) Diagnostic morphology of the third-stage larvae of *Angiostrongylus cantonensis*, *Angiostrongylus vasorum*, *Aelurostrongylus abstrusus*, and *Anafilaroides rostratus* (Nematoda: Metastrongyloidea). *Journal of Parasitology* 56, 249–253.
- Burns RE, Bicknese EJ, Qvarnstrom Y, DeLeon-Carnes M, Drew CP, Gardiner CH and Rideout BA (2014) Cerebral *Angiostrongylus cantonensis* infection in a captive African pygmy falcon (*Polihierax semitorquatus*) in southern California. *Journal of Veterinary Diagnostic Investigation* 26, 695–698.
- Čabanová V, Miterpáková M, Druga M, Hurníková Z and Valentová D (2018) GIS-based environmental analysis of fox and canine lungworm distribution: an epidemiological study of *Angiostrongylus vasorum* and *Crenosoma vulpis* in red foxes from Slovakia. *Parasitology Research* 117, 521–530.
- Caldeira RL, Mendonça CL, Goveia CO, Lenzi HL, Graeff-Teixeira C, Lima WS, Mota EM, Pecora IL, Medeiros AMZ and Carvalho ODS (2007) First record of molluscs naturally infected with *Angiostrongylus cantonensis* (Chen, 1935) (Nematoda: Metastrongylidae) in Brazil. *Memórias do Instituto Oswaldo Cruz* 102, 887–889.
- Campbell BG and Little MD (1988) The finding of *Angiostrongylus cantonensis* in rats in New Orleans. *American Journal of Tropical Medicine and Hygiene* 38, 568–573.
- Červená B, Modrý D, Fecková B, Hrazdilová K, Foronda P, Alonso AM, Lee R, Walker J, Neibuhr CR, Malik R and Šlapeta J (2019) Low diversity of *Angiostrongylus cantonensis* complete mitochondrial DNA sequences from Australia, Hawaii, French Polynesia and the Canary Islands revealed using whole genome next-generation sequencing. *Parasites & Vectors* 12, 1–13.
- Chaisiri K, Dusitsittipon S, Panitvong N, Ketboonlue T, Nuamtanong S, Thaenkham U, Morand S and Dekumyoy P (2019) Distribution of the newly invasive New Guinea flatworm *Platydemus manokwari* (Platyhelminthes: Geoplanidae) in Thailand and its potential role as a paratenic host carrying *Angiostrongylus malaysiensis* larvae. *Journal of Helminthology* 93, 711–719.
- Chen HT (1935) Un nouveau nematode Pulmonaire, *Pulmonema cantonensis*, n. g., n. sp. des rats de Canton. *Annals of Parasitology* 13, 312–317.
- Conboy G, Guselle N and Schaper R (2017) Spontaneous shedding of metastrongyloid third-stage larvae by experimentally infected *Limax maximus*. *Parasitology Research* 116, 41–54.
- Cowie RH (2017) *Angiostrongylus cantonensis*: agent of a sometimes fatal globally emerging infectious disease (rat lungworm disease). *ACS Chemical Neuroscience* 8, 2102–2104.
- Cowie R (2019) Annotated catalogue of species of *Angiostrongylus* and the related genera *Gallestrongylus*, *Rodentocaulus* and *Stefanskostrongylus* (Nematoda: Metastrongyloidea, Angiostrongylidae). *Journal of Helminthology* 93, 389–423.
- Delgado-Serra S, Sola J, Negre N and Paredes-Esquível C (2022) *Angiostrongylus cantonensis* Nematode Invasion Pathway, Mallorca, Spain. *Emerging Infectious Diseases* 28, 1163–1169.
- Diakou A, Psalla D, Migli D, Di Cesare A, Youlatos D, Marcer F and Traversa D (2016) First evidence of the European wildcat (*Felis silvestris silvestris*) as definitive host of *Angiostrongylus chabaudi*. *Parasitology Research* 115, 1235–1244.
- Eleni C, Di Cesare A, Cavicchio P, Tonnicchia MC, Meoli R, di Regalbano AF, Paoletti B, Pietrobelli M and De Liberato C (2016) Fatal *Angiostrongylus dujardini* infection in callitrichid monkeys and suricates in an Italian zoological garden. *Parasitology International* 65, 333–335.
- Ferdushy T, Kapel CMO, Webster P, Al-Sabi MNS and Grønbold J (2009) The occurrence of *Angiostrongylus vasorum* in terrestrial slugs from forests and parks in the Copenhagen area, Denmark. *Journal of Helminthology* 83, 379.
- Ferdushy T, Kapel CMO, Webster P, Al-Sabi MNS and Grønbold JR (2010) The effect of temperature and host age on the infectivity and development of *Angiostrongylus vasorum* in the slug *Arion lusitanicus*. *Parasitology Research* 107, 147–151.
- Foronda P, López-González M, Miquel J, Torres J, Segovia M, Abreu-Acosta N, Casanova JC, Valladares B, Mas-Coma S, Bargues MD and Feliu C (2010) Finding of *Parastrongylus cantonensis* (Chen, 1935) in *Rattus rattus* in Tenerife, Canary Islands (Spain). *Acta Tropica* 114, 123–127.
- Fuehrer HP, Morelli S, Unterköfler MS, Bajer A, Bakran-Lebl K, Dwuznik-Szarek D, Farkas R, Grandi G, Heddergott M, Jokelainen P, Knific T, Leschnik M, Miterpáková M, Modrý D, Petersen HH, Skírnisson K, Vergles Rataj A, Schnyder M and Strube C (2021) *Dirofilaria* spp. and *Angiostrongylus vasorum*: current risk of spreading in Central and Northern Europe. *Pathogens (Basel, Switzerland)* 10, 1268.
- Galán-Puchades MT, Gómez-Samblás M, Osuna A, Sáez-Durán S, Bueno-Marí R and Fuentes MV (2022) Autochthonous *Angiostrongylus cantonensis* lungworms in urban rats, Valencia, Spain, 2021. *Emerging Infectious Diseases* 28, 2564–2567.
- Gerichter CB (1948) Observation on the life history of lung nematodes using snails as intermediate hosts. *American Journal of Veterinary Research* 9, 109–112.
- González P, Carbonell E, Urios V and Rozhnov VV (2007) Coprology of *Panthera tigris altaica* and *Felis bengalensis euptilurus* from the Russian far east. *Journal of Parasitology* 93, 948–950.
- Graille M, Ferté H, Petit T, Ollivet Courtois F, Gauchot JY, Nougailon JL, Vitaud C, Wardzynski C and Lemberger K (2015) Fatal *Parastrongylus dujardini* infection in captive callitrichids. *Veterinary Pathology* 52, 364–368.
- Heyneman D and Lim BL (1967) *Angiostrongylus cantonensis*: proof of direct transmission with its epidemiological implications. *Science (New York, N.Y.)* 158, 1057–1058.
- Hollingsworth RG, Howe K and Jarvi SI (2013) Control measures for slug and snail hosts of *Angiostrongylus cantonensis*, with special reference to the semi-slug *Parmarion martensi*. *Hawai'i Journal of Medicine & Public Health* 72(6 Suppl 2), 75–80.
- Ishii AI (1984) Effects of temperature on the larval development of *Angiostrongylus cantonensis* in the intermediate host, *Biomphalaria glabrata*. *Zeitschrift für Parasitenkunde* 70, 375–379.
- Kim JR, Hayes KA, Yeung NW and Cowie RH (2014) Diverse gastropod hosts of *Angiostrongylus cantonensis*, the rat lungworm, globally and with a focus on the Hawaiian Islands. *PloS One* 9, e94969.
- Kuberski T and Wallace GD (1979) Clinical manifestations of eosinophilic meningitis due to *Angiostrongylus cantonensis*. *Neurology* 29, 1566–1570.

- Lange MK, Penagos-Tabares F, Hirtzmann J, Failing K, Schaper R, Van Bourgionie YR, Bäckeljaou T, Hermosilla T and Taubert A (2018) Prevalence of *Angiostrongylus vasorum*, *Aelurostrongylus abstrusus* and *Crenosoma vulpis* larvae in native slug populations in Germany. *Veterinary Parasitology* **254**, 120–130.
- Lindo JF, Escoffery CT, Reid B, Codrington G, Cunningham-Myrie C and Eberhard ML (2004) Fatal autochthonous eosinophilic meningitis in a Jamaican child caused by *Angiostrongylus cantonensis*. *American Journal of Tropical Medicine and Hygiene* **70**, 425–428.
- Lv S, Zhou XN, Zhang YI, Liu HX, Zhu D, Yin WG, Steinmann P, Wang XH and Jia TW (2006) The effect of temperature on the development of *Angiostrongylus cantonensis* (Chen 1935) in *Pomacea canaliculata* (Lamarck 1822). *Parasitology Research* **99**, 583–587.
- Lv S, Guo YH, Nguyen HM, Sinuon M, Sayasone S, Lo NC, Zhou XN and Andrews JR (2018) Invasive *Pomacea* snails as important intermediate hosts of *Angiostrongylus cantonensis* in Laos, Cambodia and Vietnam: implications for outbreaks of eosinophilic meningitis. *Acta Tropica* **183**, 32–35.
- Mackerras MJ and Sandars DF (1955) The life history of the rat lung-worm, *Angiostrongylus cantonensis* (Chen) (Nematoda: Metastrongylidae). *Australian Journal of Zoology* **3**, 1–21.
- Majoros G, Fukár O and Farkas R (2010) Autochthonous infection of dogs and slugs with *Angiostrongylus vasorum* in Hungary. *Veterinary Parasitology* **174**, 351–354.
- Modrý D, Fecková B, Putnová B, Manalo SM and Otranto D (2021) Alternative pathways in *Angiostrongylus cantonensis* (Metastrongyloidea: Angiostrongylidae) transmission. *Parasitology* **148**, 167–173.
- Monks DJ, Carlisle MS, Carrigan M, Rose K, Spratt D, Gallagher A and Prociw P (2005) *Angiostrongylus cantonensis* as a cause of cerebrospinal disease in a yellow-tailed black cockatoo (*Calyptorhynchus funereus*) and two tawny frogmouths (*Podargus strigoides*). *Journal of Avian Medicine and Surgery* **19**, 289–293.
- Morelli S, Traversa D, Colombo M, Raue K, Strube C, Pollmeier M and Di Cesare A (2020) The effect of the hibernation on the larval development of *Troglostrongylus brevior* in the land snail *Cornu aspersum*. *Veterinary Parasitology* **282**, 109123.
- Morelli S, Colombo M, Diakou A, Traversa D, Grillini M, Frangipane di Regalbono A and Di Cesare A (2021) The influence of temperature on the larval development of *Aelurostrongylus abstrusus* in the land snail *Cornu aspersum*. *Pathogens (Basel, Switzerland)* **10**, 960.
- Morgan ER, Modry D, Paredes-Esquivel C, Foronda P and Traversa D (2021) Angiostrongylosis in animals and humans in Europe. *Pathogens (Basel, Switzerland)* **10**, 1236.
- Odani J, Sox E, Coleman W, Jha R and Malik R (2021) First documented cases of canine neuroangiostrongyliasis due to *Angiostrongylus cantonensis* in Hawaii. *Journal of the American Animal Hospital Association* **57**, 42–46.
- Paredes-Esquivel C, Sola J, Delgado-Serra S, Riera MP, Negre N, Miranda MÁ and Jurado-Rivera JA (2019) *Angiostrongylus cantonensis* in North African hedgehogs as vertebrate hosts, Mallorca, Spain, October 2018. *Eurosurveillance* **24**, 1900489.
- Prociw P and Carlisle MS (2001) The spread of *Angiostrongylus cantonensis* in Australia. *Southeast Asian Journal of Tropical Medicine and Public Health* **32**, 126–128.
- Radomyos P, Tungtrongchitr A, Praewanich R, Khewwathan P, Kantangkul T, Junlananto P and Ayudhya SI (1994) Occurrence of the infective stage of *Angiostrongylus cantonensis* in the yellow tree monitor (*Varanus bengalensis*) in five provinces of Thailand. *Southeast Asian Journal of Tropical Medicine and Public Health* **25**, 498–500.
- Rollins RL, Cowie RH, Echaluse MV and Medeiros MC (2021) Host snail species exhibit differential *Angiostrongylus cantonensis* prevalence and infection intensity across an environmental gradient. *Acta Tropica* **216**, 105824.
- South A (2012) *Terrestrial Slugs: Biology, Ecology and Control*. Dordrecht: Springer Science & Business Media.
- Stockdale Walden HD, Slapcinsky JD, Roff S, Mendieta Calle J, Diaz Goodwin Z, Stern J, Corlett R, Conway J and McIntosh A (2017) Geographic distribution of *Angiostrongylus cantonensis* in wild rats (*Rattus rattus*) and terrestrial snails in Florida, USA. *PLoS One* **12**, e0177910.
- Tieri EE, Saletti MA, D'Angelo AR, Parisiani G, Pelini S, Cocco A, Teodoro G, Censo E, D'Alterio N, Latrofa MS, Otranto D and Pascucci I (2021) *Angiostrongylus vasorum* in foxes (*Vulpes vulpes*) and wolves (*Canis lupus italicus*) from Abruzzo region, Italy. *International Journal for Parasitology: Parasites and Wildlife* **15**, 184–194.
- Traversa D and Di Cesare A (2014) Cardio-pulmonary parasitic nematodes affecting cats in Europe: unraveling the past, depicting the present, and predicting the future. *Frontiers in Veterinary Science* **1**, 1–9. doi: 10.3389/fvets.2014.00011.
- Turck HC, Fox MT and Cowie RH (2022) Paratenic hosts of *Angiostrongylus cantonensis* and their relation to human neuroangiostrongyliasis globally. *One Health (Amsterdam, Netherlands)* **15**, 100426.
- Walden HDS, Slapcinsky J, Rosenberg J and Wellehan JFX (2021) *Angiostrongylus cantonensis* (rat lungworm) in Florida, USA: current status. *Parasitology* **148**, 149–152.
- Wallace GD and Rosen L (1967) Studies on eosinophilic meningitis. 4. Experimental infection of fresh-water and marine fish with *Angiostrongylus cantonensis*. *American Journal of Epidemiology* **85**, 395–402.
- Wang QP, Wu ZD, Wei J, Owen RL and Lun ZR (2012) Human *Angiostrongylus cantonensis*: an update. *European Journal of Clinical Microbiology & Infectious Diseases* **31**, 389–395.
- Wang H, Lu L, She D, Wen Z, Mo Z, Li J and Li H (2018) Eating centipedes can result in *Angiostrongylus cantonensis* infection: two case reports and pathogen investigation. *The American Journal of Tropical Medicine and Hygiene* **99**, 743.
- Yong HS and Eamsobhana P (2013) Definitive rodent hosts of the rat lung-worm *Angiostrongylus cantonensis*. *Raffles Bulletin of Zoology* **29** (Supplement 29), 111–115.
- York EM, Butler CJ and Lord WD (2014) Global decline in suitable habitat for *Angiostrongylus* (= *Parastrongylus*) *cantonensis*: the role of climate change. *PLoS One* **9**, e103831.
- York EM, Creecy JP, Lord WD and Caire W (2015) Geographic range expansion for rat lungworm in North America. *Emerging Infectious Diseases* **21**, 1234.