

Corrigendum

Bulletin of Entomological Research, Vol. 90(3), pp. 233–244

The following table replaces that which originally appeared in the above article.

Table 3. Mean lengths (\pm SE) in mm of nine characters of clones of *Myzus persicae* reared on different host-plants.

Clones	Host-plants	ant III*	base IV	pt	urs	hf	ht2	ls	mws	lc
T1	potato (n=20)	0.460 b (0.005)	0.132 ab (0.002)	0.499 ab (0.005)	0.126 a (0.001)	0.692 ab (0.006)	0.117 b (0.0006)	0.557 b (0.005)	0.057 b (0.001)	0.239 b (0.002)
	cabbage (n=20)	0.423 a (0.007)	0.123 a (0.001)	0.486 a (0.004)	0.127 a (0.001)	0.613 a (0.009)	0.113 a (0.001)	0.407 a (0.008)	0.054 a (0.001)	0.208 a (0.004)
	broad bean (n=20)	0.476 c (0.005)	0.135 b (0.001)	0.518 c (0.004)	0.126 a (0.001)	0.715 b (0.005)	0.120 c (0.0006)	0.559 b (0.004)	0.053 a (0.0006)	0.239 b (0.002)
	tobacco (n=20)	0.479 c (0.004)	0.135 b (0.001)	0.513 bc (0.006)	0.130 a (0.001)	0.732 b (0.006)	0.120 c (0.0006)	0.582 c (0.004)	0.055 a (0.001)	0.248 b (0.003)
T2	potato (n=18)	0.482 b (0.004)	0.134 a (0.001)	0.569 b (0.004)	0.128 a (0.001)	0.727 c (0.005)	0.129 c (0.001)	0.605 b (0.005)	0.054 b (0.001)	0.249 b (0.003)
	cabbage (n=20)	0.382 a (0.005)	0.131 b (0.0007)	0.512 a (0.005)	0.127 a (0.001)	0.573 b (0.006)	0.112 a (0.001)	0.456 a (0.006)	0.052 a (0.001)	0.203 a (0.003)
	broad bean (n=20)	0.483 b (0.005)	0.147 c (0.001)	0.521 a (0.008)	0.127 a (0.0005)	0.694 a (0.005)	0.118 b (0.001)	0.562 c (0.005)	0.052 ab (0.0006)	0.246 b (0.003)
	tobacco (n=20)	0.475 b (0.006)	0.143 d (0.002)	0.550 b (0.009)	0.127 a (0.001)	0.738 c (0.009)	0.123 c (0.001)	0.609 b (0.005)	0.054 b (0.0005)	0.270 c (0.003)
T3	potato (n=21)	0.450 c (0.005)	0.142 a (0.002)	0.549 b (0.006)	0.125 a (0.001)	0.651 d (0.007)	0.118 c (0.001)	0.532 d (0.005)	0.054 b (0.001)	0.244 c (0.003)
	cabbage (n=22)	0.369 a (0.006)	0.138 a (0.001)	0.515 a (0.004)	0.125 a (0.0005)	0.565 a (0.007)	0.111 a (0.001)	0.436 a (0.007)	0.050 a (0.001)	0.206 a (0.003)
	broad bean (n=19)	0.417 b (0.007)	0.139 a (0.002)	0.513 a (0.006)	0.122 b (0.001)	0.587 b (0.009)	0.109 ab (0.001)	0.474 c (0.006)	0.050 a (0.001)	0.208 a (0.004)
	tobacco (n=20)	0.424 b (0.003)	0.139 a (0.002)	0.526 a (0.009)	0.125 a (0.0005)	0.626 c (0.005)	0.113 b (0.0005)	0.497 b (0.009)	0.052 ab (0.001)	0.230 b (0.002)
P	potato (n=20)	0.470 b (0.003)	0.143 b (0.004)	0.541 c (0.004)	0.119 b (0.0006)	0.686 b (0.005)	0.119 b (0.001)	0.567 b (0.003)	0.055 b (0.0008)	0.229 b (0.001)
	cabbage (n=11)	0.376 a (0.007)	0.134 a (0.005)	0.488 a (0.008)	0.117 a (0.0006)	0.544 a (0.007)	0.111 a (0.001)	0.424 a (0.006)	0.051 a (0.001)	0.179 a (0.003)
	broad bean (n=22)	0.476 b (0.007)	0.141 b (0.006)	0.528 b (0.004)	0.119 b (0.0006)	0.680 b (0.01)	0.117 b (0.001)	0.554 b (0.009)	0.055 b (0.0006)	0.224 b (0.003)

T1 and T2: red androcyclic clones collected from tobacco from Anavra region. T3: green holocyclic clone collected from tobacco from Alexandria region. P: green holocyclic clone collected from peach, away from tobacco growing regions (Portaria, Volos).

For each clone mean values of specimens followed by different letter are significantly different ($P < 0.05$). Duncan's test was used

*See Table 2 for explanation of the characters.