

SUBJECT INDEX

| | |
|---|---------------------------------------|
| abundance gradients | 500 |
| abundances | 142, 169 |
| abundances, molecular | 179 |
| acceleration of clouds | 209 |
| acceleration of electrons | 319 |
| acceleration of cosmic rays | 75, 325 |
| adiabatic expansion | 34, 43, 69, 279, 435 |
| age estimates | 65, 353 |
| ambient density | 285 |
| amplification of magnetic field | 235, 267, 331 |
| atomic clouds | 92, 205, 245, 455 |
| AXAF | 112 |
| Balmer line emission | 162, 415 |
| barrel-shaped SNRs | 270, 321, 359 |
| blast wave models | 145 |
| blast waves | 205, 517 |
| blue supergiants | 27 |
| bow shocks | 101, 311 |
| Brackett γ emission | 391, 395 |
| bubbles, ionized | 92 |
| bubbles, stellar wind | 95, 217, 278, 477 |
| C-type shocks | 172 |
| Cambridge low-frequency synthesis telescope | 51 |
| carbon deflagration | 31, 149, 151 |
| cavity, HII region | 278 |
| cavity, interstellar HI | 245 |
| cavity, re-energized | 245 |
| cavity, stellar wind | 233 |
| CCD imaging | 197 |
| centre-filled remnants | 99, 115, 219, 269, 297, 305, 335, 347 |
| charge exchange | 162 |
| circumstellar density | 33 |
| circumstellar medium | 15, 69, 235 |
| cirrus clouds | 498 |
| cirrus infrared | 365 |
| [C I] emission | 8 |
| classification of supernovae | 1 |
| [Cl II] emission | 8 |

| | |
|--|---------------------------------------|
| cloud compression | 208 |
| cloud crushing | 495 |
| cloud evaporation | 212 |
| cloud-cloud collisions | 495 |
| cloudy interstellar medium | 231 |
| clumpy ejecta | 227 |
| clumpy medium | 223 |
| CO core | 27 |
| CO emission | 249, 257, 261, 275, 399, 473 |
| CO filaments | 257 |
| collapse, gravitational | 211, 509 |
| collisional equilibrium ionization (CEI) | 105 |
| collisional heating of dust | 367, 385 |
| collisionless shocks | 162, 312 |
| composite supernova remnants | 100, 253, 269, 335, 347 |
| Compton scattering | 15 |
| condensational instabilities | 439 |
| conduction of heat | 34 |
| conduction, thermal | 173 |
| cooling parameters | 422 |
| cooling, molecular | 168 |
| coronal line emission | 407, 411 |
| cosmic ray acceleration | 75, 325 |
| cosmic ray ionization | 179 |
| cosmic rays | 309, 315, 405 |
| Coulomb collision losses | 315 |
| Crab-like remnants | 99, 115, 219, 269, 297, 305, 335, 347 |
| density scale height | 75 |
| depletion of heavy elements | 170 |
| detection of new remnants | 197, 201, 293, 339 |
| detection of young remnants | 279 |
| dissociation of H ₂ | 168 |
| distance estimates | 5, 121, 239, 253 |
| DRAO synthesis telescope | 245, 270, 297, 301, 339 |
| dust grain cooling | 383 |
| dust grains | 363 |
| dust grains, collisionally heated | 383 |
| dust temperature | 370 |
| dust, radiatively heated | 403 |
| dust, shock heated | 387, 403 |
| Effelsberg 100-m telescope | 254, 293 |
| Einstein focal plane crystal spectrometer (FPCS) | 102 |
| Einstein observatory | 99, 115, 125, 129, 137, 142, 153, 157 |
| Einstein solid state spectrometer (SSS) | 102 |
| ejecta, stellar | 33, 59, 65, 191 |
| ejectum | 235 |
| electron acceleration | 319 |
| equilibration of ion-electron energy | 145 |
| eschelle spectroscopy | 193 |
| ESO 3.6-m telescope | 391 |
| ESO-CTIO telescope | 189 |

| | |
|--|---------------------------------------|
| evaporation | 101 |
| evaporation of clouds | 212 |
| evolution of remnants | 216, 231, 376, 419, 435 |
| EXOSAT observatory | 99, 119, 137, 142 |
| EXOSAT gas scintillation proportional counter (GSPC) | 103 |
| expansion of SNRs | 43, 51, 69, 279, 435 |
| explosion energy E_0 | 121, 142, 285 |
| Fabry-Perot imaging interferometry (TAURUS) | 443 |
| Fabry-Perot spectrophotometry | 411 |
| Faraday rotation | 47, 351, 355 |
| fast-moving knots | 6 |
| Fe abundance | 391 |
| Fe K-line emission | 102, 151 |
| Fe II absorption | 59 |
| [Fe II] emission | 391, 395 |
| [Fe X] emission | 145, 407, 411 |
| [Fe XIV] emission | 145, 411 |
| [Fe XVII] emission | 146 |
| Fe/H ratio | 494 |
| Fermi acceleration | 267, 310, 320 |
| filamentary radio emission | 301, 332 |
| filaments, CO emission | 257 |
| filaments, optical morphology | 219 |
| filaments, proper motions | 57, 65 |
| filaments, radial velocities | 56, 443 |
| filled-centre remnants | 99, 115, 219, 269, 297, 305, 335, 347 |
| firehose instability | 314 |
| Fleurs synthesis telescope (FIRST) | 272 |
| flux densities of SNRs in infrared | 380 |
| free expansion | 43, 69, 219, 279 |
| galactic halos | 461 |
| Galactic spiral arms | 485 |
| galactic winds | 493 |
| gas dynamics | 161 |
| gas-grain collisions | 368 |
| Giant HII-regions | 201 |
| Giant Molecular Clouds (GMC) | 469 |
| Ginga satellite | 151 |
| Gould's Belt | 493 |
| grain destruction | 166, 367, 397 |
| grain size distribution | 374 |
| grain-grain, grain-ion collisions | 166 |
| gravitational collapse | 211, 509 |
| gravitational instabilities | 450 |
| gravity wave instabilities | 173 |
| gyro-radius, protons | 313 |
| HI absorption | 239, 275 |
| HI clouds | 92, 205, 245, 455 |
| HI emission | 239, 245, 249, 253, 274, 341 |
| HI filaments, sheets | 92 |
| HI shells, giant | 473 |

| | |
|--|---------------------------------------|
| HII region cavity | 278 |
| HII regions | 96, 201, 485 |
| H α emission | 3, 24, 55, 74, 193, 197, 415, 443 |
| H α /[N II] ratio | 55 |
| H β emission | 391, 403 |
| H ₂ emission | 391, 395, 399 |
| Hat Creek Observatory 6-m telescope | 473 |
| HCO ⁺ emission | 399 |
| heat conduction | 34, 43 |
| hydrodynamic code | 69, 149, 235, 405, 461, 517 |
| hydrodynamic evolution | 137 |
| hydrodynamic models | 91, 133, 141 |
| hydrodynamical simulations | 435, 477 |
| InfraRed Astronomical Satellite (IRAS) | 363, 379, 383 |
| infrared cooling | 368 |
| infrared emission | 363 |
| infrared flux densities of SNRs | 380 |
| infrared morphology of SNRs | 373 |
| Infrared Space Observatory (ISO) | 387 |
| infrared spectroscopy | 366, 391, 395, 403 |
| infrared survey of SNRs | 379 |
| inhomogeneities, small scale | 426 |
| inhomogeneous medium | 205, 223 |
| initial mass function | 495, 517 |
| instabilities | 43 |
| dynamical | 450, 509 |
| firehose | 314 |
| gravitational | 450, 509 |
| gravity wave | 173 |
| Kelvin-Helmholtz | 210 |
| oscillational and condensational | 439 |
| radiative | 429 |
| Rayleigh-Taylor | 74, 173, 210, 233, 450, 463, 478, 512 |
| Vishniac | 173, 450, 461, 512 |
| interplanetary shocks | 311 |
| interstellar cavity | 245 |
| interstellar medium | 73, 91, 205, 223, 227, 231, 253, |
| interstellar pressure | 75 |
| ionization, non-equilibrium | 155 |
| IR shocked dust emission | 233 |
| IRAS observations | 227, 363, 379, 383 |
| IRAS telescope | 363, 379, 383 |
| irregularities in density | 223 |
| Isaac Newton 2.5-m telescope | 200, 443 |
| Itapetinga 13.7-m telescope | 305 |
| IUE observatory | 3, 187 |
| J-type shocks | 162, 172 |
| jets in SNRs | 273, 353 |
| knots, fast-moving | 173 |
| KPNO 2.1-m telescope | 26, 183 |
| KPNO 4-m telescope | 55, 197 |

| | |
|---|---|
| Larmor radius | 314 |
| luminosity of SNRs | 435 |
| magnetic field amplification | 235, 267, 331 |
| magnetic field, intrinsic direction | 351 |
| magnetic fields | 70, 76, 163, 215, 223, 311, 331, 355, 405, 456 |
| magnetic precursors | 170 |
| mass exchange in the ISM | 212 |
| mass injection in SNRs | 216 |
| mass loss, galactic | 517 |
| mass loss, stellar | 1, 15, 95, 217, 232 |
| metallicity | 500, 517 |
| molecular clouds | 96, 179, 205, 249, 257, 261, 265, 399, 473, 493 |
| molecular gas | 477 |
| Molonglo observatory synthesis telescope (MOST) | 270, 359 |
| morphology of SNRs | 265, 270, 331, 359, 373 |
| “mystery spot” of SN1987a | 322 |
| [N II] emission | 5, 47 |
| N-D relation | 276, 285 |
| Nagoya 4-m telescope | 259, 261 |
| NASA/Goddard 36-in telescope | 407 |
| Ne VIII emission | 146 |
| Neon plasma | 125 |
| neutrinos | 11 |
| neutron stars | 100, 157, 343 |
| [Ni II] emission | 183 |
| Nobeyama 45-m telescope | 257, 293 |
| non-equilibrium ionization (NEI) | 105, 137, 369 |
| non-luminous matter | 517 |
| non-radiative shocks | 74, 207, 415 |
| non-thermal emission | 309 |
| non-thermal X-ray spectrum | 99 |
| novae | 47 |
| numerical models of supershell dynamics | 461 |
| numerical simulations | 43, 69, 223 |
| numerical simulations of shocks | 312 |
| O I recombination lines | 8, 187 |
| [O I] emission | 411 |
| [O II] emission | 183 |
| [O III] emission | 65, 187, 195, 415, 443 |
| O VII resonance, forbidden lines | 154 |
| O VIII $\text{Ly}\alpha, \beta$ | 154 |
| O/Fe ratio | 494 |
| O-star progenitor | 231 |
| O_f stars | 95 |
| OB associations | 95, 447, 461 |
| Ooty synthesis radio telescope (OSRT) | 265 |
| oscillational instabilities | 439 |
| oxygen-rich supernova remnants | 187 |
| Palomar Observatory 60-in telescope | 415 |
| Parkes interferometer | 275 |
| particle acceleration | 235, 309, 331 |

| | |
|---------------------------------|--|
| photoevaporation | 217 |
| photoionization, UV and X-ray | 59 |
| planetary bow shocks | 311 |
| planetary nebulae | 47 |
| plerions | 99, 115, 219, 269, 297, 305, 335, 347 |
| polarization, radio | 47, 223, 274, 349, 351 |
| Population I stars | 447 |
| porosity of the ISM | 327, 454 |
| post-shock temperatures | 421 |
| Prandtl number | 212 |
| pre-supernova environments | 95 |
| pressure, cosmic ray | 77 |
| pressure, interstellar gas | 75, 513 |
| pressure, magnetic | 77 |
| progenitor bubble | 231 |
| progenitor cavities | 227 |
| progenitor processing | 217, 227 |
| proper motions of filaments | 57, 65 |
| pulsars | 100, 339, 343 |
| quasi-stationary flocculi (QSF) | 6, 96, 239 |
| radial velocities of filaments | 56, 443 |
| radiative cooling | 91, 419, 435 |
| radiative instabilities | 429 |
| radiative shocks | 74, 233, 439 |
| radiative shocks, stability | 439 |
| radiatively heated dust | 403 |
| radio continuum emission | 133, 269, 289, 293, 297, 301, 305, 339 |
| radio observations | 51, 265 |
| radio polarization | 223, 274 |
| radio supernovae | 23 |
| radioactive decay | 60 |
| Rankine-Hugoniot conditions | 163, 309, 316, 319 |
| rate of supernovae | 478, 481, 489 |
| Rayleigh-Taylor instabilities | 74, 173, 210, 233, 450, 463, 478, 512 |
| red supergiants | 15 |
| relativistic electrons | 267 |
| remnants, stellar | 157 |
| reverse shocks | 16, 35, 59, 100, 142, 151 |
| Reynold's number | 209, 211 |
| ring nebulae | 95 |
| ROSAT satellite | 99, 112 |
| rotation measure | 355 |
| SAX | 112 |
| scale height of galactic disk | 499 |
| Sedov-Taylor expansion | 34, 43, 69, 279, 435 |
| shell remnants | 100, 115, 335, 339, 347 |
| shells of ejecta | 335 |
| shock acceleration | 209 |
| shock chemistry | 179 |
| shock excitation | 161, 179, 391, 399 |
| shock heated dust | 227, 403 |

| | |
|--------------------------------|---------------------------------------|
| shock propagation | 223 |
| shock structure | 164, 309 |
| shock velocities | 5 |
| shock waves | 309, 517 |
| shocked HI | 248 |
| shocks, | |
| adiabatic | 162 |
| atomic gas | 161 |
| collisionless | 162, 312 |
| interplanetary | 311 |
| metal-rich | 172, 174 |
| MHD | 170 |
| molecular gas | 161, 179 |
| non-radiative | 74, 207, 415 |
| non-steady | 172 |
| oxygen-rich | 172 |
| planetary bow | 311 |
| quasi-parallel, -perpendicular | 309 |
| radiative | 74, 162, 233, 439 |
| reverse | 16, 35, 59, 100, 142, 151 |
| supernova remnant | 205 |
| $\Sigma - D$ relation | 276, 285, 289 |
| [S II] emission | 5, 24, 191, 193, 197 |
| [S II]/H α imagery | 203 |
| [S III] emission | 191 |
| simulations, numerical | 43, 69, 223 |
| snowplow, pressure-driven | 435 |
| spectral index, radio | 302, 305, 343, 348 |
| spectral line widths | 223 |
| Spectrosat | 112 |
| sputtering of dust grains | 166, 367 |
| stability of radiative shocks | 439 |
| star formation | 91, 456, 467, 469, 494, 509, 513 |
| star formation rate | 497, 513 |
| starburst galaxies | 203, 477 |
| statistical properties of SNRs | 276, 285, 289, 481, 485 |
| stellar ejecta | 33, 65, 191 |
| stellar remnant | 157 |
| stellar winds | 15, 95, 206, 217, 232, 337 |
| stellar wind bubbles | 95, 217, 278, 477 |
| stellar wind cavity | 233 |
| stochastic heating | 374 |
| Strömgren sphere | 217 |
| superbubbles | 73, 465 |
| superclouds | 469 |
| supernova explosions | 233, 477 |
| supernova remnants, | 1 |
| barrel morphology | 270, 321, 359 |
| composite | 100, 253, 269, 335, 347 |
| Crab-like | 99, 115, 219, 269, 297, 305, 335, 347 |
| distances | 5, 121, 239, 253 |

| | |
|---|---|
| supernova remnants | |
| distribution in Galaxy | 481, 485 |
| evolution | 31, 216, 231, 376, 419, 435 |
| newly detected | 197, 201, 293, 339 |
| searches | 201 |
| survey in infrared | 379 |
| total number in Galaxy | 483, 487 |
| supernovae, | 1 |
| collective effects | 493 |
| explosions | 27, 206, 469 |
| rate | 92, 478, 481, 489 |
| radio emission | 19, 23 |
| Type I | 1, 25, 43, 141, 489 |
| Type Ia | 31, 149 |
| Type Ib | 2, 6, 15, 31 |
| Type II | 1, 11, 15, 25, 95, 141, 447, 489 |
| Type V | 19, 23 |
| supershell dynamics | 448, 461 |
| supershells | 447, 451, 461, 465, 469 |
| synchrotron emission | 332, 405 |
| synchrotron nebulae | 115 |
| T-Tauri winds | 493 |
| TAURUS, Fabry-Perot imaging interferometer | 443 |
| temperatures, post-shock | 421 |
| Tenma satellite | 99, 103, 131, 137, 151 |
| thermal conduction | 173 |
| thermonuclear burning | 27 |
| turbulence | 70, 235 |
| United Kingdom InfraRed Telescope (UKIRT) | 395 |
| UV observations | 3 |
| Very Large Array (VLA) | 23, 47, 254, 270, 331, 335, 343, 357, 465 |
| Vishniac instabilities | 173, 450, 461, 512 |
| wave-particle interactions | 317 |
| Westerbork synthesis radio telescope (WSRT) | 239, 275, 465 |
| white dwarfs | 2 |
| wind heating | 91 |
| wind-blown bubbles | 95, 217, 278, 477 |
| winds, galactic | 493 |
| winds, stellar | 15, 95, 206, 217, 232, 337 |
| Wolf-Rayet stars | 2, 6, 32, 95 |
| X-ray emission | 133, 149, 407 |
| X-ray heating of dust | 363, 384 |
| X-ray images | 115, 157 |
| X-ray models | 145 |
| X-ray observations | 99, 125 |
| X-ray spectra | 119, 129, 137, 141, 153 |
| X-ray thermal spectrum | 100 |
| X-rays, generation of | 75 |
| XMM | 112 |
| ZEUS hydrodynamics code | 461 |
| zodiacal infrared emission | 365 |