

RESEARCH ARTICLE

Cuspidal ℓ -modular representations of $\mathrm{GL}_n(F)$ distinguished by a Galois involution

Robert Kurinczuk¹, Nadir Matringe^{2,3} and Vincent Sécherre⁴

¹School of Mathematical and Physical Sciences, University of Sheffield, Sheffield, S3 7RH, United Kingdom
E-mail: robkurinczuk@gmail.com (corresponding author).

²Institute of Mathematical Sciences, NYU Shanghai, Shanghai, China; E-mail: nrm6864@nyu.edu.

³Institut de Mathématiques de Jussieu-Paris Rive Gauche, Université Paris Cité, Paris, 75205, France;
E-mail: matringe@imj-prg.fr.

⁴Laboratoire de Mathématiques de Versailles, UVSQ, CNRS, Université Paris-Saclay, Versailles, 78035, France;
E-mail: vincent.secherre@uvsq.fr.

Received: 20 October 2023; **Revised:** 17 October 2024; **Accepted:** 6 November 2024

2020 Mathematics Subject Classification: *Primary* – 22E50; *Secondary* – 11F70, 11F85

Abstract

Let F/F_0 be a quadratic extension of non-Archimedean locally compact fields of residual characteristic $p \neq 2$ with Galois automorphism σ , and let R be an algebraically closed field of characteristic $\ell \notin \{0, p\}$. We reduce the classification of $\mathrm{GL}_n(F_0)$ -distinguished cuspidal R -representations of $\mathrm{GL}_n(F)$ to the level 0 setting. Moreover, under a parity condition, we give necessary conditions for a σ -self-dual cuspidal R -representation to be distinguished. Finally, we classify the distinguished cuspidal $\overline{\mathbb{F}}_\ell$ -representations of $\mathrm{GL}_n(F)$ having a distinguished cuspidal lift to $\overline{\mathbb{Q}}_\ell$.

Contents

1	Introduction	1
2	Notation	5
3	Basic results	7
4	Reduction to level zero	11
5	The odd case	27
6	Distinguished lift theorems	33

1. Introduction

1.1.

Let F/F_0 be a quadratic extension of non-Archimedean locally compact fields whose residual characteristic is a prime number p different from 2. Let σ be its nontrivial automorphism, and G be the general linear group $\mathrm{GL}_n(F)$ for some positive integer n . It is a totally disconnected, locally compact group, on which the involution σ acts componentwise, and the group G^σ of its σ -fixed points is equal to $\mathrm{GL}_n(F_0)$.

Now, fix an algebraically closed field R of characteristic different from p . A (smooth) representation π of G on an R -vector space V is said to be *distinguished* (by G^σ) if V carries a nonzero G^σ -invariant

linear form; more generally, if χ is a smooth character of G^σ with values in R^\times , the representation π is said to be χ -distinguished if V carries a nonzero linear form Λ such that

$$\Lambda(\pi(h)v) = \chi(h)\Lambda(v), \quad h \in G^\sigma, \quad v \in V.$$

1.2.

In the case where R is the field of complex numbers, distinguished irreducible representations of G have been extensively studied:

1. They are σ -self-dual, that is, the contragredient π^\vee of a distinguished irreducible representation π of G is isomorphic to its σ -conjugate π^σ ([16, 32, 33]) and their central character is trivial on F_0^\times ,
2. Any σ -self-dual discrete series representation of G is either distinguished or \varkappa -distinguished (\varkappa denotes the character of F_0^\times whose kernel is the subgroup of F/F_0 -norms), but not both: This is the dichotomy and disjunction theorem ([23, 2, 3]),
3. Distinguished generic irreducible representations of G are classified in terms of their cuspidal support ([4, 26, 27]),
4. Distinguished cuspidal representations of G are characterized in terms of their Galois parameter ([17]) and in terms of type theory (see [35] and below).

1.3.

Distinguished irreducible representations of G with coefficients in a field R of positive characteristic have been less well studied (see [3, 35, 25, 12]). As in the complex case, they are σ -self-dual and their central character is trivial on F_0^\times . For σ -self-dual *supercuspidal* representations, that is, irreducible representations which do not occur as subquotients of parabolically induced representations from a proper Levi subgroup, one has a dichotomy and disjunction theorem (see Paragraph 3.2). One also has a distinction criterion in terms of Galois parameters ([12] Proposition 3.15) and in terms of types ([35] Theorem 10.9). But there are explicit examples of σ -self-dual nonsupercuspidal *cuspidal* representations that are neither distinguished nor \varkappa -distinguished (as in [35] Remark 2.18) and of Steinberg representations that are both distinguished and \varkappa -distinguished ([12] Remark 1.9). Also, there is no known classification of distinguished cuspidal representations of $GL_n(F)$ for an arbitrary $n \geq 3$ (see [12] and Paragraph 4.13 below for $n = 2$).

In this paper, which can be considered as a sequel to [35], where all distinguished supercuspidal R -representations of G have been classified, we investigate the classification of distinguished cuspidal R -representations of G in terms of their supercuspidal support. We:

- reduce this classification to that of distinguished cuspidal representations of level 0, and from there to finite group theory (see Section 4),
- give a necessary condition of distinction for σ -self-dual cuspidal representations of G that satisfy a certain parity condition (see Section 5),
- classify the (distinguished, cuspidal) $\overline{\mathbb{F}}_\ell$ -representations of G having a distinguished cuspidal lift to $\overline{\mathbb{Q}}_\ell$, where $\overline{\mathbb{Q}}_\ell$ is an algebraic closure of the field of ℓ -adic numbers with residue field $\overline{\mathbb{F}}_\ell$.

Let us explain these results in more detail.

1.4.

Bushnell and Kutzko [9], in work extended to the modular setting by Vignéras [41], have given an explicit construction of a collection of pairs (\mathbf{J}, λ) called *extended maximal simple types* (which we will abbreviate to *types* here), consisting of a compact-mod-centre open subgroup \mathbf{J} of G and an irreducible R -representation λ of \mathbf{J} such that the representations $\text{ind}_{\mathbf{J}}^G(\lambda)$ are (irreducible and) cuspidal and such that every cuspidal R -representation of G appears in the collection of $\text{ind}_{\mathbf{J}}^G(\lambda)$.

We need the following invariants associated to a cuspidal R -representation of G following this explicit construction by compact induction (see Paragraphs 4.2 and 4.7):

1. the *endo-class* Θ : a fine refinement of the level introduced by Bushnell–Henniart in [6] and which applies equally well to the modular setting,
2. the *tame parameter field* T : a tamely ramified extension of F of degree dividing n , uniquely determined up to F -isomorphism by Θ ,
3. the *relative degree* m : a positive integer such that $m[T : F]$ divides n , uniquely determined by Θ and n .

Suppose further that Θ is σ -self-dual (which follows if for example the cuspidal representation itself is σ -self-dual), then there is a uniquely determined tamely ramified extension T_0 of F_0 contained in T such that T is isomorphic to $T_0 \otimes_{F_0} F$. The Galois group of T/T_0 canonically identifies with that of F/F_0 , and the unique nontrivial automorphism of T/T_0 extending σ will be denoted by σ (see §4.3). Our main theorem on reduction to the level 0 setting is then (see Theorem 4.41):

Theorem 1.1. 1. *There is a natural bijection*

$$\pi \mapsto \pi_t \tag{1.1}$$

from the set of isomorphism classes of cuspidal representations of G with endo-class Θ to the set of isomorphism classes of cuspidal representations of level 0 of $\mathrm{GL}_m(T)$.

2. *The representation π is σ -self-dual if and only if π_t is σ -self-dual.*
3. *The representation π is $\mathrm{GL}_n(F_0)$ -distinguished if and only if π_t is $\mathrm{GL}_m(T_0)$ -distinguished.*

The map (1.1) is also compatible with supercuspidal support; see Proposition 4.43 for a precise statement.

1.5.

Let us briefly explain how the map (1.1) above is defined. Let (\mathbf{J}, λ) be a type inducing a cuspidal representation π of G with σ -self-dual endo-class Θ , tame parameter field T and relative degree m . Then:

- the group \mathbf{J} has a unique maximal compact subgroup \mathbf{J}^0 , and a unique maximal normal pro- p subgroup \mathbf{J}^1 ,
- there is a group isomorphism $\mathbf{J}^0/\mathbf{J}^1 \simeq \mathrm{GL}_m(I)$, where I is the residue field of T ,
- the restriction of λ to \mathbf{J}^1 is isotypic for an irreducible representation η of \mathbf{J}^1 , and this representation η extends (noncanonically) to \mathbf{J} ,
- the choice of a representation κ of \mathbf{J} extending η determines a decomposition $\lambda \simeq \kappa \otimes \tau$, where τ is a representation of \mathbf{J} trivial on \mathbf{J}^1 , uniquely determined up to isomorphism.

The fact that Θ is σ -self-dual implies that there is a preferred choice for (\mathbf{J}, λ) : The group \mathbf{J} is fixed by σ , the representation η is σ -self-dual and there exists a natural isomorphism between the space of G^σ -invariant linear forms on π and that of $\mathbf{J} \cap G^\sigma$ -invariant linear forms on λ . Such a type is called *generic* (see Definition 4.31). We prove (see Proposition 4.16):

Proposition 1.2. *The representation η has a unique extension κ to \mathbf{J} which is both σ -self-dual and $\mathbf{J} \cap G^\sigma$ -distinguished and whose determinant has order a power of p .*

The choice of the representation κ given by Proposition 1.2 thus uniquely determines a representation τ of \mathbf{J} trivial on \mathbf{J}^1 .

Now, there is a natural choice, as explained in §4.10, of a σ -fixed maximal compact subgroup \mathbf{J}_t^0 of $\mathrm{GL}_m(T)$, with normalizer \mathbf{J}_t and pro- p -radical \mathbf{J}_t^1 , such that there is a σ -equivariant group isomorphism:

$$\mathbf{J}/\mathbf{J}^1 \simeq \mathbf{J}_t/\mathbf{J}_t^1.$$

The representation τ then defines a representation of \mathbf{J}_t trivial on \mathbf{J}_t^1 , denoted τ_t . The cuspidal representation π_t associated with π by the bijection (1.1) is then the compact induction of τ_t to $\mathrm{GL}_m(T)$.

1.6.

Having reduced the classification of distinguished cuspidal R -representations to level 0, we further reduce this classification to the finite group setting. Let π be a σ -self-dual cuspidal R -representation of G of level 0 with central character c_π and generic type (\mathbf{J}, λ) . Restricting λ to \mathbf{J}^0 defines a cuspidal R -representation V of $\mathrm{GL}_n(\mathbf{k})$, where \mathbf{k} is the residue field of F . We prove (see Theorem 4.45):

Theorem 1.3. *Suppose $n \neq 1$. The representation π is $\mathrm{GL}_n(F_0)$ -distinguished if and only if its central character c_π is trivial on F_0^\times and*

1. *if F/F_0 is unramified, then V is $\mathrm{GL}_n(\mathbf{k}_0)$ -distinguished (\mathbf{k}_0 is the residue field of F_0);*
2. *if F/F_0 is ramified, then n is even, V is $\mathrm{GL}_{n/2}(\mathbf{k}) \times \mathrm{GL}_{n/2}(\mathbf{k})$ -distinguished and*

$$s = \begin{pmatrix} 0 & \mathrm{id} \\ \mathrm{id} & 0 \end{pmatrix} \in \mathrm{GL}_n(\mathbf{k})$$

acts on the space of its $\mathrm{GL}_{n/2}(\mathbf{k}) \times \mathrm{GL}_{n/2}(\mathbf{k})$ -invariant linear forms by the sign $c_\pi(\varpi)$, where ϖ is any uniformizer of F .

As an application, we classify all distinguished cuspidal representations of $\mathrm{GL}_2(F)$ (see Paragraph 4.13).

1.7.

Let π be a cuspidal nonsupercuspidal R -representation of G . Following [29], we recall in Paragraph 3.4 that there are a uniquely determined integer $r = r(\pi) \geq 2$ and a supercuspidal R -representation ρ of $\mathrm{GL}_{n/r}(F)$ such that π is isomorphic to $\mathrm{St}_r(\rho)$, where $\mathrm{St}_r(\rho)$ denotes the unique generic subquotient of the parabolically induced representation

$$\rho \nu^{-(r-1)/2} \times \dots \times \rho \nu^{(r-1)/2}$$

(where ν denote the unramified character which is the absolute value of F composed with the determinant). The representation ρ is not unique in general, but, if π is σ -self-dual and r is odd, and if one further demands that ρ be σ -self-dual, then ρ is uniquely determined up to isomorphism (see Proposition 3.8). In this case, we obtain further necessary conditions for distinction (see Theorem 5.1):

Theorem 1.4. *Let π be a σ -self-dual cuspidal nonsupercuspidal R -representation of $\mathrm{GL}_n(F)$. Assume that the integer $r = r(\pi)$ is odd, thus π is isomorphic to $\mathrm{St}_r(\rho)$ for a uniquely determined σ -self-dual supercuspidal representation ρ of $\mathrm{GL}_{n/r}(F)$. If π is $\mathrm{GL}_n(F_0)$ -distinguished, then*

1. *the relative degree $m = m(\pi)$ and the ramification index of T/T_0 have the same parity,*
2. *the representation ρ is $\mathrm{GL}_{n/r}(F_0)$ -distinguished.*

As a corollary, we extend the disjunction theorem from the supercuspidal setting (that is, the statement that, if $\ell \neq 2$, a supercuspidal R -representation is not both distinguished and \varkappa -distinguished) to include cuspidal R -representations π with $r(\pi)$ odd.

1.8.

Say that an irreducible $\overline{\mathbb{F}}_\ell$ -representation π of G lifts to $\overline{\mathbb{Q}}_\ell$ if there exists a free $\overline{\mathbb{Z}}_\ell$ -lattice L equipped with a linear action of G such that the $\overline{\mathbb{F}}_\ell$ -representation of G on $L \otimes \overline{\mathbb{F}}_\ell$ is isomorphic to π . When this is

the case, say that the smooth $\overline{\mathbb{Q}}_\ell$ -representation of G on $L \otimes \overline{\mathbb{Q}}_\ell$ is a *lift* of π to $\overline{\mathbb{Q}}_\ell$. By [41], any cuspidal $\overline{\mathbb{F}}_\ell$ -representation of G lifts to $\overline{\mathbb{Q}}_\ell$ and any of its lifts is cuspidal.

According to [25] (see Theorem 3.3), any cuspidal $\overline{\mathbb{F}}_\ell$ -representation of G having a G^σ -distinguished lift to $\overline{\mathbb{Q}}_\ell$ is G^σ -distinguished. The converse holds for supercuspidal $\overline{\mathbb{F}}_\ell$ -representations of G (see [35] and [12]), but there are examples of G^σ -distinguished $\overline{\mathbb{F}}_\ell$ -cuspidal representations of $\mathrm{GL}_2(F)$ with no distinguished lift if $\ell = 2$ (see Paragraph 4.13 and Remark 6.4). In the final section, we classify the G^σ -distinguished cuspidal $\overline{\mathbb{F}}_\ell$ -representations of G having a G^σ -distinguished cuspidal lift to $\overline{\mathbb{Q}}_\ell$ (see Propositions 6.1 and 6.2 for a precise statement).

1.9.

Finally, let us comment on how the main results of this article could be pushed further.

Let π be a σ -self-dual cuspidal nonsupercuspidal $\overline{\mathbb{F}}_\ell$ -representation of G . In the case when $r(\pi)$ is even and $\ell > 2$, we conjecture that π is distinguished if and only if it has a distinguished lift to $\overline{\mathbb{Q}}_\ell$ (see Conjecture 6.3). We expect that this conjecture could be approached via the use of the Rankin–Selberg local factors developed in [24].

In the case when $r(\pi)$ is odd, we do not expect the necessary conditions of Theorem 1.4 to be sufficient for distinction. It would be interesting to determine whether any G^σ -distinguished cuspidal representation of G has a distinguished lift to $\overline{\mathbb{Q}}_\ell$.

Finally, this work is part of a wider programme aiming at classifying all distinguished generic R -representations of G . Such a classification might be approached by developing a theory of mod ℓ intertwining periods.

Structure of the paper

After setting some notation in Section 2, in Section 3 we collect together necessary background from the literature and prove some basic results on σ -self-dual cuspidal R -representations.

Section 4 constitutes the technical heart of the paper. It reduces the problem of classifying distinguished cuspidal R -representations to level 0.

In Section 5, under a parity condition, we provide necessary conditions for distinction, allowing us to deduce the disjunction theorem and a lifting theorem.

In Section 6, we classify the cuspidal $\overline{\mathbb{F}}_\ell$ -representations having a distinguished cuspidal lift.

2. Notation

2.1.

Given any non-Archimedean locally compact field F , we write \mathcal{O}_F for its ring of integers, \mathfrak{p}_F for the maximal ideal of \mathcal{O}_F , \mathbf{k}_F for its residue field and q_F for the cardinality of \mathbf{k}_F .

We also write val_F for the valuation of F taking any uniformizer to 1, and $|\cdot|_F$ for the absolute value of F taking any uniformizer to the inverse of q_F .

Given any finite extension L of K , we write $N_{L/K}$ and $\mathrm{tr}_{L/K}$ for the norm and trace maps.

2.2.

Given a locally compact, totally disconnected topological group G and an algebraically closed field R of characteristic different from p , we consider smooth representations of G on R -vector spaces. We will abbreviate *smooth R -representation* to *R -representation*, or even *representation* if the coefficient field R is clear from the context.

An *R -character* (or *character*) of G is a group homomorphism from G to R^\times with open kernel.

Let π be a representation of G . We write π^\vee for its contragredient. Given a character χ of G , we write π_χ for the representation $g \mapsto \chi(g)\pi(g)$ of G .

Let π be a representation of a closed subgroup H of G . Given any element $g \in G$, we write π^g for the representation $x \mapsto \pi(gxg^{-1})$ of $H^g = g^{-1}Hg$. Given any continuous involution σ of G , we write π^σ for the representation $\pi \circ \sigma$ of $\sigma(H)$. Given any character μ of $H \cap G^\sigma$, we say that π is μ -distinguished if the space $\text{Hom}_{H \cap G^\sigma}(\pi, \chi)$ is nonzero. If μ is the trivial character, we will abbreviate μ -distinguished to $H \cap G^\sigma$ -distinguished, or just distinguished.

2.3.

Let us fix a separable quadratic extension F/F_0 of non-Archimedean locally compact fields of residual characteristic p , and let σ denote its nontrivial automorphism. Let R be an algebraically closed field of characteristic different from p . Let

$$\varkappa = \varkappa_{F/F_0} : F_0^\times \rightarrow \{-1, 1\} = \mathbb{Z}^\times \tag{2.1}$$

denote the \mathbb{Z} -valued character of F_0^\times with kernel $N_{F/F_0}(F_0^\times)$. When needed, we will consider \varkappa as a character with values in R . We abbreviate $q = q_F$ and $q_0 = q_{F_0}$. We fix a square root

$$q_0^{1/2} \in R \tag{2.2}$$

of q_0 in R and define

$$q^{1/2} = \begin{cases} q_0^{1/2} & \text{if } F/F_0 \text{ is ramified,} \\ q_0 & \text{if } F/F_0 \text{ is unramified,} \end{cases} \tag{2.3}$$

which we will use to normalize parabolic induction and restriction functors (see below).

2.4.

Given a positive integer $n \geq 1$, the automorphism σ acts on the group $\text{GL}_n(F)$ componentwise, thus defines a continuous involution of $\text{GL}_n(F)$, still denoted σ . Its fixed points form the subgroup $\text{GL}_n(F_0)$.

We denote by ν the unramified character ‘absolute value of the determinant’ of $\text{GL}_n(F)$ and by $\nu^{1/2}$ the unramified character taking any element whose determinant has valuation 1 to $q^{-1/2}$. We thus have $(\nu^{1/2})^2 = \nu$. Similarly, we define the characters ν_0 and $\nu_0^{1/2}$ of $\text{GL}_n(F_0)$.

Given positive integers n_1, \dots, n_r such that $n_1 + \dots + n_r = n$ and, for each $i = 1, \dots, r$, given an R -representation π_i of $\text{GL}_{n_i}(F)$, we write

$$\pi_1 \times \dots \times \pi_r \tag{2.4}$$

for the representation of $\text{GL}_n(F)$ obtained by normalized parabolic induction from $\pi_1 \otimes \dots \otimes \pi_r$ along the parabolic subgroup generated by upper triangular matrices and the standard Levi subgroup $\text{GL}_{n_1}(F) \times \dots \times \text{GL}_{n_r}(F)$.

An irreducible R -representation of $\text{GL}_n(F)$ is said to be *cuspidal* (respectively, *supercuspidal*) if it does not occur as a subrepresentation (respectively, a subquotient) of any representation of the form (2.4) with $r \geq 2$. Any supercuspidal representation of $\text{GL}_n(F)$ is cuspidal. When R has characteristic 0, any cuspidal representation of $\text{GL}_n(F)$ is supercuspidal. When R has characteristic $\ell > 0$, the group $\text{GL}_n(F)$ may have cuspidal nonsupercuspidal representations (see §3.4).

Given a representation π of $\text{GL}_n(F)$ and a character χ of F^\times , we will write π_χ for $\pi(\chi \circ \det)$.

2.5.

Let us fix an algebraic closure $\overline{\mathbb{Q}}_\ell$ of the field of ℓ -adic numbers. Let $\overline{\mathbb{Z}}_\ell$ denote its ring of integers, and $\overline{\mathbb{F}}_\ell$ denote the residue field of $\overline{\mathbb{Z}}_\ell$.

We call an irreducible representation π of a locally compact, totally disconnected group G on a $\overline{\mathbb{Q}}_\ell$ -vector space V *integral* if it stabilizes a $\overline{\mathbb{Z}}_\ell$ -lattice L in V . In this case, we obtain a smooth $\overline{\mathbb{F}}_\ell$ -representation $L \otimes \overline{\mathbb{F}}_\ell$ of G whose isomorphism class may depend on the choice of L .

If G is either the group of rational points of a connected reductive linear algebraic F -group or a finite group (see [42, Theorem 1] and the Brauer–Nesbitt principle), the smooth $\overline{\mathbb{F}}_\ell$ -representation $L \otimes \overline{\mathbb{F}}_\ell$ has finite length, and its semisimplification is independent of the choice of L . This semisimplification is called the *reduction modulo ℓ* of π and is denoted by $r_\ell(\pi)$.

Given an irreducible $\overline{\mathbb{F}}_\ell$ -representation ρ of G , we call an irreducible integral $\overline{\mathbb{Q}}_\ell$ -representation with reduction modulo ℓ equal to ρ a *$\overline{\mathbb{Q}}_\ell$ -lift* of ρ .

3. Basic results

In this section, p is an arbitrary prime number, F/F_0 is a separable quadratic extension and R has characteristic $\ell \neq p$. We fix a positive integer $n \geq 1$.

3.1.

Fundamental results of Flicker and Prasad [16, 32, 33] on irreducible complex representations of $\mathrm{GL}_n(F)$ distinguished by $\mathrm{GL}_n(F_0)$ have been extended to irreducible R -representations in [35] Theorem 4.1.

Theorem 3.1. *Let π be an irreducible representation of $\mathrm{GL}_n(F)$ distinguished by $\mathrm{GL}_n(F_0)$.*

1. *The central character c_π of π is trivial on F_0^\times .*
2. *The R -vector space $\mathrm{Hom}_{\mathrm{GL}_n(F_0)}(\pi, R)$ has dimension 1.*
3. *The contragredient π^\vee of π is isomorphic to π^σ .*

We will say that a representation π of $\mathrm{GL}_n(F)$ is *σ -self-dual* if π^\vee is isomorphic to π^σ .

3.2.

For supercuspidal representations, we have the following dichotomy and disjunction theorem ([23] Theorem 4, [2] Corollary 1.6 if $\ell = 0$, [35] Theorem 10.8 if $p \neq 2$ and [12] Theorem 3.14 if $\ell \neq 0, 2$).

Theorem 3.2. *Let ρ be a σ -self-dual supercuspidal R -representation of $\mathrm{GL}_n(F)$.*

1. *If $\ell = 2$, then ρ is distinguished.*
2. *If $\ell \neq 2$, then ρ is either distinguished or κ -distinguished, but not both.*

3.3.

In this paragraph, ℓ is a prime number different from p and we will consider representations with coefficients in $\overline{\mathbb{Q}}_\ell$ or $\overline{\mathbb{F}}_\ell$. The following theorem is [25] Theorem 3.4.

Theorem 3.3. *Let π be an integral σ -self-dual cuspidal $\overline{\mathbb{Q}}_\ell$ -representation of $\mathrm{GL}_n(F)$. If π is distinguished by $\mathrm{GL}_n(F_0)$, then its reduction mod ℓ is (irreducible, cuspidal and) distinguished.*

It follows that any σ -self-dual cuspidal $\overline{\mathbb{F}}_\ell$ -representation of $\mathrm{GL}_n(F)$ having a distinguished lift to $\overline{\mathbb{Q}}_\ell$ is distinguished. For supercuspidal representations, one has the following converse (see [35] Theorem 10.11 if $p \neq 2$, and [12] Theorem 3.4):

Theorem 3.4. *Any $\mathrm{GL}_n(F_0)$ -distinguished supercuspidal $\overline{\mathbb{F}}_\ell$ -representation of $\mathrm{GL}_n(F)$ has a $\mathrm{GL}_n(F_0)$ -distinguished lift to $\overline{\mathbb{Q}}_\ell$.*

We also have the following distinguished lift theorem, making Theorem 3.4 more precise.

Theorem 3.5. *Let ρ be a σ -self-dual supercuspidal $\overline{\mathbb{F}}_\ell$ -representation of $\mathrm{GL}_n(F)$.*

1. The representation ρ has a σ -self-dual lift to $\overline{\mathbb{Q}}_\ell$.
2. Let μ be a σ -self-dual lift of ρ to $\overline{\mathbb{Q}}_\ell$, and suppose that $\ell \neq 2$. Then μ is distinguished if and only if ρ is distinguished.

Proof. If $p \neq 2$, this is [35] Theorem 10.11. Assume now that $p = 2$, thus $\ell \neq 2$.

By Theorem 3.2, the representation ρ is either distinguished or κ -distinguished. If it is distinguished, it has a σ -self-dual lift thanks to Theorem 3.4 and Theorem 3.1(3). If it is κ -distinguished, fix a $\overline{\mathbb{Q}}_\ell$ -character ξ of F^\times extending the canonical $\overline{\mathbb{Q}}_\ell$ -lift of κ . The reduction mod ℓ of ξ is an $\overline{\mathbb{F}}_\ell$ -character of F^\times extending κ , denoted χ . The representation $\rho\chi$ is distinguished and supercuspidal. It thus has a σ -self-dual lift π . Then $\pi\xi^{-1}$ is a distinguished lift of ρ . This proves (1).

Let μ be a σ -self-dual lift of ρ , and assume that ρ is distinguished. If μ is not distinguished, it must then be κ -distinguished. By Theorem 3.3, this implies that ρ is κ -distinguished, which contradicts the dichotomy and disjunction theorem. Conversely, if μ is distinguished, then ρ is distinguished thanks to Theorem 3.3. □

3.4.

From now on, we consider the case of cuspidal nonsupercuspidal R -representations, thus ℓ is a prime number different from p . Let us recall how they are classified in terms of their supercuspidal support.

Recall that a representation π of $\mathrm{GL}_n(F)$ on an R -vector space V is *generic* if V carries a nonzero R -linear form Θ such that $\Theta(\pi(u)v) = \theta(u)v$ for all $v \in V$ and all unipotent upper triangular matrices u , where $\theta(u) = \psi(u_{1,2} + \dots + u_{n-1,n})$ and ψ is a nontrivial R -character of F .

Let $k \geq 1$ be a positive integer, and ρ be a supercuspidal R -representation of $\mathrm{GL}_k(F)$. According to [29] 8.1, for any $r \geq 1$, the induced representation

$$\rho v^{-(r-1)/2} \times \dots \times \rho v^{(r-1)/2} \tag{3.1}$$

contains a unique generic irreducible subquotient, denoted $\mathrm{St}_r(\rho)$.

Let $e(\rho)$ be the smallest integer $i \geq 1$ such that ρv^i is isomorphic to ρ and $t(\rho)$ be the torsion number of ρ , that is, the number of unramified characters χ of F^\times such that $\rho\chi$ is isomorphic to ρ . By [31] Lemme 3.6, these integers are related by the identity

$$e(\rho) = \text{order of } q^{t(\rho)} \text{ mod } \ell. \tag{3.2}$$

By [29] Théorème 6.14, one has the following classification.

Proposition 3.6. *Let π be a cuspidal nonsupercuspidal R -representation of $\mathrm{GL}_n(F)$.*

1. *There are a unique positive integer $r = r(\pi) \geq 2$ dividing n and a supercuspidal representation ρ of $\mathrm{GL}_{n/r}(F)$ such that π is isomorphic to $\mathrm{St}_r(\rho)$.*
2. *There is a unique integer $v \geq 0$ such that $r = e(\rho)\ell^v$.*
3. *Let ρ' be a supercuspidal representation of $\mathrm{GL}_{n/r}(F)$. The representation π is isomorphic to $\mathrm{St}_r(\rho')$ if and only if ρ' is isomorphic to ρv^i for some $i \in \mathbb{Z}$.*

Note that, conversely, by the same references, if ρ is a supercuspidal representation of $\mathrm{GL}_k(F)$ and $r = e(\rho)\ell^v$ for some $v \geq 0$, the representation $\mathrm{St}_r(\rho)$ is cuspidal.

It will be convenient to set $r(\pi) = 1$ for any supercuspidal R -representation π of $\mathrm{GL}_n(F)$.

3.5.

We now classify σ -self-dual cuspidal representations.

Lemma 3.7. *Let ρ be a supercuspidal R -representation of $\mathrm{GL}_k(F)$ for some $k \geq 1$. Let $r \geq 2$ be such that $\mathrm{St}_r(\rho)$ is cuspidal, and suppose that $\mathrm{St}_r(\rho)$ is σ -self-dual. Then there is an $i \in \mathbb{Z}$, uniquely determined mod $e(\rho)$ such that $\rho^{\vee\sigma}$ is isomorphic to ρv^i .*

Proof. The representation $\text{St}_r(\rho)$ is the unique generic irreducible subquotient of (3.1). The representation $\text{St}_r(\rho)^{\vee\sigma}$ is thus generic, irreducible and it is a subquotient of

$$\rho^{\vee\sigma} \nu^{(r-1)/2} \times \dots \times \rho^{\vee\sigma} \nu^{-(r-1)/2}.$$

Equivalently (see [29] Proposition 2.6), it is a subquotient of $\rho^{\vee\sigma} \nu^{-(r-1)/2} \times \dots \times \rho^{\vee\sigma} \nu^{(r-1)/2}$. By uniqueness of the generic irreducible subquotient, we thus have

$$\text{St}_r(\rho)^{\vee\sigma} \simeq \text{St}_r(\rho^{\vee\sigma}).$$

The representation $\text{St}_r(\rho)$ is thus σ -self-dual if and only if $\text{St}_r(\rho^{\vee\sigma})$ is isomorphic to $\text{St}_r(\rho)$. The result then follows from Proposition 3.6. □

Proposition 3.8. *Let π be a cuspidal σ -self-dual representation of $\text{GL}_n(F)$. Set $r = r(\pi)$ and write $k = n/r$.*

1. *If r is odd or $\ell = 2$, there is a unique σ -self-dual supercuspidal representation ρ of $\text{GL}_k(F)$ such that π is isomorphic to $\text{St}_r(\rho)$.*
2. *Suppose that r is even and $\ell \neq 2$.*
 - (a) *There are a supercuspidal representation ρ of $\text{GL}_k(F)$ and an $i \in \{0, 1\}$ such that π is isomorphic to $\text{St}_r(\rho)$ and $\rho^{\vee\sigma} \simeq \rho \nu^i$.*
 - (b) *Let ρ' be a supercuspidal representation of $\text{GL}_k(F)$ and $j \in \{0, 1\}$ such that π is isomorphic to $\text{St}_r(\rho')$ and $\rho'^{\vee\sigma} \simeq \rho' \nu^j$. Then $j = i$, and either $\rho' \simeq \rho$ or $\rho' \simeq \rho \nu^{r/2}$.*

Proof. If $r = 1$, the result is trivial. Let us assume that $r \geq 2$. Fix a supercuspidal irreducible representation ρ of $\text{GL}_k(F)$ such that π is isomorphic to $\text{St}_r(\rho)$. By Lemma 3.7, there is an $i \in \mathbb{Z}$ such that $\rho^{\vee\sigma} \simeq \rho \nu^i$. Changing ρ to $\rho' = \rho \nu^s$ for some $s \in \mathbb{Z}$ does not change $\text{St}_r(\rho)$, but changes i to $i - 2s$. If r is odd or $\ell = 2$, then $e(\rho)$ is odd, thus $2\mathbb{Z} + e(\rho)\mathbb{Z} = \mathbb{Z}$. This proves (1). Similarly, if r is even and $\ell \neq 2$, then $e(\rho)$ is even: we thus may assume that $i \in \{0, 1\}$, proving (2.a). Moreover, if ρ' and j are as in (2.b), then $j - i$ is even, thus $j = i$. Moreover, ρ' is isomorphic to $\rho \nu^s$ for some $0 \leq s < e(\rho)$ such that $\nu^{2st(\rho)} = 1$, thus $e(\rho)$ divides $2s$. □

3.6.

We will need the finite field analogue of 3.4 (see [41] III.2.5 or [10] Theorem 19.3).

Proposition 3.9. *Let k be a finite field of characteristic p .*

1. *Let $f \geq 1$ be a positive integer and ϱ be a supercuspidal representation of $\text{GL}_f(k)$.*
 - (a) *For all $u \geq 1$, the induced representation*

$$\varrho \times \dots \times \varrho \quad (u \text{ times})$$

has a unique generic irreducible subquotient, denoted $\text{st}_u(\varrho)$.

- (b) *Let $e(\varrho)$ be the order of $q^f \pmod{\ell}$. The representation $\text{st}_u(\varrho)$ is cuspidal if and only if $u = 1$ or $u = e(\varrho)\ell^v$ for some $v \geq 0$.*
2. *Let W be a cuspidal representation of $\text{GL}_n(k)$. There exist a unique integer $u = r(W) \geq 1$ dividing n and a unique supercuspidal representation ϱ of $\text{GL}_{n/u}(k)$ such that $W \simeq \text{st}_u(\varrho)$.*

3.7.

As in the previous paragraph, k is a finite field of characteristic p . Let us recall how to parametrize cuspidal representations of $\text{GL}_n(k)$ by regular characters ([19], [14] Theorem 3.5 and [15, 21]).

Let \bar{k} be an algebraic closure of k . For any integer $s \geq 1$, let k_s be the extension of k of degree s contained in \bar{k} . Let Δ denote the group $\text{Gal}(k_n/k)$. A character of k_n^\times is Δ -regular if it is fixed by no nontrivial element of Δ .

Proposition 3.10. 1. Associated with any Δ -regular $\overline{\mathbb{Q}_\ell}$ -character ξ of k_n^\times , there is a cuspidal $\overline{\mathbb{Q}_\ell}$ -representation W_ξ of $\text{GL}_n(k)$, unique up to isomorphism, such that

$$\text{tr } W_\xi(x) = (-1)^{n-1} \cdot \sum_{\delta \in \Delta} \xi(x^\delta)$$

for all $x \in k_n^\times$ of degree n over k , where k_n^\times is considered as a maximal torus in $\text{GL}_n(k)$.
 2. The correspondence

$$\xi \mapsto W_\xi$$

induces a bijection from the set of Δ -conjugacy classes of Δ -regular $\overline{\mathbb{Q}_\ell}$ -characters of k_n^\times to that of isomorphism classes of cuspidal $\overline{\mathbb{Q}_\ell}$ -representations of $\text{GL}_n(k)$.

By reduction mod ℓ , we get the following classification.

- Proposition 3.11.** 1. Given any Δ -regular $\overline{\mathbb{Q}_\ell}$ -character ξ of k_n^\times , the reduction mod ℓ of W_ξ , denoted \overline{W}_ξ , is irreducible and cuspidal, and it only depends on the reduction mod ℓ of ξ .
 2. Reduction mod ℓ induces a bijection from the set of Δ -conjugacy classes of $\overline{\mathbb{F}_\ell}$ -characters of k_n^\times having a Δ -regular lift to $\overline{\mathbb{Q}_\ell}$ to that of isomorphism classes of cuspidal $\overline{\mathbb{F}_\ell}$ -representations of the group $\text{GL}_n(k)$.
 3. The integer $r(\overline{W}_\xi)$ is the greatest divisor r of n such that the reduction of ξ mod ℓ factorizes through a character of $k_{n/r}^\times$.

Definition 3.12. A parameter of a cuspidal representation ρ of $\text{GL}_n(k)$ is a character of k_n^\times whose Δ -conjugacy class corresponds to ρ by the bijection of either Proposition 3.10 or 3.11.

3.8.

Finally, we will need the following distinction criterion for cuspidal $\overline{\mathbb{Q}_\ell}$ -representations (see [20] Proposition 6.1 and [11] Lemme 3.4.10) of $\text{GL}_n(k)$ when p is odd.

Proposition 3.13. Assume that q is odd, n is even and write $n = 2u$. We consider the group $\text{GL}_u(k) \times \text{GL}_u(k)$ as a Levi subgroup of $\text{GL}_n(k)$. Let ξ be a Δ -regular $\overline{\mathbb{Q}_\ell}$ -character of k_n^\times .

1. The following assertions are equivalent.
 - (a) The cuspidal $\overline{\mathbb{Q}_\ell}$ -representation W_ξ is $\text{GL}_u(k) \times \text{GL}_u(k)$ -distinguished.
 - (b) The space of $\text{GL}_u(k) \times \text{GL}_u(k)$ -invariant linear forms on W_ξ has $\overline{\mathbb{Q}_\ell}$ -dimension 1.
 - (c) The cuspidal $\overline{\mathbb{Q}_\ell}$ -representation W_ξ is self-dual.
 - (d) The character ξ is trivial on k_u^\times .
2. Assume that the conditions of (1) are satisfied, and fix an element $\alpha \in k_n^\times$ such that $\alpha \notin k_u^\times$ and $\alpha^2 \in k_u^\times$. The element

$$s = \begin{pmatrix} 0 & \text{id} \\ \text{id} & 0 \end{pmatrix} \in \text{GL}_n(k), \tag{3.3}$$

where id is the identity in $\text{GL}_u(k)$, normalizes the group $\text{GL}_u(k) \times \text{GL}_u(k)$ and acts on the $\overline{\mathbb{Q}_\ell}$ -vector space of $\text{GL}_u(k) \times \text{GL}_u(k)$ -invariant linear forms on W_ξ by the sign $-\xi(\alpha)$.

Remark 3.14. Suppose that W_ξ is $\text{GL}_u(k) \times \text{GL}_u(k)$ -distinguished. By [35] Lemma 2.6, the cuspidal $\overline{\mathbb{F}_\ell}$ -representation \overline{W}_ξ is $\text{GL}_u(k) \times \text{GL}_u(k)$ -distinguished as well. More precisely, if we fix a nonzero

$GL_u(\mathbf{k}) \times GL_u(\mathbf{k})$ -invariant $\overline{\mathbb{Q}}_\ell$ -linear form Λ on W_ξ together with a $GL_n(\mathbf{k})$ -stable $\overline{\mathbb{Z}}_\ell$ -lattice $L \subseteq W_\xi$, then the associated $\overline{\mathbb{F}}_\ell$ -linear form

$$\overline{\Lambda} : L \otimes \overline{\mathbb{F}}_\ell \rightarrow \Lambda(L) \otimes \overline{\mathbb{F}}_\ell$$

is nonzero and $GL_u(\mathbf{k}) \times GL_u(\mathbf{k})$ -invariant. Moreover, if s acts on Λ by a sign $c \in \{-1, 1\} \subseteq \overline{\mathbb{Q}}_\ell^\times$, then s acts on $\overline{\Lambda}$ by the image of c in $\overline{\mathbb{F}}_\ell^\times$.

4. Reduction to level zero

In this section, p is odd, ℓ is any prime number different from p and R has characteristic 0 or ℓ . Let us fix a positive integer $n \geq 1$, and set $G = GL_n(F)$. We fix a character

$$\psi : F \rightarrow R^\times \tag{4.1}$$

which is trivial on \mathfrak{p}_F but not on \mathcal{O}_F .

4.1.

Let us recall the definitions and main results of [9, 8, 30, 3] which we will need.

Let $[\mathfrak{a}, \beta]$ be a simple stratum in the algebra $\mathbf{M}_n(F)$ of $n \times n$ matrices with entries in F . Recall that \mathfrak{a} is a hereditary \mathcal{O}_F -order of $\mathbf{M}_n(F)$ and β is an element of $\mathbf{M}_n(F)$ such that

- the F -algebra $E = F[\beta]$ is a field, and
- the multiplicative group E^\times normalizes \mathfrak{a}

(plus an extra technical condition on β which is not necessary to recall here: See [9] 1.5.5).

Let $\mathcal{K}_\mathfrak{a}$ be the normalizer of \mathfrak{a} in G and $\mathfrak{p}_\mathfrak{a}$ be its Jacobson radical, and set $U_\mathfrak{a}^1 = 1 + \mathfrak{p}_\mathfrak{a}$. Let B be the centralizer of E in $\mathbf{M}_n(F)$. The intersection $\mathfrak{b} = \mathfrak{a} \cap B$ is a hereditary order in B .

Associated with $[\mathfrak{a}, \beta]$ in [9] Chapter 3, there are compact mod centre open subgroups

$$H^1(\mathfrak{a}, \beta) \subseteq \mathbf{J}^1(\mathfrak{a}, \beta) \subseteq \mathbf{J}^0(\mathfrak{a}, \beta) \subseteq \mathbf{J}(\mathfrak{a}, \beta) \subseteq \mathcal{K}_\mathfrak{a}$$

and a nonempty finite set $\mathcal{C}(\mathfrak{a}, \beta)$ of characters of $H^1(\mathfrak{a}, \beta)$ called *simple characters*, depending on the choice of the character (4.1). We write $\mathbf{J} = \mathbf{J}(\mathfrak{a}, \beta)$, $\mathbf{J}^0 = \mathbf{J}^0(\mathfrak{a}, \beta)$, $\mathbf{J}^1 = \mathbf{J}^1(\mathfrak{a}, \beta)$ and $H^1 = H^1(\mathfrak{a}, \beta)$ for simplicity.

We will only be interested in the case where \mathfrak{b} is a maximal order in B , in which case the simple stratum $[\mathfrak{a}, \beta]$ and the simple characters in $\mathcal{C}(\mathfrak{a}, \beta)$ are said to be *maximal*. For the following result, see [8] 2.1, 3.2 and [9] 5.1.1.

Proposition 4.1. *Let $[\mathfrak{a}, \beta]$ be a maximal simple stratum.*

1. *The group \mathbf{J}^0 is the unique maximal compact subgroup of \mathbf{J} , and \mathbf{J}^1 is its unique maximal normal pro- p -subgroup.*
2. *One has $\mathbf{J} = E^\times \mathbf{J}^0 = (\mathbf{J} \cap B^\times) \mathbf{J}^1$ and*

$$\mathbf{J} \cap B^\times = \mathcal{K}_\mathfrak{b}, \quad \mathbf{J}^0 \cap B^\times = \mathfrak{b}^\times, \quad \mathbf{J}^1 \cap B^\times = U_\mathfrak{b}^1. \tag{4.2}$$

3. *There is an isomorphism of E -algebras*

$$B \simeq \mathbf{M}_m(E), \quad m = n/[E : F], \tag{4.3}$$

sending \mathfrak{b}^\times to the maximal compact open subgroup $GL_m(\mathcal{O}_E)$, which induces isomorphisms

$$\mathbf{J}^0/\mathbf{J}^1 \simeq \mathfrak{b}^\times/U_\mathfrak{b}^1 \simeq GL_m(I), \tag{4.4}$$

where I is the residue field of E .

- 4. Given any simple character $\theta \in \mathcal{C}(\mathfrak{a}, \beta)$, we have
 - (a) the normalizer of θ in G is equal to \mathbf{J} , and
 - (b) there is an irreducible representation η of \mathbf{J}^1 , unique up to isomorphism, whose restriction to H^1 contains θ , and such a representation extends to \mathbf{J} .

The representation η of (4.b) is called the *Heisenberg representation* associated with θ . If κ is a representation of \mathbf{J} extending η , any other extension of η to \mathbf{J} has the form $\kappa\xi$ for a unique character ξ of \mathbf{J} trivial on \mathbf{J}^1 .

Remark 4.2. 1. An isomorphism (4.3) as in Proposition 4.1(3) comes from the choice of an \mathcal{O}_E -basis of an \mathcal{O}_E -lattice \mathcal{L} in F^n whose endomorphism algebra is \mathfrak{b} .
 2. Changing the isomorphism (4.3), that is, changing the basis of \mathcal{L} above, has the effect of conjugating the identification (4.4) by an inner automorphism of $\mathrm{GL}_m(I)$.

A character θ of an open pro- p -subgroup H of G will be called a *maximal simple character* if there is a maximal simple stratum $[\mathfrak{a}, \beta]$ in $\mathbf{M}_n(F)$ such that $H = H^1(\mathfrak{a}, \beta)$ and $\theta \in \mathcal{C}(\mathfrak{a}, \beta)$. Given a maximal simple character θ of G , we will write H_θ^1 for the group on which θ is defined, \mathbf{J}_θ for its G -normalizer, \mathbf{J}_θ^0 for its unique maximal compact subgroup, \mathbf{J}_θ^1 for its unique maximal normal pro- p -subgroup and T for the maximal tamely ramified extension of F in $E = F[\beta]$. The following result shows how the latter depends on the choice of $[\mathfrak{a}, \beta]$ (see [8] 2.1, 2.5 and 2.6).

Proposition 4.3. Let $[\mathfrak{a}', \beta']$ be a simple stratum such that $\theta \in \mathcal{C}(\mathfrak{a}', \beta')$, and set $E' = F[\beta']$.

- 1. The orders $\mathfrak{a}, \mathfrak{a}'$ are equal and E, E' have the same degree over F .
- 2. The simple stratum $[\mathfrak{a}, \beta']$ is maximal.
- 3. The maximal tamely ramified extension of F in E' is \mathbf{J}_θ^1 -conjugate to T .

It follows that the G -conjugacy class of the simple character θ uniquely determines the integer m in equation (4.3), as well as the extension T up to G -conjugacy (or equivalently up to F -isomorphism). However, the fields E, E' need not be isomorphic (see [8] Example 2.1).

4.2.

In this paragraph only, we let n vary among all positive integers, and consider the set

$$\mathcal{C}_{\max}(F) = \bigcup_{[\mathfrak{a}, \beta]} \mathcal{C}(\mathfrak{a}, \beta),$$

where the union is taken over all maximal simple strata of $\mathbf{M}_n(F)$, for any $n \geq 1$. It is endowed with an equivalence relation called *endo-equivalence* ([6, 7]). An equivalence class for this equivalence relation is called an *endo-class* (see [3] 3.2).

Given a maximal simple character $\theta \in \mathcal{C}_{\max}(F)$ with endo-class Θ , the degree $[E : F]$ and the F -isomorphism class of its tame parameter field T only depend on Θ . They are called the *degree* and *tame parameter field* of Θ , respectively.

For a given n , any two maximal simple characters of $\mathrm{GL}_n(F)$ are endo-equivalent if and only if they are $\mathrm{GL}_n(F)$ -conjugate.

Remark 4.4. Note that endo-equivalence is defined in [6, 7] for arbitrary simple characters, not only for maximal ones, but we will not need this extra generality.

4.3.

We go back to the situation of Paragraph 4.1, assuming further that the character ψ of equation (4.1) is σ -invariant, which is possible since $p \neq 2$. As in [35], we will say that:

- o a simple stratum $[\mathfrak{a}, \beta]$ in $\mathbf{M}_n(F)$ is *σ -self-dual* if \mathfrak{a} is σ -stable and $\sigma(\beta) = -\beta$,

- a simple character θ is σ -self-dual if the group H_θ^1 is σ -stable and $\theta^{-1} \circ \sigma = \theta$,
- an endo-class Θ of (maximal) simple characters is σ -self-dual if for some (or equivalently for any) $\theta \in \Theta$, the character $\theta^{-1} \circ \sigma$ is in Θ .

Proposition 4.5. *Let θ be a σ -self-dual maximal simple character.*

1. *There is a σ -stable simple stratum $[\mathfrak{a}, \beta]$ such that $\theta \in \mathcal{C}(\mathfrak{a}, \beta)$.*
2. *Let E_0 be the σ -fixed points of E and \mathfrak{l}_0 be its residue field. There is an isomorphism (4.3) inducing an isomorphism (4.4) which transports the action of σ on $\mathbf{J}^0/\mathbf{J}^1$ to*
 - (a) *the action of the nontrivial element of $\text{Gal}(\mathfrak{l}/\mathfrak{l}_0)$ on $\text{GL}_m(\mathfrak{l})$ if E/E_0 is unramified,*
 - (b) *the adjoint action of*

$$\begin{pmatrix} -\text{id}_i & 0 \\ 0 & \text{id}_{m-i} \end{pmatrix} \in \text{GL}_m(\mathfrak{l}) \tag{4.5}$$

on $\text{GL}_m(\mathfrak{l})$ if E/E_0 is ramified, for a uniquely determined integer $i \in \{0, \dots, \lfloor m/2 \rfloor\}$.

If θ is a σ -self-dual maximal simple character, we will write T_0 for the maximal tamely ramified extension of F_0 in E_0 , that is, $T_0 = T \cap E_0$. By [3] Lemma 4.10, the canonical homomorphism

$$T_0 \otimes_{F_0} F \rightarrow T \tag{4.6}$$

is an isomorphism. Also, T/T_0 and E/E_0 have the same ramification index. By [3] Lemma 4.29, the F_0 -isomorphism class of T_0 is uniquely determined by the endo-class Θ of θ . And it follows from the isomorphism (4.6) that the F_0 -isomorphism class of T_0 determines the F -isomorphism class of T .

The following result is given by [35] Proposition 6.12, Lemma 6.20 and [36] Lemme 3.28. (The latter reference in [36] is for representations with coefficients in $R = \mathbb{C}$, but its proof is still valid in the ℓ -modular case.)

Proposition 4.6. *Let θ be a σ -self-dual maximal simple character.*

1. *The Heisenberg representation η of θ is σ -self-dual and $\mathbf{J}^1 \cap G^\sigma$ -distinguished, and the space $\text{Hom}_{\mathbf{J}^1 \cap G^\sigma}(\eta, R)$ has dimension 1.*
2. *For any representation κ of \mathbf{J} extending η , there are*
 - (a) *a unique character ξ of \mathbf{J} trivial on \mathbf{J}^1 such that $\kappa^{\vee\sigma}$ is isomorphic to $\kappa\xi$,*
 - (b) *a unique character χ of $\mathbf{J} \cap G^\sigma$ trivial on $\mathbf{J}^1 \cap G^\sigma$ such that*

$$\text{Hom}_{\mathbf{J}^1 \cap G^\sigma}(\eta, R) = \text{Hom}_{\mathbf{J} \cap G^\sigma}(\kappa, \chi^{-1}), \tag{4.7}$$

and the restriction of ξ to $\mathbf{J} \cap G^\sigma$ is equal to χ^2 .

3. *Given a representation κ as in (2) and an irreducible representation τ of \mathbf{J} trivial on \mathbf{J}^1 , the canonical linear map*

$$\text{Hom}_{\mathbf{J}^1 \cap G^\sigma}(\eta, R) \otimes \text{Hom}_{\mathbf{J} \cap G^\sigma}(\tau, \chi) \rightarrow \text{Hom}_{\mathbf{J} \cap G^\sigma}(\kappa \otimes \tau, R) \tag{4.8}$$

is an isomorphism of R -vector spaces.

4. *There exists a σ -self-dual representation of \mathbf{J} extending η .*

Conversely, let Θ be a σ -self-dual endo-class of degree dividing n . By [3] Section 4, it contains a σ -self-dual maximal simple character θ in G , and we have the following classification.

Proposition 4.7. *Let T/T_0 be the quadratic extension associated with Θ as above, and let us write $m = n/\text{deg}(\Theta)$.*

1. *If T/T_0 is unramified, the G -conjugacy class of θ contains a unique G^σ -conjugacy class of σ -self-dual simple characters.*

2. If T/T_0 is ramified, the number of G^σ -conjugacy classes of σ -self-dual simple characters in the G -conjugacy class of θ is equal to $\lfloor m/2 \rfloor + 1$, each class corresponding bijectively to an integer $i \in \{0, \dots, \lfloor m/2 \rfloor\}$ characterized by Proposition 4.5(2.b).

Remark 4.8. When T/T_0 is ramified, we define (as in [3, 35]) the *index* of a σ -self-dual maximal simple character θ to be the integer $i \in \{0, \dots, \lfloor m/2 \rfloor\}$ associated with its G^σ -conjugacy class. By [3] Remark 4.28 or [35] 5.D, if θ has index 0 and if

$$y_i = \text{diag}(\varpi, \dots, \varpi, 1, \dots, 1) \in \text{GL}_m(E) \simeq B^\times, \quad \varpi \text{ a uniformizer of } E \text{ occurring } i \text{ times,}$$

for some $i \in \{0, \dots, \lfloor m/2 \rfloor\}$, then θ^{y_i} is a σ -self-dual maximal simple character of index i .

4.4.

Let θ be a maximal simple character, and $[\alpha, \beta]$ be a simple stratum such that $\theta \in \mathcal{C}(\alpha, \beta)$. As in 4.1, write $\mathbf{J} = \mathbf{J}_\theta$, $\mathbf{J}^0 = \mathbf{J}_\theta^0$, $\mathbf{J}^1 = \mathbf{J}_\theta^1$ and $H^1 = H_\theta^1$. Let η be the Heisenberg representation of \mathbf{J}^1 associated with θ . In this paragraph, we give a slightly generalized version of [8] 3.3.

Fix an \mathcal{O}_E -lattice \mathcal{L} in $V = F^n$ whose endomorphism algebra is \mathfrak{b} . (It is uniquely determined up to homothety as \mathfrak{b} is maximal.) Fix a divisor $u \geq 1$ of m and decompositions

$$\mathcal{L} = \mathcal{L}_* \oplus \dots \oplus \mathcal{L}_*, \quad V = V_* \oplus \dots \oplus V_*, \tag{4.9}$$

in which V_* is an E -vector space of dimension m/u and \mathcal{L}_* is an \mathcal{O}_E -lattice of rank m/u in V_* . It defines a Levi subgroup M of G . Fix a pair (Q_-, Q_+) of M -opposite parabolic subgroups of G with Levi component M , and write N_-, N_+ for their unipotent radicals. Define $\mathfrak{a}_* = \text{End}_{\mathcal{O}_E}(\mathcal{L}_*)$. It is a hereditary order, and $[\mathfrak{a}_*, \beta]$ is a maximal simple stratum in $\text{End}_F(V_*)$. Write $\mathbf{J}_*, \mathbf{J}_*^0, \mathbf{J}_*^1$ and H_*^1 for the subgroups associated with it. Compare with [8] 3.3, which corresponds to the particular case where $u = m$.

The next result follows from [6] Example 10.9 (compare with Lemma 1 in [8] 3.3).

Lemma 4.9. 1. *There are Iwahori decompositions*

$$H^1 = (H^1 \cap N_-) \cdot (H^1 \cap M) \cdot (H^1 \cap N_+),$$

$$H^1 \cap M = H_*^1 \times \dots \times H_*^1$$

and

$$\mathbf{J}^1 = (\mathbf{J}^1 \cap N_-) \cdot (\mathbf{J}^1 \cap M) \cdot (\mathbf{J}^1 \cap N_+),$$

$$\mathbf{J}^1 \cap M = \mathbf{J}_*^1 \times \dots \times \mathbf{J}_*^1.$$

2. *The character θ is trivial on $H^1 \cap N_-, H^1 \cap N_+$, and there exists a unique simple character $\theta_* \in \mathcal{C}(\mathfrak{a}_*, \beta)$ such that θ agrees with $\theta_* \otimes \dots \otimes \theta_*$ on $H^1 \cap M$.*

Moreover, the map $\theta \mapsto \theta_*$ defined by Lemma 4.9(2) is a bijection from $\mathcal{C}(\alpha, \beta)$ to $\mathcal{C}(\mathfrak{a}_*, \beta)$: this is an instance of the transfer of [9] 3.6.

Let η_* denote the Heisenberg representation of \mathbf{J}_*^1 associated with θ_* . Compare the next result with Lemma 2 in [8] 3.3 and the discussion after it.

Lemma 4.10. *Let κ_* be a representation of \mathbf{J}_* extending η_* .*

1. *The set $\mathbf{J}_+ = (H^1 \cap N_-) \cdot (\mathbf{J} \cap Q_+)$ is a group, and there is a unique representation κ_+ of \mathbf{J}_+ which is trivial on $H^1 \cap N_-, \mathbf{J}^0 \cap N_+$ and agrees with $\kappa_* \otimes \dots \otimes \kappa_*$ on $\mathbf{J} \cap M$.*
2. *The representation $\tilde{\kappa}_+$ of $(\mathbf{J}^1 \cap N_-) \cdot (\mathbf{J} \cap Q_+) = \mathbf{J}^1 \mathbf{J}_+$ induced by κ_+ extends η .*
3. *There is a unique irreducible representation κ of \mathbf{J} extending $\tilde{\kappa}_+$.*

Proof. That \mathbf{J}_+ is a group follows from the fact that H^1 is normalized by \mathbf{J} , thus by $\mathbf{J} \cap Q_+$. The existence of κ_+ follows from the containment

$$(\mathbf{J}^0 \cap N_+) \cdot (H^1 \cap N_-) \subseteq (H^1 \cap N_-) \cdot (\mathbf{J}^1 \cap M) \cdot (\mathbf{J}^0 \cap N_+)$$

(see the argument of [34] 2.3). Mackey’s formula implies that the restriction of $\tilde{\kappa}_+$ to \mathbf{J}^1 is

$$\text{Ind}_{\mathbf{J}^1 \cap \mathbf{J}_+}^{\mathbf{J}^1} (\kappa_+).$$

The restriction of κ_+ to $\mathbf{J}^1 \cap \mathbf{J}_+ = (H^1 \cap N_-) \cdot (\mathbf{J}^1 \cap Q_+)$ is the unique representation η_+ which is trivial on $H^1 \cap N_-$, $\mathbf{J}^1 \cap N_+$ and agrees with $\eta_* \otimes \cdots \otimes \eta_*$ on $\mathbf{J}^1 \cap M$. The representation it induces to \mathbf{J}^1 is isomorphic to η : Indeed, this representation contains θ by Lemma 4.9(2), and it has dimension

$$\begin{aligned} \dim(\eta_* \otimes \cdots \otimes \eta_*) \cdot (\mathbf{J}^1 \cap N_- : H^1 \cap N_-) &= (\mathbf{J}^1 \cap M : H^1 \cap M)^{1/2} \cdot (\mathbf{J}^1 \cap N_- : H^1 \cap N_-) \\ &= (\mathbf{J}^1 : H^1)^{1/2} \end{aligned}$$

which is the dimension of η (see [30] 2.3).

It remains to prove (3). First, uniqueness follows from the fact that any two such extensions differ from a character of \mathbf{J} trivial on $\mathbf{J}^1 \mathbf{J}_+$, and such a character is trivial since $p \neq 2$. Existence follows from [9] 5.2.4 (see [30] 2.4 in the modular case). \square

The reader will pay attention to the fact that, although $\mathbf{J}^0 \cap M$ is equal to $\mathbf{J}_*^0 \times \cdots \times \mathbf{J}_*^0$, the group $\mathbf{J} \cap M$ is not equal to $\mathbf{J}_* \times \cdots \times \mathbf{J}_*$ in general (unless $u = 1$). It is generated by $\mathbf{J}^0 \cap M$ and E^\times , the latter being diagonal in M .

Lemma 4.11. 1. *The map*

$$\kappa_* \mapsto \kappa \tag{4.10}$$

from isomorphism classes of representations of \mathbf{J}_ extending η_* to isomorphism classes of representation of \mathbf{J} extending η is surjective.*

2. *Any two representations of \mathbf{J}_* extending η_* have the same image if and only if they are twists of each other by a character of \mathbf{J}_* trivial on \mathbf{J}_*^0 and of order dividing u .*

Proof. The case where $u = m$ is given by [8] 3.3, Corollary 1. For the general case, first note that, if κ is the image of κ_* by the map (4.10), and if ξ_* is a character of \mathbf{J}_* trivial on \mathbf{J}_*^0 , then the image of $\kappa_* \xi_*$ by the map (4.10) is equal to $\kappa \xi^u$, where ξ is the character of \mathbf{J} trivial on \mathbf{J}^0 coinciding with ξ_* on any uniformizer of E . Now, start with a representation κ extending η . Let $\tilde{\kappa}_+$ denote its restriction to $\mathbf{J}^1 \mathbf{J}_+$ and κ_+ denote the representation of \mathbf{J}_+ on the space of $\mathbf{J}^1 \cap N_-$ -invariants of $\tilde{\kappa}_+$. The restriction of κ_+ to $\mathbf{J}^0 \cap M$ has the form

$$\kappa_*^0 \otimes \cdots \otimes \kappa_*^0$$

for a uniquely determined representation κ_*^0 of \mathbf{J}_*^0 extending η_* . Since any two extensions of κ_*^0 to \mathbf{J}_* are twists of each other by a character of $\mathbf{J}_* = E^\times \mathbf{J}_*^0$ trivial on \mathbf{J}_*^0 , the result follows. \square

Remark 4.12. Suppose that u is equal to m . Let $y \in M \cap B^\times$, and write $\theta' = \theta^y \in \mathcal{C}(\mathfrak{a}^y, \beta)$. The groups associated with θ' are $\mathbf{J}' = \mathbf{J}^y$, etc. The group isomorphism $B^\times \simeq \text{GL}_m(E)$ identifies $M \cap B^\times$ with the diagonal torus $E^\times \times \cdots \times E^\times$, and E^\times normalizes θ_* . The character θ' is thus trivial on $H^{1y} \cap N_-$, $H^{1y} \cap N_+$ and agrees with $\theta_* \otimes \cdots \otimes \theta_*$ on $H^{1y} \cap M = H^1 \cap M$. If κ_* is a representation of \mathbf{J}_* extending η_* , the representation of \mathbf{J}' corresponding to it by the map (4.10) is κ^y .

4.5.

Suppose now that the simple character θ and the simple stratum $[\mathfrak{a}, \beta]$ of 4.4 are σ -self-dual. The groups $\mathbf{J}, \mathbf{J}^0, \mathbf{J}^1$ and H^1 are thus σ -stable. Suppose also that the decompositions (4.9) are σ -stable. (To obtain such decompositions, consider the vertex in the reduced building of B^\times defined by the \mathcal{O}_E -lattice \mathcal{L} (see [5]) and choose a σ -stable apartment containing this vertex, whose existence is granted by [13] since $p \neq 2$.)

The Levi subgroup M is thus σ -stable, and we may assume that Q_-, Q_+ are σ -stable as well. Also, the simple stratum $[\mathfrak{a}_*, \beta]$ and the simple character θ_* given by Lemma 4.9 are σ -self-dual. Let G_* denote the group $\text{Aut}_F(V_*)$. It is isomorphic to $\text{GL}_{n/u}(F)$.

We may also assume that our choice of basis induces an isomorphism of groups (4.4) between $\mathbf{J}^0/\mathbf{J}^1$ and $\text{GL}_m(\mathbf{I})$ as in Proposition 4.5(4), transporting the action of σ on $\mathbf{J}^0/\mathbf{J}^1$ to

- the action of the nontrivial element of $\text{Gal}(\mathbf{I}/\mathbf{I}_0)$ on $\text{GL}_m(\mathbf{I})$ if T/T_0 is unramified,
- the adjoint action of (4.5) on $\text{GL}_m(\mathbf{I})$ for some $i \in \{0, \dots, \lfloor m/2 \rfloor\}$, if T/T_0 is ramified.

Let κ_* be a representation of \mathbf{J}_* extending η_* , and let κ correspond to it by the map (4.10).

Lemma 4.13. *If κ_* is σ -self-dual, then κ is σ -self-dual.*

Proof. First, $\kappa_+^{\vee\sigma}$ is trivial on both $H^1 \cap N_-, \mathbf{J}^1 \cap N_+$ and agrees with $\kappa_*^{\vee\sigma} \otimes \dots \otimes \kappa_*^{\vee\sigma}$ on $\mathbf{J} \cap M$. If κ_* is σ -self-dual, it follows by uniqueness that κ_+ is σ -self-dual, thus $\tilde{\kappa}_+$ is σ -self-dual as well. The unique representation of \mathbf{J} extending $\tilde{\kappa}_+^{\vee\sigma}$ is $\kappa^{\vee\sigma}$, hence κ is σ -self-dual. □

We will need the following lemma. Let ϖ be a uniformizer of E such that

$$\sigma(\varpi) = \begin{cases} \varpi & \text{if } T/T_0 \text{ is unramified,} \\ -\varpi & \text{if } T/T_0 \text{ is ramified.} \end{cases} \tag{4.11}$$

Note that the group \mathbf{J} is generated by \mathbf{J}^0 and ϖ .

Lemma 4.14. *The group $\mathbf{J} \cap G^\sigma$ is generated by $\mathbf{J}^0 \cap G^\sigma$ and an element ϖ' such that*

1. $\varpi' = \varpi$ if T/T_0 is unramified,
2. $\varpi' = \varpi^2$ if T/T_0 is ramified and $m \neq 2i$,
3. $\varpi' \in \varpi \mathbf{J}^0$ if T/T_0 is ramified and $m = 2i$.

Proof. If T/T_0 is unramified: See [35] Lemma 9.1. Suppose that T/T_0 is ramified, and assume that there is an $x \in \mathbf{J} \cap G^\sigma, x \notin \langle \varpi^2, \mathbf{J}^0 \cap G^\sigma \rangle$. We have $x = \varpi^v y$, where $v \in \mathbb{Z}$ is odd and $y \in \mathbf{J}^0$ satisfies $\sigma(y) = -y$. Reducing mod \mathbf{J}^1 , we get an $\alpha \in \text{GL}_m(\mathbf{I})$ such that $\sigma(\alpha) = -\alpha$. Since the involution σ acts on $\text{GL}_m(\mathbf{I})$ by conjugacy by

$$\delta = \text{diag}(-1, \dots, -1, 1, \dots, 1)$$

(where -1 occurs i times), this implies that δ and $-\delta$ are $\text{GL}_m(\mathbf{I})$ -conjugate, thus $m = 2i$. Conversely, if $m = 2i$, then

$$w = \begin{pmatrix} 0 & \text{id}_i \\ \text{id}_i & 0 \end{pmatrix} \in \mathbf{J}^0 \cap B^\times = \text{GL}_{2i}(\mathcal{O}_E) \tag{4.12}$$

is σ -anti-invariant, and $\varpi' = \varpi w$ has the required property. □

We now investigate the behavior of the map (4.10) with respect to distinction. The case where $u = m$ will be sufficient for our purpose (see Paragraph 4.6).

Lemma 4.15. *Suppose that $u = m$ and κ_* is $\mathbf{J}_* \cap G_*^\sigma$ -distinguished.*

1. *If T/T_0 is unramified, or if T/T_0 is ramified and $m \neq 2i$, the representation κ is $\mathbf{J} \cap G^\sigma$ -distinguished.*
2. *If T/T_0 is ramified and $m = 2i$, there exists a quadratic character ξ of \mathbf{J} trivial on \mathbf{J}^0 such that $\kappa\xi$ is $\mathbf{J} \cap G^\sigma$ -distinguished.*

Proof. The representation κ_+ is $\mathbf{J}_+ \cap G^\sigma$ -distinguished, thus $\tilde{\kappa}_+$ is $\mathbf{J}^1\mathbf{J}_+ \cap G^\sigma$ -distinguished. It follows that κ is $\mathbf{J}^1\mathbf{J}_+ \cap G^\sigma$ -distinguished. Let χ be the character of $\mathbf{J} \cap G^\sigma$ associated with κ by Proposition 4.6. It is trivial on $\mathbf{J}^1\mathbf{J}_+ \cap G^\sigma$. Restricting to $\mathbf{J}^0 \cap G^\sigma$, it is a character of

$$(\mathbf{J}^0 \cap G^\sigma)/(\mathbf{J}^1 \cap G^\sigma) \simeq \text{GL}_m(\mathcal{I})^\sigma.$$

Since $\mathbf{J} \cap M \subseteq \mathbf{J}_+$, it is trivial on the image of $(\mathbf{J} \cap M) \cap (\mathbf{J}^0 \cap G^\sigma)$ in $\text{GL}_m(\mathcal{I})^\sigma$, which is made of the σ -fixed points of the diagonal torus $M = \mathcal{I}^\times \times \dots \times \mathcal{I}^\times$.

If T/T_0 is unramified, we have $\text{GL}_m(\mathcal{I})^\sigma = \text{GL}_m(\mathcal{I}_0)$ and $M^\sigma = \mathcal{I}_0^\times \times \dots \times \mathcal{I}_0^\times$, thus χ is trivial on $\mathbf{J}^0 \cap G^\sigma$. If T/T_0 is ramified, we have $\text{GL}_m(\mathcal{I})^\sigma = \text{GL}_i(\mathcal{I}) \times \text{GL}_{m-i}(\mathcal{I})$ and $M^\sigma = M$. Again, χ is trivial on $\mathbf{J}^0 \cap G^\sigma$.

By Lemma 4.14, it remains to consider the value of χ at ϖ' . If T/T_0 is unramified, or if T/T_0 is ramified and $m \neq 2i$, we have $\varpi' \in \mathbf{J}^1\mathbf{J}_+ \cap G^\sigma$, thus χ is trivial.

Now, assume that T/T_0 is ramified and $m = 2i$. Let ξ be the quadratic character of \mathbf{J} trivial on \mathbf{J}^0 defined by $\xi(\varpi') = \chi(\varpi')$. Then $\kappa\xi$ is $\mathbf{J} \cap G^\sigma$ -distinguished. □

We will prove in Paragraph 4.6 that the quadratic character ξ of Lemma 4.15(2) is always trivial: See Corollary 4.19.

4.6.

As in Paragraph 4.5, the simple character θ and the simple stratum $[\alpha, \beta]$ are both maximal σ -self-dual, and η is the Heisenberg representation of \mathbf{J}^1 associated with θ . The next proposition, which says that η has a canonical extension to \mathbf{J} , is the core of our proof of Theorem 4.41.

Proposition 4.16. *There is, up to isomorphism, a unique representation κ of \mathbf{J} extending η satisfying the following conditions:*

1. *it is both σ -self-dual and $\mathbf{J} \cap G^\sigma$ -distinguished,*
2. *its determinant has order a power of p .*

This unique representation will be denoted κ_θ .

Remark 4.17. This extends (and makes more precise) the results of [35] (see *ibid.*, Propositions 7.9, 9.4) where θ is assumed to be generic and either T/T_0 is unramified and m is odd, or T/T_0 is ramified and $m \in \{1, 2i\}$. See also [35] Remarks 9.5 and 9.9.

Proof. Suppose first that there exists a representation satisfying (1). As in the proof of [36] Corollary 6.12, one then easily proves the existence of a representation κ satisfying (1) and (2). Let us prove that such a representation is unique. Any other representation of \mathbf{J} satisfying the conditions of the proposition is of the form $\kappa\phi$ for some character ϕ of \mathbf{J} which is σ -self-dual and trivial on $(\mathbf{J} \cap G^\sigma)\mathbf{J}^1$, and whose order is a power of p . The restriction of ϕ to \mathbf{J}^0 can be considered as a character of $\text{GL}_m(\mathcal{I})$. Since the latter group is not isomorphic to $\text{GL}_2(\mathbb{F}_2)$ (for $p \neq 2$), this character factors through the determinant. Its order is thus prime to p , which implies that ϕ is trivial on \mathbf{J}^0 . It is thus a character of $\mathbf{J}/(\mathbf{J} \cap G^\sigma)\mathbf{J}^0$ which, by Lemma 4.14, has order at most 2. Uniqueness follows from the fact that $p \neq 2$.

We are now reduced to proving the existence of a representation κ satisfying (1). If $m = 1$, this follows from [35] Propositions 7.9, 9.4. (See also Remark 4.17.)

Now, consider the constructions of 4.4 and 4.5 with $u = m$. Thanks to the case where m is equal to 1, there is a representation κ_* of \mathbf{J}_* extending η_* which is both σ -self-dual and $\mathbf{J}_* \cap G_*^\sigma$ -distinguished. Let κ be the representation of \mathbf{J} extending η associated with it by the map (4.10). Lemma 4.13 implies that it

is σ -self-dual, and Lemma 4.15 implies that there is a quadratic character ξ of \mathbf{J} trivial on \mathbf{J}^0 such that $\kappa\xi$ is $\mathbf{J} \cap G^\sigma$ -distinguished. Since ξ is unramified and quadratic, $\kappa\xi$ is also σ -self-dual and extends η . \square

Remark 4.18. Notice that this gives another proof of [35] Propositions 7.9, 9.4, based on the case $m = 1$ only.

Now, we can improve Lemma 4.15. Suppose we are in the situation of Paragraphs 4.4 and 4.5, with $u = m$.

Corollary 4.19. *Suppose that $u = m$. Let κ_* be a representation of \mathbf{J}_* extending η_* and κ correspond to it by the map (4.10). If κ_* is $\mathbf{J}_* \cap G_*^\sigma$ -distinguished, then κ is $\mathbf{J} \cap G^\sigma$ -distinguished.*

Proof. The result is given by Lemma 4.15, except when T/T_0 is ramified and $m = 2i$, which we assume now. Suppose that κ_* is $\mathbf{J}_* \cap G_*^\sigma$ -distinguished. By Lemma 4.15, there is a quadratic character ξ of \mathbf{J} trivial on \mathbf{J}^0 such that $\kappa\xi$ is $\mathbf{J} \cap G^\sigma$ -distinguished. Let κ_θ be the representation given by Proposition 4.16, and write $\kappa\xi = \kappa_\theta\phi$ for some character ϕ of \mathbf{J} trivial on $(\mathbf{J} \cap G^\sigma)\mathbf{J}^1$. Restricting to \mathbf{J}^0 , the character ϕ can be seen as a character of $\mathrm{GL}_m(I)$ of the form $\alpha \circ \det$, for some character α of I^\times , which is trivial on $\mathrm{GL}_m(I)^\sigma = \mathrm{GL}_i(I) \times \mathrm{GL}_i(I)$. This implies that α is trivial, thus ϕ is trivial on \mathbf{J}^0 . Also, ϕ is trivial at $\varpi' \in \varpi\mathbf{J}^0$ by Lemma 4.14. It is thus trivial. In conclusion, we have $\kappa = \kappa_\theta\xi$. Taking determinants, we get

$$\det \kappa = \xi^{\dim \kappa_\theta} \cdot \det \kappa_\theta = \xi \cdot \det \kappa_\theta. \tag{4.13}$$

(The second equality comes from the fact that ξ is quadratic and $p \neq 2$ so $\dim \kappa_\theta$ is odd.)

There exists a $y \in M \cap B^\times$ such that $\theta' = \theta^y \in \mathcal{C}(\alpha^y, \beta)$ is a σ -self-dual maximal simple character of index 0 (in the sense of Remark 4.8). By Remark 4.12, the simple character of $\mathcal{C}(\alpha_*, \beta)$ associated with θ' by Lemma 4.9 is still θ_* , and the representation of $\mathbf{J}' = \mathbf{J}^y$ corresponding to κ_* by the map (4.10) is κ^y . Let $\kappa_{\theta'}$ be the representation associated with θ' by Proposition 4.16. By Lemma 4.15, κ' is distinguished. By the discussion above, it follows that

$$\det \kappa^y = \det \kappa_{\theta'}. \tag{4.14}$$

But the characters $\det \kappa$, $\det \kappa^y$ have the same order (since they are conjugate to each others), and the latter one has order a power of p thanks to equation (4.14). Now, equation (4.13) implies that ξ has order a power of p . Since ξ is quadratic and $p \neq 2$, this character is trivial. \square

We extract from the proof of Corollary 4.19 the following valuable corollary.

Corollary 4.20. *Suppose that $u = m$. Let κ_{θ_*} and κ_θ be the representations associated with θ_* and θ by Proposition 4.16, respectively. Then the map (4.10) takes κ_{θ_*} to κ_θ .*

We also have the following corollary, which extends [35] Lemma 7.10(3), Corollary 9.6(1).

Corollary 4.21. *Any $\mathbf{J} \cap G^\sigma$ -distinguished representation of \mathbf{J} extending η is σ -self-dual.*

Proof. Let κ be a $\mathbf{J} \cap G^\sigma$ -distinguished representation of \mathbf{J} extending η , and ξ be the unique character of \mathbf{J} trivial on \mathbf{J}^1 such that $\kappa = \kappa_\theta\xi$. We have to prove that $\xi^{-1} \circ \sigma = \xi$. The fact that κ is distinguished implies that ξ is trivial on $(\mathbf{J} \cap G^\sigma)\mathbf{J}^1$. Restricting to \mathbf{J}^0 , the character ξ can be seen as a character of $\mathrm{GL}_m(I)$ of the form $\alpha \circ \det$, for some character α of I^\times .

If T/T_0 is unramified, α is trivial on I_0^\times , thus $\xi^{-1} \circ \sigma$ and ξ coincide on \mathbf{J}^0 . They also coincide on $\varpi \in \mathbf{J} \cap G^\sigma$, thus they are equal.

If T/T_0 is ramified, α is trivial, thus $\xi^{-1} \circ \sigma$ and ξ are both trivial on \mathbf{J}^0 . Since ξ is trivial on $\varpi^2 \in \mathbf{J} \cap G^\sigma$, we get $\xi^{-1} \circ \sigma(\varpi) = \xi^{-1}(-\varpi) = \xi^{-1}(\varpi) = \xi(\varpi)$, which finishes the proof. \square

4.7.

We now come to the type theoretic description of cuspidal representations of G . The following proposition follows from [9] Theorem 8.4.1, Corollary 6.2.3, Theorem 5.7.1 (see [30] Théorèmes 3.4, 3.7 and [37] Theorem 7.2 in the modular case).

Proposition 4.22. *Let π be a cuspidal representation of G . There is, up to G -conjugacy, a unique simple character θ such that the restriction of π to H_θ^1 contains θ , and it is maximal.*

Let π be a cuspidal representation of G , and let θ be a simple character occurring in π . Associated with it, there are:

- the positive integer $m(\pi) = m \geq 1$ defined by equation (4.3), called the *relative degree* of π ,
- the G -conjugacy class (or equivalently the F -isomorphism class) of the tamely ramified extension T of F associated with θ , called the *tame parameter field* of π ,
- the endo-class Θ of θ , called the *endo-class* of π .

(Note that, when π has level 0, one has $m = n$ and $T = F$, and Θ is the null endo-class.)

Write $\mathbf{J} = \mathbf{J}_\theta$, $\mathbf{J}^0 = \mathbf{J}_\theta^0$, $\mathbf{J}^1 = \mathbf{J}_\theta^1$, and let η be the Heisenberg representation of θ . The next proposition follows from [30] Lemme 5.3, Theorem 3.11.

Proposition 4.23. *Let κ be a representation of \mathbf{J} extending η , and define a representation of \mathbf{J} on the space $\text{Hom}_{\mathbf{J}^1}(\kappa, \pi)$ by making $x \in \mathbf{J}$ act on $f \in \text{Hom}_{\mathbf{J}^1}(\kappa, \pi)$ by*

$$x \cdot f = \pi(x) \circ f \circ \kappa(x)^{-1}.$$

This representation, denoted τ , has the following properties:

1. *It is irreducible, and trivial on \mathbf{J}^1 .*
2. *If one identifies $\mathbf{J}^0/\mathbf{J}^1$ with a finite general linear group as in (4.4), its restriction to \mathbf{J}^0 is the inflation of a cuspidal representation.*
3. *The compact induction of $\kappa \otimes \tau$ from \mathbf{J} to G is isomorphic to π .*

Any two representations of \mathbf{J} extending η differ from a character of \mathbf{J} trivial on \mathbf{J}^1 . The pair

$$(\mathbf{J}, \kappa \otimes \tau) \tag{4.15}$$

thus only depends on π and the choice of θ and not on the choice of κ .

When π varies among all cuspidal representations of G and θ varies among all maximal simple characters in π , the pairs (4.15) are called *extended maximal simple types* in [9, 30], which we will abbreviate to *types* here. A given cuspidal representation of G thus contains, up to G -conjugacy, a unique type (\mathbf{J}, λ) : There is a unique maximal simple character θ such that $\mathbf{J}_\theta = \mathbf{J}$ and θ occurs in the restriction of λ to H_θ^1 , a representation κ of \mathbf{J} which restricts irreducibly to \mathbf{J}^1 and a representation τ of \mathbf{J} trivial on \mathbf{J}^1 such that λ is isomorphic to $\kappa \otimes \tau$.

Remark 4.24. If \mathfrak{a} is a maximal order in $\mathbf{M}_n(F)$, the trivial character of $U_\mathfrak{a}^1$ is a maximal simple character, with $E = T = F$ and $m = n$. The cuspidal representations of G that contain such a simple character are precisely the cuspidal representations of level 0.

Fix a representation κ of \mathbf{J} extending η , and define τ as in Proposition 4.23, and fix a simple stratum $[\mathfrak{a}, \beta]$ such that $\theta \in \mathcal{C}(\mathfrak{a}, \beta)$ and an isomorphism (4.3). This gives a field E and an isomorphism $\mathbf{J}^0/\mathbf{J}^1 \simeq \text{GL}_m(I)$, where I is the residue field of T .

By Proposition 4.23(2), the restriction of τ to \mathbf{J}^0 is the inflation of a cuspidal irreducible representation, denoted V .

On the other hand, the representation τ has a central character: It is a character of the centre $E^\times \mathbf{J}^1/\mathbf{J}^1$ of \mathbf{J}/\mathbf{J}^1 or equivalently a tamely ramified character of E^\times . Since E is totally wildly ramified over its maximal tamely ramified subextension T , any tamely ramified character of E^\times factors through the norm

$N_{E/T}$. The restriction of τ to E^\times is thus a multiple of $\omega \circ N_{E/T}$ for a uniquely determined tamely ramified character ω of T^\times .

The data V and ω are subject to the compatibility condition that the restriction of V to I^\times is a multiple of the character of I^\times whose inflation to \mathcal{O}_T^\times is the restriction of ω^{p^e} , with $p^e = [E : T]$. Associated with V by Proposition 3.9, there are a unique integer $u \geq 1$ dividing m and a unique supercuspidal representation ϱ of $GL_{m/u}(I)$ such that V is isomorphic to $st_u(\varrho)$. The next important result is [31] Lemma 3.2. The integer $r(\pi)$ has been defined in Paragraph 3.4.

Lemma 4.25. *The integer u is equal to $r(\pi)$.*

It follows that $r(\pi)$ divides m and that π is supercuspidal if and only if V is supercuspidal.

4.8.

Write $r = r(\pi)$ and $k = n/r$, and let ρ be a supercuspidal representation of $GL_k(F)$ such that π is isomorphic to $St_r(\rho)$ given by Proposition 3.6. In this paragraph, we will compare the type theoretic description of π with that of ρ . As in 4.7, we fix a representation κ of \mathbf{J} extending η . It defines an irreducible representation τ of \mathbf{J} trivial on \mathbf{J}^1 , then a cuspidal representation V of $GL_m(I)$ and a tamely ramified character ω of T^\times . There is also a (unique) supercuspidal representation ϱ of $GL_{m/r}(I)$ such that V is isomorphic to $st_r(\varrho)$.

Since r divides m , we may apply the results of 4.4 to the case where $u = r$, which we assume now. Let θ_* be the simple character associated with θ by Lemma 4.9.

Lemma 4.26. *The representation ρ contains θ_* .*

Proof. This follows from the description¹ of $St_r(\rho)$ in [29] Section 6. □

Consequently, π and ρ have the same endo-class. We have the following immediate corollary.

Corollary 4.27. *We have $m(\pi) = m(\rho)r$ and the representations π, ρ have the same tame parameter field.*

Let η_* be the Heisenberg representation associated with θ_* , and let κ_* be a representation of \mathbf{J}_* extending η_* such that the representation of \mathbf{J} associated with it by the map (4.10) is κ . It defines an irreducible representation τ_* of \mathbf{J}_* trivial on \mathbf{J}_*^1 such that the pair $(\mathbf{J}_*, \kappa_* \otimes \tau_*)$ is a type in ρ . Associated with this, there are a cuspidal representation ϱ_* of $GL_{m/r}(I)$ (which is supercuspidal thanks to the comment after Lemma 4.25) and a tamely ramified character ω_* of T^\times . The following proposition compares the pairs (ϱ, ω) and (ϱ_*, ω_*) associated with τ and τ_* .

Proposition 4.28. *We have $\varrho \simeq \varrho_*$ and $\omega = \omega_*^r$.*

Proof. Again, the fact that ϱ is isomorphic to ϱ_* follows from the description of $St_r(\rho)$ in [29] Section 6. It thus remains to prove the second equality. For this, consider the action of \mathbf{J} on

$$\text{Hom}_{\mathbf{J}^1}(\kappa, \mathcal{F}(\rho, r)), \tag{4.16}$$

where $\mathcal{F}(\rho, r)$ is the parabolically induced representation (3.1). By [38] Proposition 5.6, its restriction to \mathbf{J}^0 is the inflation of the induced representation $\varrho_* \times \cdots \times \varrho_*$ of $GL_m(I)$. By tracking the action of E^\times in the arguments of [38] Section 5, we see that it acts on the space (4.16) by the character

$$\omega_*^r \circ N_{E/T}.$$

In particular, E^\times acts through this character on the subquotient $\text{Hom}_{\mathbf{J}^1}(\kappa, \pi)$, which is τ . □

¹Warning: the representation denoted $St(\rho, r)$ in [29] corresponds to $St_r(\rho v^{(r-1)/2})$, and the one denoted $St_r(\rho)$ in [29] corresponds to $St_v(\rho v^{(v-1)/2})$ with $v = e(\rho)\ell^r$.

4.9.

Suppose that the cuspidal representation π is σ -self-dual. We say a type (\mathbf{J}, λ) is σ -self-dual if \mathbf{J} is σ -stable and $\lambda^{\vee\sigma}$ is isomorphic to λ . The next result is [3] Theorem 4.1.

Proposition 4.29. *The representation π contains a σ -self-dual type.*

A type (\mathbf{J}, λ) contains a unique simple character θ such that $\mathbf{J}_\theta = \mathbf{J}$: It follows that, if (\mathbf{J}, λ) is σ -self-dual, θ is σ -self-dual as well. In particular, π contains a σ -self-dual simple character.

Let θ be a σ -self-dual simple character occurring in π , and $[\mathfrak{a}, \beta]$ be a σ -self-dual simple stratum such that $\theta \in \mathcal{C}(\mathfrak{a}, \beta)$ (which exists by Proposition 4.5). The G^σ -conjugacy class (or equivalently the F_0 -isomorphism class) of the tamely ramified extension T_0 of E_0 associated with θ only depends on π . Associated with π , there is thus a quadratic extension T/T_0 .

Remark 4.30. When π has level 0, one has $T_0 = F_0$.

It follows from Proposition 4.7 that π contains

- only one G^σ -conjugacy class of σ -self-dual types if T/T_0 is unramified,
- $\lfloor m/2 \rfloor + 1$ different G^σ -conjugacy classes of σ -self-dual types if T/T_0 is ramified.

Among these G^σ -conjugacy classes of σ -self-dual types, one is of particular importance.

Definition 4.31. A σ -self-dual type (\mathbf{J}, λ) is said to be *generic* if either T/T_0 is unramified, or T/T_0 is ramified and the integer i of Proposition 4.5(2.b) is equal to $\lfloor m/2 \rfloor$.

A σ -self-dual cuspidal representation of G thus contains, up to G^σ -conjugacy, a unique generic σ -self-dual type. The next result is [35] Theorem 10.3 (see also [3] Section 6).

Proposition 4.32. *Let π be a σ -self-dual cuspidal representation of G and (\mathbf{J}, λ) be its generic σ -self-dual type. Then π is distinguished if and only if λ is $\mathbf{J} \cap G^\sigma$ -distinguished.*

If (\mathbf{J}, λ) is a σ -self-dual type, and if θ is the unique simple character contained in λ such that $\mathbf{J}_\theta = \mathbf{J}$, we will write λ_w for the unique representation κ_θ of \mathbf{J} extending the Heisenberg representation of θ given by Proposition 4.16. The next result extends [35] Propositions 7.9, 9.8 to the case of arbitrary σ -self-dual cuspidal representations.

Proposition 4.33. *Let π be a σ -self-dual cuspidal representation of G . Let (\mathbf{J}, λ) be a generic σ -self-dual type in π and τ be the representation of \mathbf{J} trivial on \mathbf{J}^1 such that λ is isomorphic to $\lambda_w \otimes \tau$. Then π is distinguished if and only if τ is $\mathbf{J} \cap G^\sigma$ -distinguished.*

Proof. This follows from Proposition 4.32 together with the fact that

$$\text{Hom}_{\mathbf{J} \cap G^\sigma}(\lambda, R) \simeq \text{Hom}_{\mathbf{J} \cap G^\sigma}(\lambda_w, R) \otimes \text{Hom}_{\mathbf{J} \cap G^\sigma}(\tau, R)$$

and $\text{Hom}_{\mathbf{J} \cap G^\sigma}(\lambda_w, R)$ has dimension 1 (see Proposition 4.6(4)). □

Fix isomorphisms

$$B \simeq \mathbf{M}_m(E), \quad \mathbf{J}^0/\mathbf{J}^1 \simeq \text{GL}_m(I), \tag{4.17}$$

as in Proposition 4.5(4). The restriction of τ to $\mathbf{J} \cap B^\times$ is a generic σ -self-dual type of level 0 in $B^\times \simeq \text{GL}_m(E)$ and \mathbf{J}/\mathbf{J}^1 is naturally isomorphic to $(\mathbf{J} \cap B^\times)/(\mathbf{J}^1 \cap B^\times)$. The representation τ is thus distinguished by $\mathbf{J} \cap G^\sigma$ if and only if its restriction to $\mathbf{J} \cap \text{GL}_m(E)$ is distinguished by $\mathbf{J} \cap \text{GL}_m(E_0)$. Proposition 4.33 used twice thus implies that π is distinguished by G^σ if and only if the cuspidal representation of level 0 of $\text{GL}_m(E)$ compactly induced from the restriction of τ to $\mathbf{J} \cap \text{GL}_m(E)$ is distinguished by $\text{GL}_m(E_0)$.

However, the field extension E is not canonical. In 4.10, we will canonically associate with π a σ -self-dual cuspidal representation π_t of level 0 of $\text{GL}_m(T)$, which is $\text{GL}_m(T_0)$ -distinguished if and only if

π is G^σ -distinguished, where T/T_0 is the quadratic extension associated with π . Our strategy is inspired from [8] Section 3.

The following proposition relates the parity of m/r to the ramification of T/T_0 .

Proposition 4.34. *Let π be a σ -self-dual cuspidal representation of $\mathrm{GL}_n(F)$ with quadratic extension T/T_0 , and write $m = m(\pi)$, $r = r(\pi)$. Then*

$$m/r \text{ is } \begin{cases} \text{odd if } T/T_0 \text{ is unramified,} \\ \text{either even or equal to 1 if } T/T_0 \text{ is ramified.} \end{cases}$$

Proof. Write π as $\mathrm{St}_r(\rho)$ as in Proposition 3.8 with $\rho^{\sigma^\vee} \simeq \rho \nu^i$ for some $i \in \{0, 1\}$. Then $\rho \nu^{i/2}$ is a σ -self-dual supercuspidal representation of $\mathrm{GL}_{n/r}(F)$, and the quadratic extension associated with it is T/T_0 . Applying [35] Propositions 8.1, 9.8, we get the expected result. \square

4.10.

In order to prove Theorem 4.41, it will be useful to consider the slightly more general situation where π is a cuspidal representation of G with σ -self-dual endo-class Θ . Thus π itself needs not be σ -self-dual. However, it has a relative degree m and, since Θ is σ -self-dual, there is a quadratic extension T/T_0 associated with it. Moreover, by Proposition 4.7, it contains, up to G^σ -conjugacy, a unique generic σ -self-dual maximal simple character θ . Let \mathbf{J} be its normalizer in G and κ_θ be the representation of \mathbf{J} given by Proposition 4.16. Then π contains a unique type of the form

$$(\mathbf{J}, \kappa_\theta \otimes \tau) \tag{4.18}$$

for a uniquely determined irreducible representation τ of \mathbf{J} trivial on \mathbf{J}^1 . Fix a σ -self-dual simple stratum $[\alpha, \beta]$ and isomorphisms (4.17) as in Proposition 4.5.

First, we define an open and compact mod centre subgroup \mathbf{J}_t of $\mathrm{GL}_m(T)$ as follows:

- if T/T_0 is unramified, \mathbf{J}_t is the normalizer of $\mathrm{GL}_m(\mathcal{O}_T)$ in $\mathrm{GL}_m(T)$,
- if T/T_0 is ramified, and if t is a uniformizer of T such that $\sigma(t) = -t$, then \mathbf{J}_t is the normalizer in $\mathrm{GL}_m(T)$ of the conjugate of $\mathrm{GL}_m(\mathcal{O}_T)$ by the diagonal element

$$\mathrm{diag}(t, \dots, t, 1, \dots, 1) \in \mathrm{GL}_m(T),$$

where t occurs $\lfloor m/2 \rfloor$ times.

The group \mathbf{J}_t (which does not depend on the choice of t in the ramified case) has a unique maximal compact subgroup \mathbf{J}_t^0 and a unique normal maximal pro- p -subgroup \mathbf{J}_t^1 . The natural group isomorphism

$$\mathbf{J}_t^0/\mathbf{J}_t^1 \simeq \mathrm{GL}_m(I) \tag{4.19}$$

transports the action of $\sigma \in \mathrm{Gal}(T/T_0)$ on $\mathbf{J}_t^0/\mathbf{J}_t^1$ to

- the action of the nontrivial element of $\mathrm{Gal}(I/I_0)$ on $\mathrm{GL}_m(I)$ if T/T_0 is unramified,
- the adjoint action of

$$\begin{pmatrix} -\mathrm{id}_{\lfloor m/2 \rfloor} & 0 \\ 0 & \mathrm{id}_{m-\lfloor m/2 \rfloor} \end{pmatrix} \in \mathrm{GL}_m(I), \tag{4.20}$$

on $\mathrm{GL}_m(I)$ if T/T_0 is ramified.

Remark 4.35. When T/T_0 is ramified, the isomorphism (4.19) depends on the choice of t : changing t to another uniformizer t' conjugates the isomorphism by the σ -invariant element

$$\mathrm{diag}(\alpha, \dots, \alpha, 1, \dots, 1) \in \mathrm{GL}_m(I),$$

where α (which occurs i times) is the image of $t't^{-1}$ in I^\times . This element is central in $\mathrm{GL}_m(I)^\sigma$.

We now associate to τ an irreducible representation τ_t of \mathbf{J}_t trivial on \mathbf{J}_t^1 . On the one hand, the restriction of τ to \mathbf{J}^0 is the inflation of an irreducible cuspidal representation V of $\mathrm{GL}_m(\mathbf{I})$. On the other hand, the restriction of τ to E^\times is a multiple of $\omega \circ N_{E/T}$ for a uniquely determined tamely ramified character ω of T^\times : See 4.7. Note that $[E : T] = p^e$ for some $e \geq 1$.

Lemma 4.36. *Let V and ω be as above.*

1. *There is a unique representation τ_t of \mathbf{J}_t trivial on \mathbf{J}_t^1 such that*
 - (a) *the restriction of τ_t to T^\times is a multiple of the character ω ,*
 - (b) *the restriction of τ_t to \mathbf{J}_t^0 is the inflation of $V^{(p^{-e})}$, where $V^{(p^{-e})}$ is the representation of $\mathrm{GL}_m(\mathbf{I})$ obtained from V by applying the field automorphism $x \mapsto x^{p^{-e}}$.*
2. *The pair (\mathbf{J}_t, τ_t) is a level 0 type in $\mathrm{GL}_m(T)$.*
3. *Up to isomorphism, the representation τ_t only depends on τ , and not on the choice of the σ -self-dual simple stratum $[\mathfrak{a}, \beta]$, the uniformizer t and the identification $\mathbf{J}^0/\mathbf{J}^1 \simeq \mathrm{GL}_m(\mathbf{I})$.*

Proof. Uniqueness follows from the fact that \mathbf{J}_t is generated by \mathbf{J}_t^0 and T^\times , and the existence of τ_t follows from the fact that the restriction of $V^{(p^{-e})}$ to \mathbf{I}^\times is a multiple of the character of \mathbf{I}^\times defined by the restriction of ω to the units of T^\times . Since $V^{(p^{-e})}$ is cuspidal, the pair (\mathbf{J}_t, τ_t) is a level 0 type by construction. It remains to prove (3). Since it will require techniques which are not used anywhere else in the paper, we will prove it apart, in Paragraph 4.14. □

It will be convenient to give another description of the representation τ_t .

- Lemma 4.37.** 1. *There is a unique group isomorphism $\pi : \mathbf{J}/\mathbf{J}^1 \rightarrow \mathbf{J}_t/\mathbf{J}_t^1$ such that*
- (a) *its restriction to $\mathrm{GL}_m(\mathbf{I})$ is the automorphism acting entrywise by $\phi : x \mapsto x^{p^e}$,*
 - (b) *for all $x \in E^\times$, the image of $x\mathbf{J}^1$ is $N_{E/T}(x)\mathbf{J}_t^1$.*
2. *The isomorphism π is σ -equivariant.*
3. *The representation τ_t is isomorphic to $\tau \circ \pi^{-1}$.*

Proof. Again, uniqueness follows from the fact that \mathbf{J} is generated by \mathbf{J}^0 and E^\times . Existence follows from the fact that $N_{E/T}(x) = x^{p^e}$ for all $x \in \mathcal{O}_E^\times$ of order prime to p and that $N_{E/T}$ induces a group isomorphism from $E^\times/(1 + \mathfrak{p}_E)$ to $T^\times/(1 + \mathfrak{p}_T)$.

Define π_1 to be $\sigma \circ \pi \circ \sigma^{-1}$. The restriction of π_1 to E^\times corresponds to $\sigma \circ N_{E/T} \circ \sigma^{-1}$, which is equal to $N_{E/T}$ since E and T are stable by σ . The restriction of π_1 to $\mathrm{GL}_m(\mathbf{I})$ is

- the automorphism defined by making $\sigma \circ \phi \circ \sigma^{-1} = \phi \in \mathrm{Gal}(\mathbf{I}/\mathbb{F}_p)$ act entrywise if T/T_0 is unramified,
- the automorphism $\mathrm{Ad}(\delta^{-1}\phi(\delta)) \circ \phi = \phi$ if T/T_0 is ramified, where δ is the ϕ -invariant matrix defined by equation (4.5).

The fact that π is σ -equivariant now follows from its uniqueness, and (3) is immediate. □

Now, let us describe the behavior of $\tau \mapsto \tau_t$ with respect to duality and distinction.

- Lemma 4.38.** 1. *The representation τ_t is σ -self-dual if and only if τ is σ -self-dual.*
2. *The representation τ_t is $\mathbf{J}_t \cap \mathrm{GL}_m(T_0)$ -distinguished if and only if τ is $\mathbf{J} \cap G^\sigma$ -distinguished.*

Proof. Saying that τ is σ -self-dual is equivalent to saying that the representation V and the character $\omega \circ N_{E/T}$ are σ -self-dual. Assertion (1) follows from the fact that $(V^{(p^{-e})})^{\vee\sigma}$ is isomorphic to $(V^{\vee\sigma})^{(p^{-e})}$ and $(\omega \circ N_{E/T})^{-1} \circ \sigma$ is equal to $(\omega^{-1} \circ \sigma) \circ N_{E/T}$.

Assertion (2) follows from the fact that $\tau_t \circ \pi = \tau$ and π maps $(\mathbf{J}/\mathbf{J}^1)^\sigma$ to $(\mathbf{J}_t/\mathbf{J}_t^1)^\sigma$. □

Corollary 4.39. *The pair (\mathbf{J}_t, τ_t) is a generic σ -self-dual type if and only if $(\mathbf{J}, \kappa_\theta \otimes \tau)$ is a generic σ -self-dual type.*

Proof. This follows from Lemma 4.38(1), thanks to our choice of \mathbf{J}_t (see equation (4.20)). □

4.11.

We still are in the situation of Paragraph 4.10. Consider the compactly induced representation

$$\pi_t = \text{ind}_{\mathbf{J}_t}^{\text{GL}_m(T)}(\tau_t). \tag{4.21}$$

It satisfies the following properties.

- Proposition 4.40.** 1. *The representation π_t is cuspidal, irreducible and has level 0.*
 2. *One has $m(\pi_t) = m$ and $r(\pi_t) = r$.*
 3. *The representation π_t is σ -self-dual if and only if π is σ -self-dual.*
 4. *The representation π_t is $\text{GL}_m(T_0)$ -distinguished if and only if π is $\text{GL}_n(F_0)$ -distinguished.*

Proof. Assertion (1) follows from the fact that π_t is compactly induced from a level 0 type in $\text{GL}_m(T)$ (see Lemma 4.36 and Remark 4.24). The first equality of Assertion (2) follows from Remark 4.24, and the second one from Lemma 4.25.

Suppose that π is σ -self-dual. Then τ is σ -self-dual (see 4.9). By Lemma 4.38, the representation τ_t is σ -self-dual as well. By compact induction, it follows that π_t is σ -self-dual. The argument also works the other way round, proving (3). Assertion (4) follows from Proposition 4.33 together with Lemma 4.38(2). □

Theorem 4.41. 1. *The process*

$$\pi \mapsto \pi_t \tag{4.22}$$

induces a bijection from the set of isomorphism classes of cuspidal representations of G with endo-class Θ to that of cuspidal representations of level 0 of $\text{GL}_m(T)$.

2. *The bijection (4.22) maps σ -self-dual representations onto σ -self-dual representations and G^σ -distinguished representations onto $\text{GL}_m(T_0)$ -distinguished representations.*
 3. *For any cuspidal representation π with endo-class Θ and any tamely ramified character χ of F^\times , the representation $(\pi\chi)_t$ is isomorphic to $\pi_t(\chi \circ N_{T/F})$.*

Proof. For (1), let π_0 be a cuspidal representation of level 0 of $\text{GL}_m(T)$. It contains a level 0 type (\mathbf{J}_t, τ_0) for a uniquely determined representation τ_0 of \mathbf{J}_t trivial on \mathbf{J}_t^1 . It then suffices to check that the process

$$\pi_0 \mapsto \text{ind}_{\mathbf{J}}^G(\kappa_\theta \otimes (\tau_0 \circ \pi))$$

gives the inverse bijection. For (3), notice that if π contains the type $(\mathbf{J}, \kappa_\theta \otimes \tau)$, then $\pi\chi$ contains the type $(\mathbf{J}, \kappa_\theta \otimes \tau\chi^{\mathbf{J}})$, where $\chi^{\mathbf{J}}$ is the unique character of \mathbf{J} trivial on \mathbf{J}^1 whose restriction to $\mathbf{J} \cap B^\times \simeq \text{GL}_m(E)$ is equal to $(\chi \circ N_{E/F}) \circ \det_E$ where \det_E is the determinant on $B \simeq \mathbf{M}_m(E)$. Then $(\tau\chi^{\mathbf{J}})_t$ is isomorphic to the representation τ_t twisted by the character of \mathbf{J}_t trivial on \mathbf{J}_t^1 given by $(\chi \circ N_{T/F}) \circ \det_T$, where \det_T is the determinant on $\mathbf{M}_m(T)$. Assertion (2) is given by Proposition 4.40. □

Corollary 4.42. *Let μ be a tamely ramified character of F_0^\times . A cuspidal representation π of $\text{GL}_n(F)$ with endo-class Θ is distinguished by μ if and only if π_t is distinguished by $\mu \circ N_{T_0/F_0}$.*

Proof. Fix a tamely ramified character ξ of F^\times extending μ . Then π is μ -distinguished if and only if $\pi\xi^{-1}$ is distinguished, and $(\pi\xi^{-1})_t$ is isomorphic to $\pi_t(\xi^{-1} \circ N_{T/F})$. Thus, π is μ -distinguished if and only if π_t is distinguished by the character $\xi \circ N_{T/F}|_{T_0^\times} = \mu \circ N_{T_0/F_0}$. □

Finally, let us describe the compatibility of the process (4.22) with the description of cuspidal representations in terms of supercuspidal ones of 4.8.

Proposition 4.43. *Let π be a cuspidal representation of G with endo-class Θ and $r = r(\pi)$. Let ρ be a supercuspidal representation of $\text{GL}_{n/r}(F)$ such that π is isomorphic to $\text{St}_r(\rho)$. Then π_t is isomorphic to $\text{St}_r(\rho_t)$.*

Remark 4.44. Note that this makes sense since, by Corollary 4.27, the representations π, ρ have the same endo-class Θ , thus the same quadratic extension T/T_0 .

Proof. The representation π contains a type of the form $(\mathbf{J}, \kappa_\theta \otimes \tau)$ for a unique representation τ of \mathbf{J} trivial on \mathbf{J}^1 . Fix a σ -self-dual simple stratum $[\mathfrak{a}, \beta]$ and isomorphisms (4.17) as in Proposition 4.5. Associated with τ , there are a tamely ramified character ω of T , and a cuspidal representation $V = \text{st}_r(\varrho)$ of $\text{GL}_m(I)$, for some supercuspidal representation ϱ of $\text{GL}_{m/r}(I)$. The representation τ is entirely determined by the fact that

- its restriction to \mathbf{J}^0 is the inflation of V ,
- its restriction to E^\times is a multiple of the character $\omega \circ N_{E/T}$.

We now use the results of 4.4 and 4.5 for $u = r$. Let θ_* be the simple character associated with θ by Lemma 4.9. Thanks to Corollary 4.20, the representation κ_{θ_*} corresponds to κ_θ via the map (4.10). Paragraph 4.8 says that ρ contains the type $(\mathbf{J}_*, \kappa_{\theta_*} \otimes \tau_*)$, where τ_* is the representation of \mathbf{J}_* trivial on \mathbf{J}_*^1 determined by

- its restriction to \mathbf{J}_*^0 is the inflation of ϱ ,
- its restriction to E^\times is a multiple of $\omega_* \circ N_{E/T}$, where ω_* is a tamely ramified character of T^\times such that $\omega_*^r = \omega$.

Thus, ρ_t is compactly induced from the level 0 type $(\mathbf{J}_{*,t}, \tau_{*,t})$ where $\tau_{*,t}$ is determined by

- its restriction to $\mathbf{J}_{*,t}^0$ is the inflation of $\varrho^{(p^{-e})}$,
- its restriction to T^\times is a multiple of ω_* .

Thus, $\text{St}_r(\rho_t)$ is compactly induced from the level 0 type (\mathbf{J}_t, δ) , where δ is determined by

- its restriction to \mathbf{J}_t^0 is the inflation of $\text{st}_r(\varrho^{(p^{-e})}) \simeq V^{(p^{-e})}$,
- its restriction to T^\times is a multiple of $\omega_*^r = \omega$.

It follows that δ is isomorphic to τ_t , whence $\text{St}_r(\rho_t)$ is isomorphic to π_t . □

4.12.

Finally, let π be a σ -self-dual cuspidal representation of G , of level 0. It has a central character c_π and its generic type (\mathbf{J}, λ) defines a cuspidal representation V of $\text{GL}_n(\mathbf{k})$. Assume that $n \neq 1$. In the spirit of Proposition 4.33, we give a necessary and sufficient condition for π to be distinguished by $\text{GL}_n(F_0)$ in terms of c_π and V .

Theorem 4.45. *The representation π is $\text{GL}_n(F_0)$ -distinguished if and only if its central character c_π is trivial on F_0^\times and*

1. if F/F_0 is unramified, then V is $\text{GL}_n(\mathbf{k}_0)$ -distinguished,
2. if F/F_0 is ramified, then n is even, V is $\text{GL}_{n/2}(\mathbf{k}) \times \text{GL}_{n/2}(\mathbf{k})$ -distinguished and

$$s = \begin{pmatrix} 0 & \text{id} \\ \text{id} & 0 \end{pmatrix} \in \text{GL}_n(\mathbf{k}) \tag{4.23}$$

acts on the space of its $\text{GL}_{n/2}(\mathbf{k}) \times \text{GL}_{n/2}(\mathbf{k})$ -invariant linear forms by the sign $c_\pi(\varpi)$, where ϖ is any uniformizer of F .

Note that, if π is distinguished and F/F_0 is ramified, the space of $\text{GL}_{n/2}(\mathbf{k}) \times \text{GL}_{n/2}(\mathbf{k})$ -invariant linear forms on V has dimension 1 by Proposition 3.13.

Proof. By Proposition 4.33, the representation π is $\text{GL}_n(F_0)$ -distinguished if and only if λ is $\mathbf{J} \cap G^\sigma$ -distinguished. In the unramified case, the result follows from the fact that $\mathbf{J} \cap G^\sigma$ is generated by F_0^\times and $\mathbf{J}^0 \cap G^\sigma$ (see Lemma 4.14) and $(\mathbf{J}^0 \cap G^\sigma)/(\mathbf{J}^1 \cap G^\sigma)$ identifies with $\text{GL}_n(\mathbf{k}_0)$.

Assume now that we are in the ramified case. Since c_π is trivial on F_0^\times and $1 + \mathfrak{p}_F$, its value at ϖ does not depend on the choice of a uniformizer ϖ of F . We thus may and will assume that $\sigma(\varpi) = -\varpi$, thus $c_\pi(\varpi) \in \{-1, 1\}$. By Lemma 4.14 again, $\mathbf{J} \cap G^\sigma$ is generated by F_0^\times , $\mathbf{J}^0 \cap G^\sigma$ and ϖw , where the element $w \in \mathbf{J}^0$ is defined by equation (4.12). The quotient $(\mathbf{J}^0 \cap G^\sigma)/(\mathbf{J}^1 \cap G^\sigma)$ identifies with $\mathrm{GL}_{n/2}(\mathbf{k}) \times \mathrm{GL}_{n/2}(\mathbf{k})$, the image of w in $\mathbf{J}^0/\mathbf{J}^1 \simeq \mathrm{GL}_n(\mathbf{k})$ is the element s and ϖs acts on the (one-dimensional) space of $\mathrm{GL}_{n/2}(\mathbf{k}) \times \mathrm{GL}_{n/2}(\mathbf{k})$ -invariant linear forms on V as $c_\pi(\varpi)s$. \square

Putting Theorems 4.41 and 4.45 together, we have thus reduced the problem of characterizing distinguished cuspidal representations of $\mathrm{GL}_n(F)$ to a problem about distinction of cuspidal representations of finite general linear groups.

4.13.

As an application of Theorem 4.45, we classify all distinguished cuspidal representations of $\mathrm{GL}_2(F)$. The supercuspidal case has been treated in [35] and the case where $\ell \neq 2$ has been treated in [12]. It thus remains the case of nonsupercuspidal cuspidal representations for $\ell = 2$.

Proposition 4.46. *Suppose that $\ell = 2$. The $\mathrm{GL}_2(F_0)$ -distinguished cuspidal nonsupercuspidal representations of $\mathrm{GL}_2(F)$ are the representations $\mathrm{St}_2(\chi)$, where χ is a character of F^\times trivial on F_0^\times .*

Proof. According to Lemma 3.7, and since the character ν is trivial when $\ell = 2$, the σ -self-dual nonsupercuspidal cuspidal representations of $\mathrm{GL}_2(F)$ are the representations of the form $\mathrm{St}_2(\chi)$ for a uniquely determined σ -self-dual character χ of F^\times . Since $N_{F/F_0}(F^\times)$ has index 2 in F_0^\times , any σ -self-dual character of F^\times is trivial on F_0^\times . We are thus reduced to proving that the representation $\pi = \mathrm{St}_2(1)$ is distinguished.

This representation has level 0 and trivial central character. By Theorem 4.45, it is distinguished if and only if the cuspidal representation $V = \mathrm{st}_2(1)$ of $\mathrm{GL}_2(\mathbf{k})$ is

- distinguished by $\mathrm{GL}_2(\mathbf{k}_0)$ if F/F_0 is unramified,
- distinguished by $\mathbf{k}^\times \times \mathbf{k}^\times$ if F/F_0 is ramified.

(The condition on the action of s disappears in the ramified case since $-1 \equiv 1 \pmod{\ell = 2}$.) We are thus reduced to proving that V is distinguished.

Let W denote the parabolic induction to $\mathrm{GL}_2(\mathbf{k})$ of the trivial character of the Borel subgroup B made of all upper triangular matrices. By [40], it is indecomposable of length 3, its socle and cosocle are irreducible and isomorphic to the trivial character and its remaining irreducible subquotient is V .

Let us assume first that F/F_0 is unramified. Then the action of $\mathrm{GL}_2(\mathbf{k}_0)$ on $B \backslash \mathrm{GL}_2(\mathbf{k})$ has two orbits \mathcal{O}_0 and \mathcal{O}_1 . If one identifies $B \backslash \mathrm{GL}_2(\mathbf{k})$ with the projective line $\mathbb{P}^1(\mathbf{k})$, then $\mathcal{O}_0 = \mathbb{P}^1(\mathbf{k}_0)$ and $\mathcal{O}_1 = \mathbb{P}^1(\mathbf{k}) \setminus \mathbb{P}^1(\mathbf{k}_0)$. It follows that W has a two-dimensional space of $\mathrm{GL}_2(\mathbf{k}_0)$ -invariant linear forms, generated by Λ_0 and Λ_1 with

$$\Lambda_i(f) = \sum_{x \in \mathcal{O}_i} f(x), \quad f \in W, \quad i \in \{0, 1\}.$$

These two linear forms vanish on the subspace R of constant functions. They thus define nonzero linearly independent $\mathrm{GL}_2(\mathbf{k}_0)$ -invariant linear forms on $X = W/R$. If both of them vanish on V , then the 1-dimensional quotient X/V has a two-dimensional space of invariant linear forms: contradiction. It thus follows that V is distinguished by $\mathrm{GL}_2(\mathbf{k}_0)$.

Now, assume that F/F_0 is ramified. Then the action of $\mathbf{k}^\times \times \mathbf{k}^\times$ on $B \backslash \mathrm{GL}_2(\mathbf{k})$ has three orbits \mathcal{O}_∞ , \mathcal{O}_0 and \mathcal{O}_1 , the first two orbits being single points. They define three linearly independent invariant linear forms Λ_∞ , Λ_0 and Λ_1 . The two linear forms $\Lambda_\infty - \Lambda_0$ and Λ_1 vanish on the subspace R of constant functions. From there, an argument similar to that of the unramified case shows that V is distinguished by $\mathbf{k}^\times \times \mathbf{k}^\times$. \square

4.14.

In this paragraph, we prove Lemma 4.36(3). First, by Remarks 4.2, 4.35, changing the isomorphism (4.17) and t does not affect the isomorphism class of τ_t . Let $[\alpha, \beta']$ be another σ -self-dual maximal simple stratum such that $\theta \in \mathcal{C}(\alpha, \beta')$. Conjugating by \mathbf{J}^1 , we may and will assume that the maximal tamely ramified extension of F in $E' = F[\beta']$ is T . This gives us another isomorphism π' from \mathbf{J}/\mathbf{J}^1 to $\mathbf{J}_t/\mathbf{J}_t^1$. By construction, it coincides with π on $\mathbf{J}^0/\mathbf{J}^1$ and the image of $x\mathbf{J}^1$ by π' is equal to $N_{E'/T}(x)\mathbf{J}^1$ for all $x \in E'^\times$. We are going to prove that π' is equal to π . The result will then follow from the fact that τ_t is equal to $\tau \circ \pi$. For this, it suffices to prove that π and π' take the same value at some given uniformizer of E' . Let ϖ, ϖ' be uniformizers of E, E' , respectively.

The centre of \mathbf{J}/\mathbf{J}^1 is $E^\times\mathbf{J}^1/\mathbf{J}^1 = E'^\times\mathbf{J}^1$, thus $E'^\times \subseteq E^\times\mathbf{J}^1$. We thus may write $\varpi' \in \varpi\zeta\mathbf{J}^1$ for some root of unity ζ of T^\times of order prime to p . Changing ϖ' to $\varpi'\zeta^{-1}$, we may and will assume that $\varpi' \in \varpi\mathbf{J}^1$. It suffices to prove the following claim.

Claim 4.47. We have $N_{E'/T}(\varpi') \equiv N_{E/T}(\varpi) \pmod{1 + \mathfrak{p}_T}$.

First, this is true when $m = 1$. Indeed, writing G_T for the centralizer of T in G and \det_T for the determinant on G_T , we have $\det_T(x) = N_{E/T}(x)$ for all $x \in E^\times$, thus

$$N_{E'/T}(\varpi') = \det_T(\varpi') \in \det_T(\varpi) \cdot \det_T(\mathbf{J}^1 \cap G_T) = N_{E/T}(\varpi) \cdot (1 + \mathfrak{p}_T).$$

Now, assume that $m > 1$. We use the results of 4.4 for $u = m$. Let $\theta_* \in \mathcal{C}(\alpha_*, \beta)$ denote the transfer of θ as in Lemma 4.9. Fix a T -embedding

$$\iota : E' \rightarrow \text{End}_T(V_*) \subseteq \text{End}_F(V_*)$$

such that α_* is normalized by $\iota E'^\times$, and transfer θ to $\theta_\bullet \in \mathcal{C}(\alpha_*, \iota\beta')$ in the sense of [9] 3.6. The simple character θ is in $\mathcal{C}(\alpha, \beta) \cap \mathcal{C}(\alpha, \beta')$. It follows from [6] Theorem 8.7 that θ_*, θ_\bullet intertwine in G_* , and from [9] Theorem 3.5.11 that $\theta_\bullet = \theta_*^x$ for some $x \in \mathcal{K}_{\alpha_*}$. Changing ι to $\text{Ad}(x) \circ \iota$, we thus may assume that $\theta_\bullet = \theta_* \in \mathcal{C}(\alpha_*, \beta) \cap \mathcal{C}(\alpha_*, \iota\beta')$. By using ι , we get a diagonal embedding

$$E' \rightarrow \text{End}_T(V_*) \times \cdots \times \text{End}_T(V_*) \subseteq \text{End}_T(V)$$

denoted ϕ , which is the identity on T^\times . The Skolem–Noether theorem implies that $\phi = \text{Ad}(g)$ for some $g \in G_T$. Conjugating by g , we thus may assume that E^\times and E'^\times are both diagonal in M . The identity $\varpi' \in \varpi\mathbf{J}^1$ thus implies $\varpi' \in \varpi\mathbf{J}_*^1$. We are thus reduced to the case where $m = 1$.

Remark 4.48. The fact that τ_t does not depend on the choice of β is claimed in [8] Lemma 3.6. However, Property (b) of this lemma does not hold: Using the notation of *ibid.*, the restriction of $\lambda_\xi^{\mathbf{J}}$ to P^\times is a multiple of the character $\xi \circ N_{P/T}$, whereas the restriction of $(\xi|_{T^\times})^{\mathbf{J}}$ to P^\times is $(\xi \circ N_{P/T})^s$. (Note that P corresponds to our E , and s corresponds to our m .)

5. The odd case

In this section, p is odd, ℓ is any prime number different from p and the field R has characteristic ℓ . This section is devoted to the proof of the following theorem.

Theorem 5.1. *Let π be a σ -self-dual cuspidal nonsupercuspidal R -representation of $\text{GL}_n(F)$. Assume that the integer $r = r(\pi)$ is odd, thus π is isomorphic to $\text{St}_r(\rho)$ for a uniquely determined σ -self-dual supercuspidal representation ρ of $\text{GL}_k(F)$, with $k = n/r$. If π is $\text{GL}_n(F_0)$ -distinguished, then*

1. *the relative degree $m = m(\pi)$ and the ramification index of T/T_0 have the same parity,*
2. *the representation ρ is $\text{GL}_k(F_0)$ -distinguished.*

Note that the fact that r is odd and $r \neq 1$ implies that $\ell \neq 2$.

5.1.

Before we start the proof of Theorem 5.1, let us prove the following disjunction theorem.

Corollary 5.2. *Let π be a σ -self-dual cuspidal R -representation of $GL_n(F)$. Assume that $r(\pi)$ is odd. Then π cannot be both distinguished and \varkappa -distinguished.*

Proof. Assume that π is both distinguished and \varkappa -distinguished, and let χ be a tamely ramified character of F^\times extending \varkappa . Then $\pi\chi$ is distinguished, it is isomorphic to $St_r(\rho\chi)$ and $\rho\chi$ is supercuspidal and σ -self-dual. Theorem 5.1 applied to both π and $\pi\chi$ implies that ρ is both distinguished and \varkappa -distinguished. This contradicts Theorem 3.2. □

We also have the following Distinguished Lift Theorem.

Corollary 5.3. *Let π be a $GL_n(F_0)$ -distinguished cuspidal $\overline{\mathbb{F}}_\ell$ -representation of $GL_n(F)$ with $r(\pi)$ odd. There is a $GL_n(F_0)$ -distinguished integral generic $\overline{\mathbb{Q}}_\ell$ -representation of $GL_n(F)$ whose reduction mod ℓ contains π .*

Proof. Write π as $St_r(\rho)$ with ρ distinguished. Let μ be a distinguished integral cuspidal lift of ρ , which exists by Theorem 3.5. Then the generic representation $St_r(\mu)$ satisfies the required property (see [1] Theorem 1.3 or [26] Corollary 4.2 when F has characteristic zero, and observe that the argument in [26] holds verbatim in positive characteristic thanks to [22, Theorem 4.7]). □

Remark 5.4. A $GL_n(F_0)$ -distinguished integral generic $\overline{\mathbb{Q}}_\ell$ -representation of $GL_n(F)$ as in Corollary 5.3 may not be cuspidal. See Section 6 for the classification of all distinguished cuspidal $\overline{\mathbb{F}}_\ell$ -representations of $GL_n(F)$ having a cuspidal distinguished lift to $\overline{\mathbb{Q}}_\ell$.

Finally, compare Theorem 5.1 with the following finite field analogue.

Proposition 5.5. *Let k/k_0 be a quadratic extension of finite fields of characteristic p . Let ϱ be a supercuspidal R -representation of $GL_f(k)$ for some $f \geq 1$, and r be an odd integer such that $st_r(\varrho)$ is cuspidal. If $st_r(\varrho)$ is distinguished by $GL_{f,r}(k_0)$, then ϱ is distinguished by $GL_f(k_0)$.*

Proof. First, [35] Remark 4.3 tells us that $st_r(\varrho)$ is σ -self-dual (where σ is here the nontrivial automorphism of k/k_0). Proposition 3.9 implies that ϱ is σ -self-dual. By [35] Lemma 2.5, it is distinguished by $GL_f(k_0)$. □

5.2.

Let us prove Theorem 5.1(1). Since r is odd, m has the same parity as m/r , and, since π is nonsupercuspidal, we have $r > 1$, thus $m > 1$. It follows from [35] Proposition 7.1 that, if m is odd, T/T_0 is unramified, and from Proposition 4.34 that, if m is even, T/T_0 is ramified.

5.3.

We now start the proof of Theorem 5.1(2). We thus have a distinguished cuspidal representation π of $GL_n(F)$, which we write $St_r(\rho)$ with ρ supercuspidal and σ -self-dual.

Associated with π , there are a positive divisor m of n , a quadratic extension T/T_0 and a cuspidal representation π_τ of $GL_m(T)$. By Proposition 4.40, the representation π_τ has level 0, it is distinguished by $GL_m(T_0)$ and it satisfies $r(\pi_\tau) = r$.

Similarly, associated with ρ , there is a supercuspidal σ -self-dual representation ρ_τ of $GL_{m/r}(T)$, which has level 0, and is distinguished by $GL_{m/r}(T_0)$ if and only if ρ is distinguished by $GL_k(F_0)$. By Proposition 4.43, the representation π_τ is isomorphic to $St_r(\rho_\tau)$.

It follows that, in order to prove Theorem 5.1(2), we may assume that π has level 0.

5.4.

Let π be a distinguished cuspidal representation of level 0 of $GL_n(F)$. Associated with it, there are its central character c_π and a cuspidal representation V of $GL_n(k)$ (see §4.7).

The representation π is isomorphic to $St_r(\rho)$ for a unique σ -self-dual supercuspidal representation ρ , and ρ has level 0. Associated with ρ , there are its central character c_ρ and a supercuspidal representation ϱ of $GL_k(k)$. We have the relation

$$c_\pi = (c_\rho)^r \tag{5.1}$$

and, by Proposition 4.28, the representation V is isomorphic to $st_r(\varrho)$.

Since π is distinguished, its central character is trivial on F_0^\times . Since ρ is σ -self-dual, the restriction of c_ρ to F_0^\times has order at most 2. Restricting the relation (5.1) to F_0^\times , and since r is odd, we deduce that c_ρ is trivial on F_0^\times .

5.5.

In this paragraph, we will assume that F/F_0 is unramified. By Theorem 4.45, the representation V is distinguished by $GL_n(k_0)$. By [35] Remark 4.3, it is thus σ -self-dual, that is

$$st_r(\varrho) \simeq V \simeq V^{\sigma^\vee} \simeq st_r(\varrho^{\sigma^\vee}).$$

It follows from Proposition 3.9 that ϱ is σ -self-dual. By [35] Lemma 2.5, it is thus distinguished by $GL_k(k_0)$. Applying Theorem 4.45 again, we deduce that ρ is distinguished by $GL_k(F_0)$. This proves Theorem 5.1 in the unramified case.

5.6.

From now on, and until the end of this section, we assume that F/F_0 is ramified. By Theorem 4.45, we may write $n = 2u$ for some integer $u \geq 1$. We write $G = G_n = GL_n(k)$, $H = H_n = GL_u(k) \times GL_u(k)$ and $K = K_n$ for the normalizer of H in G , which is generated by H and

$$s = s_n = \begin{pmatrix} 0 & \text{id} \\ \text{id} & 0 \end{pmatrix} \in G,$$

where id is the identity in $GL_u(k)$. It will be convenient to introduce the following definition.

Definition 5.6. Let $c \in \{-1, 1\} \subseteq R^\times$. An irreducible R -representation V of G is said to be c -distinguished by H if V is H -distinguished and s acts on the space of H -invariant linear forms on V by multiplication by c .

By Theorem 4.45, the representation V is H -distinguished and s acts on the one-dimensional vector space $\text{Hom}_H(V, R)$ by the sign $c = c_\pi(\varpi)$. In other words, V is c -distinguished by H . We are now reduced to proving the following result. (Note that k is even since n is even and r is odd.)

Proposition 5.7. *The supercuspidal representation ϱ is c -distinguished by H_k .*

Indeed, since r is odd, the identity (5.1) together with Proposition 5.7 will give us $c = c_\rho(\varpi)$. It will then follow from Theorem 4.45 that ρ is $GL_k(F_0)$ -distinguished.

5.7.

Let π be an irreducible $\overline{\mathbb{F}}_\ell$ -representation of G . The natural map

$$\text{Hom}_{\overline{\mathbb{F}}_\ell H}(\pi, \overline{\mathbb{F}}_\ell) \otimes R \rightarrow \text{Hom}_{RH}(\pi \otimes R, \overline{\mathbb{F}}_\ell \otimes R)$$

defined by $f \otimes r \mapsto r(f \otimes \text{id})$ is an isomorphism of R -vector spaces. Moreover, these spaces have dimension at most 1, and it follows from this isomorphism that π is c -distinguished by H if and only if $\pi \otimes R$ is c -distinguished by H .

Since G is finite, any irreducible R -representation of G is defined over $\overline{\mathbb{F}}_\ell$, that is, isomorphic to $\pi_0 \otimes R$ for some irreducible $\overline{\mathbb{F}}_\ell$ -representation π_0 of G . In order to prove Proposition 5.7, we thus may assume that R is equal to $\overline{\mathbb{F}}_\ell$.

5.8.

From now on, we assume that R is equal to $\overline{\mathbb{F}}_\ell$. The remaining part of the section will be devoted to the proof of Proposition 5.7.

Lemma 5.8. *There exists a c -distinguished irreducible $\overline{\mathbb{Q}}_\ell$ -representation of G whose reduction mod ℓ contains V .*

Proof. Let χ denote the unique $\overline{\mathbb{F}}_\ell$ -character of K trivial on H such that $\chi(s) = c$. Since V is c -distinguished, it embeds in $\text{Ind}_K^G(\chi)$. Equivalently, the representation $\text{Ind}_K^G(\chi)$, which is self-dual (as χ is equal to χ^{-1}), surjects onto the contragredient W of V . Let Π be a projective indecomposable $\overline{\mathbb{F}}_\ell$ -representation of G whose unique irreducible quotient is isomorphic to W . Let $\tilde{\Pi}$ be the unique projective $\overline{\mathbb{Z}}_\ell$ -representation of G such that $\tilde{\Pi} \otimes \overline{\mathbb{F}}_\ell$ is isomorphic to Π . Let Λ be a surjective homomorphism from $\text{Ind}_K^G(\chi)$ to W . By projectivity, it defines a nonzero homomorphism Λ' from Π to $\text{Ind}_K^G(\chi)$, then a nonzero homomorphism Λ'' from $\tilde{\Pi}$ to $\text{Ind}_K^G(\tilde{\chi})$, where $\tilde{\chi}$ is the canonical $\overline{\mathbb{Z}}_\ell$ -lift of χ .

By inverting ℓ , we deduce that there is an irreducible $\overline{\mathbb{Q}}_\ell$ -representation X of G occurring in each of the semi-simple representations $J = \text{Ind}_K^G(\tilde{\chi}) \otimes \overline{\mathbb{Q}}_\ell$ and $\tilde{\Pi} \otimes \overline{\mathbb{Q}}_\ell$. It is thus c -distinguished and, by [39] 15.4, its reduction mod ℓ contains W .

Now, observe that, since $\tilde{\chi}$ is quadratic, J is self-dual. The contragredient of X is thus c -distinguished and its reduction mod ℓ contains V . □

5.9.

Let τ be a c -distinguished irreducible $\overline{\mathbb{Q}}_\ell$ -representation as in Lemma 5.8. Consider its cuspidal support: There are positive integers n_1, \dots, n_t such that $n_1 + \dots + n_t = n$ and, for each i in $\{1, \dots, t\}$, a cuspidal irreducible $\overline{\mathbb{Q}}_\ell$ -representation ρ_i of $\text{GL}_{n_i}(\mathbf{k})$ such that τ occurs as a component of the parabolically induced representation $\rho_1 \times \dots \times \rho_t$, denoted W . The representation W is thus c -distinguished. We claim the following.

Claim 5.9. There is an $i \in \{1, \dots, t\}$ such that n_i is even and ρ_i is c -distinguished by H_{n_i} .

Before proving this claim in the next paragraph, let us explain how it implies Proposition 5.7.

Propositions 3.9 and 3.11 imply that, for each $i \in \{1, \dots, t\}$, the reduction mod ℓ of ρ_i is irreducible and cuspidal, of the form $\text{st}_{r_i}(\varrho_i)$ for a unique positive integer r_i and a unique supercuspidal representation ϱ_i . Since the reduction mod ℓ of τ contains V , the representation V occurs as an irreducible component of the parabolically induced representation $r_\ell(\rho_1) \times \dots \times r_\ell(\rho_t)$. Uniqueness of the supercuspidal support implies that $\varrho_i \simeq \varrho$ for all i . It follows that either $r_i = 1$ or $r_i = e(\varrho)\ell^{v_i}$ for some $v_i \geq 0$. Observe that, as $r = e(\varrho)\ell^v$ for some $v \geq 0$ and r is odd, the integer $e(\varrho)$ is odd, thus r_i is odd in any case, for all i .

Fix an integer i as in Claim 5.9, and let ξ_i be a parameter for ρ_i in the sense of Definition 3.12. It is a $\text{Gal}(\mathbf{k}_{n_i}/\mathbf{k})$ -regular $\overline{\mathbb{Z}}_\ell$ -character of $\mathbf{k}_{n_i}^\times$. By Proposition 3.13, it is trivial on $\mathbf{k}_{u_i}^\times$, where u_i is defined by $n_i = 2u_i$, and it takes the unique element of $\mathbf{k}_{n_i}^\times/\mathbf{k}_{u_i}^\times$ of order 2 to $-c$.

Since the reduction mod ℓ of ρ_i is $\text{st}_{r_i}(\varrho)$, the reduction mod ℓ of ξ_i takes the form $\vartheta \circ N_{\mathbf{k}_{n_i}/\mathbf{k}}$, where ϑ is a parameter for ϱ .

Since $n_i = r_i k$ and r_i is odd, ϑ is trivial on \mathbf{k}_l^\times (where $k = 2l$) and takes the element of $\mathbf{k}_k^\times/\mathbf{k}_l^\times$ of order 2 to $-c$.

By Proposition 3.13, the canonical $\overline{\mathbb{Z}}_\ell$ -lift of ϑ is the parameter of a c -distinguished $\overline{\mathbb{Q}}_\ell$ -lift of ϱ , which implies that ϱ is c -distinguished (see Remark 3.14). This proves Proposition 5.7.

5.10.

The remaining part of this section will be devoted to the proof of Claim 5.9. We follow the argument of [28] Section 3, which simplifies in our situation since we deal with finite groups. Let \mathbf{A} denote the set of t -uples $\alpha = (\alpha_1, \dots, \alpha_t)$ where

- for each i , the element α_i is a family of $t + 1$ nonnegative integers of the form

$$\alpha_i = (n_{i,1}, \dots, n_{i,i-1}, n_{i,i}^+, n_{i,i}^-, n_{i,i+1}, \dots, n_{i,t})$$

of sum n_i ,

- one has $n_{1,1}^+ + \dots + n_{t,t}^+ = n_{1,1}^- \dots + n_{t,t}^-$ and $n_{i,j} = n_{j,i}$ for all $i \neq j$.

For an $\alpha \in \mathbf{A}$, it will be convenient to set $n_{i,i} = n_{i,i}^+ + n_{i,i}^-$ for each integer $i \in \{1, \dots, t\}$.

As in [28] 3.1, the set \mathbf{A} parametrizes the set of (P, H) -double cosets in G , where P is the parabolic subgroup of G generated by upper triangular matrices and the standard Levi subgroup M isomorphic to $G_{n_1} \times \dots \times G_{n_t}$. Let us explain how this parametrization works. Associated with any $\alpha \in \mathbf{A}$, there are

- a standard Levi subgroup

$$M_\alpha = (G_{n_{1,1}} \times G_{n_{1,2}} \times \dots \times G_{n_{1,t}}) \times \dots \times (G_{n_{t,1}} \times G_{n_{t,2}} \times \dots \times G_{n_{t,t}}) \subseteq M,$$

- a diagonal element

$$d_\alpha = \text{diag} \left(\begin{pmatrix} \text{id}_{n_{1,1}^+} & \\ & -\text{id}_{n_{1,1}^-} \end{pmatrix}, \text{id}_{n_{1,2}}, \dots, \text{id}_{n_{1,t}}, \dots, \text{id}_{n_{t,1}}, \text{id}_{n_{t,2}}, \dots, \begin{pmatrix} \text{id}_{n_{t,t}^+} & \\ & -\text{id}_{n_{t,t}^-} \end{pmatrix} \right) \in M_\alpha,$$

- a permutation matrix $w_\alpha \in G$ defined as follows: Decompose $\{1, \dots, n\}$ as the disjoint union of intervals $J_{i,j} = \{a_{i,j}, a_{i,j} + 1, \dots, b_{i,j}\}$ of length $n_{i,j}$, for each $i, j \in \{1, \dots, t\}$, where $a_{1,1} = 1$, $a_{i,j+1} = b_{i,j} + 1$ if $j \neq t$ and $a_{i+1,1} = b_{i,t} + 1$ if $i \neq t$; then w_α is the involution which
 - restricts to the identity on $J_{i,i}$ for each i ,
 - exchanges the intervals $J_{i,j}$ and $J_{j,i}$ if $i \neq j$, and sends the k th element of any of these intervals to the k th element of the other one, for all $k \in \{1, \dots, n_{i,j}\}$.

A system of representatives $(x_\alpha)_{\alpha \in \mathbf{A}}$ of (P, H) -double cosets in G is then obtained by any choice of $x_\alpha \in G$ such that

$$x_\alpha \begin{pmatrix} \text{id}_u & \\ & -\text{id}_u \end{pmatrix} x_\alpha^{-1} = e_\alpha, \tag{5.2}$$

where $e_\alpha = d_\alpha w_\alpha$.

Definition 5.10. An $\alpha \in \mathbf{A}$ is called *admissible* if, for any i , there exists a unique j such that $n_{i,j} \neq 0$. This defines an involution $\sigma_\alpha : i \mapsto j$ on $\{1, \dots, t\}$.

When this is the case, let us write H_α for the subgroup of M made of the $\text{diag}(g_1, \dots, g_t) \in M$ such that

$$\begin{aligned} g_{\sigma_\alpha(i)} &= g_i && \text{for all } i \in \{1, \dots, t\}, \\ g_i &\in \text{GL}_{n_{i,i}^+}(\mathbf{k}) \times \text{GL}_{n_{i,i}^-}(\mathbf{k}) && \text{for all } i \text{ fixed by } \sigma_\alpha. \end{aligned}$$

Moreover, if $n_{i,i}^+ = n_{i,i}^-$ for all $i \in \{1, \dots, t\}$, we define a matrix $k_\alpha = \text{diag}(k_1, \dots, k_t) \in M$ by

$$k_i = \text{id}_{n_i} = -k_{\sigma_\alpha(i)} \text{ if } i < \sigma_\alpha(i), \quad k_i = s_{n_i} \text{ if } i = \sigma_\alpha(i).$$

This matrix normalizes H_α , and we write K_α for the group generated by H_α and k_α .

We denote by θ_α the inner automorphism of the group $\text{PGL}_n(\mathbf{k})$ induced by conjugacy by e_α (which normalizes M_α). It is not hard to check that:

Lemma 5.11. *Let Z denote the centre of G .*

1. *An $\alpha \in \mathbf{A}$ is admissible if and only if M/Z is θ_α -stable in $G/Z = \text{PGL}_n(\mathbf{k})$.*
2. *Suppose that $\alpha \in \mathbf{A}$ is admissible. The preimage of $(M/Z)^{\theta_\alpha}$ in G , denoted by L_α , is*

$$\begin{cases} K_\alpha & \text{if } n_{ii}^+ = n_{ii}^- \text{ for all } i, \\ H_\alpha & \text{otherwise.} \end{cases}$$

When $L_\alpha = K_\alpha$, we denote by χ_α the character of K_α trivial on H_α and sending k_α to c . Otherwise, we set χ_α to be the trivial character of $L_\alpha = H_\alpha$. We have the following lemma.

Lemma 5.12. *Suppose that α is admissible and $L_\alpha = K_\alpha$. Then there is a system of representatives $(x_\alpha)_{\alpha \in \mathbf{A}}$ of (P, H) -double cosets of G satisfying both equation (5.2) and $x_\alpha s_n x_\alpha^{-1} = k_\alpha$.*

Proof. Let us set $m_i = n_{i,i}^+ = n_{i,i}^- = n_{i,i}/2$ for any integer $i \in \{1, \dots, t\}$ such that $\sigma_\alpha(i) = i$. For each $\alpha \in \mathbf{A}$, we look for a matrix $x_\alpha \in G$ such that

$$x_\alpha \begin{pmatrix} \text{id}_u & \\ & -\text{id}_u \end{pmatrix} x_\alpha^{-1} = e_\alpha \quad \text{and} \quad x_\alpha \begin{pmatrix} & \text{id}_u \\ \text{id}_u & \end{pmatrix} x_\alpha^{-1} = k_\alpha.$$

To make an explicit choice of $x_\alpha \in G$, it will be convenient to introduce the matrix $v_\alpha \in G$ defined as follows: For all integers $i, j \in \{1, \dots, t\}$, the (i, j) -block of v_α in $\mathbf{M}_{n_i, n_j}(\mathbf{k})$ is

- o the identity matrix id_{n_i} if $j = i$ or $j = \sigma_\alpha(i) < i$,
- o its opposite $-\text{id}_{n_i}$ if $j = \sigma_\alpha(i) > i$,
- o and 0 otherwise.

Then we choose y_α the permutation matrix corresponding to the permutation of minimal length (with the usual generators of the symmetric group) satisfying

$$y_\alpha \begin{pmatrix} \text{id}_u & \\ & -\text{id}_u \end{pmatrix} y_\alpha^{-1} = l_\alpha,$$

where $l_\alpha = \text{diag}(l_1, \dots, l_t) \in M$ is defined by

$$l_i = \text{id}_{n_i} = -l_{\sigma_\alpha(i)} \text{ if } i < \sigma_\alpha(i), \quad l_i = \begin{pmatrix} \text{id}_{m_i} & \\ & -\text{id}_{m_i} \end{pmatrix} \text{ if } i = \sigma_\alpha(i).$$

Finally, we put $x_\alpha = v_\alpha y_\alpha$, which has the desired property thanks to the equality

$$\begin{pmatrix} \text{id}_k & -\text{id}_k \\ \text{id}_k & \text{id}_k \end{pmatrix} \begin{pmatrix} \text{id}_k & \\ & -\text{id}_k \end{pmatrix} \begin{pmatrix} \text{id}_k & -\text{id}_k \\ \text{id}_k & \text{id}_k \end{pmatrix}^{-1} = s_{2k} \tag{5.3}$$

valid for any $k \geq 1$. With this choice, the careful reader checks by a computation relying again on Equality (5.3), that $y_\alpha s_n y_\alpha^{-1} = v_\alpha^{-1} k_\alpha v_\alpha$, which is the desired equality. \square

Now, we have the following lemma.

Lemma 5.13. *There is an admissible $\alpha \in \mathbf{A}$ such that $\text{Hom}_{L_\alpha}(\rho_1 \otimes \dots \otimes \rho_t, \chi_\alpha)$ is nonzero.*

Proof. Given any subgroup X of G , we will write \overline{X} for its image in $G/Z = \text{PGL}_n(\mathbf{k})$. In particular, we have $\overline{G} = G/Z$. Note that $\overline{K} = K/Z$ is the subgroup of \overline{G} made of all elements fixed by conjugacy by

$$\begin{pmatrix} \text{id}_u & 0 \\ 0 & -\text{id}_u \end{pmatrix} \text{ mod } Z.$$

Let χ be the unique character of K trivial on H such that $\chi(s) = c$. The character that it induces on \overline{K} will still be denoted by χ . Since W is c -distinguished, Mackey’s formula implies that there is an $x \in G$ such that ρ , the representation of P inflated from $\rho_1 \otimes \cdots \otimes \rho_t$, is distinguished by the character $\chi^x|_{P \cap K^x}$. We derive from ρ a representation $\overline{\rho}$ of \overline{P} distinguished by $\chi^x|_{\overline{P} \cap \overline{K}^x}$.

In fact, because H is a subgroup of K , we can chose x to be some x_α for $\alpha \in \mathbf{A}$. Now, we claim that for all non admissible $\alpha \in \mathbf{A}$, the space

$$\text{Hom}_{\overline{P} \cap \overline{K}^{x_\alpha}}(\overline{\rho}, \chi^{x_\alpha}) \tag{5.4}$$

is zero, so in particular x can only be of the form x_α for admissible α . Indeed, it follows from [28] Proposition 3.5 that, for a nonadmissible α , the group $P \cap H^{x_\alpha}$ contains a nontrivial unipotent radical U_α of some parabolic subgroup of M , but the character χ_α is trivial on U_α , so if the space (5.4) were not reduced to zero, we would deduce that $\text{Hom}_{U_\alpha}(\rho, R)$ is nonzero, contradicting the cuspidality of ρ . Hence, we deduce $x = x_\alpha$ for an α which is admissible. In this case, $\overline{M} \cap \overline{K}^{x_\alpha}$ is equal to $\overline{M}^{\theta_\alpha}$ so that the space

$$\text{Hom}_{\overline{M}^{\theta_\alpha}}(\overline{\rho}, \chi^{x_\alpha}) = \text{Hom}_{L_\alpha}(\rho, \chi^{x_\alpha})$$

is nonzero. If $L_\alpha = K_\alpha$, then χ^{x_α} is equal to χ_α thanks to Lemma 5.12. Otherwise, χ^{x_α} and χ_α are trivial, thus equal. The statement now follows. \square

Recall that, for any $i \in \{1, \dots, t\}$, either $r_i = 1$ or $r_i = e(\varrho)\ell^{v_i}$ for some $v_i \geq 0$.

Lemma 5.14. *Let $\alpha \in \mathbf{A}$ be as in Lemma 5.13. Then the involution σ_α has a fixed point.*

Proof. Let I_1 be the set of $i \in \{1, \dots, t\}$ such that $r_i > 1$, let t_1 be the cardinality of this set and define $t_0 = t - t_1$. The identity $r = r_1 + \cdots + r_t$ implies

$$r = t_0 + e(\varrho) \cdot \sum_{i \in I_1} \ell^{v_i}.$$

Since r , $e(\varrho)$ and ℓ are odd, it follows that $t_0 + t_1 = t$ is odd. Thus, σ_α has a fixed point. \square

Claim 5.9 now follows from Lemmas 5.13, 5.14. Indeed, by Lemma 5.13, there is an admissible $\alpha \in \mathbf{A}$ such that $\text{Hom}_{L_\alpha}(\rho_1 \otimes \cdots \otimes \rho_t, \chi_\alpha)$ is nonzero. Since L_α contains H_α , the representation ρ_i is, for all i fixed by σ_α , distinguished by the Levi subgroup

$$\text{GL}_{n_{i,i}^+}(\mathbf{k}) \times \text{GL}_{n_{i,i}^-}(\mathbf{k}).$$

By [35] Proposition 2.14, this implies that $n_{i,i}^+ = n_{i,i}^-$ for all i fixed by σ_α , thus $L_\alpha = K_\alpha$. By Lemma 5.14, there is an integer $i \in \{1, \dots, t\}$ fixed by σ_α . The i th block of $L_\alpha = K_\alpha$ is K_{n_i} and $\chi_\alpha(k_i) = c$. Thus, ρ_i is c -distinguished.

6. Distinguished lift theorems

In this section, p is odd and ℓ is a prime number different from p . We look for a necessary and sufficient condition for an \mathbb{F}_ℓ -cuspidal representation of $\text{GL}_n(F)$ to have a $\text{GL}_n(F_0)$ -distinguished lift to \mathbb{Q}_ℓ . Since the case of supercuspidal representations is treated by Theorems 3.3 and 3.4, we will concentrate on nonsupercuspidal cuspidal representations.

6.1.

We will prove the following two propositions.

Proposition 6.1. *Let π be a σ -self-dual cuspidal $\overline{\mathbb{F}}_\ell$ -representation of $\mathrm{GL}_n(F)$ with quadratic extension T/T_0 and $m = m(\pi)$. Assume that $r = r(\pi) > 1$ is odd. Then π has a distinguished lift to $\overline{\mathbb{Q}}_\ell$ if and only if*

1. *the representation π is isomorphic to $\mathrm{St}_r(\rho)$ for some $\mathrm{GL}_{n/r}(F_0)$ -distinguished supercuspidal representation ρ of $\mathrm{GL}_{n/r}(F)$,*
2. *if e, e_0 are the orders of the cardinalities of the residue fields $\mathfrak{l}, \mathfrak{l}_0$ of $T, T_0 \bmod \ell$, then*
 - (a) *either T/T_0 is unramified and e_0 is even,*
 - (b) *or T/T_0 is ramified, m is even and m/e is odd.*

Note that the assumption ‘ $r > 1$ is odd’ in Proposition 6.1 implies that $\ell \neq 2$.

Proposition 6.2. *Let π be a σ -self-dual cuspidal $\overline{\mathbb{F}}_\ell$ -representation of $\mathrm{GL}_n(F)$ with quadratic extension T/T_0 and $m = m(\pi)$. Assume that $r = r(\pi)$ is even. Then π has a distinguished lift to $\overline{\mathbb{Q}}_\ell$ if and only if*

1. *the extension T/T_0 is ramified,*
2. *one has $m = r$,*
3. *if we denote by v_0 the normalized absolute value of F_0 , then*
 - (a) *either $\ell \neq 2$ and π is isomorphic to $\mathrm{St}_m(\rho)$ for some supercuspidal representation ρ of $\mathrm{GL}_{n/m}(F)$ which is either κ -distinguished or v_0^{-1} -distinguished,*
 - (b) *or $\ell = m = r = 2$, the cardinality of the residue field of T is congruent to $-1 \bmod 4$ and π is isomorphic to $\mathrm{St}_2(\rho)$ where ρ is a $\mathrm{GL}_{n/2}(F_0)$ -distinguished supercuspidal representation of $\mathrm{GL}_{n/2}(F)$.*

We also formulate the following conjecture making Proposition 6.2 more precise when $\ell > 2$.

Conjecture 6.3. *Assume that $\ell \neq 2$. Let π be a σ -self-dual cuspidal $\overline{\mathbb{F}}_\ell$ -representation of the group $\mathrm{GL}_n(F)$ such that $r(\pi)$ is even. The following assertions are equivalent:*

1. *The representation π is distinguished,*
2. *The representation π has a distinguished lift to $\overline{\mathbb{Q}}_\ell$,*
3. *The three conditions of Proposition 6.2 hold.*

By Proposition 6.2, Theorem 3.3, we know that (2) implies (1) and is equivalent to (3). We thus conjecture that (1) implies (3). See [12] Theorem 4.6 for the case $n = r = 2$.

Remark 6.4. When $\ell = 2$, there are distinguished cuspidal $\overline{\mathbb{F}}_\ell$ -representations of $\mathrm{GL}_2(F)$ with no distinguished lift. This is the case exactly when either F/F_0 is unramified, or F/F_0 is ramified and q is congruent to 1 mod 4. See Paragraph 4.13.

6.2.

Let π be a σ -self-dual cuspidal $\overline{\mathbb{F}}_\ell$ -representation of $G = \mathrm{GL}_n(F)$. Let (\mathbf{J}, λ) be a generic σ -self-dual type in π , and let λ_w be the representation of \mathbf{J} given by Proposition 4.16 (see Paragraph 4.9) and τ be the representation of \mathbf{J} trivial on \mathbf{J}^1 such that λ is isomorphic to $\lambda_w \otimes \tau$. Associated with π by equation (4.21), there is also a σ -self-dual cuspidal $\overline{\mathbb{F}}_\ell$ -representation π_1 of $\mathrm{GL}_m(T)$.

Lemma 6.5. *The following assertions are equivalent.*

1. *The representation π has a $\mathrm{GL}_n(F_0)$ -distinguished lift to $\overline{\mathbb{Q}}_\ell$.*
2. *The representation λ has a $\mathbf{J} \cap \mathrm{GL}_n(F_0)$ -distinguished lift to $\overline{\mathbb{Q}}_\ell$.*
3. *The representation τ has a $\mathbf{J} \cap \mathrm{GL}_n(F_0)$ -distinguished lift to $\overline{\mathbb{Q}}_\ell$.*
4. *The representation π_1 has a $\mathrm{GL}_m(T_0)$ -distinguished lift to $\overline{\mathbb{Q}}_\ell$.*

Proof. Fix a σ -self-dual simple stratum $[\mathfrak{a}, \beta]$ as well as isomorphisms (4.17) as in Proposition 4.5. Let $\theta \in \mathcal{C}(\mathfrak{a}, \beta)$ be the σ -self-dual maximal simple character associated with λ , and $\hat{\theta}$ be its unique $\overline{\mathbb{Q}}_\ell$ -lift:

this is a σ -self-dual maximal simple character (with respect to the unique $\overline{\mathbb{Q}}_\ell$ -lift $\tilde{\psi}$ of the character ψ given by equation (4.1)) having the same G -normalizer \mathbf{J} as θ .

Let λ_w be the $\overline{\mathbb{Q}}_\ell$ -representation of \mathbf{J} associated with θ by Proposition 4.16. It is $\mathbf{J} \cap G^\sigma$ -distinguished and σ -self-dual, and its determinant has order a power of p . It is thus integral. Let us consider its reduction mod ℓ . On the one hand, it is $\mathbf{J} \cap G^\sigma$ -distinguished, σ -self-dual, and its determinant has order a power of p . On the other hand, [30] Proposition 2.37 implies that it is an irreducible representation extending the Heisenberg representation associated with θ . By uniqueness, we deduce that λ_w is a $\overline{\mathbb{Q}}_\ell$ -lift of λ_w .

Suppose that π has a G^σ -distinguished $\overline{\mathbb{Q}}_\ell$ -lift $\tilde{\pi}$. Thus, $\tilde{\pi}$ is a σ -self-dual and cuspidal representation of G containing the maximal simple character $\tilde{\theta}$. By Proposition 4.32, this representation $\tilde{\pi}$ contains a distinguished generic σ -self-dual type, which we may assume to be of the form $(\mathbf{J}, \tilde{\lambda})$ with $\tilde{\lambda} = \tilde{\lambda}_w \otimes \tilde{\tau}$ and the representation $\tilde{\tau}$ is $\mathbf{J} \cap G^\sigma$ -distinguished. Reducing mod ℓ , we deduce that π contains the type $\lambda_w \otimes \delta$, where δ is the reduction mod ℓ of $\tilde{\tau}$. But π also contains the type $\lambda_w \otimes \tau$, thus δ is isomorphic to τ , and the reduction mod ℓ of $\tilde{\lambda}$ is isomorphic to λ . Thus, (1) implies both (2) and (3).

Conversely, suppose that τ has a distinguished $\overline{\mathbb{Q}}_\ell$ -lift $\tilde{\tau}$. Then the pair $(\mathbf{J}, \lambda_w \otimes \tilde{\tau})$ is a distinguished type whose compact induction to G is a G^σ -distinguished $\overline{\mathbb{Q}}_\ell$ -lift of π , and whose reduction mod ℓ is isomorphic to $\lambda_w \otimes \tau \simeq \lambda$. Thus, (3) implies both (1) and (2).

Applying these results to the representation π_t , we get that π_t has a distinguished lift to $\overline{\mathbb{Q}}_\ell$ if and only if τ_t has a distinguished lift to $\overline{\mathbb{Q}}_\ell$. The fact that τ is isomorphic to $\tau_t \circ \pi$ (by Lemma 4.37) thus implies that (4) is equivalent to (3). □

It follows from Lemma 6.5, together with Corollary 4.42 and Proposition 4.43, that, in order to prove Propositions 6.1 and 6.2, it suffices to prove them for σ -self-dual cuspidal $\overline{\mathbb{F}}_\ell$ -representations of level 0. (For Proposition 6.2(3.a), this also follows from the fact that $\varkappa_{F/F_0} \circ N_{T_0/F_0} = \varkappa_{T/T_0}$ and $\nu_{F_0} \circ N_{T_0/F_0} = \nu_{T_0}$.)

6.3.

We continue with the situation of Paragraph 6.2, assuming further that π has level 0. Thus, π is a σ -self-dual cuspidal $\overline{\mathbb{F}}_\ell$ -representation of G of level 0. We will also assume that π is nonsupercuspidal, that is, $r = r(\pi) > 1$. Let (\mathbf{J}, λ) be a generic σ -self-dual type in π . Associated with it in Paragraph 4.7, there are

- a σ -self-dual tamely ramified character ω of F^\times , which is the central character c_π of π ,
- and a σ -self-dual cuspidal representation V of $GL_n(\mathbf{k})$ of the form $\text{st}_r(\varrho)$ for some supercuspidal representation ϱ of $GL_{n/r}(\mathbf{k})$, uniquely determined up to isomorphism (thus V is nonsupercuspidal).

Recall that the restriction of λ to \mathbf{J}^0 is the inflation of V , and that its restriction to F^\times is a multiple of ω . Since V is σ -self-dual, Proposition 3.9 implies that ϱ is σ -self-dual.

The action of σ on $GL_n(\mathbf{k})$ is described in Proposition 4.5: This is the action of the nontrivial automorphism of \mathbf{k}/\mathbf{k}_0 if F/F_0 is unramified, and the adjoint action of the element (4.5) with $i = \lfloor m/2 \rfloor$ otherwise.

Let us fix a uniformizer ϖ of F such that $\varpi \in F_0$ if F/F_0 is unramified, and $\varpi^2 \in F_0$ if F/F_0 is ramified. (One thus has $\sigma(\varpi) = -\varpi$ in the ramified case.)

Lemma 6.6. *The representation π has a $GL_n(F_0)$ -distinguished lift to $\overline{\mathbb{Q}}_\ell$ if and only if V has a $GL_n(\mathbf{k})^\sigma$ -distinguished lift \tilde{V} to $\overline{\mathbb{Q}}_\ell$ such that*

1. if F/F_0 is unramified, then $\omega(\varpi) = 1$,
2. if F/F_0 is ramified, then n is even and the element (4.23) acts on the space of $GL_{n/2}(\mathbf{k}) \times GL_{n/2}(\mathbf{k})$ -invariant linear forms on \tilde{V} by a sign whose reduction mod ℓ is equal to $\omega(\varpi)$.

Proof. By Lemma 6.5, the representation π has a $GL_n(F_0)$ -distinguished lift if and only if the type λ has a $\mathbf{J} \cap GL_n(F_0)$ -distinguished lift to $\overline{\mathbb{Q}}_\ell$. Suppose λ has a distinguished lift $\tilde{\lambda}$. Then the pair (\mathbf{J}, λ) is the generic type of a distinguished cuspidal $\overline{\mathbb{Q}}_\ell$ -representation $\tilde{\pi}$, compactly induced from $\tilde{\lambda}$. Associated with it, there are

- a cuspidal $\overline{\mathbb{Q}}_\ell$ -representation \widetilde{V} of $GL_n(\mathbf{k})$ lifting V ,
- a tamely ramified $\overline{\mathbb{Q}}_\ell$ -character $\widetilde{\omega}$ of F^\times lifting ω .

By Theorem 4.45, the character $\widetilde{\omega}$ is trivial on F_0^\times and \widetilde{V} is distinguished by $GL_n(\mathbf{k})^\sigma$. If F/F_0 is unramified, then $\widetilde{\omega}(\varpi) = 1$, thus $\omega(\varpi) = 1$. If F/F_0 is ramified, then $n = 2u$ for some $u \geq 1$ and \widetilde{V} is $\widetilde{\omega}(\varpi)$ -distinguished (in the sense of Definition 5.6), and the reduction mod ℓ of $\widetilde{\omega}(\varpi)$ is equal to $\omega(\varpi)$.

Conversely, suppose that V has a $GL_n(\mathbf{k})^\sigma$ -distinguished lift \widetilde{V} satisfying the conditions of the lemma. Let $\widetilde{\omega}$ be a $\overline{\mathbb{Q}}_\ell$ -lift of ω coinciding on the units of F with the inflation of the central character of \widetilde{V} , and

1. if F/F_0 is unramified, then $\widetilde{\omega}(\varpi) = 1$,
2. if F/F_0 is ramified, then $\widetilde{\omega}(\varpi) \in \{-1, 1\}$ and the representation \widetilde{V} is $\widetilde{\omega}(\varpi)$ -distinguished.

Inflate \widetilde{V} to \mathbf{J}^0 , and extend it to a representation $\widetilde{\lambda}$ of \mathbf{J} by demanding that the restriction of $\widetilde{\lambda}$ to F^\times is a multiple of $\widetilde{\omega}$. The representation $\widetilde{\lambda}$ is then a $\mathbf{J} \cap GL_n(F_0)$ -distinguished lift of λ . □

6.4.

In this paragraph, we assume that F/F_0 is unramified. Remind that q denotes the cardinality of \mathbf{k} and q_0 denotes that of \mathbf{k}_0 .

Lemma 6.7. *Let W be a σ -self-dual cuspidal $\overline{\mathbb{F}}_\ell$ -representation of $GL_n(\mathbf{k})$. It has a $GL_n(\mathbf{k}_0)$ -distinguished lift to $\overline{\mathbb{Q}}_\ell$ if and only if n is odd and*

1. either W is supercuspidal,
2. or W is nonsupercuspidal and the order of $q_0 \bmod \ell$ is even (thus $\ell \neq 2$).

Proof. By [18] Theorem 3.6, an irreducible $\overline{\mathbb{Q}}_\ell$ -representation of $GL_n(\mathbf{k})$ is $GL_n(\mathbf{k}_0)$ -distinguished if and only if it is σ -self-dual.

First, the condition on the parity of n is necessary: See [35] Lemma 2.3 for instance. Now, assume that n is odd. If $\ell \neq 2$, the result is given by [25] Proposition 4.6. If $\ell = 2$, then W has the form $st_r(\varrho)$, where ϱ is a supercuspidal representation of $GL_{n/r}(\mathbf{k})$ and $r = 2^\nu$ for some $\nu \geq 0$. Since n is odd, W must be supercuspidal, and the result is given by [35] Remark 2.7. □

Remark 6.8. Let e and e_0 be the orders of q and $q_0 \bmod \ell$, respectively. Note that $r = e(\varrho)\ell^\nu$ for some $\nu \geq 0$, where $e(\varrho)$ is the order of $q^f \bmod \ell$ with $f = n/r$. If n is odd, then f and r are odd, thus $e(\varrho)$ is odd. But $e(\varrho) = e/(e, f)$. It follows that $e = e_0/(e_0, 2)$ is odd. Thus, e_0 is not divisible by 4.

Example 6.9. Let W be the σ -self-dual cuspidal $\overline{\mathbb{F}}_\ell$ -representation $st_e(1)$ of $GL_e(\mathbf{k})$. We have $e = e_0/(e_0, 2)$, which is odd if and only if e_0 is not divisible by 4. Thus, W has a $GL_e(\mathbf{k}_0)$ -distinguished lift to $\overline{\mathbb{Q}}_\ell$ if and only if e_0 is divisible by 2 but not by 4.

Suppose first that π has a distinguished lift to $\overline{\mathbb{Q}}_\ell$. On the one hand, the generic type of such a lift defines a σ -self-dual cuspidal $\overline{\mathbb{Q}}_\ell$ -representation of $GL_n(\mathbf{k})$, and [35] Lemma 2.3 implies that n is odd, thus r is odd. On the other hand, Theorem 3.3 implies that π is distinguished. It thus follows from Theorem 5.1 that π is isomorphic to $St_r(\rho)$ for some distinguished supercuspidal representation ρ of $GL_{n/r}(F)$. Finally, Lemma 6.6 says that V has a distinguished lift. It follows from Lemma 6.7 that the order e_0 of the cardinality of $\mathbf{k}_0 \bmod \ell$ is even.

We thus proved that, when F/F_0 is unramified, if π has a distinguished lift, then r is odd and Conditions (1), (2.a) of Proposition 6.1 are satisfied.

Conversely, suppose that the conditions (1), (2.a) of Proposition 6.1 are satisfied. Then V has a distinguished lift \widetilde{V} . By Lemma 6.6, the representation π has a $GL_n(F_0)$ -distinguished lift to $\overline{\mathbb{Q}}_\ell$ if and only if $\omega(\varpi) = 1$. By Paragraph 4.8, the central character ω_* of ρ satisfies $\omega_*^r = \omega$. Since ρ is distinguished, we have $\omega_*(\varpi) = 1$, thus $\omega(\varpi) = \omega_*(\varpi)^r = 1$.

We proved Proposition 6.1 in the case when F/F_0 is unramified.

6.5.

In this paragraph, we assume that F/F_0 is ramified. Let q denote the cardinality of \mathbf{k} , and let e denote the order of $q \bmod \ell$.

Lemma 6.10. *Let W be a self-dual cuspidal $\overline{\mathbb{F}}_\ell$ -representation of $GL_n(\mathbf{k})$, isomorphic to $st_r(\varrho)$ for some self-dual supercuspidal representation ϱ of $GL_{n/r}(\mathbf{k})$. Write $u = \lfloor n/2 \rfloor$. Then W has a lift to $\overline{\mathbb{Q}}_\ell$ which is distinguished by $GL_u(\mathbf{k}) \times GL_{n-u}(\mathbf{k})$ if and only if*

1. either W is supercuspidal,
2. or W is nonsupercuspidal, n is even and
 - (a) either $\ell \neq 2$ and $r, n/e$ are odd,
 - (b) or $\ell \neq 2$ and $r = n$,
 - (c) or $\ell = n = r = 2$ and q is congruent to $-1 \pmod 4$, and ϱ is trivial.

Proof. First note that n must be either even or equal to 1: See [35] Lemma 2.17 for instance. Also, the supercuspidal case is given by [35] Remark 2.21. Let us assume that W is nonsupercuspidal (thus n is even, and we will write $n = 2u$). We use the notation of Paragraph 3.7.

Set $f = n/r$, and let α be a $\text{Gal}(\mathbf{k}_f/\mathbf{k})$ -regular $\overline{\mathbb{F}}_\ell$ -character of \mathbf{k}_f^\times of order A which is a parameter of ϱ in the sense of Definition 3.12. Let $\tilde{\nu}$ be the canonical $\overline{\mathbb{Q}}_\ell$ -lift of

$$\nu = \alpha \circ N_{\mathbf{k}_n/\mathbf{k}_f},$$

that is, its unique lift of order A . Let \tilde{W} be a cuspidal lift of W . It is parametrized by a $\text{Gal}(\mathbf{k}_n/\mathbf{k})$ -regular character of \mathbf{k}_n^\times lifting ν , that is, of the form $\tilde{\nu}\phi$, where ϕ is a $\overline{\mathbb{Q}}_\ell$ -character of \mathbf{k}_n^\times of order ℓ^s for some $s \geq 0$. Since W is not supercuspidal, one has $s \geq 1$. The character $\tilde{\nu}\phi$ has order $A\ell^s$.

By Proposition 3.13, the representation \tilde{W} is distinguished by $GL_u(\mathbf{k}) \times GL_u(\mathbf{k})$ if and only if it is self-dual, which is also equivalent (see, for instance, [35] (2.7)) to $A\ell^s$ dividing $q^u + 1$. Similarly, the fact that ϱ is self-dual is equivalent to

- either $f = 1$ and ϱ is a quadratic character (thus, A is equal to 1 or 2),
- or f is even and A divides $q^{f/2} + 1$ (thus, $A > 2$ since q has order $f \geq 2 \pmod A$).

Suppose that $\ell \neq 2$ and f is even. If \tilde{W} is distinguished, then A divides $q^{f/2} + 1$ and $q^u + 1$. Since $u = rf/2$, we have

$$q^u = (q^{f/2})^r \equiv (-1)^r \pmod A,$$

thus A divides $1 + (-1)^r$. Since $A > 2$, it follows that r is odd. Also, ℓ divides $q^u + 1$, that is, the order of $q^u \bmod \ell$ is $e/(e, u) = 2$, which implies that n/e is odd. Conversely, suppose that r and n/e are odd. The fact that A divides $q^{f/2} + 1$ and r is odd implies that A divides $q^u + 1$. Now, ℓ^s divides $q^n - 1 = (q^u + 1)(q^u - 1)$. If ℓ divides $q^u - 1$, then e divides $u = n/2$, thus n/e is even: contradiction. Thus, ℓ^s divides $q^u + 1$, thus \tilde{W} is distinguished.

Suppose that $\ell \neq 2$ and $f = 1$. Then ϱ is a character of \mathbf{k}^\times , thus $r = e\ell^v$ for some $v \geq 0$. This gives $n/e = \ell^v$, which is odd. The same argument as above implies that $q^u + 1$ is a multiple of ℓ^s . It is also a multiple of $A \in \{1, 2\}$ since it is even. Thus, \tilde{W} is distinguished.

Now, suppose that $\ell = 2$. If \tilde{W} is distinguished and f is even, then, as in the case where $\ell \neq 2$, the integer $A > 2$ divides $q^{f/2} + 1$ and $q^{rf/2} + 1$, thus r is odd. But the fact that W is cuspidal implies that r is a power of 2. It follows that $r = 1$: contradiction. Thus, $f = 1$, that is W is the representation $st_n(1)$ with $n = 2^t$ for some $t \geq 1$. Moreover, q has order $m \pmod{2^s}$, that is, 2^s divides $q^n - 1$ but not $q^u - 1$. Set

$$a = v_2(q^u + 1), \quad b = v_2(q^u - 1).$$

We have $b < s \leq a + b$ and $\min(a, b) = 1$. The fact that \tilde{W} is distinguished implies $s \leq a$, which gives $b = 1 < a$, that is 4 divides $q^u + 1$. Since u is a power of 2, we deduce that 4 divides $q + 1$ and $u = 1$.

Conversely, suppose that $\ell = n = r = 2$ and 4 divides $q + 1$ (hence $b = 1 < a$). Then any $\overline{\mathbb{Q}}_2$ -character of k_2^\times of order 2^a parametrizes a distinguished cuspidal $\overline{\mathbb{Q}}_2$ -representation of $GL_2(k)$ lifting $W = st_2(1)$. □

Example 6.11. The fact that $GL_f(k)$ has a self-dual supercuspidal $\overline{\mathbb{F}}_\ell$ -representation is equivalent to the fact that there is an k -regular $\overline{\mathbb{F}}_\ell$ -character of k_f^\times which is trivial on $k_{f/2}^\times$, that is, there exists an integer A with the following properties:

1. A is prime to ℓ and the order of $q \bmod A$ is equal to f ,
2. A divides $q^{f/2} + 1$.

Now, suppose that $\ell > 2$ and $f = 2$. Thus, $GL_2(k)$ has a self-dual supercuspidal $\overline{\mathbb{F}}_\ell$ -representation if and only if there exists an integer A prime to ℓ dividing $q + 1$ but not $q - 1$, that is, if and only if $q + 1$ has a prime divisor different from 2 and ℓ . Assume this is the case, and let ϱ be a self-dual supercuspidal $\overline{\mathbb{F}}_\ell$ -representation of $GL_2(k)$. Let W be the self-dual cuspidal $\overline{\mathbb{F}}_\ell$ -representation $st_r(\varrho)$ of $GL_n(k)$ with $r = e/(e, 2)$ and $m = 2r$. Then r is odd if and only if e is not divisible by 4, and $n/e = 2/(e, 2)$ is odd if and only if e is even. If we take $q = 9$ and $\ell = 7$, we get $r = 3$ and $n/e = 2$. If we take $q = 5$, we get $q - 1 = 4$ and $q^6 - 1 = 1953 \times 8$. Thus, if ℓ is a prime divisor of 1953, we get $r = 3$ and $n/e = 6$.

In addition, we have the following result. We assume that $n = 2u$ for some $u \geq 1$.

Lemma 6.12. *Let W be a self-dual cuspidal $\overline{\mathbb{F}}_\ell$ -representation of $GL_n(k)$ of the form $st_n(\varrho)$ for some quadratic character ϱ of k^\times . Assume W is distinguished by $GL_u(k) \times GL_u(k)$. Then the element (4.23) acts on the space of $GL_u(k) \times GL_u(k)$ -invariant linear forms on W by*

$$\begin{cases} -1 & \text{if } \varrho \text{ is trivial,} \\ (-1)^{u(q-1)/2} & \text{if } \varrho \text{ is nontrivial.} \end{cases}$$

Proof. Let c be the sign such that W is c -distinguished. If $\ell = 2$, the result is immediate since the only sign is 1. Assume that $\ell \neq 2$. By Lemma 6.10, the representation W has a distinguished cuspidal $\overline{\mathbb{Q}}_\ell$ -lift. Let \widetilde{W} be such a $\overline{\mathbb{Q}}_\ell$ -lift and ξ be a parameter for \widetilde{W} . Let α be an element of k_n such that $\alpha \notin k_u$ and $\alpha^2 \in k_u$. By Proposition 3.13, the representation \widetilde{W} is $-\xi(\alpha)$ -distinguished by $GL_u(k) \times GL_u(k)$. Since \widetilde{W} lifts W , we have

- the reduction mod ℓ of the parameter ξ is equal to $(\varrho \circ N_{k_n/k})\phi$ where ϕ is a character whose order is a power of ℓ (see Proposition 3.11),
- the reduction mod ℓ of $-\xi(\alpha)$ is equal to c (see Remark 3.14).

On the one hand, the character ξ is trivial on k_u^\times since \widetilde{W} is self-dual (see Proposition 3.13). On the other hand, $\varrho \circ N_{k_n/k}$ is trivial on k_u^\times since ϱ is quadratic and the index of k_u^\times in k_n^\times is even. We deduce that ϕ is trivial on k_u^\times , thus $\phi(\alpha)$ is a sign. Since it has order a power of $\ell \neq 2$, it is trivial. It follows that

$$c = -\varrho(N_{k_n/k}(\alpha)).$$

If ϱ is trivial, this gives $c = -1$, as expected. Assume now that ϱ is nontrivial. It thus coincides with \varkappa on k^\times . Since α^2 is not a square in k_u^\times , its k_u/k -norm is not a square in k^\times . Thus,

$$c = -\varkappa(N_{k_u/k}(\alpha^{q^u+1})) = -(-1)^{(q^u+1)/2},$$

and one verifies that this is equal to $\varkappa(-1)^u = (-1)^{u(q-1)/2}$ as expected. □

6.6.

Let us prove Proposition 6.1 when F/F_0 is ramified. Assume that r is odd, and suppose that π has a distinguished lift to $\overline{\mathbb{Q}}_\ell$. By Theorem 3.3, it is distinguished. Thus, Theorem 5.1 implies that n/r is

even and π is isomorphic to $\text{St}_r(\rho)$ for some distinguished supercuspidal representation ρ of $\text{GL}_{n/r}(F)$. Lemma 6.6 says that V has a distinguished lift. It follows from Lemma 6.10 that n/e is odd.

Conversely, assume that r is odd, π is isomorphic to $\text{St}_r(\rho)$ for some distinguished supercuspidal representation ρ of $\text{GL}_{n/r}(F)$ of level 0, n is even and n/e is odd. It follows from Lemma 6.10 that V has a distinguished lift \tilde{V} . Let $\varepsilon \in \{-1, 1\}$ be the unique sign such that \tilde{V} is ε -distinguished by $\text{GL}_u(\mathbf{k}) \times \text{GL}_u(\mathbf{k})$ in the sense of Definition 5.6, with $n = 2u$. By Lemma 6.6, the representation π has a distinguished lift if and only if $\omega(\varpi)$ is equal to the image of ε in $\overline{\mathbb{F}}_\ell^\times$, denoted c . We are going to prove that this is the case. Let ω_* be the central character of ρ . By Theorem 4.45, we have

- the representation ϱ is $\omega_*(\varpi)$ -distinguished by $\text{GL}_{k/2}(\mathbf{k}) \times \text{GL}_{k/2}(\mathbf{k})$.

(Note that k is even since n is even and r is odd.) By Proposition 4.28, we have

- the sign $\omega(\varpi)$ is equal to $\omega_*(\varpi)^r = \omega_*(\varpi)$.

Let α be the unique sign such that V is α -distinguished by $\text{GL}_u(\mathbf{k}) \times \text{GL}_u(\mathbf{k})$. By Remark 3.14, we have $\alpha = c$. On the other hand, we have $\alpha = \omega_*(\varpi)$ by Proposition 5.7. Putting these facts together, we get $\omega(\varpi) = \omega_*(\varpi) = \alpha = c$ as expected. This proves Proposition 6.1 if F/F_0 is ramified. Together with Paragraph 6.4, this finishes the proof of Proposition 6.1.

6.7.

In this paragraph and the next one, we prove Proposition 6.2. Assume that r is even, and let q be the cardinality of \mathbf{k} . Since r divides n , we have $n = 2u$ for some $u \geq 1$.

Suppose that π has a distinguished lift. By Paragraph 6.4, this implies that F/F_0 is ramified. By Lemma 6.6, the representation V has a distinguished $\overline{\mathbb{Q}}_\ell$ -lift. By Lemma 6.10, one has $r = n$, thus V is isomorphic to $\text{st}_n(\varrho)$ for a character ϱ of \mathbf{k}^\times of order at most 2. Besides, if $\ell = 2$, then $n = 2$ and q is congruent to $-1 \pmod 4$. Since $r = n$, the representation π is isomorphic to $\text{St}_n(\rho)$ for a tamely ramified character ρ of F^\times whose restriction to the units of F is the inflation of ϱ .

Suppose first that $\ell = 2$. Since π is distinguished (by Theorem 3.3), it is σ -self-dual (by Theorem 3.1). It follows from Proposition 3.8 that the representation ρ is σ -self-dual and from Theorem 3.2 that it is F_0^\times -distinguished, as expected.

Suppose now that $\ell \neq 2$. By Proposition 3.8, we may choose ρ so that $\rho^{-1} \circ \sigma = \rho v^i$ for some $i \in \{0, 1\}$, that is, $\rho \circ \text{N}_{F/F_0} = v^{-i}$. It remains to prove that the restriction of ρ to F_0^\times is either \varkappa or v_0^{-1} .

Let c be the sign by which the element (4.23) acts on the space of $\text{GL}_u(\mathbf{k}) \times \text{GL}_u(\mathbf{k})$ -invariant linear forms on V , which is given by Lemma 6.12. Remind that we have fixed a uniformizer ϖ of F such that $\sigma(\varpi) = -\varpi$, thus $\varpi_0 = \varpi^2$ is a uniformizer of F_0 . The representation π is distinguished (by Theorem 3.3) and it follows from Theorem 4.45 that $c = c_\pi(\varpi)$. We have

$$c_\pi(\varpi) = \rho(\varpi)^n = \rho(\varpi_0)^u. \tag{6.1}$$

On the other hand, the identity $\rho \circ \text{N}_{F/F_0} = v^{-i}$ implies that $\rho(-\varpi_0) = q^i$.

Lemma 6.13. *We have $q^u \equiv -1 \pmod \ell$.*

Proof. Since $r \geq 2$ and π is cuspidal, r has the form $e(\rho)\ell^v$ for some $v \geq 0$, where $e(\rho)$ is the order of $q^k \pmod \ell$ by equation (3.2). In particular, $(q^k)^r = q^n$ is congruent to 1 $\pmod \ell$. Moreover, since ℓ is odd, one has $q^u \equiv -1 \not\equiv 1 \pmod \ell$. □

It follows from Lemma 6.13 and equation (6.1) that

$$c = \begin{cases} (-1)^i & \text{if } \varrho \text{ is trivial,} \\ (-1)^i \cdot \varkappa(-1)^u & \text{otherwise (that is, if } \varrho = \varkappa). \end{cases}$$

Comparing with Lemma 6.12, we get the following corollary.

Corollary 6.14. *We have $i = 1$ if ϱ is trivial, and $i = 0$ if ϱ is nontrivial.*

If $i = 0$, then ρ is self-dual. By Theorem 3.2, it is either distinguished or \varkappa -distinguished. Since its restriction to the units of F is the inflation of $\varrho = \varkappa$, we deduce that ρ is \varkappa -distinguished.

If $i = 1$, then $\rho v^{1/2}$ is unramified and self-dual. By Theorem 3.2, it is distinguished. Thus, the restriction of ρ to F_0^\times is equal to $v^{-1/2}|_{F_0^\times} = v_0^{-1}$.

6.8.

Let us finish the proof of Proposition 6.2. Assume that $n = r = 2u$ for some $u \geq 1$, the extension F/F_0 is ramified and π is isomorphic to $\text{St}_n(\rho)$ for some tamely ramified character ρ of F^\times . We also assume that

- either $\ell \neq 2$ and the restriction of ρ to F_0^\times is either \varkappa or v_0^{-1} ,
- or $\ell = n = r = 2$, q is congruent to $-1 \pmod 4$ and ρ is trivial on F_0^\times .

It follows from Lemma 6.10 that V has a distinguished $\overline{\mathbb{Q}}_\ell$ -lift \widetilde{V} , which is ε -distinguished for some sign $\varepsilon \in \{-1, 1\}$. By Lemma 6.6, the representation π has a distinguished lift to $\overline{\mathbb{Q}}_\ell$ if and only if the reduction of $\varepsilon \pmod \ell$, denoted c , is equal to $\omega(\varpi)$. Let us prove that this is the case. On the one hand, we have $\omega(\varpi) = \rho(\varpi)^n = \rho(\varpi_0)^u$. If $\ell = 2$, we have $\omega(\varpi) = 1$. Otherwise, we have

$$\omega(\varpi) = \begin{cases} q^u & \text{if the restriction of } \rho \text{ to } F_0^\times \text{ is } v_0^{-1}, \\ \varkappa(-1)^u & \text{if the restriction of } \rho \text{ to } F_0^\times \text{ is } \varkappa. \end{cases}$$

On the other hand, V is distinguished, and it is isomorphic to $\text{st}_n(\varrho)$, where ϱ is the character of k^\times defined by the restriction of ρ to the units of F . One thus may apply Lemma 6.12, which says that V is α -distinguished, with

$$\alpha = \begin{cases} -1 & \text{if } \varrho \text{ is trivial,} \\ (-1)^{u(q-1)/2} & \text{if } \varrho \text{ is nontrivial.} \end{cases}$$

Note that $\alpha = c$ by Remark 3.14, and that ϱ is trivial if and only if the restriction of ρ to F_0^\times is equal to v_0^{-1} . Together with Lemma 6.13, this gives $\omega(\varpi) = c$ as expected.

Acknowledgements. We thank Alberto Mínguez and Shaun Stevens for their interest and useful conversations.

Competing interest. The authors have no competing interest to declare.

Financial support. The first author was supported by EPSRC grant EP/V001930/1 and the Heilbronn Institute for Mathematical Research. The third author was partially supported by the Institut Universitaire de France.

This work was partially supported by the Erwin Schrödinger Institute in Vienna, where we benefitted from the Research in Teams grant ‘ ℓ -modular Langlands Quotient Theorem and Applications’. We thank the institute for hospitality and for excellent working conditions.

References

- [1] U. K. Anandavardhanan, ‘Root numbers of Asai L-functions’, *Int. Math. Res. Not. IMRN.* (2008), **25**:Art. ID rnn125.
- [2] U. K. Anandavardhanan, A. C. Kable and R. Tandon, ‘Distinguished representations and poles of twisted tensor L-functions’, *Proc. Amer. Math. Soc.*, **132**(10) (2004), 2875–2883.
- [3] U. K. Anandavardhanan, R. Kurinczuk, N. Matringe, V. Sécherre and S. Stevens, ‘Galois self-dual cuspidal types and Asai local factors’, *J. Eur. Math. Soc. (JEMS)*, **23**(9) (2021), 3129–3191.
- [4] U. K. Anandavardhanan and C. S. Rajan, ‘Distinguished representations, base change, and reducibility for unitary groups’, *Int. Math. Res. Not.* (14) (2005), 841–854.
- [5] F. Bruhat and J. Tits, ‘Schémas en groupes et immeubles des groupes classiques sur un corps local’, *Bull. Soc. Math. France.* **112**(2) (1984), 259–301.
- [6] C. J. Bushnell and G. Henniart, ‘Local tame lifting for $GL(N)$ I. Simple characters’, *Inst. Hautes Études Sci. Publ. Math.* **83** (1996), 105–233.
- [7] C. J. Bushnell and G. Henniart, ‘Intertwining of simple characters in $GL(n)$ ’, *Int. Math. Res. Not. IMRN.* (17) (2013), 3977–3987.

- [8] C. J. Bushnell and G. Henniart, 'To an effective local Langlands correspondence', *Mem. Amer. Math. Soc.* **231**(1087), (2014), v+88.
- [9] C. J. Bushnell and P. C. Kutzko, 'The admissible dual of $GL(N)$ via compact open subgroups', volume 129 of *Annals of Mathematics Studies* (Princeton University Press, Princeton, NJ, 1993).
- [10] M. Cabanes, 'Modular representations of finite reductive groups', in *Modular Representation Theory of Finite and p -adic Groups*. Volume 30 of *Lect. Notes Ser. Inst. Math. Sci. Natl. Univ. Singap.* (World Sci. Publ., Hackensack, NJ, 2015), 1–46.
- [11] C. Coniglio-Guilloton, 'Correspondance de Jacquet-Langlands et distinction: cas des représentations cuspidales de niveau 0', *Bull. Soc. Math. France.* **144**(2) (2016), 163–216.
- [12] P. Cui, T. Lanard and H. Lu, 'Modulo ℓ distinction problems', *Compos. Math.* **160**(10) (2024), 2285–2321.
- [13] P. Delorme and V. Sécherre, 'An analogue of the Cartan decomposition for p -adic symmetric spaces of split p -adic reductive groups', *Pacific J. Math.* **251**(1) (2011), 1–21.
- [14] R. Dipper, 'On the decomposition numbers of the finite general linear groups. II', *Trans. Amer. Math. Soc.* **292**(1), (1985), 123–133.
- [15] R. Dipper and G. James, 'Identification of the irreducible modular representations of $GL_n(q)$ ', *J. Algebra.* **104**(2) (1986), 266–288.
- [16] Y. Z. Flicker, 'On distinguished representations', *J. Reine Angew. Math.* **418** (1991), 139–172.
- [17] W. T. Gan and A. Raghuram, 'Arithmeticity for periods of automorphic forms', in *Automorphic Representations and L -functions*, Volume 22 of *Tata Inst. Fundam. Res. Stud. Math.* (Tata Inst. Fund. Res., Mumbai, 2013), 187–229.
- [18] R. Gow, 'Two multiplicity-free permutation representations of the general linear group', *Math. Z.* **188**(1) (1984), 45–54.
- [19] J. A. Green, 'The characters of the finite general linear groups', *Trans. Amer. Math. Soc.* **80** (1955), 402–447.
- [20] J. Hakim and F. Murnaghan, 'Two types of distinguished supercuspidal representations', *Int. Math. Res. Not.* (35) (2002), 1857–1889.
- [21] G. James, 'The irreducible representations of the finite general linear groups', *Proc. London Math. Soc. (3)*, **52**(2) (1986), 236–268.
- [22] Y. Jo, 'Local exterior square and Asai L-functions for in odd characteristic', *Pacific J. Math.* **322**(2) (2023), 301–340.
- [23] A. C. Kable, 'Asai L-functions and Jacquet's conjecture', *Amer. J. Math.* **126**(4) (2004), 789–820.
- [24] R. Kurinczuk and N. Matringe, 'Rankin-Selberg local factors modulo ℓ ', *Selecta Math. (N.S.)*. **23**(1) (2017), 767–811.
- [25] R. Kurinczuk and N. Matringe, 'Characterisation of the poles of the ℓ -modular Asai L-factor', *Bull. Soc. Math. France.* **148**(3) (2020), 481–514.
- [26] N. Matringe, 'Conjectures about distinction and local Asai L-functions', *Int. Math. Res. Not. IMRN.* (9) (2009), 1699–1741.
- [27] N. Matringe, 'Distinguished generic representations of $GL(n)$ over p -adic fields', *Int. Math. Res. Not. IMRN.* (1) (2011), 74–95.
- [28] N. Matringe, 'On the local Bump–Friedberg L-function', *J. Reine Angew. Math.* **709**, (2015), 119–170.
- [29] A. Mínguez and V. Sécherre, 'Représentations lisses modulo ℓ de $GL_m(D)$ ', *Duke Math. J.* **163**(4) (2014), 795–887.
- [30] A. Mínguez and V. Sécherre, 'Types modulo ℓ pour les formes intérieures de GL_n sur un corps local non archimédien', *Proc. Lond. Math. Soc. (3)*, **109**(4) (2014), 823–891. With an appendix by Vincent Sécherre et Shaun Stevens.
- [31] A. Mínguez and V. Sécherre, 'Correspondance de Jacquet-Langlands locale et congruences modulo ℓ ', *Invent. Math.* **208**(2), 2017, 553–631.
- [32] D. Prasad, 'Trilinear forms for representations of and local ϵ -factors', *Compositio Math.* **75**(1) (1990), 1–46.
- [33] D. Prasad, 'On a conjecture of Jacquet about distinguished representations of $GL(n)$ ', *Duke Math. J.* **109**(1) (2001), 67–78.
- [34] V. Sécherre, 'Représentations lisses de $GL(m, D)$ II. β -extensions', *Compos. Math.* **141**(6) (2005), 1531–1550.
- [35] V. Sécherre, 'Supercuspidal representations of $GL_n(F)$ distinguished by a Galois involution', *Algebra Number Theory* **13**(7) (2019), 1677–1733.
- [36] V. Sécherre, 'Représentations cuspidales de $GL_r(D)$ distinguées par une involution intérieure', *Ann. Sci. Éc. Norm. Supér. (4)*, **57**(4) (2024), 961–1038.
- [37] V. Sécherre and S. Stevens, 'Smooth representations of $GL_m(D)$ VI: semisimple types', *Int. Math. Res. Not. IMRN.* (13) (2012), 2994–3039.
- [38] V. Sécherre and S. Stevens, 'Block decomposition of the category of ℓ -modular smooth representations of $GL_n(F)$ and its inner forms', *Ann. Sci. Éc. Norm. Supér. (4)*, **49**(3) (2016), 669–709.
- [39] J.-P. Serre, *Représentations Linéaires des Groupes Finis*. Revised edition. (Hermann, Paris, revised edition, 1978).
- [40] M.-F. Vignéras, 'Représentations modulaires de en caractéristique l , F corps p -adique, $p \neq l$ ', *Compositio. Math.* **72**(1) (1989), 33–66.
- [41] M.-F. Vignéras, *Représentations l -modulaires d'un groupe réductif p -adique avec $l \neq p$* , volume 137 of *Progress in Mathematics* (Birkhäuser Boston, Inc., Boston, MA, 1996).
- [42] M.-F. Vignéras, 'On highest Whittaker models and integral structures', in *Contributions to Automorphic Forms, Geometry, and Number Theory* (Johns Hopkins Univ. Press, Baltimore, MD, 2004), 773–801.