Index of Subjects

Acinetobacter baumannii, genetic diversity and clonal relationships of, strains isolated in a neonatal ward: epidemiological investigations by allozyme, whole-cell protein and antibiotic resistance analysis, 491

Badger, Mycobacterium bovis in the European, (Meles meles): epidemiological findings in the tuberculous badgers from a naturally infected population, 9

BCG, protective efficacy of, vaccine against leprosy in southern Malawi, 21

Campylobacter, epidemiological investigation of risk factors for, colonization in Norwegian broiler flocks, 245

Campylobacter jejuni, molecular subtyping by genome and plasmid analysis of, serogroups O1 and O2 (Penner) from sporadic and outbreak cases of human diarrhoea, 415

Chlamydial, the influence of local antichlamydial antibody on the acquisition and persistence of human ocular, infection: IgG antibodies are not protective, 315

Clostridium difficile, a molecular characterization of, isolates from humans, animals and their environments, 257

Colds, signs and symptoms in common, 143

Crimean—Congo haemorrhagic fever virus, enzyme-linked immunosorbent assays for the detection of antibody to, in the sera of livestock and wild vertebrates, 547

Dengue epidemic in the state of Rio de Janeiro. Brazil, 1990–1: co-circulation of dengue 1 and dengue 2 serotypes, 163

Escherichia coli. a miniaturized biotyping system for strain discrimination in. 81

 $Escherichia\ coli.\ {\it cattle\ as\ a\ possible\ source\ of\ verocytotoxin-producing.\ O157\ infections\ in\ man.}$

Escherichia coli, replicon typing characterization of plasmids encoding resistance to gentamicin and apramycin in, and Salmonella typhimurium isolated from human and animal sources in Belgium, 229

Escherichia coli, verotoxin producing, O157 infections associated with the consumption of yoghurt, 181

Escherichia coli, virulence properties of, strains belonging to serogroups O26, O55, O111 and O128 isolated in the United Kingdom in 1991 from patients with diarrhoea, 429

Faecal flora, characterization of human, by means of an improved fluoro-morphometrical method, 265

Faecal flora, the prevalence of antimicrobial resistance in human, in South Africa, 221

Faecal microflora, healthy individuals possess circulating antibodies against their indigenous, as well as against allogenous faecal microflora; an immunomorphometrical study, 273

Foot-and-mouth disease, the role of wild animals, other than buffalo, in the current epidemiology of, in Zimbabwe, 559

Foot-and-mouth disease virus, characterization of monoclonal antibodies against a type SAT 2, 391

Haemophilus parainfluenzae, an analysis of the diversity of, in the adult human respiratory tract by genomic DNA fingerprinting. 89

Hantavirus, the clinical spectrum of human, infection in Somerset, UK, 171

Helicobacter pylori, social and geographical risk factors in, infection, 63

Helicobacter pylori, the coccoid forms of. Criteria for their viability, 483

Influenza A, the haemagglutinins of. (H1N1) viruses in the 'O' or 'D' phases exhibit biological and antigenic differences. 135

Influenza immunization; vaccine offer, request and uptake in high-risk patients during the 1991/2 season, 347

Influenza type B, antigenic and genetic analyses of, viruses isolated in Russia, 1987-91, 539

Legionella pneumophila, intracellular growth of, serogroup 1 monoclonal antibody type 2 positive and negative bacteria, 499

Legionnaires' disease, a model for autumn outbreaks of, associated with cooling towers, linked to system operation and size, 287

Leishmaniasis, the epidemiology and surveillance of visceral, in the Campania region of Italy. The value of zymodeme typing, 297

Leprosy, protective efficacy of BCG vaccine against, in southern Malawi. 21

Listeria monocytogenes, comparison of multilocus enzyme electrophoresis (MEE), ribotyping, restriction enzyme analysis (REA) and phage typing for typing of, 71

Malaria prevalence amongst Brazilian Indians assessed by a new mathematical model. 525 Malaria transmission rates estimated from serological data, 503

MRSA-strains, outbreak of nosocomial infections with two different, involved; significance of genomic DNA fragment patterns in strains otherwise difficult to type, 55

Mycobacterium bovis in the European badger (Meles meles): epidemiological findings in tuberculous badgers from a naturally infected population. 9

Mycoplasma pulmonis infection of the murine oropharynx protects against subsequent vaginal colonization, 307

Pacific oysters, bacteriophage as models for virus removal from. (Crassostrea gigus) during relaying. 325

Paracoccidiodes brasiliensis, factors associated with, infection among permanent residents of three endemic areas in Colombia, 121

Pertussis, age-specific efficacy of, vaccine during epidemic and non-epidemic periods. 41 Pneumonoccal, levels of anti-, antibodies in young children in Papua New Guinea, 109

Respiratory virus infection, impact of, in patients with chronic disease, 337

Rotavirus, distinct yearly change of serotype distribution of human, in Thailand as determined by ELISA and PCR, 407

Salmonella enteritidis. interrelationships between strains of, 1

Salmonella enteritidis, egg age and the growth of, PT4 in egg contents. 209

Salmonella enteritidis, outer membrane characteristics of, phage type 4 growing in chickens. 449 Salmonella enteritidis, plasmid analysis of, isolated from human gastroenteritis cases and from epidemiologically associated poultry flocks. 239

Salmonella enteritidis, variations in biochemical phenotypes and phage types of. in Germany 1980–92, 199

Salmonella mikawasima, an outbreak of, associated with doner kebabs. 473

 $Salmonella\ typhimurium$, replicon typing characterization of plasmids encoding resistance to gentamicin and apramycin in $Escherichia\ coli$ and, isolated from human and animal sources in Belgium, 229

Salmonella typhimurium, the problems of tracing a geographically widespread outbreak of salmonellosis from a commonly eaten food: DT193 in North West England and North Wales in 1991, 465

Salmonellas, a comparison of multiple drug resistance in from humans and food animals in England and Wales, 1981 and 1990, 189

Salmonellas, a quantitative polymerase chain reaction method for the detection in avian faeces of, carrying the spvR gene, 455

Serratia marcescens, comparison of serotype, biotype and bacteriocin type with rDNA RFLP patterns for the type identification of, 99

SRSV, outbreak of, gastroenteritis at an international conference traced to food handled by a post-symptomatic caterer, 157

Streptoccous pneumoniae, acquisition and invasiveness of different serotypes of, in young children, 27

 $\label{like toxin} \emph{Vibrio mimicus}. \ \emph{high prevalence of thermostable direct hemolysin (TDH)-like toxin in. strains isolated from diarrhoeal patients. 49$

Viral meningitis, laboratory surveillance of, by examination of cerebrospinal fluid in Cape Town, $1981-9,\ 357$

Western equine encephalitis, weather factors in the prediction of, epidemics in Manitoba, 373

Yoghurt, verotoxin producing $\textit{Escherichia coli}\ \text{O157}$ infections associated with the consumption of, 481