




Hyperbolic geometry and real moduli of five points on the line

Olivier de Gaay Fortman 

ABSTRACT

We show that each connected component of the moduli space of smooth real binary quintics is isomorphic to an open subset of an arithmetic quotient of the real hyperbolic plane. Moreover, our main result says that the induced metric on this moduli space extends to a complete real hyperbolic orbifold structure on the space of stable real binary quintics. This turns the moduli space of stable real binary quintics into the quotient of the real hyperbolic plane by an explicit non-arithmetic triangle group.

1. Introduction

For interesting classes of complex varieties, there is a period map that identifies the moduli space with an open subset of an arithmetic quotient of a hermitian symmetric domain. Classical examples include abelian varieties, K3-surfaces, and configurations of points on the line. To study moduli of real algebraic varieties, several authors have analyzed the equivariance of the complex period map with respect to the action of complex conjugation on cohomology [Kha76, Nik79, Kha84, SS89, Yos98, AY98, DIK00]. An important difference between the complex and the real case is that moduli spaces of smooth real varieties are often not connected. This implies that a real period map has to be defined on each connected component of the moduli space separately; in favorable cases, this defines an isomorphism between any such component and the quotient of a Riemannian manifold by a discrete group of isometries (see e.g. [GH81, ACT10, Chu11, HR18]).

To salvage the non-connectedness of the real moduli space, one can sometimes define a slightly larger moduli space by allowing mild singularities. The idea is that, in such a larger space, the smooth varieties of one topological type do now deform into smooth varieties of another topological type, making the moduli space *connected*. In their beautiful paper [ACT10], Allcock, Carlson and Toledo showed that, for cubic surfaces, the real period maps defined on the various connected components of the moduli space of smooth surfaces extend to a global period map, defined on the moduli space of stable real cubic surfaces. In this way, they identified the latter with a single non-arithmetic real hyperbolic quotient. They proved analogous results for moduli of stable binary sextics, and stable binary sextics with a double root at infinity [ACT06; ACT07].

Received 4 August 2023, accepted 10 March 2025.

2020 Mathematics Subject Classification: 14J15, 14P99 (primary), 22E40 (secondary).

Keywords: Binary quintics, moduli spaces, period maps, real algebraic geometry, hyperbolic geometry.

© The Author(s), 2025. Published by Cambridge University Press on behalf of Foundation Compositio Mathematica, in partnership with the London Mathematical Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

It turns out that binary quintics provide a new example of this phenomenon. Let $X \cong \mathbf{A}_{\mathbf{R}}^6$ be the real algebraic variety that parametrizes homogeneous polynomials $F \in \mathbf{R}[x, y]$ of degree five. Let $X_0 \subset X$ parametrize polynomials with distinct roots, and $X_s \subset X$ polynomials with roots of multiplicity at most two (i.e. stable in the sense of geometric invariant theory). The principal goal of this paper is to study the geometry of the moduli space of stable *real binary quintics*

$$\mathcal{M}_s(\mathbf{R}) := \mathrm{GL}_2(\mathbf{R}) \setminus X_s(\mathbf{R}) \supset \mathrm{GL}_2(\mathbf{R}) \setminus X_0(\mathbf{R}) =: \mathcal{M}_0(\mathbf{R}).$$

Let $P_s \subset \mathbf{P}^1(\mathbf{C})^5$ be the set of five-tuples (x_1, \dots, x_5) such that no three $x_i \in \mathbf{P}^1(\mathbf{C})$ coincide (cf. [MS72]), and let $P_0 \subset P_s$ be the subset of five-tuples whose coordinates are distinct. These spaces are naturally acted upon by \mathfrak{S}_5 , the symmetric group on five letters. Moreover, complex conjugation $\sigma: \mathbf{P}^1(\mathbf{C})^5 \rightarrow \mathbf{P}^1(\mathbf{C})^5$ induces an anti-holomorphic involution $\sigma: P_s/\mathfrak{S}_5 \rightarrow P_s/\mathfrak{S}_5$ that preserves P_0/\mathfrak{S}_5 . Let $(P_0/\mathfrak{S}_5)(\mathbf{R})$ and $(P_s/\mathfrak{S}_5)(\mathbf{R})$ denote the respective fixed loci. Then

$$\mathcal{M}_0(\mathbf{R}) \cong \mathrm{PGL}_2(\mathbf{R}) \setminus (P_0/\mathfrak{S}_5)(\mathbf{R}) \quad \text{and} \quad \mathcal{M}_s(\mathbf{R}) \cong \mathrm{PGL}_2(\mathbf{R}) \setminus (P_s/\mathfrak{S}_5)(\mathbf{R}).$$

In other words, $\mathcal{M}_0(\mathbf{R})$ is the space of subsets $S \subset \mathbf{P}^1(\mathbf{C})$ of cardinality $|S| = 5$ that are stable under complex conjugation modulo real projective transformations; in $\mathcal{M}_s(\mathbf{R})$ one or two pairs of points are allowed to collapse.

By the Deligne–Mostow theory, the coarse moduli space $\mathcal{M}_s(\mathbf{C}) = \mathrm{GL}_2(\mathbf{C}) \setminus X_s(\mathbf{C})$ of stable complex binary quintics has a complex hyperbolic orbifold structure. Indeed, for five distinct points $u_1, \dots, u_5 \in \mathbf{A}^1(\mathbf{C}) \subset \mathbf{P}^1(\mathbf{C})$, the projective model of the normalization of the affine curve $z^5 = (x - u_1)^2 \cdots (x - u_5)^2$ is a smooth curve C of genus six; this curve C carries an automorphism of order five that induces an automorphism on the space $H^0(C, \Omega_C^1)$ of holomorphic one-forms on C whose $e^{2\pi i/5}$ -eigenspace defines a line in the corresponding eigenspace in $H^1(C(\mathbf{C}), \mathbf{C})$. This line is negative for a natural hermitian form, and hence one can associate to $\{u_1, \dots, u_5\}$ a point in a certain two-dimensional complex ball quotient $P\Gamma \setminus \mathbf{CH}^2$. This construction was already known to Shimura; see [Shi63; Shi64]. By varying the subset $\{u_1, \dots, u_5\}$ of points on $\mathbf{P}^1(\mathbf{C})$, or rather the associated complex binary quintic, one obtains a period map $\mathcal{M}_0(\mathbf{C}) = \mathrm{GL}_2(\mathbf{C}) \setminus X_0(\mathbf{C}) \rightarrow P\Gamma \setminus \mathbf{CH}^2$ (see Section 4.1 for details). The results of Deligne and Mostow [DM86] imply that this period map extends to an isomorphism of complex analytic spaces $\mathcal{M}_s(\mathbf{C}) = \mathrm{GL}_2(\mathbf{C}) \setminus X_s(\mathbf{C}) \xrightarrow{\sim} P\Gamma \setminus \mathbf{CH}^2$; see Theorem 4.1. Since strictly stable quintics correspond to points in a hyperplane arrangement $\mathcal{H} \subset \mathbf{CH}^2$ (see Proposition 4.3), one thus obtains an isomorphism

$$\mathcal{M}_0(\mathbf{C}) = \mathrm{GL}_2(\mathbf{C}) \setminus X_0(\mathbf{C}) \xrightarrow{\sim} P\Gamma \setminus (\mathbf{CH}^2 - \mathcal{H}). \quad (1)$$

By investigating the equivariance of the period map with respect to suitable anti-holomorphic involutions $\alpha_j: \mathbf{CH}^2 \rightarrow \mathbf{CH}^2$, we obtain the following real analogue.

THEOREM 1.1. *For $j \in \{0, 1, 2\}$, let \mathcal{M}_j be the connected component of $\mathcal{M}_0(\mathbf{R})$ parametrizing $\mathrm{Gal}(\mathbf{C}/\mathbf{R})$ -stable subsets $S \subset \mathbf{P}^1(\mathbf{C})$ with $2j$ complex and $5 - 2j$ real points. The period map induces an isomorphism of real analytic orbifolds*

$$\mathcal{M}_j \xrightarrow{\sim} P\Gamma_j \setminus (\mathbf{RH}^2 - \mathcal{H}_j). \quad (2)$$

Here \mathbf{RH}^2 is the real hyperbolic plane, \mathcal{H}_j a union of geodesic subspaces in \mathbf{RH}^2 , and $P\Gamma_j$ an arithmetic lattice in $\mathrm{PO}(2, 1)$. Moreover, the lattices $P\Gamma_j$ are projective orthogonal groups attached to explicit quadratic forms over $\mathbf{Z}[\lambda]$, $\lambda = \zeta_5 + \zeta_5^{-1} = (\sqrt{5} - 1)/2$; see (41).

In particular, Theorem 1.1 endows each connected component $\mathcal{M}_j \subset \mathcal{M}_0(\mathbf{R})$ with a hyperbolic metric. Since one can deform the topological type of a $\mathrm{Gal}(\mathbf{C}/\mathbf{R})$ -stable five-element subset of $\mathbf{P}^1(\mathbf{C})$ by allowing two points to collide, the compactification $\mathcal{M}_s(\mathbf{R}) \supset \mathcal{M}_0(\mathbf{R})$ is connected.




FIGURE 1. The moduli space of stable real binary quintics as the hyperbolic triangle $\Delta_{3,5,10} \subset \mathbf{RH}^2$. Here $\lambda = \zeta_5 + \zeta_5^{-1} = (\sqrt{5} - 1)/2$ and $\omega = \zeta_3$ (where $\zeta_n = e^{2\pi i/n} \in \mathbf{C}$ for $n \in \mathbf{Z}_{\geq 3}$).

One may wonder whether the metrics on the components \mathcal{M}_j extend to a metric on the whole of $\mathcal{M}_s(\mathbf{R})$. If so, what does the resulting space look like at the boundary? Our main result answers these questions in the following way.

THEOREM 1.2. *There exists a complete hyperbolic metric on $\mathcal{M}_s(\mathbf{R})$ that restricts to the metrics on \mathcal{M}_j induced by (2). Let $\overline{\mathcal{M}}_{\mathbf{R}}$ denote the resulting metric space, and define $\Gamma_{3,5,10}$ as the group*

$$\Gamma_{3,5,10} = \langle a, b, c \mid a^2 = b^2 = c^2 = (ab)^3 = (ac)^5 = (bc)^{10} = 1 \rangle. \quad (3)$$

There exist an open embedding $\coprod_j P\Gamma_j \setminus (\mathbf{RH}^2 - \mathcal{H}_j) \hookrightarrow \Gamma_{3,5,10} \setminus \mathbf{RH}^2$ and an isometry

$$\overline{\mathcal{M}}_{\mathbf{R}} \cong \Gamma_{3,5,10} \setminus \mathbf{RH}^2 \quad (4)$$

that extend the real analytic orbifold isomorphisms (2) in Theorem 1.1. In particular, $\overline{\mathcal{M}}_{\mathbf{R}}$ is isometric to the hyperbolic triangle $\Delta_{3,5,10}$ with angles $\pi/3, \pi/5, \pi/10$; see Figure 1.

Note that the closure

$$\overline{\mathcal{M}}_0 \subset \overline{\mathcal{M}}_{\mathbf{R}}$$

of \mathcal{M}_0 in $\overline{\mathcal{M}}_{\mathbf{R}}$ is the moduli space of stable configurations of five real points on $\mathbf{P}_{\mathbf{R}}^1$. This moduli space was investigated by Apéry and Yoshida in [AY98], who proved that $\overline{\mathcal{M}}_0$ is the hyperbolic

triangle with angles $\pi/2, \pi/4$ and $\pi/5$. From this, together with the knowledge of the angles of $\overline{\mathcal{M}}_{\mathbf{R}}$ and the fact that the two hyperplanes in Figure 1 intersect orthogonally, one can deduce the remaining angles of the closures $\overline{\mathcal{M}}_j \subset \overline{\mathcal{M}}_{\mathbf{R}}$ of the subsets $\mathcal{M}_j \subset \overline{\mathcal{M}}_{\mathbf{R}}$ ($j \in \{0, 1, 2\}$).

THEOREM 1.3. Consider Figure 1. For $j = 0, 1, 2$, let $\overline{\mathcal{M}}_j \subset \overline{\mathcal{M}}_{\mathbf{R}}$ be the closure of $\mathcal{M}_j \subset \overline{\mathcal{M}}_{\mathbf{R}}$.

- (i) The angle of $\overline{\mathcal{M}}_0$ at $(0, -1, \infty, \infty, 1)$ is $\pi/2$, and its angle at $(0, -1, -1, \infty, \infty)$ is $\pi/4$.
- (ii) The angle of $\overline{\mathcal{M}}_1$ at $(0, -1, \infty, \infty, 1)$ is $\pi/2$, and its angle at $(0, -i, \infty, \infty, i)$ is $\pi/2$.
- (iii) The angle of $\overline{\mathcal{M}}_2$ at $(0, -1, -1, \infty, \infty)$ is $\pi/4$, and its angle at $(0, -i, \infty, \infty, i)$ is $\pi/2$.

Remark 1.4.

- (i) The lattice $\Gamma_{3,5,10} \subset \mathrm{PO}(2, 1)$ is non-arithmetic; see [Tak77].
 - (ii) The topological space $\mathcal{M}_s(\mathbf{R})$ underlies two topological orbifold structures: the natural orbifold structure on $\mathrm{GL}_2(\mathbf{R}) \setminus X_s(\mathbf{R})$ and the orbifold structure on $\overline{\mathcal{M}}_{\mathbf{R}}$ induced by (4). These orbifold structures only differ at one point of $\mathcal{M}_s(\mathbf{R})$, which is $(\infty, i, i, -i, -i)$ (see Figure 1). The stabilizer group of $\mathcal{M}_s(\mathbf{R})$ at $(\infty, i, i, -i, -i)$ is isomorphic to $\mathbf{Z}/2$, whereas the stabilizer group of $\overline{\mathcal{M}}_{\mathbf{R}}$ at $(\infty, i, i, -i, -i)$ is isomorphic to the dihedral group of order 20.
 - (iii) Important ingredients in the proof of Theorem 1.2 are:
 - (a) the fact that under the complex period map (1), moduli of singular binary quintics correspond to points in the quotient of a certain hyperplane arrangement $\mathcal{H} \subset \mathbf{CH}^2$; and
 - (b) the fact that the hyperplane arrangement $\mathcal{H} \subset \mathbf{CH}^2$ is an orthogonal arrangement in the sense of [ACT02a].
- We prove (a) in Proposition 4.3, and (b) holds by [GF24, Theorem 2.5 and Example 2.12].

Remark 1.5. Let $PX_0(\mathbf{C})$ denote the space of \mathbf{C}^* -equivalence classes of smooth complex binary quintics $F \in \mathbf{C}[x, y]$. The natural map $P_0 \rightarrow PX_0(\mathbf{C})$ induces a $\mathrm{PGL}_2(\mathbf{C})$ -equivariant isomorphism $P_0/\mathfrak{S}_5 \xrightarrow{\sim} PX_0(\mathbf{C})$, and the quotient $\mathcal{M}_0(\mathbf{C}) = \mathrm{PGL}_2(\mathbf{C}) \setminus PX_0(\mathbf{C})$ is the moduli space of smooth complex binary quintics. It turns out that neither $\pi_1(PX_0(\mathbf{C}))$ nor $\pi_1^{\mathrm{orb}}(\mathcal{M}_0(\mathbf{C}))$ is a lattice in any Lie group with finitely many connected components. In view of [ACT02a, Theorem 1.2], this follows from the isomorphism $\mathcal{M}_0(\mathbf{C}) \cong P\Gamma \setminus (\mathbf{CH}^2 - \mathcal{H})$ (see (1) above) and the orthogonality of the hyperplane arrangement $\mathcal{H} \subset \mathbf{CH}^2$ (see Remark 1.4(iii)(b) above).

1.1 Overview of this paper

In Section 3, we recall known results on families of quintic covers of the complex projective line, branched along a binary quintic. We consider moduli of complex binary quintics in Section 4. In particular, we show that the Deligne–Mostow theory provides an isomorphism between the space of stable complex binary quintics and an arithmetic ball quotient. In Section 5, we prove that moduli of stable real binary quintics are in one-to-one correspondence with points in the real hyperbolic quotient space $P\Gamma_{\mathbf{R}} \setminus \mathbf{RH}^2$ defined by a lattice $P\Gamma_{\mathbf{R}} \subset \mathrm{PO}(2, 1)$. We calculate $P\Gamma_{\mathbf{R}}$ in Section 6: it is conjugate to the lattice $\Gamma_{3,5,10}$ defined in (3). In Section 7, we study monodromy groups of moduli spaces of smooth binary quintics over \mathbf{C} and over \mathbf{R} , and prove Theorems 1.2 and 1.3. In Section 8, we use [GF24, Theorem 1.8] and the main results of this paper to provide an explicit sequence $\{\Gamma_n \subset \mathrm{PO}(n, 1)\}_{n \geq 2}$ of non-arithmetic lattices Γ_n , with $\Gamma_2 = P\Gamma_{\mathbf{R}}$.

2. Notation

Let K be the cyclotomic field $\mathbf{Q}(\zeta)$, where

$$\zeta := \zeta_5 = e^{2\pi i/5} \in \mathbf{C}.$$

The ring of integers \mathcal{O}_K of K is $\mathbf{Z}[\zeta]$ (see e.g. [Neu99, Chapter I, Proposition 10.2]). Let $\mu_K \subset \mathcal{O}_K^*$ be the torsion subgroup of the unit group \mathcal{O}_K^* , and recall that μ_K is cyclic of order ten, generated by $-\zeta$. Define an involution $\rho: K \rightarrow K$ by $\rho(\zeta) = \zeta^{-1}$, and let $F = K^\rho$ be the maximal totally real subfield of K . Recall that F is generated over \mathbf{Q} by the element

$$\lambda := \zeta + \zeta^{-1} = (\sqrt{5} - 1)/2.$$

Define

$$\eta = \zeta^2 - \zeta^{-2} \in \mathcal{O}_K,$$

and consider the different ideal $\mathfrak{D}_K \subset \mathcal{O}_K$ (see e.g. [Neu99, Chapter III, Section 2, Definition 2.1] or [Ser79, Chapter III, Section 3]). We have $(\zeta - \zeta^{-1}) \cdot (\zeta^2 - \zeta^{-2}) \cdot (\zeta^3 - \zeta^{-3}) \cdot (\zeta^4 - \zeta^{-4}) = 5$, and hence $(\eta)^4 = (5)$ as ideals of \mathcal{O}_K . This implies (see e.g. [Neu99, Chapter III, Theorem 2.6]) that

$$\mathfrak{D}_K = (5/\eta) = (\eta)^3.$$

3. Preliminaries on quintic covers of the projective line branched along five points

In this section, we recollect some known results on quintic covers $C \rightarrow \mathbf{P}_{\mathbf{C}}^1$ ramified along five points with local monodromy $\exp(4\pi i/5)$ around each point. Some of these results are well known but hard to find in the literature; we state and prove these for the convenience of the reader. Other results of this section are available in the literature, but we formulate them in a different manner.

Recall from the introduction that $X \cong \mathbf{A}_{\mathbf{R}}^6$ is the real algebraic variety that parametrizes homogeneous polynomials of degree five, X_0 the subvariety of polynomials with distinct roots, and $X_s \subset X$ the subvariety of polynomials with roots of multiplicity at most two (i.e. non-zero polynomials whose class in the associated projective space is stable in the sense of geometric invariant theory [MFK94] for the action of $\mathrm{SL}_{2,\mathbf{R}}$ on it).

3.1 Refined Hodge numbers of a quintic cover of \mathbf{P}^1 branched along five points

Let $F \in X_0(\mathbf{C})$ be a smooth binary quintic; thus $F = F(x, y) \in \mathbf{C}[x, y]$ is a homogeneous polynomial of degree five whose zeros in $\mathbf{P}^1(\mathbf{C})$ all have multiplicity one. Let $t_1, \dots, t_5 \in \mathbf{P}^1(\mathbf{C})$ be the zeros of F , with $t_j = [u_j : v_j]$ in homogeneous coordinates of $\mathbf{P}^1(\mathbf{C})$, for $j = 1, \dots, 5$. Let

$$C_F \rightarrow \mathbf{P}_{\mathbf{C}}^1 \tag{5}$$

be the cyclic quintic cover with branch points $t_1, \dots, t_5 \in \mathbf{P}^1(\mathbf{C})$ and local monodromy

$$\exp(2\pi i \cdot 2/5) \in \mu_5 \tag{6}$$

around t_j for each $j \in \{1, \dots, 5\}$. If the points $t_j = [u_j : 1]$ are all in $\mathbf{A}^1(\mathbf{C})$, the cover (5) is, in affine coordinates, given by the normalization of the curve defined by the equation

$$z^5 = (x - u_1)^2 \cdot (x - u_2)^2 \cdots (x - u_5)^2,$$

with $\zeta \in \mu_5 \subset \mathbf{C}$ acting by $(x, z) \mapsto (x, \zeta \cdot z)$. We have $g = 6$ for the genus $g = g(C_F)$ of the curve C_F . Let JC_F be the Jacobian of the curve C_F , so that

$$JC_F(\mathbf{C}) = H^1(C_F(\mathbf{C}), \mathcal{O}_{C_F})/H^1(C_F(\mathbf{C}), \mathbf{Z})(1),$$

with weight -1 Hodge decomposition

$$H_1(JC_F(\mathbf{C}), \mathbf{C}) = H^1(C_F(\mathbf{C}), \mathbf{Z})(1) \otimes \mathbf{C} = H^{-1,0}(JC_F) \oplus H^{0,-1}(JC_F). \quad (7)$$

Note that $H^{0,-1}(JC_F)$ is naturally isomorphic to the space $H^0(C_F, \Omega^1)$ of global holomorphic differentials on the curve C_F . The order five automorphism

$$\zeta: C_F \xrightarrow{\sim} C_F$$

defined above induces an embedding of rings

$$\varphi: \mathbf{Z}[\zeta] \rightarrow \text{End}(JC_F), \quad \varphi(\zeta) = (\zeta^{-1})^*,$$

which is compatible with the Hodge decomposition (7). For $k \in \{1, 2, 3, 4\}$, define

$$H^{0,-1}(JC_F)_{\zeta^k} = \left\{ x \in H^{0,-1}(JC_F) \mid \varphi(\zeta)x = \zeta^k x \right\} \subset H^{0,-1}(JC_F) = H^{1,0}(C_F) = H^0(C_F, \Omega^1),$$

and define $H^{-1,0}(JC_F)_{\zeta^k} \subset H^{-1,0}(JC_F)$ in a similar way.

The way to calculate the refined Hodge numbers $h^{0,-1}(JC_F)_{\zeta^k} = \dim H^{0,-1}(JC_F)_{\zeta^k}$ and $h^{-1,0}(JC_F)_{\zeta^k} = \dim H^{-1,0}(JC_F)_{\zeta^k}$ for $k = 1, 2, 3, 4$ is well known; the result is as follows.

LEMMA 3.1. *Let $F \in X_0(\mathbf{C})$ be a smooth binary quintic, and let JC_F be the Jacobian of the cyclic cover $C_F \rightarrow \mathbf{P}^1(\mathbf{C})$ associated to F as in (5). One has the following refined Hodge numbers:*

$$\begin{aligned} h^{0,-1}(JC_F)_{\zeta} &= 1, & h^{0,-1}(JC_F)_{\zeta^2} &= 3, & h^{0,-1}(JC_F)_{\zeta^3} &= 0, & h^{0,-1}(JC_F)_{\zeta^4} &= 2, \\ h^{-1,0}(JC_F)_{\zeta} &= 2, & h^{-1,0}(JC_F)_{\zeta^2} &= 0, & h^{-1,0}(JC_F)_{\zeta^3} &= 3, & h^{-1,0}(JC_F)_{\zeta^4} &= 1. \end{aligned}$$

Proof. This follows from the Hurwitz–Chevalley–Weil formula (see [MO13, Proposition 5.9]). Alternatively, see [Loo07, Lemma 4.2]. \square

3.2 The hermitian lattice

We fix a smooth binary quintic $F_0 \in X_0(\mathbf{C})$. Let $C \rightarrow \mathbf{P}_{\mathbf{C}}^1$ be the cyclic cover of $\mathbf{P}_{\mathbf{C}}^1$ associated to F_0 as in (5), and consider the Jacobian variety with \mathcal{O}_K -action

$$(JC, \quad \varphi: \mathcal{O}_K = \mathbf{Z}[\zeta] \rightarrow \text{End}(A).$$

See Section 3.1 above. Define Λ as the free \mathcal{O}_K -module

$$\Lambda = H^1(C(\mathbf{C}), \mathbf{Z}).$$

The canonical principal polarization of the Jacobian JC is given by a symplectic pairing

$$E: \Lambda \times \Lambda = H^1(C(\mathbf{C}), \mathbf{Z}) \times H^1(C(\mathbf{C}), \mathbf{Z}) \rightarrow H^2(C(\mathbf{C}), \mathbf{Z}) = \mathbf{Z}$$

which satisfies $E(\varphi(a)x, y) = E(x, \varphi(\rho(a))y)$ for each $a \in \mathcal{O}_K$ and $x, y \in \Lambda$, where ρ is the automorphism $K \xrightarrow{\sim} K, \zeta \mapsto \zeta^{-1}$ (see Section 2). Consider the different ideal $\mathfrak{D}_K \subset \mathcal{O}_K$, and define a skew-hermitian form T on Λ as follows:

$$T: \Lambda \times \Lambda \rightarrow \mathfrak{D}_K^{-1}, \quad T(x, y) = \frac{1}{5} \sum_{j=0}^4 \zeta^j E(x, \varphi(\zeta)^j y).$$

By [GF24, Example 11.2.2], this is the skew-hermitian form corresponding to E , via [GF24, Lemma 11.1]. Let $\eta \in \mathcal{O}_K$ be the purely imaginary element

$$\eta = \zeta^2 - \zeta^{-2} \in \mathcal{O}_K,$$

and note that the different ideal $\mathfrak{D}_K \subset \mathcal{O}_K$ is generated by $5/\eta$ (see Section 2). We obtain a hermitian form on the free \mathcal{O}_K -module Λ as follows:

$$\mathfrak{h}: \Lambda \times \Lambda \rightarrow \mathcal{O}_K, \quad \mathfrak{h}(x, y) = \frac{5}{\eta} \cdot T(x, y) = \frac{1}{\zeta^2 - \zeta^{-2}} \cdot \sum_{j=0}^4 \zeta^j E(x, \varphi(\zeta)^j y). \quad (8)$$

By [GF24, Lemma 11.1], we have that (Λ, \mathfrak{h}) is unimodular as a hermitian lattice over \mathcal{O}_K , because (Λ, E) is unimodular as an alternating lattice over \mathbf{Z} .

3.3 The signature

Recall that a CM type on a CM field L of degree $2g$ over \mathbf{Q} is a set $\Phi \subset \text{Hom}(L, \mathbf{C})$ of g embeddings of L into \mathbf{C} , such that for each embedding $\varphi: L \rightarrow \mathbf{C}$, either φ lies in Φ or the complex conjugate of φ does.

Define a CM type $\Phi \subset \text{Hom}(K, \mathbf{C})$ as

$$\Phi = \{\sigma_1, \sigma_2: K \rightarrow \mathbf{C}\}, \quad \sigma_k(\zeta) = \zeta^k \quad \text{for } k = 1, 2. \quad (9)$$

Observe that, for $k = 1, 2$, we have

$$\Lambda \otimes_{\mathcal{O}_K, \sigma_k} \mathbf{C} = (\Lambda \otimes_{\mathbf{Z}} \mathbf{C})_{\zeta^k} = H^1(C(\mathbf{C}), \mathbf{C})_{\zeta^k}.$$

LEMMA 3.2. For $k \in \{1, 2\}$, consider the hermitian form

$$\mathfrak{h}^{\sigma_k}: H^1(C(\mathbf{C}), \mathbf{C})_{\zeta^k} \times H^1(C(\mathbf{C}), \mathbf{C})_{\zeta^k} \rightarrow \mathbf{C}.$$

Then $\text{sign}(\mathfrak{h}^{\sigma_1}) = (2, 1)$ and $\text{sign}(\mathfrak{h}^{\sigma_2}) = (3, 0)$, where $\text{sign}(\mathfrak{h}^{\sigma_k})$ is the signature of \mathfrak{h}^{σ_k} .

Proof. Write $\Lambda_{\mathbf{C}} = \Lambda \otimes_{\mathbf{Z}} \mathbf{C} = H^1(C(\mathbf{C}), \mathbf{C})$. For each embedding $\phi: K \rightarrow \mathbf{C}$, the restriction of the hermitian form

$$\phi(5/\eta) \cdot E_{\mathbf{C}}(x, \bar{y}): \Lambda_{\mathbf{C}} \times \Lambda_{\mathbf{C}} \rightarrow \mathbf{C}$$

to $(\Lambda_{\mathbf{C}})_{\phi} \subset \Lambda_{\mathbf{C}}$ coincides with \mathfrak{h}^{ϕ} by [GF24, Lemma 11.3]. Moreover, the hermitian form

$$i \cdot E_{\mathbf{C}}(x, \bar{y}): H^1(C(\mathbf{C}), \mathbf{C}) \times H^1(C(\mathbf{C}), \mathbf{C}) \rightarrow \mathbf{C}$$

is positive definite on $H^0(C, \Omega^1) = H^{1,0}(C) = H^{0,-1}(JC)$ and negative definite on $H^{-1,0}(JC) = H^{0,1}(C)$ (see [Voi02, Théorème 6.32]). As $\Im(\sigma_1(\eta)) > 0$ and $\Im(\sigma_2(\eta)) < 0$, we have $\Im(\sigma_1(5/\eta)) < 0$ and $\Im(\sigma_2(5/\eta)) > 0$. Consequently, the hermitian form

$$\sigma_1(5/\eta) \cdot E_{\mathbf{C}}(x, \bar{y}) = \mathfrak{h}^{\sigma_1}(x, y): H^1(C(\mathbf{C}), \mathbf{C})_{\zeta} \times H^1(C(\mathbf{C}), \mathbf{C})_{\zeta} \rightarrow \mathbf{C}$$

is negative definite on $H^{0,-1}(JC)_{\zeta}$ and positive definite on $H^{-1,0}(JC)_{\zeta}$, so that $\text{sign}(\mathfrak{h}^{\sigma_1}) = (2, 1)$ by Lemma 3.1. Similarly, the hermitian form

$$\sigma_2(5/\eta) \cdot E_{\mathbf{C}}(x, \bar{y}) = \mathfrak{h}^{\sigma_2}(x, y): H^1(C(\mathbf{C}), \mathbf{C})_{\zeta^2} \times H^1(C(\mathbf{C}), \mathbf{C})_{\zeta^2} \rightarrow \mathbf{C}$$

is positive definite on $H^{0,-1}(JC)_{\zeta^2}$ and negative definite on $H^{-1,0}(JC)_{\zeta^2}$. Hence, using Lemma 3.1 again, we conclude that $\text{sign}(\mathfrak{h}^{\sigma_2}) = (3, 0)$. \square

3.4 The monodromy representation

Consider the real algebraic variety X_0 introduced in Section 1. Let $D \subset \mathrm{GL}_2(\mathbf{C})$ be the subgroup $D = \{\zeta^j \cdot \mathrm{Id}\} \subset \mathrm{GL}_2(\mathbf{C})$ of scalar matrices of the form $\zeta^j \cdot \mathrm{Id}$ with $j \in \{0, 1, 2, 3, 4\}$ and $\mathrm{Id} \in \mathrm{GL}_2(\mathbf{C})$ the identity two-by-two matrix. Define

$$G(\mathbf{C}) = \mathrm{GL}_2(\mathbf{C})/D. \quad (10)$$

The group $G(\mathbf{C})$ acts from the left on $X_0(\mathbf{C})$ in the following way: if

$$F = F(x, y) \in \mathbf{C}[x, y]$$

is a binary quintic, we may view F as a function $\mathbf{C}^2 \rightarrow \mathbf{C}$, and define $g \cdot F = F(g^{-1})$ for $g \in G(\mathbf{C})$. This gives a canonical isomorphism of complex analytic orbifolds

$$\mathcal{M}_0(\mathbf{C}) = G(\mathbf{C}) \backslash X_0(\mathbf{C}),$$

where \mathcal{M}_0 is the moduli stack of smooth binary quintics. Define

$$g: \mathcal{C} \rightarrow X_0 \quad (11)$$

as the universal family of cyclic quintic covers $C \rightarrow \mathbf{P}^1$ ramified along a smooth binary quintic $\{F = 0\} \subset \mathbf{P}^1$ with local monodromy given by (6), and let

$$J = J(\mathcal{C}) \rightarrow X_0$$

be the relative Jacobian of \mathcal{C}/X_0 . By Lemma 3.2, J is a polarized abelian scheme of relative dimension six over X_0 , equipped with \mathcal{O}_K -action of signature $\{(2, 1), (3, 0)\}$ with respect to $\Phi = \{\sigma_1, \sigma_2\}$. Let $\mathbf{V} = R^1 g_* \mathbf{Z}$ be the local system of hermitian \mathcal{O}_K -modules underlying the abelian scheme J/X_0 . Attached to \mathbf{V} and the base point $F_0 \in X_0(\mathbf{C})$, we have a representation

$$\rho_\Gamma: \pi_1(X_0(\mathbf{C}), F_0) \rightarrow \Gamma := \mathrm{Aut}_{\mathcal{O}_K}(\Lambda, \mathfrak{h}) \quad (12)$$

whose composition with the quotient map $\Gamma \rightarrow P\Gamma = \Gamma/\mu_K$ defines a homomorphism

$$P(\rho_\Gamma): \pi_1(X_0(\mathbf{C}), F_0) \rightarrow P\Gamma. \quad (13)$$

3.5 Marked binary quintics and points on the projective line

Let $F \in X_0(\mathbf{C})$ and consider the hypersurface $Z_F := \{F = 0\} \subset \mathbf{P}_{\mathbf{C}}^1$. A *marking* of F is an ordering $m: Z_F(\mathbf{C}) \xrightarrow{\sim} \{1, 2, 3, 4, 5\}$ of the five-element set $Z_F(\mathbf{C}) = \{x \in \mathbf{P}^1(\mathbf{C}) \mid F(x) = 0\}$.

Remark 3.3. Let $F \in X_0(\mathbf{C})$. To give a marking of F is to give an isomorphism of rings $H^0(Z_F(\mathbf{C}), \mathbf{Z}) \xrightarrow{\sim} \mathbf{Z}^5$. Moreover, sending a ring automorphism of \mathbf{Z}^5 to the induced permutation of the canonical basis of $\{e_1, \dots, e_5\} \subset \mathbf{Z}^5$ defines a group isomorphism $\mathrm{Aut}_{\mathrm{ring}}(\mathbf{Z}^5) \cong \mathfrak{S}_5$.

Define \mathcal{N}_0 as the set of marked smooth complex binary quintics (F, m) , and consider the map that forgets the marking:

$$\mathcal{N}_0 \rightarrow X_0(\mathbf{C}), \quad (F, m) \mapsto F. \quad (14)$$

To provide \mathcal{N}_0 with a complex manifold structure such that (14) is a finite covering map, we define \mathcal{N}_0 in a slightly different but equivalent way. Define an algebraic subgroup

$$T := \{(\lambda_1, \dots, \lambda_5) \in (\mathbf{C}^*)^5 \mid \lambda_1 \cdots \lambda_5 = 1\} \subset (\mathbf{C}^*)^5.$$

The group T acts freely on $(\mathbf{C}^2 - \{0\})^5$ in the usual way, i.e. by letting $(\lambda_1, \dots, \lambda_5) \in T$ act as

$$((a_1, b_1), \dots, (a_5, b_5)) \mapsto (\lambda_1 \cdot (a_1, b_1), \dots, \lambda_5 \cdot (a_5, b_5)).$$

As $\dim(T) = 4$, the quotient variety $\mathcal{N} := (\mathbf{C}^2 - \{0\})^5 / T$ is a complex manifold of dimension six, equipped with a holomorphic principal \mathbf{C}^* -bundle map

$$f: \mathcal{N} \rightarrow \mathbf{P}^1(\mathbf{C})^5, \quad (v_1, \dots, v_5) \mapsto ([v_1], \dots, [v_5]), \quad (15)$$

where \mathbf{C}^* acts on \mathcal{N} by $\lambda \cdot (v_1, v_2, \dots, v_5) = (\lambda v_1, v_2, \dots, v_5) = (v_1, \lambda v_2, \dots, v_5) = \dots = (v_1, v_2, \dots, \lambda v_5) \in \mathcal{N}$ for $\lambda \in \mathbf{C}^*$ and $(v_1, v_2, \dots, v_5) \in \mathcal{N}$. For $i, j \in \{1, \dots, 5\}$ with $i < j$, define $\overline{\Delta}_{ij} = \{(x_1, \dots, x_5) \in \mathbf{P}^1(\mathbf{C})^5 \mid x_i = x_j\}$, and let

$$\Delta_{ij} := f^{-1}(\overline{\Delta}_{ij}),$$

where f is the principal bundle map (15). Since $\overline{\Delta}_{ij} \subset \mathbf{P}^1(\mathbf{C})^5$ is Zariski closed in $\mathbf{P}^1(\mathbf{C})^5$, we see that Δ_{ij} is Zariski closed in \mathcal{N} . Define a Zariski open subset $\mathcal{N}_0 \subset \mathcal{N}$ as follows:

$$\mathcal{N}_0 := \mathcal{N} - \bigcup_{i < j} \Delta_{ij}.$$

LEMMA 3.4. *The complex manifold \mathcal{N}_0 is connected and there is a canonical bijection between \mathcal{N}_0 and the set of marked smooth complex binary quintics (F, m) . Under this bijection, the map*

$$\mathcal{N}_0 \rightarrow X_0(\mathbf{C}), \quad ((a_1, b_1), (a_2, b_2), \dots, (a_5, b_5)) \mapsto \prod_{j=1}^5 (b_j \cdot x - a_j \cdot y) \quad (16)$$

corresponds to the map (14) that forgets the marking, and (16) is a Galois covering with Galois group \mathfrak{S}_5 .

Proof. Note that \mathcal{N}_0 is connected because $(\mathbf{C}^2 - \{0\})^2$ is connected. For $v = ((a_1, b_1), \dots, (a_5, b_5)) \in \mathcal{N}_0$, define $F_v = \prod_{j=1}^5 (b_j \cdot x - a_j \cdot y)$. The polynomial F_v is a smooth binary quintic, equipped with the marking m_v defined by the ordering $\{[a_1 : b_1], \dots, [a_5 : b_5]\}$ of the roots $[a_j : b_j] \in \mathbf{P}^1(\mathbf{C})$ of F_v . The association $v \mapsto (F_v, m_v)$ defines the bijection alluded to in the first statement of the lemma. The rest is clear. \square

In a similar way, we define, for $i, j, k \in \{1, \dots, 5\}$ with $i < j < k$, closed subsets

$$\overline{\Delta}_{ijk} := \{(x_1, \dots, x_5) \in \mathbf{P}^1(\mathbf{C})^5 \mid x_i = x_j = x_k\}, \quad \Delta_{ijk} := f^{-1}(\overline{\Delta}_{ijk}) \subset \mathcal{N},$$

and put

$$\mathcal{N}_s := \mathcal{N} - \bigcup_{i < j < k} \Delta_{ijk}.$$

The space \mathcal{N}_s is equipped with a finite ramified covering map

$$\mathcal{N}_s \rightarrow X_s(\mathbf{C}), \quad ((a_1, b_1), (a_2, b_2), \dots, (a_5, b_5)) \mapsto \prod_{j=1}^5 (b_j \cdot x - a_j \cdot y) \in \mathbf{C}[x, y],$$

that commutes with (16) and the natural open embedding $\mathcal{N}_0 \subset \mathcal{N}_s$. Observe that $\mathcal{N}_s \rightarrow X_s(\mathbf{C})$ is the Fox completion (cf. [Fox57] or [DM86, Section 8.1]) of the spread $\mathcal{N}_0 \rightarrow X_0(\mathbf{C}) \hookrightarrow X_s(\mathbf{C})$.

3.6 Monodromy

Choose a marking m_0 lying over our base point $F_0 \in X_0(\mathbf{C})$. By Lemma 3.4, we have a surjective homomorphism

$$\rho_5: \pi_1(X_0(\mathbf{C}), F_0) \twoheadrightarrow \mathfrak{S}_5 \quad (17)$$

whose kernel is given by the image of the natural embedding $\pi_1(\mathcal{N}_0, m_0) \hookrightarrow \pi_1(X_0(\mathbf{C}), F_0)$. Composing the latter with the maps ρ_Γ and $P(\rho_\Gamma)$ of (12) and (13) yields homomorphisms

$$\mu: \pi_1(\mathcal{N}_0, m_0) \rightarrow \Gamma \quad \text{and} \quad P(\mu): \pi_1(\mathcal{N}_0, m_0) \rightarrow P\Gamma. \quad (18)$$

Consider the three-dimensional \mathbf{F}_5 vector space $\Lambda/(1-\zeta)\Lambda$, as well as the quadratic space

$$W := (\Lambda/(1-\zeta)\Lambda, \mathfrak{q})$$

over \mathbf{F}_5 . Here, \mathfrak{q} is the quadratic form obtained by reducing \mathfrak{h} modulo $(1-\zeta)\Lambda$. Define two groups Γ_θ and $P\Gamma_\theta$ as follows:

$$\Gamma_\theta = \text{Ker}(\Gamma \rightarrow \text{Aut}(W)), \quad P\Gamma_\theta = \text{Ker}(P\Gamma \rightarrow P\text{Aut}(W)).$$

The following proposition seems to be due to Terada [Ter83] and Yamazaki and Yoshida [YY84].

PROPOSITION 3.5. *The image of the map $P(\mu)$ defined in (18) is the group $P\Gamma_\theta$. Moreover, the map $P\Gamma \rightarrow P\text{Aut}(W)$ is surjective, and the induced homomorphism*

$$\bar{\rho}_\Gamma: \mathfrak{S}_5 = \pi_1(X_0(\mathbf{C}), F_0)/\pi_1(\mathcal{N}_0, m_0) \rightarrow P\Gamma/P\Gamma_\theta = P\text{Aut}(W)$$

is an isomorphism. As a consequence, we obtain the following commutative diagram with exact rows.

$$\begin{array}{ccccccc} 0 & \longrightarrow & \pi_1(\mathcal{N}_0, m_0) & \longrightarrow & \pi_1(X_0(\mathbf{C}), F_0) & \xrightarrow{\rho_5} & \mathfrak{S}_5 \longrightarrow 0 \\ & & \downarrow P(\mu) & & \downarrow P(\rho_\Gamma) & & \downarrow \wr \bar{\rho}_\Gamma \\ 0 & \longrightarrow & P\Gamma_\theta & \longrightarrow & P\Gamma & \longrightarrow & P\text{Aut}(W) \longrightarrow 0 \end{array} \quad (19)$$

Proof. The equality $P(\mu)(\pi_1(\mathcal{N}_0, m_0)) = P\Gamma_\theta$ follows from $\mu(\pi_1(\mathcal{N}_0, m_0)) = \Gamma_\theta$. For the latter, see [YY84, Propositions 4.2 and 4.3]. The group $P\text{Aut}(W) \cong \text{PO}_3(\mathbf{F}_5)$ is isomorphic to \mathfrak{S}_5 , and $\bar{\rho}_\Gamma: \mathfrak{S}_5 \rightarrow P\Gamma/P\Gamma_\theta$ is an isomorphism (cf. [YY84, Propositions 4.2 and 4.3] and [AY98, p. 10]). \square

COROLLARY 3.6. *The monodromy representation $P(\rho_\Gamma): \pi_1(X_0(\mathbf{C}), F_0) \rightarrow P\Gamma$ is surjective.*

3.7 Framed binary quintics, nodal binary quintics and local monodromy

By a *framing* of a point $F \in X_0(\mathbf{C})$ we mean a projective equivalence class $[f]$, where

$$f: \mathbf{V}_F = H^1(C_F(\mathbf{C}), \mathbf{Z}) \xrightarrow{\sim} \Lambda$$

is an \mathcal{O}_K -linear isometry: two such isometries are in the same class if and only if they differ by an element in $\mu_K = (\mathcal{O}_K^*)_{\text{tors}}$. Let \mathcal{F}_0 be the collection of all framings of all points $F \in X_0(\mathbf{C})$. The set \mathcal{F}_0 can naturally be given the structure of a complex manifold, in a way similar to the procedure described in [ACT02b, (3.9)]. In what follows, we consider \mathcal{F}_0 as a complex manifold.

LEMMA 3.7.

(i) *The complex manifold \mathcal{F}_0 is connected and the map*

$$\mathcal{F}_0 \rightarrow X_0(\mathbf{C}), \quad (F, [f]) \mapsto F \quad (20)$$

that forgets the framing is a Galois covering map, with Galois group $P\Gamma$.

(ii) *The spaces $P\Gamma_\theta \backslash \mathcal{F}_0$ and \mathcal{N}_0 are isomorphic as covering spaces of $X_0(\mathbf{C})$. In particular, there is a covering map $\mathcal{F}_0 \rightarrow \mathcal{N}_0$ with Galois group $P\Gamma_\theta$.*

Proof. Indeed, \mathcal{F}_0 is connected by Corollary 3.6. The isomorphism $PT_\theta \setminus \mathcal{F}_0 \cong \mathcal{N}_0$ of covering spaces of $X_0(\mathbf{C})$ follows from the isomorphism $PT/PT_\theta \cong \mathfrak{S}_5$ as quotients of $\pi_1(X_0(\mathbf{C}), F_0)$, which was shown in Proposition 3.5. The lemma follows. \square

LEMMA 3.8. *The subvariety $\Delta := X_s - X_0$ is an irreducible normal crossings divisor in X_s .*

Proof. The irreducibility of Δ is well known; see e.g. [Voi02, Section 14.1.1]. Let (p, F) be a point on the incidence variety $\mathcal{I} = \{(p, F) \in \mathbf{P}^1 \times X_s \mid p \in \text{Sing}(F)\}$, choose a hyperplane away from p , and view $p \in \mathbf{C}$ and F as a polynomial $f(x)$. Since $f''(p)$ is non-zero, the tangent map

$$Tq: T_{(p,f)}(\mathcal{I}) \rightarrow T_f(X_s) = X(\mathbf{C})$$

of the projection $q: \mathcal{I} \rightarrow X_s$ is injective, and its image consists of the linear subspace $X_p \subset X$ of binary quintics that contain p as a root. If f has k double points p_j , where $k \in \{1, 2\}$, then Δ is locally isomorphic to the union $\cup_{j=1}^k X_{p_j}$. These X_{p_j} intersect transversally, and we are done. \square

DEFINITION 3.9. A node $p \in Z_F(\mathbf{C}) \subset \mathbf{P}^1(\mathbf{C})$ of a binary quintic F is a double point, i.e. a zero with multiplicity two. For $k = 1, 2$, let $\Delta_k \subset \Delta = X_s - X_0$ be the locus of stable binary quintics with exactly k nodes.

The following result is due to Deligne and Mostow [DM86].

LEMMA 3.10. *The local monodromy transformations of $\mathcal{F}_0 \rightarrow X_0(\mathbf{C})$ around every $F \in \Delta$ are of finite order. More precisely, if $F \in \Delta_k$, then the local monodromy group around F is isomorphic to $(\mathbf{Z}/10)^k$.*

Proof. Let $F_1 \in \Delta_1$ be a binary quintic with one node. Consider the universal family $\mathcal{C} \rightarrow X_s$ of quintic covers of \mathbf{P}^1 ramified along a stable binary quintic, whose restriction to X_0 is (11). Let $D \subset X_s(\mathbf{C})$ be an open disc transverse to Δ_1 at F_1 . For $F \in D^*$, one obtains a monodromy transformation $T: H^1(C_F(\mathbf{C}), \mathbf{Z}) \rightarrow H^1(C_F(\mathbf{C}), \mathbf{Z})$ induced by a vanishing cycle, and T has order ten by [DM86, Proposition 9.2]. Similarly, if $F_2 \in \Delta_2$ has two nodes, we may choose an embedding $D^2 \subset X_s(\mathbf{C})$ of the polydisc $D^2 = D \times D$ transversal to Δ_2 at F_2 . Since distinct nodes have orthogonal vanishing cycles, the local monodromy transformations commute. \square

In the following corollary, we let $D = \{z \in \mathbf{C} : |z| < 1\}$ denote the open unit disc, and $D^* = D - \{0\}$ the punctured open unit disc.

COROLLARY 3.11. *There is an essentially unique branched cover*

$$\pi: \mathcal{F}_s \rightarrow X_s(\mathbf{C}),$$

with \mathcal{F}_s a complex manifold, such that for any $x \in \Delta$, any open $x \in U \subset X_s(\mathbf{C})$ with

$$U \cong D^k \times D^{6-k} \quad \text{and} \quad U \cap X_0(\mathbf{C}) \cong (D^*)^k \times D^{6-k},$$

and any component U' of $\pi^{-1}(U) \subset \mathcal{F}_s$, there is an isomorphism $U' \cong D^k \times D^{6-k}$ such that the composition

$$D^k \times D^{6-k} \cong U' \rightarrow U \cong D^6 \text{ is the map } (z_1, \dots, z_6) \mapsto (z_1^{10}, \dots, z_k^{10}, z_{k+1}, \dots, z_6).$$

Proof. In view of [Bea09, Lemma 7.2] (see also [Fox57] and [DM86, Section 8.1]), this follows from Lemma 3.10. \square

The group $G(\mathbf{C}) = \text{GL}_2(\mathbf{C})/D$ (see (10)) acts on \mathcal{F}_0 over its action on X_0 . Explicitly, if $g \in G(\mathbf{C})$ and if $([\phi], \phi: \mathbf{V}_F \xrightarrow{\sim} \Lambda)$ is a framing of $F \in X_0(\mathbf{C})$, then

$$([\phi \circ g^*], \phi \circ g^*: \mathbf{V}_{g \cdot F} \xrightarrow{\sim} \Lambda)$$

is a framing of $g \cdot F \in X_0(\mathbf{C})$. This is a left action. The group $P\Gamma$ also acts on \mathcal{F}_0 from the left, and the actions of $P\Gamma$ and $G(\mathbf{C})$ on \mathcal{F}_0 commute. By functoriality of the Fox completion, the action of $G(\mathbf{C})$ on \mathcal{F}_0 extends to an action of $G(\mathbf{C})$ on \mathcal{F}_s .

LEMMA 3.12. *The group $G(\mathbf{C}) = \mathrm{GL}_2(\mathbf{C})/D$ acts freely on \mathcal{F}_s .*

Proof. Consider the natural action of $G(\mathbf{C})$ on \mathcal{N}_s , and the action of $G(\mathbf{C})$ on $P\Gamma_\theta \setminus \mathcal{F}_s$ induced by the action of $G(\mathbf{C})$ on \mathcal{F}_s . With respect to these actions, the isomorphism of ramified covering spaces $P\Gamma_\theta \setminus \mathcal{F}_s \cong \mathcal{N}_s$ of $X_s(\mathbf{C})$ that results from Lemma 3.7(ii) is $G(\mathbf{C})$ -equivariant. In particular, the natural ramified covering map $\mathcal{F}_s \rightarrow \mathcal{N}_s$ is $G(\mathbf{C})$ -equivariant, and so it suffices to show that $G(\mathbf{C})$ acts freely on \mathcal{N}_s .

To this end, note that \mathcal{N}_s admits a natural \mathbf{C}^* -quotient map

$$\mathcal{N}_s \rightarrow P_s, \quad (21)$$

where $P_s \subset \mathbf{P}^1(\mathbf{C})^5$ is the space of stable ordered five-tuples in $\mathbf{P}^1(\mathbf{C})$ introduced in Section 1, and where \mathbf{C}^* acts on \mathcal{N}_s by $\lambda \cdot (v_1, v_2, \dots, v_5) = (\lambda v_1, v_2, \dots, v_5) = (v_1, \lambda v_2, \dots, v_5) = \dots = (v_1, v_2, \dots, \lambda v_5) \in \mathcal{N}_s$ for $\lambda \in \mathbf{C}^*$ and $(v_1, v_2, \dots, v_5) \in \mathcal{N}_s$ (see the description of \mathcal{N}_s in Section 3.5), and (21) is equivariant for the natural homomorphism $G(\mathbf{C}) \rightarrow \mathrm{PGL}_2(\mathbf{C})$. Let $g \in \mathrm{GL}_2(\mathbf{C})$ and $x \in \mathcal{N}_s$ such that $gx = x$. Since any element of $\mathrm{PGL}_2(\mathbf{C})$ that fixes three distinct points on $\mathbf{P}^1(\mathbf{C})$ is the identity, we have that $\mathrm{PGL}_2(\mathbf{C})$ acts freely on P_s . Therefore, $g \in \mathbf{C}^* \subset \mathrm{GL}_2(\mathbf{C})$. Let $F \in X_s(\mathbf{C})$ be the image of $x \in \mathcal{N}_s$; then

$$F(x, y) = gF(x, y) = F(g^{-1}(x, y)) = F(g^{-1}x, g^{-1}y) = g^{-5}F(x, y).$$

Thus, we have $g^5 = 1 \in \mathbf{C}^*$, and we conclude that $g \in D \subset \mathrm{GL}_2(\mathbf{C})$. \square

4. Deligne–Mostow uniformization of the moduli space of complex binary quintics

In this section, we show that the results of Deligne and Mostow [DM86] yield an isomorphism of complex analytic spaces $\mathcal{M}_s(\mathbf{C}) \cong P\Gamma \setminus \mathbf{CH}^2$. This map is induced by the Riemann extension $\mathcal{P}_s: \mathcal{F}_s \rightarrow \mathbf{CH}^2$ of a holomorphic map $\mathcal{P}: \mathcal{F}_0 \rightarrow \mathbf{CH}^2$ whose definition follows rather directly from the set-up and results of the previous section. We also show that this isomorphism induces an isomorphism between the divisor $G(\mathbf{C}) \setminus \Delta(\mathbf{C}) = \mathcal{M}_s(\mathbf{C}) - \mathcal{M}_0(\mathbf{C})$ and the divisor $P\Gamma \setminus \mathcal{H} \subset P\Gamma \setminus \mathbf{CH}^2$ defined by a certain hyperplane arrangement $\mathcal{H} \subset \mathbf{CH}^2$, and prove that \mathcal{P}_s identifies binary quintics with k nodes ($k = 1, 2$) with points in \mathcal{H} where exactly k hyperplanes meet.

4.1 The period map

Define a complex hermitian vector space V as

$$V = (\Lambda \otimes_{\mathcal{O}_K, \sigma_1} \mathbf{C}, \mathfrak{h}^{\sigma_1}),$$

where \mathfrak{h}^{σ_1} is the hermitian form defined in Section 3.3. Let \mathbf{CH}^2 be the space of negative lines in V . Using [GF24, Proposition 11.7] and Lemma 3.2, we see that the abelian scheme $J \rightarrow X_0$ induces a holomorphic map, the *period map*:

$$\mathcal{P}: \mathcal{F}_0 \rightarrow \mathbf{CH}^2. \quad (22)$$

Explicitly, if $(F, [f]) \in \mathcal{F}_0$ is the framing $[f: \mathrm{H}^1(C_F(\mathbf{C}), \mathbf{Z}) \xrightarrow{\sim} \Lambda]$ of the binary quintic $F \in X_0(\mathbf{C})$, and if JC_F is the Jacobian of the curve C_F , then

$$f(\mathrm{H}^{0,-1}(JC_F)_\zeta) = f(\mathrm{H}^{1,0}(C_F)_\zeta) \subset \mathrm{H}^1(C(\mathbf{C}), \mathbf{C})_\zeta = \Lambda \otimes_{\mathcal{O}_K, \sigma_1} \mathbf{C} = V$$

is a negative line in V , and we have $\mathcal{P}(F, [f]) = f(H^{1,0}(C_F)_\zeta) \in \mathbf{CH}^2$. The map \mathcal{P} is holomorphic, and descends to a morphism of complex analytic spaces

$$\mathcal{M}_0(\mathbf{C}) = G(\mathbf{C}) \setminus X_0(\mathbf{C}) \rightarrow P\Gamma \setminus \mathbf{CH}^2.$$

Moreover, by Riemann extension, (22) extends to a $G(\mathbf{C})$ -equivariant holomorphic map

$$\mathcal{P}_s: \mathcal{F}_s \rightarrow \mathbf{CH}^2. \quad (23)$$

THEOREM 4.1 (Deligne–Mostow). *The period map (23) induces an isomorphism of complex manifolds*

$$G(\mathbf{C}) \setminus \mathcal{F}_s \cong \mathbf{CH}^2. \quad (24)$$

Taking $P\Gamma$ -quotients gives an isomorphism of complex analytic spaces

$$\mathcal{M}_s(\mathbf{C}) = G(\mathbf{C}) \setminus X_s(\mathbf{C}) \cong P\Gamma \setminus \mathbf{CH}^2. \quad (25)$$

Proof. Recall that $P_0 \subset \mathbf{P}^1(\mathbf{C})^5$ is the set of $(x_1, \dots, x_5) \in \mathbf{P}^1(\mathbf{C})^5$ such that all x_i are distinct. In accordance with [DM86], define $Q = G(\mathbf{C}) \setminus \mathcal{N}_0 = \mathrm{PGL}_2(\mathbf{C}) \setminus P_0$ and $Q_{\mathrm{st}} := G(\mathbf{C}) \setminus \mathcal{N}_s = \mathrm{PGL}_2(\mathbf{C}) \setminus P_s$. Fix a base point $0 \in Q$ whose image in $\mathfrak{S}_5 \setminus Q$ coincides with the image of $F_0 \in X_0(\mathbf{C})$ under the canonical map $X_0(\mathbf{C}) \rightarrow \mathcal{M}_0(\mathbf{C}) = \mathfrak{S}_5 \setminus Q$.

By Lemma 3.7, we have that \mathcal{F}_0 is a covering space of \mathcal{N}_0 , with Galois group $P\Gamma_\theta$. In [DM86], Deligne and Mostow define $\tilde{Q} \rightarrow Q$ to be the covering space corresponding to the monodromy representation $\pi_1(Q, 0) \rightarrow P\Gamma$; since the image of this homomorphism is $P\Gamma_\theta$ (see Proposition 3.5), it follows that $G(\mathbf{C}) \setminus \mathcal{F}_0 \cong \tilde{Q}$ as covering spaces of Q . Consequently, if $\tilde{Q}_{\mathrm{st}} \rightarrow Q_{\mathrm{st}}$ denotes the Fox completion (cf. [Fox57], [DM86, Section 8.1]) of the spread

$$\tilde{Q} \rightarrow Q \hookrightarrow Q_{\mathrm{st}},$$

then there is an isomorphism $G(\mathbf{C}) \setminus \mathcal{F}_s \cong \tilde{Q}_{\mathrm{st}}$ of branched covering spaces of Q_{st} . We obtain the following commutative diagram, in which the horizontal arrows on the left are isomorphisms.

$$\begin{array}{ccccc} G(\mathbf{C}) \setminus \mathcal{F}_s & \xrightarrow{\sim} & \tilde{Q}_{\mathrm{st}} & \longrightarrow & \mathbf{CH}^2 \\ \downarrow & & \downarrow & & \downarrow \\ G(\mathbf{C}) \setminus \mathcal{N}_s & \xrightarrow{\sim} & Q_{\mathrm{st}} & \longrightarrow & P\Gamma_\theta \setminus \mathbf{CH}^2 \\ \downarrow & & \downarrow & & \downarrow \\ G(\mathbf{C}) \setminus X_s(\mathbf{C}) & \xrightarrow{\sim} & \mathfrak{S}_5 \setminus Q_{\mathrm{st}} & \longrightarrow & P\Gamma \setminus \mathbf{CH}^2 \end{array}$$

The map $\tilde{Q}_{\mathrm{st}} \rightarrow \mathbf{CH}^2$ is an isomorphism by [DM86, (3.11)]. It follows that $Q_{\mathrm{st}} \rightarrow P\Gamma_\theta \setminus \mathbf{CH}^2$ and $\mathfrak{S}_5 \setminus Q_{\mathrm{st}} \rightarrow P\Gamma \setminus \mathbf{CH}^2$ are isomorphisms as well. Therefore, we are done if we can show that the composition $\mathcal{F}_0 \rightarrow \tilde{Q} \rightarrow \mathbf{CH}^2$ agrees with the period map \mathcal{P} of equation (22). This follows from [DM86, (2.23) and (12.9)]. \square

4.2 Nodal binary quintics and orthogonal hyperplanes

Consider the CM type $\Phi = \{\sigma_1, \sigma_2\} \subset \mathrm{Hom}(K, \mathbf{C})$ defined in (9), the hermitian \mathcal{O}_K -lattice (Λ, \mathfrak{h}) defined in (8), and the following sets (cf. [GF24, Sections 2.2 and 2.3]):

$$\mathcal{H} = \{H_r \subset \mathbf{CH}^2 \mid r \in \mathcal{R}\} \quad \text{and} \quad \mathcal{H} = \bigcup_{H \in \mathcal{H}} H \subset \mathbf{CH}^2. \quad (26)$$

Here, $\mathcal{R} \subset \Lambda$ is the set of short roots, i.e. the set of $r \in \Lambda$ with $\mathfrak{h}(r, r) = 1$, and for each $r \in \mathcal{R}$, $H_r \subset \mathbf{CH}^2$ is the hyperplane of elements $x \in \mathbf{CH}^2$ that are orthogonal to r .

PROPOSITION 4.2. *The hyperplane arrangement $\mathcal{H} \subset \mathbf{CH}^2$ is an orthogonal arrangement in the sense of [ACT02a]. In other words, any two different hyperplanes $H_1, H_2 \in \mathcal{H}$ either meet orthogonally, or not at all.*

Proof. By Lemma 3.2, we have that \mathfrak{h} has signature $(2, 1)$ with respect to the embedding $\sigma_1: K \hookrightarrow \mathbf{C}$, and signature $(3, 0)$ with respect to the embedding $\sigma_2: K \hookrightarrow \mathbf{C}$, where σ_1 and σ_2 are defined in (9). Therefore, the result follows from [GF24, Theorem 2.5] and [GF24, Example 2.12]. \square

PROPOSITION 4.3. *The map (25) induces an isomorphism of complex analytic spaces*

$$\mathcal{M}_0(\mathbf{C}) = G(\mathbf{C}) \setminus X_0(\mathbf{C}) \cong P\Gamma \setminus (\mathbf{CH}^2 - \mathcal{H}).$$

Proof. We have $\mathcal{P}_s(\mathcal{F}_0) \subset \mathbf{CH}^2 - \mathcal{H}$ by [GF24, Proposition 11.12], because the Jacobian of a smooth curve cannot contain a non-trivial abelian subvariety whose induced polarization is principal. Therefore, we have $\mathcal{P}_s^{-1}(\mathcal{H}) \subset \mathcal{F}_s - \mathcal{F}_0$. Since \mathcal{F}_s is irreducible (it is smooth by Corollary 3.11 and connected by Lemma 3.7(i)), the analytic space $\mathcal{P}_s^{-1}(\mathcal{H})$ is a divisor. Since $\mathcal{F}_s - \mathcal{F}_0$ is also a divisor by Corollary 3.11, we have $\mathcal{P}_s^{-1}(\mathcal{H}) = \mathcal{F}_s - \mathcal{F}_0$. \square

Define $\tilde{\Delta} = \mathcal{F}_s - \mathcal{F}_0$, and for $k \in \{1, 2\}$, let $\tilde{\Delta}_k \subset \tilde{\Delta}$ be the inverse image of Δ_k in $\tilde{\Delta}$ under the map $\tilde{\Delta} \rightarrow \Delta$. Here, $\Delta_k \subset \Delta$ is the subvariety defined in Definition 3.9. Moreover, for $k = 1, 2$, define $\mathcal{H}_k \subset \mathcal{H}$ as the set

$$\mathcal{H}_k := \{x \in \mathbf{CH}^2: |\mathcal{H}(x)| = k\}.$$

Thus, this is the locus of points in \mathcal{H} where exactly k hyperplanes meet. For $r \in \mathcal{R}$, define an isometry

$$\phi_r: \Lambda \rightarrow \Lambda, \quad \phi_r(x) = x - (1 - \zeta)\mathfrak{h}(x, r) \cdot r.$$

Let $[\phi_r] \in P\Gamma$ be the image of $\phi_r \in \Gamma$ in the group $P\Gamma = \Gamma/\mu_K$, and for $x \in \mathbf{CH}^2$, define

$$G(x) := \langle [\phi_r] \mid r \in \mathcal{R} \mid \mathfrak{h}(x, r) = 0 \rangle \subset P\Gamma. \quad (27)$$

LEMMA 4.4.

- (i) *The period map \mathcal{P}_s of (23) satisfies $\mathcal{P}_s(\tilde{\Delta}_k) \subset \mathcal{H}_k$.*
- (ii) *Let $\tilde{F} \in \tilde{\Delta}_k \subset \mathcal{F}_s$ and $x = \mathcal{P}_s(\tilde{F}) \in \mathcal{H}_k \subset \mathbf{CH}^2$. Let $PG_{\tilde{F}} \subset P\Gamma$ be the stabilizer of \tilde{F} in $P\Gamma$. Then $PG_{\tilde{F}} = G(x)$, where $G(x) \cong (\mathbf{Z}/10)^k$ is as in (27).*

Proof.

- (i) We know that \mathcal{P}_s induces an isomorphism $G(\mathbf{C}) \setminus \tilde{\Delta} \xrightarrow{\sim} \mathcal{H}$ by Theorem 4.1 and Proposition 4.3. This map must identify the smooth (respectively singular) locus of one analytic variety with the smooth (respectively singular) locus of the other, from which the result follows.
- (ii) This follows from Lemma 3.10 and Corollary 3.11 (compare [ACT02b, (3.10)] and [ACT10, Lemma 10.3]). \square

5. The moduli space of real binary quintics

With the period map for complex binary quintics in place, we turn to the construction of the period map for real binary quintics. Define κ as the anti-holomorphic involution

$$\kappa: X_0(\mathbf{C}) \rightarrow X_0(\mathbf{C}), \quad F(x, y) = \sum_{i+j=5} a_{ij}x^i y^j \mapsto \overline{F(x, y)} = \sum_{i+j=5} \overline{a_{ij}}x^i y^j.$$

DEFINITION 5.1. (Compare [GF24, Sections 3.1 and 3.2].)

- (i) An \mathcal{O}_F -linear bijection $\phi: \Lambda \xrightarrow{\sim} \Lambda$ is called anti-unitary if $\phi(\mu \cdot x) = \bar{\mu} \cdot \phi(x)$ and $\mathfrak{h}(\phi(x), \phi(y)) = \overline{\mathfrak{h}(x, y)}$ for $x, y \in \Lambda$, $\mu \in \mathcal{O}_K$.
- (ii) Let Γ' be the group of unitary and anti-unitary \mathcal{O}_F -linear bijections of Λ . Let $P\Gamma' = \Gamma' / \mu_K$.
- (iii) Let \mathcal{A} be the set of anti-unitary involutions $\alpha: \Lambda \rightarrow \Lambda$, and define $P\mathcal{A} = \mu_K \backslash \mathcal{A}$.
- (iv) For $\alpha \in P\mathcal{A}$, define $\mathbf{R}H_\alpha^2 \subset \mathbf{C}H^2$ as the fixed space of α , i.e. $\mathbf{R}H_\alpha^2 = (\mathbf{C}H^2)^\alpha$.
- (v) For $\alpha \in P\mathcal{A}$, let $P\Gamma_\alpha \subset P\Gamma$ be the stabilizer of the subspace $\mathbf{R}H_\alpha^2 = (\mathbf{C}H^2)^\alpha \subset \mathbf{C}H^2$.

For $\alpha \in P\mathcal{A}$, the notation $\mathbf{R}H_\alpha^2$ reflects the fact that $\mathbf{R}H_\alpha^2$ is isometric to the real hyperbolic plane $\mathbf{R}H^2$; see [GF24, Lemma 3.4].

For each $\alpha \in P\mathcal{A}$, there is a natural anti-holomorphic involution $\alpha: \mathcal{F}_0 \rightarrow \mathcal{F}_0$ lying over the anti-holomorphic involution $\kappa: X_0(\mathbf{C}) \rightarrow X_0(\mathbf{C})$. To define α , consider a framed binary quintic $(F, [f]) \in \mathcal{F}_0$, where $f: \mathbf{V}_F \rightarrow \Lambda$ is an \mathcal{O}_K -linear isometry. Let $C_F \rightarrow \mathbf{P}_{\mathbf{C}}^1$ be the induced quintic cover of $\mathbf{P}_{\mathbf{C}}^1$. Complex conjugation $\mathbf{P}^1(\mathbf{C}) \rightarrow \mathbf{P}^1(\mathbf{C})$ induces a bijection $Z_F(\mathbf{C}) \xrightarrow{\sim} Z_{\kappa(F)}(\mathbf{C})$, and extends to an anti-holomorphic diffeomorphism $\sigma_F: C_F(\mathbf{C}) \rightarrow C_{\kappa(F)}(\mathbf{C})$ with pull-back $\sigma_F^*: \mathbf{V}_{\kappa(F)} \rightarrow \mathbf{V}_F$. The composition $\alpha \circ f \circ \sigma_F^*: \mathbf{V}_{\kappa(F)} \rightarrow \Lambda$ induces a framing of $\kappa(F) \in X_0(\mathbf{C})$, and we define

$$\alpha(F, [f]) := (\kappa(F), [\alpha \circ f \circ \sigma_F^*]) \in \mathcal{F}_0.$$

Although we have chosen a representative $\alpha \in \mathcal{A}$ of the class $\alpha \in P\mathcal{A}$, the element $\alpha(F, [f]) \in \mathcal{F}_0$ does not depend on this choice.

Consider the covering map $\mathcal{F}_0 \rightarrow X_0(\mathbf{C})$ defined in (20) in Lemma 3.7, and define $\mathcal{F}_0(\mathbf{R})$ as the preimage of $X_0(\mathbf{R})$ in the space \mathcal{F}_0 . Then

$$\mathcal{F}_0(\mathbf{R}) = \coprod_{\alpha \in P\mathcal{A}} \mathcal{F}_0^\alpha \subset \mathcal{F}_0. \quad (28)$$

To see why the union in (28) is disjoint, let $\alpha \in P\mathcal{A}$; then

$$\mathcal{F}_0^\alpha = \{(F, [f]) \in \mathcal{F}_0 : \kappa(F) = F \text{ and } [f \circ \sigma_F^* \circ f^{-1}] = \alpha\}.$$

Thus, for $\alpha, \beta \in P\mathcal{A}$ and $(F, [f]) \in \mathcal{F}_0^\alpha \cap \mathcal{F}_0^\beta$, we have $\alpha = [f \circ \sigma_F^* \circ f^{-1}] = \beta$.

LEMMA 5.2. Let $\alpha \in P\mathcal{A}$. The anti-holomorphic involution $\alpha: \mathcal{F}_0 \rightarrow \mathcal{F}_0$ commutes with the period map $\mathcal{P}: \mathcal{F}_0 \rightarrow \mathbf{C}H^2$ and the anti-holomorphic involution $\alpha: \mathbf{C}H^2 \rightarrow \mathbf{C}H^2$.

Proof. If $\text{conj}: \mathbf{C} \rightarrow \mathbf{C}$ is complex conjugation, then for any $F \in X_0(\mathbf{C})$, the induced map $\sigma_F^* \otimes \text{conj}: \mathbf{V}_{\kappa(F)} \otimes_{\mathbf{Z}} \mathbf{C} \rightarrow \mathbf{V}_F \otimes_{\mathbf{Z}} \mathbf{C}$ is anti-linear and preserves the Hodge decompositions [Sil89, Chapter I, Lemma 2.4] as well as the eigenspace decompositions. \square

By Lemma 5.2, we obtain a real period map

$$\mathcal{P}^{\mathbf{R}} : \mathcal{F}_0(\mathbf{R}) = \coprod_{\alpha \in P\mathcal{A}} \mathcal{F}_0^\alpha \longrightarrow \coprod_{\alpha \in P\mathcal{A}} \mathbf{R}H_\alpha^2 =: \widetilde{Y}. \quad (29)$$

Let $\sigma: \text{GL}_2(\mathbf{C}) \rightarrow \text{GL}_2(\mathbf{C})$ be the anti-holomorphic involution that sends a matrix to its complex conjugate, and note that σ descends to an anti-holomorphic involution $\sigma: G(\mathbf{C}) \rightarrow G(\mathbf{C})$; define

$$G(\mathbf{R}) := G(\mathbf{C})^\sigma.$$

LEMMA 5.3. *The natural map $\mathrm{GL}_2(\mathbf{R}) \rightarrow G(\mathbf{R})$ is an isomorphism.*

Proof. Indeed, if $M \in \mathrm{GL}_2(\mathbf{C})$ is a matrix such that $\sigma(M) = \zeta^j \cdot M$ for some $j \in \{0, \dots, 4\}$, then we can write $\zeta^j = \zeta^{2k}$ for some $k \in \{0, \dots, 4\}$, and hence $\sigma(\zeta^k \cdot M) = \zeta^{-k} \zeta^j \cdot M = \zeta^k \cdot M$, so that $\zeta^k \cdot M \in \mathrm{GL}_2(\mathbf{R})$. This proves that $\mathrm{GL}_2(\mathbf{R}) \rightarrow G(\mathbf{R})$ is surjective. Injectivity follows from the fact that the kernel of $\mathrm{GL}_2(\mathbf{R}) \rightarrow G(\mathbf{R})$ is spanned by the elements of $D = \{1, \zeta, \dots, \zeta^4\}$ that are invariant under complex conjugation; the only such element is $1 \in D$. \square

The map (29) is constant on $G(\mathbf{R})$ -orbits, as the same is true for $\mathcal{P}: \mathcal{F}_0 \rightarrow \mathbf{CH}^2$. By abuse of notation, we write $\mathbf{R}H_\alpha^2 - \mathcal{H} = \mathbf{R}H_\alpha^2 - (\mathcal{H} \cap \mathbf{R}H_\alpha^2)$ for $\alpha \in P\mathcal{A}$.

PROPOSITION 5.4. *The period map (29) descends to a PT -equivariant diffeomorphism*

$$\mathcal{P}^{\mathbf{R}}: \mathcal{M}_0(\mathbf{R})^f := G(\mathbf{R}) \setminus \mathcal{F}_0(\mathbf{R}) \cong \coprod_{\alpha \in P\mathcal{A}} (\mathbf{R}H_\alpha^2 - \mathcal{H}). \quad (30)$$

Let $C\mathcal{A} \subset P\mathcal{A}$ be a set of representatives for the action of PT on $P\mathcal{A}$. By PT -equivariance, the map (30) induces an isomorphism of real-analytic orbifolds

$$\mathcal{P}^{\mathbf{R}}: \mathcal{M}_0(\mathbf{R}) = G(\mathbf{R}) \setminus X_0(\mathbf{R}) \cong \coprod_{\alpha \in C\mathcal{A}} PT_\alpha \setminus (\mathbf{R}H_\alpha^2 - \mathcal{H}). \quad (31)$$

Proof. We follow the proof of [ACT10, Theorem 3.3]. The first thing to observe is that the map

$$\mathcal{P}^{\mathbf{R}}: G(\mathbf{R}) \setminus \mathcal{F}_0(\mathbf{R}) \rightarrow \coprod_{\alpha \in P\mathcal{A}} (\mathbf{R}H_\alpha^2 - \mathcal{H})$$

is a local diffeomorphism, as the same holds for $\mathcal{P}: G(\mathbf{C}) \setminus \mathcal{F}_0 \rightarrow \mathbf{CH}^2 - \mathcal{H}$ by Theorem 4.1. To prove the surjectivity and injectivity of $\mathcal{P}^{\mathbf{R}}$, notice that the arguments used in [ACT10] to prove the analogous claims for cubic surfaces readily carry over to our situation. \square

Our next goal is to prove the real analogue of Theorem 4.1. By the naturality of the Fox completion, for every $\alpha \in P\mathcal{A}$ the involution $\alpha: \mathcal{F}_0 \rightarrow \mathcal{F}_0$ extends to an involution α on \mathcal{F}_s .

LEMMA 5.5. *The restriction of the map $\mathcal{P}_s: \mathcal{F}_s \rightarrow \mathbf{CH}^2$ to \mathcal{F}_s^α induces a diffeomorphism*

$$\mathcal{P}_s^\alpha: G(\mathbf{R}) \setminus \mathcal{F}_s^\alpha \cong \mathbf{R}H_\alpha^2.$$

Proof. The map $\mathcal{P}_s^\alpha: G(\mathbf{R}) \setminus \mathcal{F}_s^\alpha \rightarrow \mathbf{R}H_\alpha^2$ is a local diffeomorphism because its differential at any point is an isomorphism by Theorem 4.1. Let us prove that \mathcal{P}_s^α is injective. Apply [ACT10, Lemma 3.5] with $X = \mathcal{F}_0$, $G = G(\mathbf{C})$ and $\phi = \alpha$; then $X^\phi = \mathcal{F}_s^\alpha$ and $Z = G(\mathbf{R})$. Note that we may apply this lemma because $G(\mathbf{C})$ acts freely on \mathcal{F}_s (see Lemma 3.12). The conclusion is that the map

$$G(\mathbf{R}) \setminus \mathcal{F}_0^\alpha \rightarrow G(\mathbf{C}) \setminus \mathcal{F}_0 = \mathbf{CH}^2 - \mathcal{H}$$

is injective. To prove the surjectivity of \mathcal{P}_s^α , one uses [ACT10, Lemma 11.2] to see that the map $G(\mathbf{R}) \setminus \mathcal{F}_s^\alpha \rightarrow G(\mathbf{C}) \setminus \mathbf{CH}^2$ is proper. By Proposition 5.4, its image contains the dense open subset $\mathcal{P}_s(\mathcal{F}_0^\alpha) = \mathbf{R}H_\alpha^2 - \mathcal{H}$, so \mathcal{P}_s^α is surjective. \square

DEFINITION 5.6. (See [GF24, Definition 1.1].) Define an equivalence relation \sim on the disjoint union $\coprod_{\alpha \in P\mathcal{A}} \mathbf{R}H_\alpha^2$ in the following way. Consider two points $(x, \beta) \in \mathbf{R}H_\beta^2 \subset \coprod_{\alpha \in P\mathcal{A}} \mathbf{R}H_\alpha^2$ and $(y, \gamma) \in \mathbf{R}H_\gamma^2 \subset \coprod_{\alpha \in P\mathcal{A}} \mathbf{R}H_\alpha^2$. Then $(x, \beta) \sim (y, \gamma)$ if $x = y \in \mathbf{CH}^2$ and $\gamma \circ \beta \in G(x)$.

By [GF24, Lemma 4.4], the action of $P\Gamma$ on $\coprod_{\alpha \in P\mathcal{A}} \mathbf{R}H_\alpha^2$ is compatible with this equivalence relation; define

$$Y := \left(\coprod_{\alpha \in P\mathcal{A}} \mathbf{R}H_\alpha^2 \right) / \sim, \quad M := P\Gamma \backslash Y.$$

THEOREM 5.7. *Let $C\mathcal{A} \subset P\mathcal{A}$ be a set of representatives for the action of $P\Gamma$ on $P\mathcal{A}$. There exists a canonical real hyperbolic orbifold structure on the topological space $M = P\Gamma \backslash Y$ together with a natural open immersion of hyperbolic orbifolds*

$$\coprod_{\alpha \in C\mathcal{A}} P\Gamma_\alpha \backslash (\mathbf{R}H_\alpha^2 - \mathcal{H}) \hookrightarrow M.$$

Moreover, for each connected component $M_i \subset M$ there exists a lattice $P\Gamma_{\mathbf{R}}^i \subset \mathrm{PO}(2, 1)$ and an isomorphism of real hyperbolic orbifolds $M_i \cong P\Gamma_{\mathbf{R}}^i \backslash \mathbf{R}H^2$.

Proof. By Lemma 3.2 and [GF24, Example 2.12], this is a special case of [GF24, Theorem 1.2]. \square

Let $p: \coprod_{\alpha \in P\mathcal{A}} \mathbf{R}H_\alpha^2 \rightarrow Y$ be the quotient map, consider the map $\pi: \mathcal{F}_s \rightarrow X_s(\mathbf{C})$ (see Corollary 3.11), and define a union of embedded real submanifolds of \mathcal{F}_s as follows:

$$\mathcal{F}_s(\mathbf{R}) = \bigcup_{\alpha \in P\mathcal{A}} \mathcal{F}_s^\alpha = \pi^{-1}(X_s(\mathbf{R})).$$

We arrive at the main theorem of Section 5.

THEOREM 5.8. *There is a smooth map*

$$\mathcal{P}_s^{\mathbf{R}}: \coprod_{\alpha \in P\mathcal{A}} \mathcal{F}_s^\alpha \rightarrow \coprod_{\alpha \in P\mathcal{A}} \mathbf{R}H_\alpha^2 \quad (32)$$

that extends the real period map (29). The map (32) induces the following commutative diagram of topological spaces, in which $\mathcal{P}_s^{\mathbf{R}}$ and $\mathcal{T}_s^{\mathbf{R}}$ are homeomorphisms.

$$\begin{array}{ccccc} \coprod_{\alpha \in P\mathcal{A}} \mathcal{F}_s^\alpha & \xrightarrow{\mathcal{P}_s^{\mathbf{R}}} & \coprod_{\alpha \in P\mathcal{A}} \mathbf{R}H_\alpha^2 & & \\ \downarrow & & \downarrow p & & \\ \mathcal{F}_s(\mathbf{R}) & \xrightarrow{\mathcal{P}_s^{\mathbf{R}}} & Y & & \\ \downarrow & & \downarrow \parallel & & \\ \mathcal{M}_s(\mathbf{R})^f & \xlongequal{\quad} & G(\mathbf{R}) \backslash \mathcal{F}_s(\mathbf{R}) & \xrightarrow[\sim]{\mathcal{P}_s^{\mathbf{R}}} & Y \\ \downarrow & & \downarrow & & \downarrow \\ \mathcal{M}_s(\mathbf{R}) & \xlongequal{\quad} & G(\mathbf{R}) \backslash X_s(\mathbf{R}) & \xrightarrow[\sim]{\mathcal{T}_s^{\mathbf{R}}} & P\Gamma \backslash Y \end{array}$$

Proof. The existence of $\mathcal{P}_s^{\mathbf{R}}$ follows from the compatibility between \mathcal{P}_s and the involutions $\alpha \in P\mathcal{A}$. We claim that the composition $p \circ \mathcal{P}_s^{\mathbf{R}}$ factors through $\mathcal{F}_s(\mathbf{R})$. To prove this, let (f, α) and (g, β) be elements of the disjoint union $\coprod_{\gamma \in P\mathcal{A}} \mathcal{F}_s^\gamma$, with $f \in \mathbf{R}H_\alpha^2$ and $g \in \mathbf{R}H_\beta^2$. Then (f, α) and (g, β) have the same image in $\mathcal{F}_s(\mathbf{R})$ if and only if $f = g \in \mathcal{F}_s^\alpha \cap \mathcal{F}_s^\beta$, in which case $\mathcal{P}_s(f) = \mathcal{P}_s(g) =: x \in \mathbf{R}H_\alpha^2 \cap \mathbf{R}H_\beta^2$. Let (x, α) and (y, β) be elements of the disjoint union $\tilde{Y} =$

$\coprod_{\gamma \in P\mathcal{A}} \mathbf{R}H_\gamma^2$, with $x \in \mathbf{R}H_\alpha^2$ and $y = x \in \mathbf{R}H_\beta^2$. We need to prove that $(x, \alpha) \sim (x, \beta) \in \tilde{Y}$, for the equivalence relation \sim on \tilde{Y} defined in Definition 5.6. Note that $\alpha\beta \in P\Gamma_f$, and that \mathcal{P}_s induces an isomorphism $P\Gamma_f \cong G(x)$ (see Lemma 4.4). Hence $\alpha\beta \in G(x)$ so that $(x, \alpha) \sim (x, \beta)$, proving what we want. We conclude that the composition $p \circ \mathcal{P}_s^{\mathbf{R}}$ factors through a map $\mathcal{P}_s^{\mathbf{R}}: \mathcal{F}_s(\mathbf{R}) \rightarrow Y$.

Next, we prove the $G(\mathbf{R})$ -equivariance of $\mathcal{P}_s^{\mathbf{R}}$. Suppose that $f \in \mathcal{F}_s^\alpha, g \in \mathcal{F}_s^\beta$ such that $a \cdot f = g \in \mathcal{F}_s(\mathbf{R})$ for some $a \in G(\mathbf{R})$. Then $x := \mathcal{P}_s(f) = \mathcal{P}_s(g) \in \mathbf{C}H^2$, so we need to show that $\alpha\beta \in G(x)$. The actions of $G(\mathbf{C})$ and $P\Gamma$ on $\mathbf{C}H^2$ commute, and the same holds for the actions of $G(\mathbf{R})$ and $P\Gamma'$ on $\mathcal{F}_s^{\mathbf{R}}$, where $P\Gamma'$ is the group defined in Definition 5.1. It follows that $\alpha(g) = \alpha(a \cdot f) = a \cdot \alpha(f) = a \cdot f = g$, and hence $g \in \mathcal{F}_s^\alpha \cap \mathcal{F}_s^\beta$. This implies in turn that $(\alpha \circ \beta)(g) = g$, and hence $\alpha\beta \in P\Gamma_g \cong G(x)$ (see Lemma 4.4). Therefore $(x, \alpha) \sim (x, \beta)$, so that $\alpha\beta \in G(x)$ as desired.

Let us prove that $\mathcal{P}_s^{\mathbf{R}}$ is injective. To do so, let $f \in \mathcal{F}_s^\alpha$ and $g \in \mathcal{F}_s^\beta$ and suppose that these elements have the same image in Y . Thus, $x := \mathcal{P}_s(f) = \mathcal{P}_s(g) \in \mathbf{R}H_\alpha^2 \cap \mathbf{R}H_\beta^2$, and $\beta = \phi \circ \alpha$ for some $\phi \in G(x)$. We have $\phi \in G(x) \cong P\Gamma_f$ (Lemma 4.4), and hence $\beta(f) = \phi(\alpha(f)) = \phi(f) = f$. Therefore $f, g \in \mathcal{F}_s^\beta$; since $\mathcal{P}_s(f) = \mathcal{P}_s(g)$, it follows from Lemma 5.5 that there exists $a \in G(\mathbf{R})$ such that $a \cdot f = g$. This proves the injectivity of $\mathcal{P}_s^{\mathbf{R}}$. The surjectivity of $\mathcal{P}_s^{\mathbf{R}}: G(\mathbf{R}) \setminus \mathcal{F}_s(\mathbf{R}) \rightarrow Y$ is straightforward: it follows from the surjectivity of $\mathcal{P}_s^{\mathbf{R}}$ (see Lemma 5.5). Finally, we claim that $\mathcal{P}_s^{\mathbf{R}}$ is open. Let $U \subset G(\mathbf{R}) \setminus \mathcal{F}_s^{\mathbf{R}}$ be open. Let V be the preimage of U in $\coprod_{\alpha \in P\mathcal{A}} \mathcal{F}_s^\alpha$. Then $V = (\mathcal{P}_s^{\mathbf{R}})^{-1}(p^{-1}(\mathcal{P}_s^{\mathbf{R}}(U)))$, hence $\mathcal{P}_s^{\mathbf{R}}(V) = p^{-1}(\mathcal{P}_s^{\mathbf{R}}(U))$, so that it suffices to show that $\mathcal{P}_s^{\mathbf{R}}(V)$ is an open subset of $\coprod_{\alpha \in P\mathcal{A}} \mathbf{R}H_\alpha^2$. This follows, because $\mathcal{P}_s^{\mathbf{R}}$ is open, being the coproduct of the maps $\mathcal{F}_s^\alpha \rightarrow \mathbf{R}H_\alpha^2$, which are open since they have surjective differential at each point. \square

COROLLARY 5.9. *Let $C\mathcal{A} \subset P\mathcal{A}$ be a set of representatives for the action of $P\Gamma$ on $P\mathcal{A}$. Then there is a lattice $P\Gamma_{\mathbf{R}} \subset \mathrm{PO}(2, 1)$, an open immersion of hyperbolic orbifolds*

$$\coprod_{\alpha \in C\mathcal{A}} P\Gamma_\alpha \setminus (\mathbf{R}H_\alpha^2 - \mathcal{H}) \hookrightarrow P\Gamma_{\mathbf{R}} \setminus \mathbf{R}H^2, \quad (33)$$

and a homeomorphism

$$\mathcal{P}_s^{\mathbf{R}}: \mathcal{M}_s(\mathbf{R}) = G(\mathbf{R}) \setminus X_s(\mathbf{R}) \cong P\Gamma_{\mathbf{R}} \setminus \mathbf{R}H^2, \quad (34)$$

such that the restriction of (34) to $\mathcal{M}_0(\mathbf{R}) \subset \mathcal{M}_s(\mathbf{R})$ coincides with the isomorphism (31).

Proof. This follows directly from Theorems 5.7 and 5.8. \square

Remark 5.10. The proof of Corollary 5.9 also shows that $\mathcal{M}_s(\mathbf{R})$ is homeomorphic to a complete hyperbolic orbifold in the cases where \mathcal{M}_s is the stack of cubic surfaces or of binary sextics over \mathbf{R} . This strategy to uniformize the real moduli space differs from the one used in [ACT06, ACT07, ACT10], since we first glue the real ball quotients together (by using the general construction of [GF24]) and then prove that the real moduli space is homeomorphic to the resulting glued space.

6. The moduli space of real binary quintics as a hyperbolic triangle

Consider the moduli space $\mathcal{M}_s(\mathbf{R}) = \mathrm{GL}_2(\mathbf{R}) \setminus X_s(\mathbf{R})$ of stable real binary quintics. Let $|\mathcal{M}_s(\mathbf{R})|$ be the underlying topological space of $\mathcal{M}_s(\mathbf{R})$.

DEFINITION 6.1. Let $\overline{\mathcal{M}}_{\mathbf{R}}$ be the orbifold with $|\mathcal{M}_s(\mathbf{R})|$ as underlying space whose orbifold structure is induced by the homeomorphism (34) and the natural orbifold structure of $P\Gamma_{\mathbf{R}} \setminus \mathbf{R}H^2$.

The goal of Section 6 is to prove the following result.

THEOREM 6.2. Consider the lattice $P\Gamma_{\mathbf{R}} \subset \mathrm{PO}(2, 1)$ (see Corollary 5.9) and the hyperbolic orbifold $\overline{\mathcal{M}}_{\mathbf{R}} \cong P\Gamma_{\mathbf{R}} \setminus \mathbf{R}H^2$ (see Definition 6.1). Then $\overline{\mathcal{M}}_{\mathbf{R}}$ is isometric to the hyperbolic triangle with angles $\pi/3, \pi/5, \pi/10$. In particular, $P\Gamma_{\mathbf{R}}$ is conjugate to the lattice $\Gamma_{3,5,10}$ defined in (3).

To prove Theorem 6.2, we need to understand the orbifold structure of $\mathcal{M}_s(\mathbf{R})$ and how this structure differs from the orbifold structure of the quotient space $P\Gamma_{\mathbf{R}} \setminus \mathbf{R}H^2$ (see Corollary 5.9). To this end, we will first analyze the orbifold structure of $\mathcal{M}_s(\mathbf{R})$ by listing its stabilizer groups.

6.1 Automorphism groups of stable real binary quintics

Recall that there is a canonical orbifold isomorphism

$$\mathcal{M}_s(\mathbf{R}) = G(\mathbf{R}) \setminus X_s(\mathbf{R}) = \mathrm{PGL}_2(\mathbf{R}) \setminus (P_s/\mathfrak{S}_5)(\mathbf{R}).$$

Thus, to list those groups that occur as the automorphism groups of a binary quintic is to classify the stabilizer groups $\mathrm{PGL}_2(\mathbf{R})_x$ of points $x = (x_1, \dots, x_5) \in (P_s/\mathfrak{S}_5)(\mathbf{R})$.

PROPOSITION 6.3. Consider the stabilizer group $\mathrm{PGL}_2(\mathbf{C})_x$ of a point $x \in P_0/\mathfrak{S}_5$. If $\mathrm{PGL}_2(\mathbf{C})_x$ is non-trivial, then $\mathrm{PGL}_2(\mathbf{C})_x$ is isomorphic to one of $\mathbf{Z}/2, \mathbf{Z}/4, D_3$ or D_5 . Moreover, the conjugacy class of each such subgroup of $\mathrm{PGL}_2(\mathbf{C})$ is unique. If H equals any of the subgroups $\mathbf{Z}/4, D_3$ or D_5 of $\mathrm{PGL}_2(\mathbf{C})$, then there is a unique $\mathrm{PGL}_2(\mathbf{C})$ -orbit in P_0/\mathfrak{S}_5 with stabilizer group H .

Proof. By [Bea10, Theorem 4.2], any finite subgroup of $\mathrm{PGL}_2(\mathbf{C})$ is isomorphic to $\mathbf{Z}/n, D_n$ (the dihedral group of order $2n$), $\mathfrak{A}_4, \mathfrak{S}_4$ or \mathfrak{A}_5 , and there is only one conjugacy class for each of these groups. Let H be any of these groups, considered as a subgroup of $\mathrm{PGL}_2(\mathbf{C})$. Assume that, with respect to the action of H on the finite subsets of $\mathbf{P}^1(\mathbf{C})$, one has

$$H \cdot \{z_1, \dots, z_5\} = \{z_1, \dots, z_5\} \subset \mathbf{P}^1(\mathbf{C}) \quad \text{for } z_1, \dots, z_5 \in \mathbf{P}^1(\mathbf{C}) \text{ distinct.}$$

This gives a homomorphism $\rho: H \rightarrow \mathfrak{S}_5$ as follows: for an element $j \in \{1, 2, 3, 4, 5\}$, we let $\rho(h)(j) \in \{1, 2, 3, 4, 5\}$ be the element with $z_{\rho(h)(j)} = h \cdot z_j$.

Note that ρ is injective, as $h \cdot z_i = z_i$ for each i implies $h = \mathrm{id}$. Therefore,

$$H \in \{\mathbf{Z}/2, \mathbf{Z}/3, \mathbf{Z}/4, \mathbf{Z}/5, \mathbf{Z}/6, D_3, D_4, D_5, \mathfrak{A}_4, \mathfrak{S}_4, \mathfrak{A}_5\}.$$

Next, assume that $H = \mathrm{Stab}_{\mathrm{PGL}_2(\mathbf{C})}(x)$ for the five-element subset $x = \{z_1, \dots, z_5\} \subset \mathbf{P}^1(\mathbf{C})$. Suppose that $\phi \in H$ is an element of order three. Note that there must be three distinct elements $z_i \in x$ with $\phi(z_i) \neq z_i$. We may assume that these are z_1, z_2 and z_3 . Moreover, we may assume that $\phi(z_1) = z_2, \phi(z_2) = z_3$ and $\phi(z_3) = z_1$. By replacing ϕ by $g\phi g^{-1}$ for some $g \in \mathrm{PGL}_2(\mathbf{C})$, we may assume that $z_1 = 1, z_2 = \zeta_3$ and $z_3 = \zeta_3^2$, and that $\phi(z) = \zeta_3 \cdot z$ for $z \in \mathbf{P}^1(\mathbf{C})$. This gives $x = \{1, \zeta_3, \zeta_3^2, z_4, z_5\} \subset \mathbf{P}^1(\mathbf{C})$. As $\phi(z_4) \neq z_5$, we have $\phi(z_4) = z_4$ and $\phi(z_5) = z_5$, so that

$$x = \{1, \zeta_3, \zeta_3^2, 0, \infty\} \subset \mathbf{P}^1(\mathbf{C}). \quad (35)$$

Let $\nu \in \mathrm{PGL}_2(\mathbf{C})$ be the element with $\nu(z) = 1/z$ for $z \in \mathbf{P}^1(\mathbf{C})$. Then (35) implies that ν is contained in H , and one readily observes that $H = D_3$. We conclude that

$$H = \mathrm{Stab}_{\mathrm{PGL}_2(\mathbf{C})}(x) \in \{\mathbf{Z}/2, \mathbf{Z}/4, \mathbf{Z}/5, D_3, D_4, D_5\}.$$

It remains to exclude $\mathbf{Z}/5$ and D_4 . Suppose that H contains an element ϕ of order five. As above, one can readily show that one may assume that

$$x = \{1, \zeta_5, \zeta_5^2, \zeta_5^3, \zeta_5^4\} \subset \mathbf{P}^1(\mathbf{C}), \quad \phi(z) = \zeta_5 \cdot z \text{ for } z \in \mathbf{P}^1(\mathbf{C}).$$

This implies that the element $\nu \in \mathrm{PGL}_2(\mathbf{C})$ as defined above is contained in H , and $H = D_5$. Finally, assume H contains an element ϕ of order four. We may assume that

$$x = \{1, i, -1, -i, 0\}, \quad \phi(z) = i \cdot z \text{ for } z \in \mathbf{P}^1(\mathbf{C}).$$

As a consequence, we have $H = \mathbf{Z}/4$, and the proof is finished. \square

We proceed to prove the real analogue of Proposition 6.3.

PROPOSITION 6.4. *Let $x \in (P_s/\mathfrak{S}_5)(\mathbf{R})$ such that the stabilizer group $\mathrm{PGL}_2(\mathbf{R})_x \subset \mathrm{PGL}_2(\mathbf{R})$ of x is non-trivial. Then its stabilizer group $\mathrm{PGL}_2(\mathbf{R})_x$ is isomorphic to $\mathbf{Z}/2$, D_3 or D_5 . For each $n \in \{3, 5\}$, there is a unique $\mathrm{PGL}_2(\mathbf{R})$ -orbit of points x in $(P_s/\mathfrak{S}_5)(\mathbf{R})$ with stabilizer D_n .*

Proof. We have an injection $(P_s/\mathfrak{S}_5)(\mathbf{R}) \hookrightarrow P_s/\mathfrak{S}_5$ which is equivariant for the embedding $\mathrm{PGL}_2(\mathbf{R}) \hookrightarrow \mathrm{PGL}_2(\mathbf{C})$. In particular, $\mathrm{PGL}_2(\mathbf{R})_x \subset \mathrm{PGL}_2(\mathbf{C})_x$ for every $x \in (P_s/\mathfrak{S}_5)(\mathbf{R})$. Note that none of the groups appearing in Proposition 6.3 have subgroups isomorphic to $D_2 = \mathbf{Z}/2 \rtimes \mathbf{Z}/2$ or $D_4 = \mathbf{Z}/2 \rtimes \mathbf{Z}/4$. Consider the involution $\nu = (z \mapsto 1/z) \in \mathrm{PGL}_2(\mathbf{R})$. We will prove the proposition by using the following steps.

Step 1: Let $\tau \in \mathrm{PGL}_2(\mathbf{R})$. Consider a subset $S = \{x, y, z\} \subset \mathbf{P}^1(\mathbf{C})$ stabilized by complex conjugation, such that $\tau(x) = x$, $\tau(y) = z$ and $\tau(z) = y$. There is a transformation $g \in \mathrm{PGL}_2(\mathbf{R})$ that maps S to either $\{-1, 0, \infty\}$ or $\{-1, i, -i\}$, and that satisfies $g\tau g^{-1} = \nu = (z \mapsto 1/z) \in \mathrm{PGL}_2(\mathbf{R})$. In particular, $\tau^2 = \mathrm{id}$.

Proof of Step 1. Indeed, two transformations $g, h \in \mathrm{PGL}_2(\mathbf{C})$ that satisfy $g(x_i) = h(x_i)$ for three different points $x_1, x_2, x_3 \in \mathbf{P}^1(\mathbf{C})$ are necessarily equal. \square

Step 2: There is no $\phi \in \mathrm{PGL}_2(\mathbf{R})$ of order four that fixes a point $x \in (P_s/\mathfrak{S}_5)(\mathbf{R})$.

Proof of Step 2. By [Bea10, Theorem 4.2], all subgroups $G \subset \mathrm{PGL}_2(\mathbf{R})$ that are isomorphic to $\mathbf{Z}/4$ are conjugate to each other. Since the transformation $I: z \mapsto (z-1)/(z+1)$ is of order four, it gives a representative $G_I = \langle I \rangle$ of this conjugacy class. It is easily shown that I cannot fix any point $x \in (P_s/\mathfrak{S}_5)(\mathbf{R})$. \square

Step 3: Define $\rho \in \mathrm{PGL}_2(\mathbf{R})$ by $\rho(z) = \frac{-1}{z+1}$. Let $x = (x_1, \dots, x_5) \in (P_s/\mathfrak{S}_5)(\mathbf{R})$. Let $\phi \in \mathrm{PGL}_2(\mathbf{R})$ be of order three, with $\phi(x) = x$. There is a transformation $g \in \mathrm{PGL}_2(\mathbf{R})$ mapping x to $(-1, \infty, 0, \omega, \omega^2)$ with ω a primitive third root of unity. The stabilizer of x is the subgroup of $\mathrm{PGL}_2(\mathbf{R})$ generated by ρ and ν . In particular, we have $\mathrm{PGL}_2(\mathbf{R})_x \cong D_3$.

Proof of Step 3. By Step 1, there are elements x_1, x_2, x_3 which form an orbit under ϕ . Since complex conjugation preserves this orbit, one element in it is real; since g is defined over \mathbf{R} , they are all real. Let $g \in \mathrm{PGL}_2(\mathbf{R})$ such that $g(x_1) = -1$, $g(x_2) = \infty$ and $g(x_3) = 0$. Define $\kappa = g\phi g^{-1}$. Then $\kappa^3 = \mathrm{id}$, and κ preserves $\{-1, \infty, 0\}$ and sends -1 to ∞ and ∞ to 0 . Consequently, $\kappa(0) = -1$, and it follows that $\kappa = \rho$. Hence x is equivalent to an element of the form $(-1, \infty, 0, \alpha, \beta)$ with $\beta = \bar{\alpha}$ and $\alpha^2 + \alpha + 1 = 0$. \square

Recall that $\lambda = \zeta_5 + \zeta_5^{-1} \in \mathbf{R}$. Define $\gamma \in \mathrm{PGL}_2(\mathbf{R})$ by $\gamma(z) = \frac{(\lambda+1)z-1}{z+1}$ for $z \in \mathbf{P}^1(\mathbf{C})$.

Step 4: Let $x = (x_1, \dots, x_5) \in (P_s/\mathfrak{S}_5)(\mathbf{R})$. Suppose x is stabilized by a subgroup of $\mathrm{PGL}_2(\mathbf{R})$ of order five. There is a transformation $g \in \mathrm{PGL}_2(\mathbf{R})$ mapping x to $z = (0, -1, \infty, \lambda+1, \lambda)$ and identifying the stabilizer of x with the subgroup of $\mathrm{PGL}_2(\mathbf{R})$ generated by γ and ν . In particular, the stabilizer $\mathrm{PGL}_2(\mathbf{R})_x$ of x is isomorphic to D_5 .

Proof of Step 4. Let $\phi \in \mathrm{PGL}_2(\mathbf{R})_x$ be an element of order five. Using Step 1, one shows that the x_i are pairwise distinct, and we may assume that $x_i = \phi^{i-1}(x_1)$ for $i = 2, \dots, 5$. Since there is one real x_i and ϕ is defined over \mathbf{R} , all x_i are real.

Let $z = \{0, -1, \infty, \lambda+1, \lambda\}$. Note that z is the orbit of 0 under $\gamma: z \mapsto ((\lambda+1)z-1)/(z+1)$. The reflection $\nu: z \mapsto 1/z$ preserves z as well: we have $\lambda+1 = -(\zeta_5^2 + \zeta_5^{-2}) = -\lambda^2 + 2$, so that $\lambda(\lambda+1) = 1$. We conclude that $\mathrm{PGL}_2(\mathbf{R})_z \cong D_5$. Thus, by Proposition 6.3, there exists $g \in \mathrm{PGL}_2(\mathbf{C})$ such that $g(x_1) = 0$, $g(x_2) = -1$, $g(x_3) = \infty$, $g(x_4) = \lambda+1$ and $g(x_5) = \lambda$, and such that $g\mathrm{PGL}_2(\mathbf{R})_x g^{-1} = \mathrm{PGL}_2(\mathbf{R})_z$. Since all x_i and $z_i \in z$ are real, we have $\bar{g}(x_i) = z_i$ for each i . Hence, g and \bar{g} coincide on three points, which implies that $g = \bar{g}$, i.e. $g \in \mathrm{PGL}_2(\mathbf{R})$. \square

By Steps 1–4 above, together with Proposition 6.3, we are done. \square

6.2 Comparing the orbifold structures

There are two orbifold structures on the space $|\mathcal{M}_s(\mathbf{R})|$. On the one hand, one has the natural orbifold structure on $\mathcal{M}_s(\mathbf{R})$ by considering it as the real locus of a smooth separated Deligne–Mumford stack over \mathbf{R} (see [GF22, Proposition 2.12]); this is the orbifold structure of the quotient $G(\mathbf{R}) \backslash X_s(\mathbf{R})$. On the other hand, one has the orbifold structure $\overline{\mathcal{M}}_{\mathbf{R}}$ introduced in Definition 6.1. The goal of Section 6.2 is to calculate the difference between these orbifold structures.

We first show that there are no cone points in the orbifold $\mathrm{PGL}_2(\mathbf{R}) \backslash (P_s/\mathfrak{S}_5)(\mathbf{R})$. These are orbifold points whose stabilizer group is \mathbf{Z}/n ($n \geq 2$) acting on the orbifold chart by rotations. By Proposition 6.4, the fact that $\mathrm{PGL}_2(\mathbf{R}) \backslash (P_s/\mathfrak{S}_5)(\mathbf{R})$ has no cone points follows from the following lemma.

LEMMA 6.5. *Let $x = (x_1, \dots, x_5) \in (P_s/\mathfrak{S}_5)(\mathbf{R})$ such that $\mathrm{PGL}_2(\mathbf{R})_x = \langle \tau \rangle$ has order two. There is a $\mathrm{PGL}_2(\mathbf{R})_x$ -stable open neighborhood $U \subset (P_s/\mathfrak{S}_5)(\mathbf{R})$ of x such that $\mathrm{PGL}_2(\mathbf{R})_x \backslash U \rightarrow \mathcal{M}_s(\mathbf{R})$ is injective, and a homeomorphism $\phi: (U, x) \rightarrow (B, 0)$ for $0 \in B \subset \mathbf{R}^2$ an open ball, such that ϕ identifies $\mathrm{PGL}_2(\mathbf{R})_x$ with $\mathbf{Z}/2$ acting on B by reflections in a line through 0.*

Proof. Using Step 1 in the proof of Proposition 6.4, one checks that the only possibilities for the element $x = (x_1, \dots, x_5) \in (P_s/\mathfrak{S}_5)(\mathbf{R})$ are $(-1, 0, \infty, \beta, \beta^{-1})$, $(-1, i, -i, \beta, \beta^{-1})$, $(-1, -1, \beta, 0, \infty)$, $(-1, -1, \beta, i, -i)$, $(0, 0, \infty, \infty, -1)$ and $(-1, i, i, -i, -i)$ for some $\beta \in \mathbf{P}^1(\mathbf{R})$. \square

To analyze the difference between the two orbifolds $\mathcal{M}_s(\mathbf{R})$ and $\overline{\mathcal{M}}_{\mathbf{R}}$, we also need the following general lemma.

LEMMA 6.6. *Let X be a set and let Γ and G be groups with commuting actions on X . Let $x \in X$ with images $\bar{x} \in G \backslash X$ and $[x] \in \Gamma \backslash X$. Let $\Gamma_{[x]}$ be the stabilizer of $[x] \in \Gamma \backslash X$ in Γ , and let $G_{\bar{x}}$ be the stabilizer of \bar{x} in G . Then for each $\gamma \in \Gamma_{[x]}$ there exists an element $\phi(\gamma) \in G_{\bar{x}}$, unique up to multiplication by an element of G_x , such that $\gamma \cdot x = \phi(\gamma) \cdot x$; moreover, the map*

$$\Gamma_{[x]}/\Gamma_x \rightarrow G_{\bar{x}}/G_x, \quad \gamma \mapsto \phi(\gamma) \quad (36)$$

is an isomorphism.

Proof. The map (36) is well-defined because if $g, g' \in G$ are such that $\gamma \cdot x = g \cdot x = g' \cdot x$ then $(g')^{-1}g \in G_x$. Since the construction is symmetric in Γ and G , the analogous map $G_{\bar{x}}/G_x \rightarrow \Gamma_{[x]}/\Gamma_x$ is also well-defined. The latter is a left and right inverse of (36), and we are done. \square

Recall (see [Thu80, Proposition 13.3.1]), that the singular locus of a two-dimensional orbifold has the following types of local models: (i) $\mathbf{R}^2/(\mathbf{Z}/2)$, where $\mathbf{Z}/2$ acts on \mathbf{R}^2 by reflection in the y -axis (*mirror points*); (ii) $\mathbf{R}^2/(\mathbf{Z}/n)$, with \mathbf{Z}/n acting by rotations (*cone points of order n*); and (iii) \mathbf{R}^2/D_n , with D_n the dihedral group of order $2n$ (*corner reflectors of order n*).

PROPOSITION 6.7. Consider the orbifold structures $\mathcal{M}_s(\mathbf{R})$ and $\overline{\mathcal{M}}_{\mathbf{R}}$ on the space $|\mathcal{M}_s(\mathbf{R})|$.

- (i) For $x_0 \in \mathcal{M}_s(\mathbf{R})$, the isomorphism class of stabilizer groups of $\mathcal{M}_s(\mathbf{R})$ and $\overline{\mathcal{M}}_{\mathbf{R}}$ at x_0 differ if and only if $x_0 \in \mathcal{M}_s(\mathbf{R})$ is the moduli point attached to the five-tuple $(\infty, i, i, -i, -i)$.
- (ii) The stabilizer group of $\mathcal{M}_s(\mathbf{R})$ at the point x_0 is isomorphic to $\mathbf{Z}/2$, whereas the stabilizer group of $\overline{\mathcal{M}}_{\mathbf{R}}$ at x_0 is isomorphic to the dihedral group D_{10} of order 20.
- (iii) The orbifold $\overline{\mathcal{M}}_{\mathbf{R}}$ has no cone points and three corner reflectors, whose angles are $\pi/3, \pi/5$ and $\pi/10$.

Proof. The statements can be deduced from Proposition 6.4, Lemmas 6.5 and 6.6, and [GF24, Proposition 5.19]. To show how this works, let us introduce some notation. Let $\tilde{F} \in \mathcal{F}_s(\mathbf{R})$ with image $F \in X_s(\mathbf{R})$. Let $f \in Y$ be the image of $[\tilde{F}] \in G(\mathbf{R}) \setminus \mathcal{F}_s(\mathbf{R})$ under the isomorphism $G(\mathbf{R}) \setminus \mathcal{F}_s(\mathbf{R}) \xrightarrow{\sim} Y$ of Theorem 5.8. Let $PT_f \subset PT$ be the stabilizer of $f \in Y$ in the group PT . Let $k \in \{0, 1, 2\}$ be the number of nodes of F , and write $k = 2a + b$, where a is the number of pairs of complex conjugate nodes and b the number of real nodes. Then the image $x \in \mathbf{CH}^2$ of \tilde{F} under (23) lies on exactly k distinct mutually orthogonal hyperplanes $H \in \mathcal{H}$, with \mathcal{H} the set defined in (26). Since $F \in X_s(\mathbf{R})$, we have that $\tilde{F} \in \mathcal{F}_s^\alpha$ for some $\alpha \in P\mathcal{A}$. We get $x \in \mathbf{RH}_\alpha^n$.

Let $\mathcal{H}(x) \subset \mathcal{H}$ be the set of hyperplanes $H \in \mathcal{H}$ such that $x \in H$. Then a equals the number of pairs of hyperplanes $H_1, H_2 \in \mathcal{H}(x)$ with $\alpha(H_1) = H_2$, and b equals the number of hyperplanes $H \in \mathcal{H}(x)$ with $\alpha(H) = H$. Define $B_f \subset PT_f$ as the group generated by reflections in all $H \in \mathcal{H}(x)$ such that $\alpha(H) = H$. Consider the quotient map $p: \sqcup_{\alpha \in P\mathcal{A}} \mathbf{RH}_\alpha^2 \rightarrow Y$, let $\alpha_1, \dots, \alpha_\ell \in P\mathcal{A}$ be the elements such that $(x, \alpha_i) \sim (x, \alpha)$, and define $Y_f = \cup_{i=1}^\ell p(\mathbf{RH}_{\alpha_i}^2) \subset Y$. The subgroup $PT_f \subset PT$ preserves the subset $Y_f \subset Y$ by [GF24, Lemma 5.9]. Moreover, by [GF24, Proposition 5.19.4], there is an isometry between $B_f \setminus Y_f$ and the union of 10^a copies of $\mathbf{B}^2(\mathbf{R})$. Let \mathbf{B} be any one of these copies of $\mathbf{B}^2(\mathbf{R})$, and define

$$S_f := \text{Stab}_{PT_f/B_f}(\mathbf{B}),$$

the stabilizer of \mathbf{B} in the group PT_f/B_f . By construction of the orbifold $\overline{\mathcal{M}}_{\mathbf{R}}$ (see [GF24, Propositions 5.1 and 5.19]), the group S_f is a representative of the isomorphism class of stabilizer groups of the orbifold $\overline{\mathcal{M}}_{\mathbf{R}}$ at the moduli point $[F] \in \mathcal{M}_s(\mathbf{R})$ induced by F . Clearly, the stabilizer $G(\mathbf{R})_F \subset G(\mathbf{R})$ of $F \in X_s(\mathbf{R})$ yields the isomorphism class of stabilizer groups of the orbifold $\mathcal{M}_s(\mathbf{R})$ at the moduli point $[F] \in \mathcal{M}_s(\mathbf{R})$. In particular, we need to compare the isomorphism classes of the groups S_f and $G(\mathbf{R})_F$. To do so, we claim that there is a canonical isomorphism

$$G(\mathbf{R})_F \xrightarrow{\sim} PT_f/G(x). \quad (37)$$

Indeed, the actions of the groups PT and $G(\mathbf{R})$ on $\mathcal{F}_s(\mathbf{R})$ commute, so we can apply Lemma 6.6. Recall that $\tilde{F} \in \mathcal{F}_s(\mathbf{R})$ has images $F \in X_s(\mathbf{R})$ and $f \in Y$, and that the map $\mathcal{F}_s(\mathbf{R}) \rightarrow Y$ factors through an isomorphism $G(\mathbf{R}) \setminus \mathcal{F}_s(\mathbf{R}) \xrightarrow{\sim} Y$. Moreover, if $PT_{\tilde{F}} \subset PT$ denotes the stabilizer of

\tilde{F} in $P\Gamma$, then $P\Gamma_{\tilde{F}} = G(x)$ by Lemma 4.4. As the group $G(\mathbf{C})$ acts freely on \mathcal{F}_s by Lemma 3.12, the group $G(\mathbf{R})$ acts freely on $\mathcal{F}_s(\mathbf{R})$. Thus, (37) follows from Lemma 6.6.

If F has no nodes ($k=0$), then $G(x)$ is trivial by [GF24, Proposition 5.19.1], and $P\Gamma_f = S_f$. Thus, $G(\mathbf{R})_F \cong S_f$ in view of (37).

If F has only real nodes, then $B_f = G(x)$ and $P\Gamma_f/G(x) = S_f$. Thus, $G(\mathbf{R})_F \cong S_f$ by (37).

Finally, suppose that F has a pair of complex nodes ($a=1$ and $b=0$). The zero set of F defines a five-tuple $\underline{z} = (z_1, \dots, z_5) \in \mathbf{P}^1(\mathbf{C})^5$, well defined up to the $\mathrm{PGL}_2(\mathbf{R}) \times \mathfrak{S}_5$ action on $\mathbf{P}^1(\mathbf{C})$, where $z_1 \in \mathbf{P}^1(\mathbf{R})$ and $z_3 = \bar{z}_2 = z_5 = \bar{z}_4 \in \mathbf{P}^1(\mathbf{C}) \setminus \mathbf{P}^1(\mathbf{R})$. Write $\underline{z} = (w, z, \bar{z}, z, \bar{z})$ with $w \in \mathbf{P}^1(\mathbf{R})$ and $z \in \mathbf{P}^1(\mathbf{C}) \setminus \mathbf{P}^1(\mathbf{R})$. There is a unique $T \in \mathrm{PGL}_2(\mathbf{R})$ such that $T(w) = \infty$ and $T(z) = i$. This gives $T(\underline{z}) = (\infty, i, -i, i, -i)$. In particular, F is unique up to isomorphism.

We have $G(x) \cong (\mathbf{Z}/10)^2$, and as there are no real nodes, B_f is trivial. With respect to the isometry $\mathrm{CH}^2(\mathbf{C}) \xrightarrow{\sim} \mathbf{B}^2(\mathbf{C})$ of [GF24, Lemma 5.16], the anti-holomorphic involutions $\alpha_j: \mathbf{B}^2(\mathbf{C}) \rightarrow \mathbf{B}^2(\mathbf{C})$ induced by the $\alpha_j \in P\mathcal{A}$ are $(t_1, t_2) \mapsto (\bar{t}_2 \zeta^j, \bar{t}_1 \zeta^j)$, for $j \in \mathbf{Z}/10$. The fixed point sets are given by $\mathbf{B}^2(\mathbf{R})_{\alpha_j} = \{t_2 = \bar{t}_1 \zeta^j\} \subset \mathbf{B}^2(\mathbf{C})$. The subgroup $E_f \subset G(x)$ that stabilizes $\mathbf{B}^2(\mathbf{R})_{\alpha_j}$ is the group $E_f \cong \mathbf{Z}/10$ generated by the transformations $(t_1, t_2) \mapsto (\zeta t_1, \zeta^{-1} t_2)$. There is only one non-trivial $T \in \mathrm{PGL}_2(\mathbf{R})$ fixing ∞ and preserving $\{i, -i\} \subset \mathbf{P}^1(\mathbf{C})$, and T has order two, so $G(\mathbf{R})_F = \mathbf{Z}/2$. Therefore, by (37), we have an exact sequence of groups $0 \rightarrow G(x) \cong (\mathbf{Z}/10)^2 \rightarrow P\Gamma_f \rightarrow \mathbf{Z}/2 \rightarrow 0$ inducing an exact sequence $0 \rightarrow E_f \cong \mathbf{Z}/10 \rightarrow S_f \rightarrow \mathbf{Z}/2 \rightarrow 0$. By Proposition 6.4 and Lemma 6.5, the proposition follows. \square

6.3 The real moduli space as a hyperbolic triangle

The goal of Section 6.3 is to prove Theorem 6.2.

The results in the above Sections 6.1 and 6.2 give the orbifold singularities of $\overline{\mathcal{M}}_{\mathbf{R}}$ together with their stabilizer groups. In order to determine the hyperbolic orbifold structure of $\overline{\mathcal{M}}_{\mathbf{R}}$, we also need to know the underlying topological space $|\mathcal{M}_s(\mathbf{R})|$ of $\overline{\mathcal{M}}_{\mathbf{R}}$. The first observation is that $\mathcal{M}_s(\mathbf{R})$ is compact. Indeed, it is classical that the topological space $\mathcal{M}_s(\mathbf{C}) = G(\mathbf{C}) \setminus X_s(\mathbf{C})$, parametrizing complex stable binary quintics, is compact. This follows from the proper surjective map $\overline{M}_{0,5}(\mathbf{C})/\mathfrak{S}_5 \rightarrow \mathcal{M}_s(\mathbf{C})$ and the properness of the stack of stable five-pointed curves $\overline{M}_{0,5}$ [Knu83], or from the fact that $\mathcal{M}_s(\mathbf{C})$ is homeomorphic to a compact ball quotient [Shi64]. Moreover, the map $\mathcal{M}_s(\mathbf{R}) \rightarrow \mathcal{M}_s(\mathbf{C})$ is proper, which proves the compactness of $\mathcal{M}_s(\mathbf{R})$. The second observation is that $\mathcal{M}_s(\mathbf{R})$ is connected, since $X_s(\mathbf{R})$ is obtained from the euclidean space $X(\mathbf{R}) = \{F \in \mathbf{R}[x, y]: F \text{ homogeneous and } \deg(F) = 5\}$ by removing a subspace of codimension two. In the following lemma we generalize both of these observations.

LEMMA 6.8. *The moduli space $\mathcal{M}_s(\mathbf{R})$ is homeomorphic to a closed disc $\overline{D} \subset \mathbf{R}^2$.*

Proof. The idea is to show that the following holds.

- (i) For each $j \in \{0, 1, 2\}$, the embedding $\mathcal{M}_j \hookrightarrow \overline{\mathcal{M}}_j \subset \mathcal{M}_s(\mathbf{R})$ of the connected component \mathcal{M}_j of $\mathcal{M}_0(\mathbf{R})$ into its closure in $\mathcal{M}_s(\mathbf{R})$ is homeomorphic to the embedding $D \hookrightarrow \overline{D}$ of the open unit disc into the closed unit disc in \mathbf{R}^2 .
- (ii) We have $\mathcal{M}_s(\mathbf{R}) = \overline{\mathcal{M}}_0 \cup \overline{\mathcal{M}}_1 \cup \overline{\mathcal{M}}_2$, and this gluing corresponds, up to homeomorphism, to the gluing of three closed discs $\overline{D}_i \subset \mathbf{R}^2$ as in Figure 1.

To prove this, one considers the moduli spaces of real smooth (respectively stable) genus zero curves with five real marked points, as well as twists of this space. Define two anti-holomorphic involutions $\sigma_i: \mathbf{P}^1(\mathbf{C})^5 \rightarrow \mathbf{P}^1(\mathbf{C})^5$ by $\sigma_1(x_1, x_2, x_3, x_4, x_5) = (\bar{x}_1, \bar{x}_2, \bar{x}_3, \bar{x}_5, \bar{x}_4)$, and

$\sigma(x_1, x_2, x_3, x_4, x_5) = (\bar{x}_1, \bar{x}_3, \bar{x}_2, \bar{x}_5, \bar{x}_4)$. Then define

$$P_0^1(\mathbf{R}) = P_0^{\sigma_1}, \quad P_s^1(\mathbf{R}) = P_1(\mathbf{C})^{\sigma_1}, \quad P_0^2(\mathbf{R}) = P_0^{\sigma_2}, \quad P_s^2(\mathbf{R}) = P_1(\mathbf{C})^{\sigma_2}.$$

It is clear that $\mathcal{M}_0 = \mathrm{PGL}_2(\mathbf{R}) \setminus P_0(\mathbf{R})/\mathfrak{S}_5$. Similarly, we have

$$\mathcal{M}_1 = \mathrm{PGL}_2(\mathbf{R}) \setminus P_0^1(\mathbf{R})/\mathfrak{S}_3 \times \mathfrak{S}_2 \quad \text{and} \quad \mathcal{M}_2 = \mathrm{PGL}_2(\mathbf{R}) \setminus P_0^2(\mathbf{R})/\mathfrak{S}_2 \times \mathfrak{S}_2.$$

Moreover, we have $\overline{\mathcal{M}}_0 = \mathrm{PGL}_2(\mathbf{R}) \setminus P_s(\mathbf{R})/\mathfrak{S}_5$. We define

$$\overline{\mathcal{M}}_1 = \mathrm{PGL}_2(\mathbf{R}) \setminus P_s^1(\mathbf{R})/\mathfrak{S}_3 \times \mathfrak{S}_2, \quad \text{and} \quad \overline{\mathcal{M}}_2 = \mathrm{PGL}_2(\mathbf{R}) \setminus P_s^2(\mathbf{R})/\mathfrak{S}_2 \times \mathfrak{S}_2.$$

Each $\overline{\mathcal{M}}_j$ is homeomorphic to a closed disc in \mathbf{R}^2 . Moreover, the natural maps $\overline{\mathcal{M}}_j \rightarrow \mathcal{M}_s(\mathbf{R})$ are closed embeddings of topological spaces, and one can check that the images glue to form $\mathcal{M}_s(\mathbf{R})$ in the prescribed way. We leave the details to the reader. \square

Proof of Theorem 6.2. To any closed two-dimensional orbifold O one can associate a set of natural numbers $S_O = \{n_1, \dots, n_k; m_1, \dots, m_l\}$ by letting k be the number of cone points of X_O , l the number of corner reflectors, n_i the order of the i th cone point and m_j the order of the j th corner reflector (see [Thu80, Proposition 13.3.1]). A closed two-dimensional orbifold O is determined, up to orbifold-structure preserving homeomorphism, by its underlying space X_O and the set S_O [Thu80]. By Lemma 6.8, $\overline{\mathcal{M}}_{\mathbf{R}}$ is homeomorphic to a closed disc in \mathbf{R}^2 . By Proposition 6.7, $\overline{\mathcal{M}}_{\mathbf{R}}$ has no cone points and three corner reflectors whose angles are $\pi/3, \pi/5$ and $\pi/10$. This implies that $\overline{\mathcal{M}}_{\mathbf{R}}$ and $\Delta_{3,5,10}$ are isomorphic as topological orbifolds. Consequently, the orbifold fundamental group of $\overline{\mathcal{M}}_{\mathbf{R}}$ is abstractly isomorphic to the group $\Gamma_{3,5,10}$ defined in (3).

Let $\phi: \Gamma_{3,5,10} \hookrightarrow \mathrm{PSL}_2(\mathbf{R})$ be any embedding such that $X := \phi(\Gamma_{3,5,10}) \setminus \mathbf{R}H^2$ is a hyperbolic orbifold; we claim that there is a fundamental domain Δ for X isometric to $\Delta_{3,5,10}$. To see this, consider the generator $a \in \Gamma_{3,5,10}$. Since $\phi(a)^2 = 1$, there exists a geodesic $L_1 \subset \mathbf{R}H^2$ such that $\phi(a) \in \mathrm{PSL}_2(\mathbf{R}) = \mathrm{Isom}(\mathbf{R}H^2)$ is the reflection across L_1 . Next, consider the generator $b \in \Gamma_{3,5,10}$. There exists a geodesic $L_2 \subset \mathbf{R}H^2$ such that $\phi(b)$ is the reflection across L_2 , and we have $L_2 \cap L_1 \neq \emptyset$. Let $x \in L_1 \cap L_2$. Then $\phi(a)\phi(b)$ is an element of order three that fixes x , and hence is a rotation around x . Therefore, one of the angles between L_1 and L_2 must be $\pi/3$. Finally, we know that $\phi(c)$ is an element of order two in $\mathrm{PSL}_2(\mathbf{R})$, and hence a reflection across a line L_3 . As $L_3 \cap L_2 \neq \emptyset$ and $L_3 \cap L_1 \neq \emptyset$, the three geodesics $L_i \subset \mathbf{R}H^2$ enclose a hyperbolic triangle. As the orders of the three elements $\phi(a)\phi(b)$, $\phi(a)\phi(c)$ and $\phi(b)\phi(c)$ are respectively three, five and ten, the three interior angles of the triangle are $\pi/3, \pi/5$ and $\pi/10$. Thus, X is isometric to $\Delta_{3,5,10}$.

Consequently, $P\Gamma_{\mathbf{R}} \setminus \mathbf{R}H^2$ is isometric to $\Gamma_{3,5,10} \setminus \mathbf{R}H^2$. It follows that the lattices $P\Gamma_{\mathbf{R}}$ and $\Gamma_{3,5,10}$ are conjugate in $\mathrm{PO}(2, 1)$ (see e.g. [Rat99, Lemma 1]). \square

7. The monodromy groups

In this section, we describe the monodromy group $P\Gamma$ attached to the moduli space $X_0(\mathbf{C})$, as well as the groups $P\Gamma_{\alpha}$ appearing in Proposition 5.4. As for the lattice (Λ, \mathfrak{h}) (see (8)), we have the following.

THEOREM 7.1 (Shimura). *There is an isomorphism of hermitian \mathcal{O}_K -lattices*

$$(\Lambda, \mathfrak{h}) \cong (\mathcal{O}_K^3, \mathrm{diag}(-\lambda, 1, 1)), \quad \lambda = \zeta_5 + \zeta_5^{-1} = \frac{\sqrt{5} - 1}{2}.$$

Proof. See [Shi64, Section 6] as well as item (5) in the table on page 1 of that paper. \square

Write $\Lambda = \mathcal{O}_K^3$ and $\mathfrak{h} = \text{diag}(-\lambda, 1, 1)$. Consider the \mathbf{F}_5 -vector space $W = \Lambda/(1 - \zeta_5)\Lambda$ equipped with the quadratic form $q = \mathfrak{h} \bmod \theta$. Define three anti-isometric involutions as follows:

$$\begin{aligned}\alpha_0 &: (x_0, x_1, x_2) \mapsto (\bar{x}_0, \bar{x}_1, \bar{x}_2), \\ \alpha_1 &: (x_0, x_1, x_2) \mapsto (\bar{x}_0, -\bar{x}_1, \bar{x}_2), \\ \alpha_2 &: (x_0, x_1, x_2) \mapsto (\bar{x}_0, -\bar{x}_1, -\bar{x}_2).\end{aligned}\tag{38}$$

LEMMA 7.2. *An anti-unitary involution of Λ is Γ -conjugate to exactly one of the $\pm\alpha_j$. In particular, $|P\Gamma \setminus P\mathcal{A}| = 3$ and the α_i defined in (38) form a set of representatives for $P\Gamma \setminus P\mathcal{A}$.*

Proof. For isometries $\alpha: W \rightarrow W$, the dimension and determinant of the fixed space $(W^\alpha, q|_{W^\alpha})$ are conjugacy-invariant. Using this, one can show that the elements $\pm\alpha_i$ are pairwise non Γ -conjugate. Moreover, $|P\Gamma \setminus P\mathcal{A}| = |\pi_0(\mathcal{M}_0(\mathbf{R}))| = 3$ by Proposition 5.4 and Theorem 7.1. \square

Define

$$\theta = \zeta_5 - \zeta_5^{-1} \in \mathcal{O}_K.$$

Note that $|\theta|^2 = \frac{\sqrt{5}+5}{2}$. The fixed lattices of the anti-unitary involutions α_i defined in (38) are

$$\begin{aligned}\Lambda^{\alpha_0} &= \mathbf{Z}[\lambda] \oplus \mathbf{Z}[\lambda] \oplus \mathbf{Z}[\lambda], \\ \Lambda^{\alpha_1} &= \mathbf{Z}[\lambda] \oplus \theta\mathbf{Z}[\lambda] \oplus \mathbf{Z}[\lambda], \\ \Lambda^{\alpha_2} &= \mathbf{Z}[\lambda] \oplus \theta\mathbf{Z}[\lambda] \oplus \theta\mathbf{Z}[\lambda],\end{aligned}\tag{39}$$

where

$$\lambda = \zeta_5 + \zeta_5^{-1} = \frac{\sqrt{5} - 1}{2}.\tag{40}$$

Restricting \mathfrak{h} to the Λ^{α_j} yields quadratic forms q_0, q_1 and q_2 on $\mathbf{Z}[\lambda]^3$ defined as follows:

$$\begin{aligned}q_0(x_0, x_1, x_2) &= -\lambda x_0^2 + x_1^2 + x_2^2, \\ q_1(x_0, x_1, x_2) &= -\lambda x_0^2 + \left(\frac{\sqrt{5}+5}{2}\right) \cdot x_1^2 + x_2^2, \\ q_2(x_0, x_1, x_2) &= -\lambda x_0^2 + \left(\frac{\sqrt{5}+5}{2}\right) \cdot x_1^2 + \left(\frac{\sqrt{5}+5}{2}\right) \cdot x_2^2,\end{aligned}\tag{41}$$

We consider $\mathbf{Z}[\lambda]$ (with λ as in (40)) as a subring of \mathbf{R} via the embedding that sends λ to a positive element.

THEOREM 7.3. *Consider the quadratic forms q_j defined in (41), with λ as in (40). There is a union of geodesic subspaces $\mathcal{H}_j \subset \mathbf{R}H^2$ ($j \in \{0, 1, 2\}$) and an isomorphism of hyperbolic orbifolds*

$$\mathcal{M}_0(\mathbf{R}) \cong \prod_{j=0}^2 \text{PO}(q_j, \mathbf{Z}[\lambda]) \setminus (\mathbf{R}H^2 - \mathcal{H}_j).\tag{42}$$

Proof. By Proposition 5.4 and Lemma 7.2, we obtain an isomorphism

$$\mathcal{M}_0(\mathbf{R}) \cong \prod_{j=0}^2 P\Gamma_{\alpha_j} \setminus (\mathbf{R}H_{\alpha_j}^2 - \mathcal{H}_j).$$

Note that $P\Gamma_{\alpha_j} = N_{P\Gamma}(\alpha_j)$ for the normalizer $N_{P\Gamma}(\alpha_j)$ of α_j in $P\Gamma$. If h_j denotes the restriction of \mathfrak{h} to Λ^{α_j} , there is a natural embedding

$$\iota_j: N_{P\Gamma}(\alpha_j) \hookrightarrow \mathrm{PO}(\Lambda^{\alpha_j}, h_j, \mathbf{Z}[\lambda]).$$

We claim that ι_j is an isomorphism. This holds because the natural homomorphism $\pi_j: N_{\Gamma}(\alpha_j) \rightarrow \mathrm{O}(\Lambda^{\alpha_j}, h_j)$ is surjective, where $N_{\Gamma}(\alpha_j) = \{g \in \Gamma: g \circ \alpha_j = \alpha_j \circ g\}$ is the normalizer of α_j in Γ . The surjectivity of π_j follows in turn from the equality

$$\Lambda = \mathcal{O}_K \cdot \Lambda^{\alpha_j} + \mathcal{O}_K \cdot \theta(\Lambda^{\alpha_j})^\vee \subset K^3, \quad (43)$$

and (43) follows from (39). Since $\mathrm{PO}(\Lambda^{\alpha_j}, h_j, \mathbf{Z}[\lambda]) = \mathrm{PO}(q_j, \mathbf{Z}[\lambda])$, we are done. \square

Proof of Theorem 1.2. This follows from Corollary 5.9, Theorem 6.2 and Theorem 7.3. \square

Proof of Theorem 1.3. In [AY98], Apéry and Yoshida proved that $\overline{\mathcal{M}}_0$ is the hyperbolic triangle with angles $\pi/2, \pi/4$ and $\pi/5$. As the two hyperplanes in Figure 1 intersect orthogonally, this implies that the bottom angle of the triangle $\overline{\mathcal{M}}_0$ in Figure 1 (i.e. its angle at $(0, -1, \infty, \infty, 1)$) equals $\pi/2$, and that the angle of $\overline{\mathcal{M}}_0$ at $(0, -1, -1, \infty, \infty)$ equals $\pi/4$. One deduces that the left angle of $\overline{\mathcal{M}}_1$ is $\pi/2$, and that the angle of $\overline{\mathcal{M}}_2$ at $(0, -1, -1, \infty, \infty)$ equals $\pi/4$.

For a hyperbolic triangle with angles α, β, γ and sides a, b, c such that a is the side opposite to α , b the side opposite to β and γ the side opposite to c , one has the hyperbolic law of cosines

$$\cosh(c) = \frac{\cos(\alpha) \cos(\beta) + \cos(\gamma)}{\sin(\alpha) \sin(\beta)}. \quad (44)$$

Applying (44) to the triangles $\overline{\mathcal{M}}_0$ and $\overline{\mathcal{M}}_{\mathbf{R}}$, one can calculate the length of the side of $\overline{\mathcal{M}}_2$ that connects $(0, -1, -1, \infty, \infty)$ and $(\infty, i, i, -i, -i)$. Applying (44) again, it follows that the angle of $\overline{\mathcal{M}}_2$ at the point $(0, -i, \infty, \infty, i)$ is $\pi/2$. Thus, the angle of $\overline{\mathcal{M}}_1$ at $(0, -i, \infty, \infty, i)$ is also $\pi/2$. \square

8. Non-arithmetic lattices in the projective orthogonal group

In a previous paper we proved a result, see [GF24, Theorem 1.8], that has the following consequence. For $n \geq 2$, define

$$\mathcal{L}_{\zeta_5}^n(\lambda) := \left(\mathbf{Q}(\zeta_5), \mathbf{Z}[\zeta_5]_{\lambda}^{n,1} \right), \quad \lambda = \zeta_5 + \zeta_5^{-1} = (\sqrt{5} - 1)/2.$$

Here, $\mathbf{Z}[\zeta_5]_{\lambda}^{n,1}$ is the free $\mathbf{Z}[\zeta_5]$ -module of rank $n + 1$ equipped with the hermitian form h defined as $h(x, y) = -\lambda \cdot x_0 \bar{y}_0 + \cdots + x_n \bar{y}_n$. Then $\mathcal{L}_{\zeta_5}^n(\lambda)$ is a hermitian lattice of rank $n + 1$ in the sense of [GF24, Definition 2.2] (indeed, this follows from [GF24, Example 2.12]). For each $n \geq 2$, perform the gluing construction of [GF24, Definition 1.1] to associate to the hermitian lattice $\mathcal{L}_{\zeta_5}^n(\lambda)$ a topological space $M(\mathcal{L}_{\zeta_5}^n(\lambda))$. By [GF24, Theorem 1.2], there exists a canonical real hyperbolic orbifold structure on $M(\mathcal{L}_{\zeta_5}^n(\lambda))$ such that each connected component of $M(\mathcal{L}_{\zeta_5}^n(\lambda))$ is isomorphic to the quotient of real hyperbolic n -space $\mathbf{R}H^n$ by a lattice in $\mathrm{PO}(n, 1)$. Define an anti-unitary involution $\alpha_0: \mathbf{Z}[\zeta_5]_{\lambda}^{n,1} \rightarrow \mathbf{Z}[\zeta_5]_{\lambda}^{n,1}$ by $\alpha_0(x) = \bar{x}$, let

$$M(\mathcal{L}_{\zeta_5}^n(\lambda), \alpha_0) \subset M(\mathcal{L}_{\zeta_5}^n(\lambda))$$

be the connected component that contains the image of the natural map $\mathbf{R}H_{\alpha_0}^n \rightarrow M(\mathcal{L}_{\zeta_5}^n(\lambda))$, and let

$$\Gamma_{\zeta_5}^n(\lambda) \subset \mathrm{PO}(n, 1)$$

be a lattice such that $M(\mathcal{L}_{\zeta_5}^n(\lambda), \alpha_0) \cong \Gamma_{\zeta_5}^n(\lambda) \setminus \mathbf{R}H^n$ (compare [GF24, p. 7]).

By combining [GF24, Theorem 1.8] with the main results of this paper, one can prove the following result.

THEOREM 8.1. *For each $n \geq 2$, the lattice $\Gamma_{\zeta_5}^n(\lambda) \subset \mathrm{PO}(n, 1)$ is non-arithmetic.*


Proof. Write $\mathbf{Z}[\zeta_5]_{\lambda}^{2,1} = \Lambda$; this abuse of notation is harmless in view of Theorem 7.1. Let $\Gamma = \mathrm{Aut}(\Lambda)$. By Lemma 7.2, an anti-unitary involution of Λ is Γ -conjugate to exactly one of the involutions $\pm\alpha_j$ defined in (38). We can therefore apply [GF24, Theorem 1.8], which implies that $\Gamma_{\zeta_5}^n(\lambda) \subset \mathrm{PO}(n, 1)$ is non-arithmetic for each $n \geq 2$ provided that $\Gamma_{\zeta_5}^2(\lambda) \subset \mathrm{PO}(2, 1)$ is non-arithmetic. In other words, we are reduced to the case $n = 2$. By Theorem 7.1, the lattice $\Gamma_{\zeta_5}^2(\lambda) \subset \mathrm{PO}(2, 1)$ is conjugate to the lattice $P\Gamma_{\mathbf{R}} \subset \mathrm{PO}(2, 1)$ defined in Corollary 5.9. Moreover, by Theorem 6.2, the lattice $P\Gamma_{\mathbf{R}}$ is conjugate to the lattice $\Gamma_{3,5,10} \subset \mathrm{PO}(2, 1)$ defined in (3). Finally, by Takeuchi's classification of arithmetic triangle groups (see [Tak77]), the subgroup $\Gamma_{3,5,10} \subset \mathrm{PO}(2, 1)$ is non-arithmetic. Thus, $\Gamma_{\zeta_5}^2(\lambda)$ is non-arithmetic, and the theorem follows. \square

ACKNOWLEDGEMENTS

This project was carried out partly at the ENS in Paris and partly at the Leibniz University in Hannover. I thank my former PhD advisor Olivier Benoist for his guidance and support. I thank Romain Branchereau, Samuel Bronstein, Nicolas Tholozan and Frans Oort for useful discussions. I thank Nicolas Bergeron for pointing me to Takeuchi's paper on hyperbolic triangle groups.

I would like to thank the referee for his or her valuable comments on this paper.

FUNDING STATEMENT

This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 754362  and from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme under grant agreement No. 948066 (ERC-StG RationAlgie).

CONFLICTS OF INTEREST

None.

JOURNAL INFORMATION

Moduli is published as a joint venture of the Foundation Compositio Mathematica and the London Mathematical Society. As not-for-profit organisations, the Foundation and Society reinvest 100% of any surplus generated from their publications back into mathematics through their charitable activities.

REFERENCES

- [ACT02a] D. Allcock, J. Carlson and D. Toledo, *Orthogonal complex hyperbolic arrangements*, in Symposium in honor of C. H. Clemens, Contemporary Mathematics, vol. 312 (American Mathematical Society, 2002), 1–8.
- [ACT02b] D. Allcock, J. Carlson and D. Toledo, *The complex hyperbolic geometry of the moduli space of cubic surfaces*, J. Algebr. Geom. **11** (2002), 659–724.

- [ACT06] D. Allcock, J. Carlson and D. Toledo, *Nonarithmetic uniformization of some real moduli spaces*, Geom. Dedic. **122** (2006), 159–169.
- [ACT07] D. Allcock, J. Carlson and D. Toledo, *Hyperbolic geometry and the moduli space of real binary sextics*, in Arithmetic and geometry around hypergeometric functions, Progress in Mathematics, vol. 260 (Birkhäuser, 2007), 1–22.
- [ACT10] D. Allcock, J. Carlson and D. Toledo, *Hyperbolic geometry and moduli of real cubic surfaces*, Ann. Sci. Éc. Norm. Supér. **43** (2010), 69–115.
- [AY98] F. Apéry and M. Yoshida, *Pentagonal structure of the configuration space of five points in the real projective line*, Kyushu J. Math. **52** (1998), 1–14.
- [Bea09] A. Beauville, *Moduli of cubic surfaces and Hodge theory (after Allcock, Carlson, Toledo)*, in Géométries à courbure négative ou nulle, groupes discrets et rigidités, Séminaires et Congrès, vol. 18 (Société Mathématique de France, 2009), 445–466.
- [Bea10] A. Beauville, *Finite subgroups of $\mathrm{PGL}_2(K)$* , in Vector bundles and complex geometry, Contemporary Mathematics, vol. 522 (American Mathematical Society, 2010), 23–29.
- [Chu11] K. Chu, *On the geometry of the moduli space of real binary octics*, Can. J. Math. **63** (2011), 755–797.
- [DIK00] A. Degtyarev, I. Itenberg and V. Kharlamov, *Real Enriques surfaces*, Lecture Notes in Mathematics, vol. 1746 (Springer-Verlag, Berlin, 2000).
- [DM86] P. Deligne and G. Mostow, *Monodromy of hypergeometric functions and nonlattice integral monodromy*. in: Institut des Hautes Études Scientifiques, Publ. Math. **63** (1986), 5–89.
- [Fox57] R. Fox, *Covering spaces with singularities*, in Algebraic geometry and topology. A symposium in honor of S. Lefschetz (Princeton University Press, 1957), 243–257.
- [GF22] O. de Gaay Fortman. *Moduli spaces and algebraic cycles in real algebraic geometry*, PhD thesis (École normale supérieure de Paris, 2022).
- [GF24] O. de Gaay Fortman, *Non-arithmetic hyperbolic orbifolds attached to unitary Shimura varieties*, Preprint (2024), arXiv:2301.01598v4.
- [GH81] B. Gross and J. Harris, *Real algebraic curves*, Ann. Sci. Éc. Norm. Supér. **14** (1981), 157–182.
- [HR18] G. Heckman and S. Rieken, *Hyperbolic geometry and moduli of real curves of genus three*, Math. Ann. **370** (2018), 1321–1360.
- [Kha76] V. Kharlamov, *Topological types of nonsingular surfaces of degree 4 in \mathbf{RP}^3* , Akademija Nauk SSSR. Funkcional’nyi Analiz i ego Prilozheniya **10** (1976), 55–68.
- [Kha84] V. Kharlamov, *On the classification of nonsingular surfaces of degree 4 in \mathbf{RP}^3 with respect to rigid isotopies*, Akademiya Nauk SSSR. Funktsional’nyi Analiz i ego Prilozheniya **18** (1984), 49–56.
- [Knu83] F. Knudsen, *The projectivity of the moduli space of stable curves. II. The stacks $M_{g,n}$* , Math. Scand. **52** (1983), 161–199.
- [Loo07] E. Looijenga, *Uniformization by Lauricella functions—an overview of the theory of Deligne–Mostow*, in Arithmetic and geometry around hypergeometric functions, Progress in Mathematics, vol. 260 (Birkhäuser, Basel, 2007), 207–244.
- [MO13] B. Moonen and F. Oort, *The Torelli locus and special subvarieties*, in Handbook of moduli, Vol. II, Advanced Lectures in Mathematics, vol. 25 (International Press of Boston, 2013), 549–594.
- [MFK94] D. Mumford, J. Fogarty and F. Kirwan, *Geometric invariant theory*, Third edition, Ergebnisse der Mathematik und ihrer Grenzgebiete, vol. 34 (Springer, 1994).
- [MS72] D. Mumford and K. Suominen, *Introduction to the theory of moduli*, in Algebraic geometry, Oslo 1970 (Wolters-Noordhoff, 1972), 171–222.
- [Neu99] J. Neukirch, *Algebraic number theory*, Grundlehren der mathematischen Wissenschaften, vol. 322 (Springer, 1999).
- [Nik79] V. Nikulin, *Integer symmetric bilinear forms and some of their geometric applications*, Izvestiya Akademii Nauk SSSR. Seriya Matematicheskaya **43** (1979), 111–177, 238.

- [Rat99] J. Ratcliffe, *On the isometry groups of hyperbolic orbifolds*, *Geom. Dedicata* **78** (1999), 63–67.
- [Ser79] J.-P. Serre, *Local fields*, *Graduate Texts in Mathematics*, vol. 67 (Springer, 1979).
- [SS89] M. Seppälä and R. Silhol, *Moduli spaces for real algebraic curves and real abelian varieties*, *Math. Z.* **201** (1989), 151–165.
- [Shi63] G. Shimura, *On analytic families of polarized abelian varieties and automorphic functions*, *Ann. Math.* **78** (1963), 149–192.
- [Shi64] G. Shimura, *On purely transcendental fields automorphic functions of several variable*, *Osaka Math. J.* **1** (1964), 1–14.
- [Sil89] R. Silhol, *Real algebraic surfaces*, *Lecture Notes in Mathematics*, vol. 1392 (Springer, 1989).
- [Tak77] K. Takeuchi, *Arithmetic triangle groups*, *J. Math. Soc. Jpn.* **29** (1977), 91–106.
- [Ter83] T. Terada, *Fonctions hypergéométriques F_1 et fonctions automorphes. I*, *J. Math. Soc. Jpn.* **35** (1983), 451–475.
- [Thu80] W. Thurston, *The geometry and topology of three-manifolds* (Princeton University Press, 1980).
- [Voi02] C. Voisin, *Théorie de Hodge et Géométrie Algébrique Complexe, Cours Spécialisés*, vol. 10 (Société Mathématique de France, Paris, 2002).
- [YY84] T. Yamazaki and M. Yoshida, *On Hirzebruch's examples of surfaces with $c_1^2 = 3c_2$* , *Math. Ann.* **266** (1984), 421–431.
- [Yos98] M. Yoshida, *The real loci of the configuration space of six points on the projective line and a Picard modular 3-fold*, *Kumamoto J. Math.* **11** (1998), 43–67.

Olivier de Gaay Fortman a.o.d.degaayfortman@uu.nl

Department of Mathematics, Utrecht University, Utrecht, The Netherlands