

The Full-Sky Astrometric Mapping Explorer – Distances and Photometry of 40 Million Stars

Scott D. Horner¹, Marvin E. Germain², Thomas P. Greene³, Fred H. Harris², Mark S. Johnson⁴, Kenneth J. Johnston¹, David G. Monet², Marc A. Murison¹, James D. Phillips⁵, Robert D. Reasenber⁵, P. Kenneth Seidelmann¹, Sean E. Urban¹, Richard H. Vassar⁶

The Full-sky Astrometric Mapping Explorer (FAME) is designed to perform an all-sky, astrometric survey with unprecedented accuracy. It will create a rigid astrometric catalog of 4×10^7 stars with $5 < m_V < 15$. For bright stars, $5 < m_V < 9$, FAME will determine positions and parallaxes accurate to $< 50 \mu\text{as}$, with proper motion errors $< 50 \mu\text{as/yr}$. For fainter stars, $9 < m_V < 15$, FAME will determine positions and parallaxes accurate to $< 500 \mu\text{as}$, with proper motion errors $< 500 \mu\text{as/yr}$. It will also collect photometric data on these 4×10^7 stars in four Sloan Digital Sky Survey colors. NASA selected FAME to be one of five MIDEX missions funded for a concept study. In October 1999, NASA selected FAME for launch in 2004 as the MIDEX-4 mission in its Explorer program.

The greatest strength of FAME will be the large number of stars observed, enabling diverse studies of stellar and galactic evolution. FAME will provide a meaningful statistical sample of stars for studies of the frequency of stellar companions with $m > M_{\text{Jupiter}}$. This will provide accurate statistics on the frequency of occurrence of multiple star systems as a function of spectral type, as well as the frequency of occurrence of giant planets. It will explore the possible transition region between giant planets and brown dwarfs, improving our understanding of star system formation. By determining accurate parallaxes for 4×10^7 stars, FAME will calibrate the absolute luminosities of solar neighborhood stars. This will determine the distances and ages of open and globular clusters, refine stellar structure models, and calibrate the distances to Cepheid and RR Lyrae stars. FAME will also monitor $\sim 40,000$ solar-like stars for photometric variations at the 0.1% level to search for evidence of magnetic activity cycles analogous to the 11-yr solar activity cycle.

FAME (<http://www.usno.navy.mil/fame>) will revolutionize variable star research by determining accurate distances to a large number of Cepheid and RR Lyrae stars, identifying a large number of variable stars, and improving our understanding of stellar structure and evolution.

¹U.S. Naval Observatory, 3450 Massachusetts Ave. NW, Washington, DC 20392-5420, USA

²U.S. Naval Observatory, PO Box 1149, Flagstaff, AZ 86002-1149, USA

³NASA Ames Research Center, M.S. 245-6, Moffett Field, CA 94035-1000, USA

⁴Naval Research Laboratory, Code 8100, 4555 Overlook Ave. SW, Washington, DC 20375-5000, USA

⁵Smithsonian Astrophysical Observatory, 60 Garden St., Cambridge, MA 02138, USA

⁶Lockheed Martin Missiles and Space Advanced Technology Center, 3251 Hanover St., Palo Alto, CA 94304-1191, USA




Figure 1. FAME will map our quadrant of the galaxy out to 2 kpc from the Sun providing the information needed to calibrate the standard candles that define the extragalactic distance scale, calibrate the absolute luminosities of stars of all spectral types for studies of stellar structure and evolution, and detect orbital motions caused by brown dwarfs and giant planets.




Figure 2. Comparison of the astrometric capabilities of FAME to SIM and Hipparcos. The lines indicate where achieved parallax accuracy will be 10% error or better. Known Cepheids are indicated by circles and RR Lyrae stars with 'x's. Distances to clusters and OB associations are indicated at the right of the figure.