

A new inventory of mountain glaciers and ice caps for the Antarctic periphery

Andrew BLISS,¹ Regine HOCK,^{1,2} J. Graham COGLEY³

¹*Geophysical Institute, University of Alaska Fairbanks, Fairbanks, AK, USA
E-mail: andybliss@gmail.com*

²*Department of Geosciences, Uppsala University, Uppsala, Sweden*

³*Department of Geography, Trent University, Peterborough, Ontario, Canada*

ABSTRACT. Although the glaciers in the Antarctic periphery make up a large fraction of all mountain glaciers and ice caps on Earth, a detailed glacier inventory of the region is lacking. We compile such an inventory, recording areas, area–altitude distributions, terminus characteristics and volume estimates. Glaciers on the mainland are excluded. The inventory is derived from the Antarctic Digital Database and some manual digitization. We additionally rely on satellite imagery, digital elevation models and a flowshed algorithm to classify ice bodies. We find 1133 ice caps and 1619 mountain glaciers covering a total of $132\,867 \pm 6643 \text{ km}^2$. Estimated total volume corresponds to $0.121 \pm 0.010 \text{ m}$ sea-level equivalent. Of the total glacier area, 99% drains either into ice shelves (63%) or into the ocean (36%). The inventory will provide a database for glacier mass-balance assessments, modelling and projections, and help to reduce the uncertainties in previous studies.

1. INTRODUCTION

Mountain glaciers and ice caps (henceforth referred to as glaciers) cover $\sim 740\,000 \text{ km}^2$ of the Earth's surface (Radić and Hock, 2010) and have been identified as a major contributor to sea-level rise (Meier and others, 2007). A large fraction of these glaciers are located in the Antarctic and sub-Antarctic, mostly around the Antarctic Peninsula. This region has undergone rapid environmental changes in recent decades. Exceptional rates of temperature increase (Vaughan and others, 2001; Turner and others, 2005; Steig and others, 2009) are concurrent with retreat of glacier fronts (Cook and others, 2005), an increase in melt areas (Vaughan, 2006), surface lowering (Pritchard and others, 2009) and rapid retreat, break-up and disintegration of ice shelves (Cook and Vaughan, 2010). Chen and others (2009) and Ivins and others (2011) estimated the mass changes of the entire Antarctic Peninsula based on the Gravity Recovery and Climate Experiment (GRACE) and found mass losses of $\sim 40 \text{ Gt a}^{-1}$ ($0.111 \text{ mm sea-level equivalent (SLE) a}^{-1}$) between 2003 and 2009. Hock and others (2009) found the rate of sea-level rise due to changes in the Antarctic periphery glaciers to be $0.22 \pm 0.16 \text{ mm SLE a}^{-1}$ between 1964 and 2004, or 28% of the estimated global contribution of glaciers excluding the ice sheets.

An extensive and detailed inventory of the glaciers on the peripheral islands of Antarctica, resolving individual glaciers and allowing an accurate estimate of total glacierized area, is essential for investigations of glacier mass balance at regional and larger scales. For lack of such an inventory, and for lack of mass-balance data, a number of studies have neglected the Antarctic periphery in their global-scale mass-balance assessments and projections (Cogley, 2005; Raper and Braithwaite, 2006).

Previous inventories have been limited either in spatial scope or breadth of data. Vinogradov and Psareva (1990) compiled a list of Antarctic islands with ice cover, but the list is incomplete and provides little beyond ice area and roughly estimated ice thickness. Rabassa and others (1982) compiled an inventory of James Ross Island and Vega Island,

off the northeast coast of the Antarctic Peninsula. Davies and others (2011) updated the inventory of James Ross Island and added the northern part of the Antarctic Peninsula. The Global Land Ice Measurements from Space project contains only the outlines from Davies and others (2011) and ice outlines for Kerguelen, and hence covers <3% of all glacier area in the periphery. Drawing on the Antarctic Digital Database (ADD; Fox and Cooper, 1994; ADD Consortium, 2000), Hock and others (2009) compiled a $1^\circ \times 1^\circ$ grid representing the glacierized areas of the Antarctic periphery. Distinguishing between glaciers and the ice sheet is not straightforward, either conceptually or practically, and depending on the purpose of a study different definitions may be appropriate. Hock and others (2009) defined 'Antarctic periphery' to include all islands between the mainland coast of Antarctica and the latitude of Kerguelen (49° S), excluding those near New Zealand and the mainland of South America (here we adopt this definition; see red and orange polygons in Fig. 1).

Published estimates of the area of glaciers in the Antarctic periphery vary between $132\,000 \text{ km}^2$ (Hock and others, 2009) and $169\,000 \text{ km}^2$ (Shumskiy, 1969). Differences are due to different reference dates but also different definitions of the domain. Based on Shumskiy's $169\,000 \text{ km}^2$, Radić and Hock (2010) estimated the volume of all Antarctic glaciers as $0.147 \pm 0.064 \text{ m SLE}$, representing 25% of global glacier ice volume (excluding the ice sheets). However, uncertainties are large since the estimate was derived by upscaling size distributions of data from the World Glacier Inventory (Cogley, 2008), which covered only a few percent of the glacierized area in the Antarctic periphery. These upscaled Antarctic size distributions were also used in global-scale mass change projections (Radić and Hock, 2011).

Here we compile an inventory of all glaciers in the Antarctic periphery. We distinguish between ice caps (dome-shaped ice bodies with radial flow, largely obscuring the subsurface topography) and mountain glaciers (glaciers that are confined by surrounding mountain terrain; Cogley and others, 2011) and present glacier size distributions,




Fig. 1. Classification of ice cover based on the Antarctic Digital Database (ADD), including the mountain glacier and ice-cap categories not distinguished in the ADD. Box 1 shows more detail for the Antarctic Peninsula. Boxes 2 (South Georgia and the South Sandwich Islands), 3 (Bouvet Island) and 4 (Kerguelen and Heard Island) are displaced from their proper positions but are at the same scale as the main map. The circled letters correspond to locations mentioned in the text: A. King George Island; B. Joinville Island; C. Vega Island; D. James Ross Island; E. Deception Island; F. Brabant Island; G. Anvers Island; H. Doumer Island; I. Liard Island (Fig. 3); J. Adelaide Island; K. South Orkney Islands; L. South Shetland Islands; M. Peter the First Island; N. Thurston Island (Fig. 2); O. Berkner Island; P. Roosevelt Island; Q. Cray Ice Rise; R. Ross Island; S. Coulman Island; T. Balleny Islands; and U. Znamenskiy Island. Scott Island (67.4° S, 179.9° W) and Marion Island (46.9° S, 37.7° E) do not show up at this scale. The map is drawn on a Lambert azimuthal equal-area projection centred at the South Pole.

area–altitude distributions, properties related to calving and an updated volume estimate based on volume–area scaling. This dataset has many possible uses. In particular, the improved area estimates and area–altitude distributions will lead to more accurate large-scale mass-balance assessments (of the past) and projections (for the future), since uncertainties are much less than in previous studies. This will also lead to better estimates of the glacier contribution to future sea-level change. Our total volume estimate can serve as a starting point for future volume change predictions. The new inventory could also be used to improve the representation of ice cover in climate models. The inventory shapefiles are available from <http://add.scar.org/> and hypsometry data are in the supplementary material hosted at <http://www.igsoc.org/hyperlink/63a377.html>

2. DATA AND METHODS

2.1. Glacier classification

Our new inventory (Fig. 1) is derived from many sources (Table 1). The main source is the ADD (Thomson and Cooper, 1993; Fox and Cooper, 1994; ADD Consortium, 2000). The ADD presents data at a number of spatial resolutions; we use ‘Scale0’, with the resolution of the best available source material. The data consist of polygons (shapefiles) representing coastlines and rock areas including nunataks south of 60° S. These shapefiles were digitized from paper maps, aerial photos and Landsat, MODIS

(Moderate Resolution Imaging Spectroradiometer) and ASTER (Advanced Spaceborne Thermal Emission and Reflection Radiometer) imagery with a range of dates from 1957 to 2005. Coastline polygons are tagged with one of the following surface-type attributes: land (encompassing both grounded ice and rock outcrops), ice shelf, ice tongue and ice rumple. We extract and identify all mountain glaciers and ice caps from the dataset as follows:

Polygons classified as ice shelves, ice tongues and ice rumples are not included in our inventory.

Following Hock and others (2009) we exclude all polygons on the Antarctic mainland. Many mainland glaciers resemble mountain glaciers, in particular along the Antarctic Peninsula, and may respond very differently to climate than the large Antarctic ice sheet. However, inspection of satellite images indicated that mountain glaciers on the mainland tend to share boundaries with the ice sheet, making classification difficult and highly subjective.

From the remaining coastline polygons, we erase all rock areas as defined by the rock polygons in the ADD. We assume non-rock areas are ice-covered, thus redefining the land class as a land ice class. The land ice class is further subdivided into ice rise, ice cap and mountain glaciers according to the steps below.

Ice rises, which are excluded from our inventory, are areas in the middle or along the seaward side of ice shelves




Fig. 2. Data used in classification for the Thurston Island region (a–c) and resulting surface classification (d), with the same colours as in Figure 1. (a) Landsat image from the Landsat Image Mosaic of Antarctica. (b) Topography from the RAMP DEM. (c) Flow direction from InSAR. See Figure 1 for location.

where the ice impinging on a grounded area appears to flow up and over the bedrock, rather than around it. Usually, ice rises do not have exposed rock nearby and their surface elevations are only slightly higher than the surrounding ice shelf. We used three datasets to identify ice rises (Fig. 2 shows an example). Flow and rock features were examined in Landsat imagery (Bindschadler and others, 2008), with a spatial resolution of 15 m. A velocity map (Rignot and others, 2011) also helped distinguish local flow around bedrock bumps. It was derived from interferometric synthetic aperture radar (InSAR) data acquired between 2007 and 2009 and optimized for display at the continental scale (spatial resolution 900 m), but it still proved useful for identifying ice rises. If the first two datasets did not yield a clear distinction, we considered the RADARSAT Antarctic Mapping Project digital elevation model version 2 (RAMP DEM; Liu and others, 2001). It has a grid resolution of 200 m and was constructed from radar altimetry and topographic maps with a variety of scales. Source data spanned the 1940s to 2001, and were mostly from the 1980s and 1990s. We

calculated the elevation difference from the summit of each candidate ice body to the surrounding ice shelf. If the difference was <200 m, the ice body was classified as an ice rise. Figure 2 shows examples of these cases.

Although their summits are >200 m above the surrounding ice-shelf surface, we exclude from our inventory all islands in the interior of the Ross, Filchner and Ronne ice shelves (e.g. Berkner Island, Roosevelt Island, Cray Ice Rise).

Ice is classified as mountain glacier if Landsat imagery revealed exposed-rock ridgelines subdividing the ice body. These ice bodies are separated into individual mountain glaciers, as described in Section 2.2. Some of the glaciers thus divided are reclassified as ice caps, based on their morphology.

The remaining land ice bodies are classified as ice caps. Diagnostic features include simple map geometry and ice flow outward from a central high elevation, as seen in the velocity map and RAMP DEM.

Table 1. Characteristics of the datasets we use to derive glacier outlines and area–altitude distributions. The ADD and RAMP cover a large percentage of the new inventory's area

Dataset	Resolution	Period	Source	Area %
<i>Glacier outlines</i>				
ADD	15 m to 1:1 000 000	1957–2005	ADD Consortium (2000)	94.3
Maps	1:15 000 to 1:200 000	1956–2000	See supplementary material	3.8
South Georgia GIS	1:200 000	2002, 2003	Cook and others (2010)	1.7
Système Pour l'Observation de la Terre (SPOT) image	20 m	1988	Not applicable	0.2
<i>Elevation*</i>				
RADARSAT Antarctic Mapping Project (RAMP)	200 m	1940s–2001	Liu and others (2001)	84.0
Updated ASTER Global Digital Elevation Model version 1 (GDEM)	100 m	2000–09	Cook and others (2012)	10.1
Shuttle Radar Topography Mission (SRTM) data	90 m	2000	Reuter and others (2007)	2.4
Topographic maps	1:20 000 to 1:200 000	1956–2000	See supplementary material	1.6
ASTER Global Digital Elevation Model version 2 (GDEM2)	30 m	2000–11	Fujisada and others (2012)	0.6
Composite Gazetteer of Antarctica [†]	Not applicable	Not applicable	Cervellati and others (2000)	0.5

*For the remaining area (0.8%) no data exist, so we substitute elevations from neighbouring islands. See Section 2.4 for details.

[†]These are simply maximum and minimum elevations. See Section 2.4 for details.




Fig. 3. Data used to create glacier divides for Liard Island (see Fig. 1 for location). (a) Landsat image with the ADD coastline (orange) and manually defined glacier terminus segments (between pairs of green dots). (b) RAMP DEM. (c) Glacier divides from the flowshed algorithm. White areas within the glacier are rock outcrops. The algorithm merged some adjacent glaciers.

For South Georgia, we use ice outlines from the South Georgia GIS (e.g. Cook and others, 2010). For the other sub-Antarctic islands (Peter the First Island, the South Shetland Islands, the South Orkney Islands, the South Sandwich Islands, Bouvet Island, Kerguelen, Heard Island and the Balleny Islands) we do not use the ADD but manually digitize the outlines of all glaciers, mainly from maps but in some cases from satellite imagery. The total area of these sub-Antarctic islands is 7540 km², ~6% of the inventory by area. Ice bodies are classified as ice caps or mountain glaciers following the same criteria as above. The geography of some of the sub-Antarctic islands remains poorly known even today. For example, the best map of the Balleny Islands is an explorer's sketch, and the islands are not covered adequately by any modern DEM, while maps of the South Sandwich Islands indicate elevations by crude form lines rather than contours.

2.2. Delineation of glacier divides

Where needed, we subdivide the ADD polygons into individual glaciers using a tool for automatic delineation of glacier flow basins (C. Kienholz and others, unpublished information). The tool uses a watershed algorithm commonly available in GIS software to calculate the area that is tributary to a manually defined set of points along each glacier's terminus (Fig. 3). In practice, the set of points is a segment of the whole polygon, and the end points of the segment are chosen visually. The DEM inputs to the tool are the same as described in Section 2.4. Manual correction of small polygon artefacts is required for some basins.

Glacier divides on South Georgia are derived from a more automated version of the same GIS tool described above. Where outlet glacier tongues are well defined, the tool is able to choose the terminus end points automatically.

Divides on the remaining sub-Antarctic islands are identified visually, relying on the contours on digitized maps. For some islands the best published maps have no contours; here the mapped ice cover is left undivided.

2.3. Terminus characteristics

A large fraction of the glaciers in our inventory terminate in the ocean, suggesting that frontal ablation (the sum of mass losses by calving and submarine melting) plays a large role in the mass budget of Antarctic periphery glaciers. As a step

towards quantifying frontal ablation, we assign all termini to one of five classes: no significant calving; terrestrial 'dry' calving; marine calving; lacustrine calving; and drainage into ice shelf. The first four correspond to the classification in Cogley (2008); the last is unique to Antarctica. Classification is done visually using imagery from Google EarthTM (as of 21 March 2012). If more than one category applies to a glacier, it is assigned to the class representing the longest part of its perimeter.

2.4. Hypsometries

Often, mass-balance models are elevation-dependent (e.g. Radić and Hock, 2011) and hence require the area–altitude distribution (hypsometry) of the glacier as input. To calculate hypsometries we primarily use digital elevation data from the RAMP DEM (Table 1). We supplement that source with Shuttle Radar Topography Mission data (Reuter and others, 2007) for South Georgia, Bouvet Island, Marion Island, Kerguelen and Heard Island. The ASTER Global Digital Elevation Model version 2 (GDEM2; Fujisada and others, 2012) covers the South Sandwich Islands and the South Orkneys. An improved version of GDEM (Cook and others, 2012) covers James Ross Island, the Joinville Islands, Brabant Island, Anvers Island, Adelaide Island and many smaller islands in the vicinity of the Antarctic Peninsula. We calculate the hypsometry of each glacier using 20 m elevation bins.

The RAMP DEM covers King George Island and Deception Island, but for the rest of the South Shetland Islands and Peter the First Island we generate hypsometries from map contours, interpolating to match the 20 m elevation bins of the rest of the dataset.

The remaining glaciers either have no DEM or the GDEM2 is too noisy to use. We assign hypsometries to these glaciers assuming their geometries can be approximated by a circular ice cap (with the measured area) with a surface profile defined by assuming ice to be a perfect plastic with yield stress of 50 kPa (Raper and Braithwaite, 2006; Cuffey and Paterson, 2010, eqn 8.133). While we use the shape of that surface profile, we adjust the maximum and minimum elevations to match the elevation data available. We take maximum and minimum elevations for the Balleny Islands (Young, Sturge, Buckle and Borradaile) from the Composite Gazetteer of Antarctica (Cervellati and

Table 2. Area of Antarctic periphery glaciers, as estimated in different studies

Study	Area km ²	Error estimate
Shumskiy (1969)	169 000 ± 68 000	Radić and Hock (2010)
Hock and others (2009)	132 000 ± 11 000	assuming an error of 8%
This study	132 867 ± 6643	assuming an error of 5%

others, 2000), a database of place names and mountain peak elevations. The total area of these islands is 661 km². For islands with no DEM data and no other elevation information, we estimate the maximum elevation based on nearby islands with data. The total area of these 20 islands is 984 km². Combined, glaciers without DEM data represent only 1.2% of the total area in the inventory (Table 1).

For comparison with other studies, we also calculate synthetic hypsometries for all the glaciers and ice caps in the inventory. For ice caps, we use the formulation described in the previous paragraph. Mountain glaciers typically have the most area in the middle of the glacier's elevation range. Following Raper and Braithwaite (2006), we give the area–altitude distribution a triangular shape, with maximum area at the elevation halfway between the minimum and maximum elevations. To evaluate the appropriateness of these shapes for the Antarctic periphery, we also normalize the elevation and area of each ice body's hypsometry and calculate the mean shape of all mountain glaciers in the inventory, and, separately, the mean shape of the ice caps.

2.5. Glacier area

We use a GIS to calculate the area of glaciers in the inventory, after projecting the outlines into a Lambert azimuthal equal-area projection centred at the South Pole. Our area estimate contains errors from several sources including the underlying maps and the digitizing process. Fox and Cooper (1994) suggest that individual outlines deviate from their true locations by <300 m, and that the total area of Antarctica calculated from the ADD has an error of <3%. The <3% value is based on an earlier estimate by Drewry and others (1982) who report <1% error in digitizing paper maps from both machine and operator random errors. Drewry and others (1982) estimate the total area error to be <3% due to the quality of underlying maps, and highly mobile boundaries in many areas. It is difficult to evaluate the error of the underlying maps, without expending considerable effort to create independent outlines from satellite imagery. Hock and others (2009) estimate area for the Antarctic periphery using the GGHYDRO 2.3 global hydrographic dataset (Cogley, 2003), which gives the percentage of glacierization in a 1° × 1° global grid. They adopted Cogley's single-cell accuracy error estimate of 8% for the error of the total area. The definition of the domain – whether to count certain ice bodies as part of the periphery – also impacts the total area estimate, but we do not consider this in our error estimate. Lacking detailed information for a more rigorous approach, here we assume a 5% error for the total area of our inventory (in line with, e.g., Paul and Andreassen, 2009; Bolch and others, 2010).

2.6. Glacier volume

Total ice volume was computed using volume–area scaling (Bahr and others, 1997):

$$V = cA^\gamma$$

where V is the volume and A the area of a single glacier and c and γ are scaling parameters. Following Radić and Hock (2010) we use $c = 0.2055 \text{ m}^{3-2\gamma}$ and $\gamma = 1.375$ for mountain glaciers in the inventory. Assuming a parabolic form of thickness–length relation, the volume of each ice cap is computed using $c = 1.7026 \text{ m}^{3-2\gamma}$ and $\gamma = 1.25$. We calculate the error of volume using error propagation for the equation above, assuming random independent errors for each term. We assume the error in each glacier's area is 5%. Following Radić and Hock (2010), we assume ±40% error in c and ±0.015 error in γ .

For comparison, we calculate volumes from the BEDMAP ice thickness dataset (Lythe and others, 2001) for 157 of the largest glaciers in the inventory. All these glaciers are larger than 50 km² and they account for ~90% of the inventory's total area. BEDMAP is constructed from airborne and ground-based radar surveys, seismic reflection and gravity measurements. BEDMAP has a coarse grid resolution (5 km), and the data that went into it are quite sparse for some glaciers, but it is currently the best available ice thickness dataset that covers the area of the inventory. Lythe and others (2001) estimate the root-mean-square error of grid elevations for the Antarctic Peninsula to be 163 m, which we apply to all gridcells for our error analysis. We use volume–area scaling to fill in the volume of glaciers not covered by BEDMAP. The scaling parameters are tuned to match the BEDMAP measurements with no distinction between mountain glaciers and ice caps: $c = 0.2187 \text{ m}^{3-2\gamma}$ and $\gamma = 1.341$.

3. RESULTS AND DISCUSSION

3.1. Number and area of glaciers

In total we identify 1133 ice caps covering 91 042 ± 4552 km² and 1619 mountain glaciers covering 41 825 ± 2091 km². Hence our glacier inventory includes 132 867 ± 6643 km² of ice on Antarctic and sub-Antarctic islands. Our estimate of the total area of glaciers in the Antarctic periphery is similar to that of Hock and others (2009; Table 2). This is not surprising as they also used the ADD as a basis for their study. The difference is due to slightly different definitions of the domain, mainly with regard to ice rises. Our estimate is less than that of Shumskiy (1969), whose methods are not explained and are thus not reproducible. We also made an independent estimate of area using island coastlines extracted from the MODIS Mosaic of Antarctica (Scambos and others, 2007), again excluding the large islands in the Ross and Ronne–Filchner Ice Shelves. The resulting area, 132 272 km², is close to the estimate from our new inventory, giving us confidence in our area estimate.

Our estimate for number of glaciers must be treated with caution since it strongly depends on the definition of an individual glacier. Separating contiguous glaciers is complicated in the Antarctic periphery. Indeed, it is often a matter of subjective judgment. Some ice bodies resembling ice caps have pronounced outlet glaciers; their longitudinal profiles are convexo-concave, unlike the convex-up profiles of ice caps with well-developed domal morphology or the concave-up profiles of many mountain glaciers. Our classification does not contain a separate class of outlet glaciers, nor




Fig. 4. Histogram showing the size distributions of ice caps and mountain glaciers in the new inventory: (a) number of glaciers and (b) fraction of total area per size bin of $0.2 \log_{10}(\text{km}^2)$ vs area. The bars are stacked.

does it rest on an objective criterion for distinguishing between ice caps and mountain glaciers. Nevertheless, our estimate combined with the area of each ice body provides a first approximation for studies applying volume–area scaling for assessments of ice volume or projections of mass changes. For these purposes, the power law that relates volume to area requires that the area of each individual glacier, or at least the frequency distribution of glacier sizes, be known.

3.2. Size distribution

Figure 4 shows the size distribution of all glaciers in the inventory. Glacier sizes range from 0.01 to 6000 km². The eight largest glaciers in the inventory (>3000 km² each) cover 25% of the total area. Most of the glaciers larger than 100 km² have a surface morphology that led us to classify them as ice caps. In some cases, a single island has more than one ice cap. There are more mountain glaciers than ice caps for areas between about 0.5 and 100 km². The inventory contains a large number of small (<1 km²) ice-covered islands, most of which are observed to have an ice-cap morphology, but their combined area is only 359 km².

3.3. Terminus characteristics

More than 99% of the glacierized area drains through marine-terminating termini or into ice shelves (Table 3). In terms of area (and implicitly volume), more of the ice from the peripheral islands is discharging into ice shelves than directly into the ocean. By number of glaciers, marine calving is the most common style, but this category includes many small islands, so the total area is less than for ice

shelves. Some of the marine-calving termini are of classical type, with glacier ice standing in water, but a common form of terminus is one in which the glacier ends either at the high-water mark or sufficiently close to it that calved ice may float away. An example is the Costa Recta on Deception Island, illustrated by López-Martinez and Serrano (2002). Over 1200 glaciers end on land with no significant calving; these glaciers also tend to be small. The terrestrial calving glaciers, found only on Coulman Island, Znamenskiy Island and Bouvet Island, usually end above rocky bluffs, such that the calved ice quickly falls into the ocean, but the ocean does not contact the glacier. Of glaciers calving into lakes, there is one on Signy Island in the South Orkneys, six on Kerguelen and four on South Georgia. There are also three glaciers terminating in coastal freshwater lagoons on Heard Island and three more on South Georgia. Except for Ampère Glacier on Kerguelen (Berthier and others, 2009), there is no information on the significance of lacustrine calving as a component of the mass budget of these glaciers.

3.4. Hypsometries

Altitudes of all glaciers in the inventory range from 0 to 3600 m a.s.l. The highest point in the region is on Mount Erebus, Ross Island. By numbers, 66% of the glaciers terminate within 50 m of sea level. Approximately 94% of the glacierized area in the Antarctic periphery is within 1000 m of sea level, and 74% is within 500 m (Fig. 5a). Ice caps have 88% of their total area below 500 m a.s.l. while the corresponding number for mountain glaciers is 44% (Fig. 5b). Above 1000 m a.s.l., ice caps have only 1584 km² (2% of their total area), while mountain glaciers have 6858 km² (16%). Mountain glaciers tend to have a larger proportion of their area at higher elevations, because the ice is partially supported by drag along valley walls (Cuffey and Paterson, 2010) and the confining walls are not conducive to the ice-cap morphology.

Due to lack of inventory data, Radić and Hock (2011) used synthetic hypsometries and glacier areas to model future glacier response to climate change. The synthetic hypsometries we calculate for our inventory make use of the maximum and minimum elevation data that we generate, so they are an improvement over Radić and Hock (2011), but still do not exactly match the DEM-derived hypsometries (Fig. 5b). Within 1000 m of sea level, the measured ice-cap hypsometries have 6% more area than the synthetic (23%

Table 3. Terminus characteristics of Antarctic periphery glaciers

Terminus characteristics	Number of glaciers	Area	
		km ²	%
Discharge into ice shelf	165	83 138	62.6
Marine calving	1339	48 053	36.2
No calving	1220	1111	0.8
Lacustrine calving	17	316	0.2
Terrestrial ('dry') calving	11	249	0.2
Total	2752	132 867	100




Fig. 5. (a) Cumulative area with increasing elevation. (b) Combined hypsometry for all glaciers and ice caps in the inventory, with 20 m elevation bands. Solid lines show the hypsometry calculated from DEM data; dashed lines show the synthetic hypsometry calculated just from maximum and minimum elevations of individual ice bodies. The DEM hypsometry has spikes every 250 m as a result of imperfect interpolation when the RAMP DEM was created from map contours.

for mountain glaciers). Above 1000 m, the synthetic hypsometries have 4.2 times more area than the DEM-derived hypsometries for ice caps (2.2 times for glaciers). The greater frequency of high elevations in the synthetic dataset would bias the results of mass-balance modelling towards higher values.

The mean shape of the normalized measured hypsometries is considerably different from the standard synthetic hypsometries (Fig. 6). The observed hypsometries of mountain glaciers and ice caps are similar. Both have a relative excess of area in the lowest 20% of the elevation range, fairly constant area between 20% and 90%, and diminishing area above 90%. The extra area at low elevation is often an ice piedmont – where a glacier flows out of the mountains onto a coastal plain. The reduced area at high elevations is a function of steep mountain topography in the glacier case, or the small radius of the summit of an ice cap. Antarctic glaciers differ from the patterns assumed by Raper and Braithwaite (2006), perhaps because many terminate in the sea, instead of on land as would be typical of glaciers elsewhere in the world.




Fig. 6. Normalized hypsometries averaged over all ice caps and mountain glaciers based on DEMs and synthetic hypsometries following Raper and Braithwaite (2006). There are 100 elevation bins, so a rectangular sloping plane would plot as a vertical line at 1%.

3.5. Glacier volumes

Our best estimate of total glacier volume for the Antarctic periphery is $48\,636 \pm 3849 \text{ km}^3$, corresponding to $0.121 \pm 0.010 \text{ m SLE}$. We assume a glacier density of 900 kg m^{-3} and an oceanic area of $3.625 \times 10^8 \text{ km}^2$ (Cogley, 2012). The total volume is the sum of $17\,428 \pm 2254 \text{ km}^3$ in glaciers and $31\,208 \pm 3121 \text{ km}^3$ in ice caps. If all the ice bodies were ice caps, volume–area scaling would give $42\,194 \text{ km}^3$. If they were all mountain glaciers, the volume would be $72\,297 \text{ km}^3$. Using the BEDMAP thickness data, we find a volume of $38\,869 \pm 2639 \text{ km}^3$. We consider the BEDMAP volume to be an underestimate because ice thickness measurements are sparse while rock outcrops (assumed to have zero ice thickness) are completely mapped. Our volume estimates correspond to a domain-average thickness (total volume divided by total area) of 417 m for mountain glaciers and 343 m for ice caps. Mean thicknesses differ for glaciers and ice caps due to different parameters in the scaling relationship.

Our best volume estimate is 18% lower than the estimate by Radić and Hock (2010; $59\,521 \pm 25\,829 \text{ km}^3$), largely because of our 21% lower area estimate. Additionally, unlike Radić and Hock, we know the number of glaciers in different size ranges. Our error estimate has been reduced accordingly.

4. CONCLUSIONS

Our new inventory of Antarctic periphery glaciers contains $132\,867 \pm 6643 \text{ km}^2$ of ice. The inventory includes grounded ice that is separated from the mainland by open water or ice shelf, but does not include the islands contained within the Ross and Filchner–Ronne Ice Shelves. We have classified ice bodies by their morphology (ice cap vs mountain glacier). Based on volume–area scaling, we estimate the ice volume in the Antarctic periphery to be $48\,636 \pm 3849 \text{ km}^3$. We find that 63% of the glacier area in the inventory discharges into ice shelves, while 36% calves ice directly into the ocean. The remaining 1% terminates either on land or in a lake. In contrast to glaciers in other regions, we find that Antarctic glaciers tend to have a rather uniform distribution of area across elevations, but a disproportionately large fraction in the lowest 10% of their

elevation range. The inventory is available on the ADD website (<http://add.scar.org>). The associated hypsometry files and a list of data sources used are in the supplementary material at <http://www.igsoc.org/hyperlink/63a377.html>

The inventory is a major advance over what was previously available. Further improvements to the inventory could be made by digitizing a consistent set of outlines from satellite imagery. A time series of outlines would allow studies of glacier change. To improve the hypsometries, better DEMs are needed. Careful separation of adjacent glaciers will increase the accuracy of volume estimates from volume–area scaling. Additional ice thickness measurements from the peripheral islands are needed to better estimate the total ice volume and validate the volume–area scaling results.

ACKNOWLEDGEMENTS

This work was supported by grants from NASA (NNX11AO23G) and the US National Science Foundation (ANT-1043649). We thank Christian Kienholz for providing his glacier delineation code and assisting with its application, Justin Rich for providing code to calculate hypsometries, and Valentina Radić for helpful discussions.

REFERENCES

- ADD Consortium (2000) *Antarctic Digital Database, Version 3.0, database, manual and bibliography*. Scientific Committee on Antarctic Research, Cambridge
- Bahr DB, Meier MF and Peckham SD (1997) The physical basis of glacier volume–area scaling. *J. Geophys. Res.*, **102**(B9), 20 355–20 362 (doi: 10.1029/97JB01696)
- Berthier E, Le Bris R, Mabileau L, Testut L and Rémy F (2009) Ice wastage on the Kerguelen Islands (49°S, 69°E) between 1963 and 2006. *J. Geophys. Res.*, **114**(F3), F03005 (doi: 10.1029/2008JF001192)
- Bindschadler R and 8 others (2008) The Landsat image mosaic of Antarctica. *Remote Sens. Environ.*, **112**(12), 4214–4226 (doi: 10.1016/j.rse.2008.07.006)
- Bolch T, Menounos B and Wheate R (2010) Landsat-based inventory of glaciers in western Canada, 1985–2005. *Remote Sens. Environ.*, **114**(1), 127–137 (doi: 10.1016/j.rse.2009.08.015)
- Cervellati R, Ramorino C, Sievers J, Thomson J and Clarke D (2000) A composite gazetteer of Antarctica. *Polar Rec.*, **36**(198), 278–285 (doi: 10.1017/S0032247400016739)
- Chen JL, Wilson CR, Blankenship D and Tapley BD (2009) Accelerated Antarctic ice loss from satellite gravity measurements. *Nature Geosci.*, **2**(12), 859–862 (doi: 10.1038/ngeo694)
- Cogley JG (2003) *GGHYDRO: Global Hydrographic Data, Release 2.3*. Department of Geography, Trent University, Peterborough, Ont. (Trent Technical Note 2003-1) <http://people.trentu.ca/~gcogley/glaciology/gghrls231.pdf>
- Cogley JG (2005) Mass and energy balances of glaciers and ice sheets. In Anderson MG ed. *Encyclopaedia of hydrological sciences. Part 14. Snow and glacier hydrology*. Wiley, New York, 2555–2573
- Cogley JG (2008) *Extended format for the World Glacier Inventory, revised edn*. Department of Geography, Trent University, Peterborough, Ont. (Trent Technical Note 2005-1) <http://people.trentu.ca/~geogley/glaciology/ttn200501.pdf>
- Cogley JG (2012) Area of the ocean. *Mar. Geod.*, **35**(4), 379–388 (doi: 10.1080/01490419.2012.709476)
- Cogley JG and 10 others (2011) *Glossary of glacier mass balance and related terms*. UNESCO–International Hydrological Programme, Paris (IHP-VII Technical Documents in Hydrology 86)
- Cook AJ and Vaughan DG (2010) Overview of areal changes of the ice shelves on the Antarctic Peninsula over the past 50 years. *Cryosphere*, **4**(1), 77–98 (doi: 10.5194/tc-4-77-2010)
- Cook AJ, Fox AJ, Vaughan DG and Ferrigno JG (2005) Retreating glacier fronts on the Antarctic Peninsula over the past half-century. *Science*, **308**(5721), 541–544 (doi: 10.1126/science.1104235)
- Cook AJ, Poncet S, Cooper APR, Herbert DJ and Christie D (2010) Glacier retreat on South Georgia and implications for the spread of rats. *Antarct. Sci.*, **22**(3), 255–263 (doi: 10.1017/S0954102010000064)
- Cook AJ, Murray T, Luckman A, Vaughan DG and Barrand NE (2012) A new 100-m Digital Elevation Model of the Antarctic Peninsula derived from ASTER Global DEM: methods and accuracy assessment. *Earth Syst. Sci. Data Discuss.*, **5**(1), 365–403 (doi: 10.5194/essdd-5-365-2012)
- Cuffey KM and Paterson WSB (2010) *The physics of glaciers*, 4th edn. Butterworth-Heinemann, Oxford
- Davies BJ, Carrivick JL, Glasser NF, Hambrey MJ and Smellie JL (2011) A new glacier inventory for 2009 reveals spatial and temporal variability in glacier response to atmospheric warming in the Northern Antarctic Peninsula, 1988–2009. *Cryos. Discuss.*, **5**(6), 3541–3595 (doi: 10.5194/tcd-5-3541-2011)
- Drewry DJ, Jordan SR and Jankowski E (1982) Measured properties of the Antarctic ice sheet: surface configuration, ice thickness, volume and bedrock characteristics. *Ann. Glaciol.*, **3**, 83–91
- Fox AJ and Cooper APR (1994) Measured properties of the Antarctic ice sheet derived from the SCAR Antarctic digital database. *Polar Rec.*, **30**(174), 201–206
- Fujisada H, Urai M and Iwasaki A (2012) Technical methodology for ASTER Global DEM. *IEEE Trans. Geosci. Remote Sens.*, **50**(10), 3725–3736 (doi: 10.1109/TGRS.2012.2187300)
- Hock R, de Woul M and Radić V (2009) Mountain glaciers and ice caps around Antarctica make a large sea-level rise contribution. *Geophys. Res. Lett.*, **36**(7), L07501 (doi: 10.1029/2008GL037020)
- Ivins ER, Watkins MM, Yuan D-N, Dietrich R, Casassa G and Rülke A (2011) On-land ice loss and glacial isostatic adjustment at the Drake Passage: 2003–2009. *J. Geophys. Res.*, **116**(B2), B02403 (doi: 10.1029/2010JB007607)
- Liu H, Jezek K, Li B and Zhao Z (2001) *Radarsat Antarctic Mapping Project digital elevation model version 2*. National Snow and Ice Data Center, Boulder, CO. Digital media: <http://nsidc.org/data/nsidc-0082.html>.
- López-Martínez J and Serrano E (2002) Geomorphology. In Smellie JL and 9 others eds. *Geology and geomorphology of Deception Island*. British Antarctic Survey, Cambridge, 31–39 (BAS Geomap Series Sheets 6-A and 6-B)
- Lythe MB, Vaughan DG and BEDMAP consortium (2001) BEDMAP: a new ice thickness and subglacial topographic model of Antarctica. *J. Geophys. Res.*, **106**(B6), 11 335–11 351 (doi: 10.1029/2000JB900449)
- Meier MF and 7 others (2007) Glaciers dominate eustatic sea-level rise in the 21st century. *Science*, **317**(5841), 1064–1067 (doi: 10.1126/science.1143906)
- Paul F and Andreassen LM (2009) A new glacier inventory for the Svartisen region, Norway, from Landsat ETM+ data: challenges and change assessment. *J. Glaciol.*, **55**(192), 607–618 (doi: 10.3189/002214309789471003)
- Pritchard HD, Arthern RJ, Vaughan DG and Edwards LA (2009) Extensive dynamic thinning on the margins of the Greenland and Antarctic ice sheets. *Nature*, **461**(7266), 971–975 (doi: 10.1038/nature08471)
- Rabassa J, Skvarca P, Bertani L and Mazzoni E (1982) Glacier inventory of James Ross and Vega Islands, Antarctic Peninsula. *Ann. Glaciol.*, **3**, 260–264
- Radić V and Hock R (2010) Regional and global volumes of glaciers derived from statistical upscaling of glacier inventory data. *J. Geophys. Res.*, **115**(F1), F01010 (doi: 10.1029/2009JF001373)

- Radić V and Hock R (2011) Regionally differentiated contribution of mountain glaciers and ice caps to future sea-level rise. *Nature Geosci.*, **4**(2), 91–94 (doi: 10.1038/ngeo1052)
- Raper SCB and Braithwaite RJ (2006) Low sea level rise projections from mountain glaciers and icecaps under global warming. *Nature*, **439**(7074), 311–313 (doi: 10.1038/nature04448)
- Reuter HI, Nelson A and Jarvis A (2007) An evaluation of void-filling interpolation methods for SRTM data. *Int. J. Geogr. Inf. Sci.*, **21**(9), 983–1008 (doi: 10.1080/13658810601169899)
- Rignot E, Mouginot J and Scheuchl B (2011) Ice flow of the Antarctic Ice Sheet. *Science*, **333**(6048), 1427–1430 (doi: 10.1126/science.1208336)
- Scambos TA, Haran TM, Fahnestock MA, Painter TH and Bohlander J (2007) MODIS-based Mosaic of Antarctica (MOA) data sets: continent-wide surface morphology and snow grain size. *Remote Sens. Environ.*, **111**(2–3), 242–257 (doi: 10.1016/j.rse.2006.12.020)
- Shumskiy PA (1969) Oledenenie [Glaciation]. In Tolstikov EI ed. *Atlas Antarktiki II [Atlas of Antarctica, vol. II]*. Gidrometeoizdat, Leningrad, 367–400 [in Russian]
- Steig EJ, Schneider DP, Rutherford SD, Mann ME, Comiso JC and Shindell DT (2009) Warming of the Antarctic ice-sheet surface since the 1957 International Geophysical Year. *Nature*, **457**(7228), 459–462 (doi: 10.1038/nature07669)
- Thomson JW and Cooper APR (1993) The SCAR Antarctic digital topographic database. *Antarct. Sci.*, **5**(3), 239–244
- Turner J and 8 others (2005) Antarctic climate change during the last 50 years. *Int. J. Climatol.*, **25**(3), 279–294 (doi: 10.1002/joc.1130)
- Vaughan D (2006) Recent trends in melting conditions on the Antarctic Peninsula and their implications for ice-sheet mass balance and sea level. *Arct. Antarct. Alp. Res.*, **38**(1), 147–152 (doi: 10.1657/1523-0430(2006)038[0147:RTIM-CO]2.0.CO;2)
- Vaughan DG, Marshall GJ, Connolley WM, King JC and Mulvaney R (2001) Climate change: devil in the detail. *Science*, **293**(5536), 1777–1779 (doi: 10.1126/science.1065116)
- Vinogradov ON and Psareva TV (1990) Present-day glacier cover on the peri-Antarctic Islands. *Polar Geogr. Geol.*, **14**(4), 249–260 (doi: 10.1080/10889379009377437)