
INSTRUCTION TO CONTRIBUTORS

The *Journal of Helminthology* publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance.

Manuscripts, which must be in English or French (with an English summary), should be addressed to:

The Editor, Journal of Helminthology,
London School of Hygiene and Tropical Medicine,
Winches Farm Field Station,
395 Hatfield Road,
St Albans, Herts AL4 0XQ,
England.

Two copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief Introduction; Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Research Notes should also be preceded by a brief abstract. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration may be printed is 12.0 × 18.5 cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space, should be indicated. Some indication of scale should normally be given on the figure. Photocopies of illustrations should be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints of the same size as they are to appear in the Journal (maximum size 12.0 × 18.5 cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

DUKE, B. O. L. (1971) The ecology of onchocerciasis in man and animals. In: *Ecology and Physiology of parasites* (editor, A. M. Fallis) pp. 213–222. Adam Hilger Ltd.: London.

JAMES, C. & WEBBE, G. (1973) A comparison of Egyptian and East African strains of *Schistosoma haematobium*. *Journal of Helminthology*, 47, 49–59.

25 offprints are provided free of charge; additional copies may be ordered at the proof stage.

Contents

	Pages
Epidemiological observations on stomach worms of horses in Morocco. V. S. PANDEY, H. OUHELLI and A. ELKHALFANE	155-160
The transmission of <i>Onchocerca tarsicola</i> (Filarioidea: Onchocercidae) by <i>Odagmia ornata</i> and <i>Prosimulium nigripes</i> (Diptera: Simuliidae). H. SCHULZ-KEY and P. WENK	161-166
Pathological phenomena associated with <i>Mesocestoides corti</i> infection in mice. (Research Note) T. R. WHITE, R. C. A. THOMPSON, D. A. PASS, W. J. PENHALE and J. STEPHANSON	167-171
Researches on Pseudophyllidea (Carus, 1813) in the south of Chile. IV Occurrence of <i>Diphyllobothrium dendriticum</i> (Nitzsch). PATRICIO TORRES, RENE FRANJOLA, LUIS FIGUEROA, ROBERTO SCHLATTER, HECTOR GONZALEZ, BORIS CONTRERAS and RODOLGO MARTIN	173-187
Limited fly load and development of <i>Onchocerca volvulus</i> microfilariae in Guatemalan <i>Simulium ochraceum</i> . Y. HASHIGUCHI, M. KAWABATA, S. ITO and M. M. RECINOS C.	189-196
Experience in Northern Nigeria with countercurrent immunoelectrophoresis, double diffusion and indirect haemagglutination tests for diagnosis of hydatid cyst in camels. D. J. O. DADA, D. S. ADEGBOYE and A. N. MOHAMMED	197-202
The behaviour of infective <i>Ancylostoma caninum</i> larvae in serum gradients. M. A. ZIETSE, J. C. M. KLAVER-WESSELING and J. C. M. VETTER	203-207
The growth and antigenicity of the metacestodes of <i>Taenia crassiceps</i> in several different strains of rats. J. CHERNIN	209-222
Cytochemical characteristics of the neurosecretory cells of <i>Ceylonocotyle scoliocoelium</i> (Trematoda: Digenea). P. N. SHARMA and A. N. SHARMA	223-229
Stereoscan observations on the surface topography of <i>Gastrothylax crumenifer</i> (Creplin, 1847) Poirier, 1883 and <i>Paramphistomum epiclitum</i> Fischoeder, 1904 (Trematoda: Digenea). VEENA TANDON and S. C. MAITRA	231-237