



On Hadwiger’s covering problem in small dimensions

Andrii V. Arman, Andriy Viktorovych Bondarenko, and Andriy Prymak

Abstract. Let H_n be the minimal number such that any n -dimensional convex body can be covered by H_n translates of the interior of that body. Similarly H_n^s is the corresponding quantity for symmetric bodies. It is possible to define H_n and H_n^s in terms of illumination of the boundary of the body using external light sources, and the famous Hadwiger’s covering conjecture (illumination conjecture) states that $H_n = H_n^s = 2^n$. In this note, we obtain new upper bounds on H_n and H_n^s for small dimensions n . Our main idea is to cover the body by translates of John’s ellipsoid (the inscribed ellipsoid of the largest volume). Using specific lattice coverings, estimates of quermassintegrals for convex bodies in John’s position, and calculations of mean widths of regular simplexes, we prove the following new upper bounds on H_n and H_n^s : $H_5 \leq 933$, $H_6 \leq 6137$, $H_7 \leq 41377$, $H_8 \leq 284096$, $H_4^s \leq 72$, $H_5^s \leq 305$, and $H_6^s \leq 1292$. For larger n , we describe how the general asymptotic bounds $H_n \leq \binom{2n}{n}n(\ln n + \ln \ln n + 5)$ and $H_n^s \leq 2^n n(\ln n + \ln \ln n + 5)$ due to Rogers, Shephard and Roger, Zong, respectively, can be improved for specific values of n .

1 Introduction

Let \mathcal{K}_n be the family of all convex bodies in \mathbb{R}^n , i.e., all convex compact sets $K \subset \mathbb{R}^n$ with a nonempty interior ($\text{int}(K) \neq \emptyset$). For $A, B \subset \mathbb{R}^n$, we denote by

$$C(A, B) := \min \left\{ N : \exists t_1, \dots, t_N \in \mathbb{R}^n \text{ satisfying } A \subseteq \bigcup_{j=1}^N (t_j + B) \right\},$$

the minimal number of translates of B needed to cover A .

Hadwiger [18] raised the question of determining the value of

$$H_n = \max \{ C(K, \text{int}(K)) : K \in \mathcal{K}_n \}$$

for all $n \geq 3$.

Received by the editors March 30, 2024; revised December 13, 2024; accepted March 28, 2025.

Published online on Cambridge Core April 4, 2025.

The first author was supported by the University of Manitoba Research Grant Program project “Covering problems with applications in Euclidean Ramsey theory and convex geometry” and PIMS PDF. The second author was supported by grant 334466 of the Research Council of Norway “Fourier Methods and Multiplicative Analysis.” The third author was supported by NSERC of Canada Discovery Grant RGPIN-2020-05357.

AMS subject classification: 52A20, 52A39, 52A40, 52C07, 52C17.

Keywords: Illumination problem, illumination number, covering by smaller homothetic copies, John’s position, quermassintegrals.



Considering an n -cube, one immediately sees that $H_n \geq 2^n$, and the well-known Hadwiger's covering conjecture states that $H_n = 2^n$ for $n \geq 3$ with equality only for parallelepipeds. It was shown that $H_2 = 4$ by Levi [24]. Aside from Levi and Hadwiger, the conjecture may be associated with the names of Boltyanski, who in [8] has established an equivalent formulation in terms of illumination of the boundary of the body by external light sources, and with Gohberg–Markus [16] who asked the question in terms of the minimal number of smaller homothetic copies of K required to cover K . As of today, the conjecture, which is also known as the illumination conjecture, is wide open. For details about the history and partial results for special classes of convex bodies see, e.g., [6]. For general background on convex geometry one can refer to, e.g., [33] or to the introductory sections of [2] or [9].

In what follows, we will outline the current approaches in obtaining the upper bounds on H_n and our modifications allowing to obtain new bounds.

To this end, let \mathcal{K}_n^s be the subfamily of \mathcal{K}_n consisting of centrally symmetric convex bodies, and define

$$H_n^s = \max\{C(K, \text{int}(K)) : K \in \mathcal{K}_n^s\}.$$

The best known *explicit* upper bound on H_n in high dimensions follows from the results of Rogers and Zong [30] (which, in turn, use the Rogers–Shephard inequality [29]):

$$(1.1) \quad H_n \leq \binom{2n}{n} n (\ln n + \ln \ln n + 5).$$

This bound is valid for each dimension $n \geq 3$, so it can be used for our settings of specific small n . We remark that a similar estimate was first obtained by Erdős and Rogers [13], who showed (1.1) with $4 + \frac{1}{n}$ in place of 5, but only for sufficiently large n . Now the asymptotic behavior of $\binom{2n}{n}$ is $\frac{4^n}{\sqrt{2\pi n}}$, so this upper bound is $(4 + o(1))^n$ while the conjecture states $H_n = 2^n$. Remarkable sub-exponential improvements of (1.1) have been obtained only recently by Huang, Slomka, Tkocz, and Vritsiou [19]. Namely, using “thin-shell” volume estimates they obtained

$$(1.2) \quad H_n \leq \exp(-c\sqrt{n})4^n.$$

This was later improved, using the new breakthrough bounds on the isotropic constant, by Campos, van Hintum, Morris, and Tiba [10] to

$$(1.3) \quad H_n \leq \exp\left(\frac{-cn}{\ln n}\right)4^n.$$

Both constants c in (1.2) and (1.3) are independent of n but not given explicitly. For centrally symmetric bodies, one has (see [13] and [30])

$$(1.4) \quad H_n^s \leq 2^n n (\ln n + \ln \ln n + 5),$$

which is asymptotically close to the conjectured 2^n .

The first step in obtaining any of the inequalities (1.1)–(1.4) is the result of Rogers [28] on the covering density of \mathbb{R}^n by translates of a body L . Namely, let

$$\theta(L) = \lim_{R \rightarrow \infty} \inf_{\Lambda} \left\{ \frac{|L| \# \Lambda}{(2R)^n} : [-R, R]^n \subseteq \bigcup_{\lambda \in \Lambda} (x_\lambda + L) \right\}$$

denote the covering density of space by translates of L , where $|L|$ is the volume of L , and $\# \Lambda$ is the cardinality of Λ . Rogers [28] showed that for any body $L \in \mathcal{K}_n$,

$$(1.5) \quad \theta(L) \leq n(\ln n + \ln \ln n + 5).$$

The main idea of [28] is that an appropriate number of random translates of L covers “most” of the space, and the leftover can be handled using a maximum packing argument.

The second step is the use of the inequality

$$(1.6) \quad C(K, L) \leq \frac{|K - L|}{|L|} \theta(L),$$

which is valid for any $K, L \in \mathcal{K}^n$ and was proved by Rogers and Zong [30] using certain averaging argument. (Here $K - L = \{x - y : x \in K, y \in L\}$ is the Minkowski difference.)

For symmetric bodies, one obtains (1.4) from (1.6) by choosing $L = (1 - \varepsilon)K$ with $\varepsilon \rightarrow 0+$ and using the fact that $|K - K| = |2K| = 2^n |K|$ for $K \in \mathcal{K}_n^s$. For general bodies, (1.1) follows from (1.6) by applying the Rogers–Shephard inequality (see [29]), which, for any $K \in \mathcal{K}_n$, estimates the volume of the difference body $K - K$ by $|K - K| \leq \binom{2n}{n} |K|$.

The improvements (1.2) and (1.3) are both based on choosing L to be the largest (by volume) centrally symmetric subset of K . With such choice, we have $|K - L|/|L| \leq |2K|/|L| \leq 2^n \Delta_{KB}(K)^{-1}$, where

$$\Delta_{KB}(K) := \max_{x \in \mathbb{R}^n} \frac{|K \cap (x - K)|}{|K|}$$

is the Kövner–Besicovitch measure of symmetry of K . Simple averaging gives $\Delta_{KB}(K) \geq 2^{-n}$ for any $K \in \mathcal{K}_n$, while [19] and [10] obtain the corresponding sub-exponential improvements to the lower bound on $\Delta_{KB}(K)$, yielding (1.2) and (1.3).

For low dimensions, Lassak [22] showed that

$$(1.7) \quad H_n \leq (n+1)n^{n-1} - (n-1)(n-2)^{n-1},$$

which outperforms (1.1) for $n \leq 5$. For $n = 3$ this gives $H_3 \leq 34$, which was improved to $H_3 \leq 20$ by Lassak [23], then to $H_3 \leq 16$ by Papadoperakis [25], and then to $H_3 \leq 14$ by Prymak [26]. For slightly larger dimensions, it was shown in [27] that $H_4 \leq 96$, $H_5 \leq 1091$ and $H_6 \leq 15373$ improving both (1.1) and (1.7) for $n = 4, 5, 6$. Then Diao [12] obtained $H_5 \leq 1002$ and $H_6 \leq 14140$. All of these results in low dimensions were based on comparing the body with a suitable parallelepiped. For the symmetric case, Lassak [21] obtained the sharp result $H_3^s = 8$, but for $n \geq 4$ no estimate better than (1.4) or than the corresponding bound on H_n (obviously, $H_n^s \leq H_n$) was known.

Our main result is the following new bounds.

Theorem 1.1 $H_5 \leq 933$, $H_6 \leq 6137$, $H_7 \leq 41377$, $H_8 \leq 284096$, $H_4^s \leq 72$, $H_5^s \leq 305$, and $H_6^s \leq 1292$.

Table 1: Best known upper bounds on H_n for $3 \leq n \leq 14$.

n	$H_n \leq$	Reference	n	$H_n \leq$	Reference
3	14	[26]	9	2 064 332	Prop. 5.1
4	96	[27]	10	8 950 599	Prop. 5.1
5	933	Th. 1.1	11	38 482 394	Prop. 5.1
6	6 137	Th. 1.1	12	164 319 569	Prop. 5.1
7	41 377	Th. 1.1	13	697 656 132	Prop. 5.1
8	284 096	Th. 1.1	14	2 947 865 482	Prop. 5.1

Table 2: Best known upper bounds on H_n^s for $3 \leq n \leq 14$.

n	$H_n^s \leq$	Reference	n	$H_n^s \leq$	Reference
3	8	[21]	9	21 738	Prop. 5.1
4	72	Th. 1.1	10	49 608	Prop. 5.1
5	305	Th. 1.1	11	111 721	Prop. 5.1
6	1 292	Th. 1.1	12	248 895	Prop. 5.1
7	3 954	Prop. 5.1	13	549 506	Prop. 5.1
8	9 370	Prop. 5.1	14	1 203 936	Prop. 5.1

The main idea of the proof is to utilize (1.6) with L being the maximal (by volume) inscribed ellipsoid into K (such ellipsoids were characterized by John [20]). After an appropriate affine transform, we can assume that L is the unit ball B_2^n in \mathbb{R}^n , while K is in the so-called John's position, so applying (1.6) we get

$$(1.8) \quad C(K, \text{int}K) \leq \frac{|K + B_2^n|}{|B_2^n|} \theta(B_2^n).$$

Next we make use of several geometric results which allow us to obtain an upper bound on $|K + B_2^n|$. One ingredient in such estimates is the fact that the mean width and the volume of a body in John's position are largest for the regular simplex (general case) or for the cube (symmetric case), the results due to Ball [3], Barthe [5], Schechtman, and Schmuckenschläger [32]. Another ingredient in our estimates is a Bonnesen-type inequality by Bokowski and Heil [7] on quermassintegrals of K . Finally, we use upper bounds on $\theta(B_2^n)$ for specific small n which arise from known lattice coverings.

Theorem 1.1 is proved in Sections 2–4. Of possibly independent interest are estimates of the mean width of the regular simplex in dimensions $5 \leq n \leq 8$ given in Section 3.

In Section 5, we show how one can improve (1.5) for each fixed n by optimizing choices of certain parameters in the original proof of Rogers [28] (the original proof provides a succinct bound valid for all n). Consequently, (1.1) and (1.4) can be somewhat improved for n larger than those covered by Theorem 1.1.

To finalize, in Tables 1 and 2 we provide the best known upper bounds on H_n and H_n^s for $3 \leq n \leq 14$.

2 Preliminaries

In what follows, $B_2^n = \{x \in \mathbb{R}^n : \|x\| \leq 1\}$, and $\mathbb{S}^{n-1} = \{x \in \mathbb{R}^n : \|x\| = 1\}$. The n -dimensional volume is denoted by $|\cdot|_n$, and subscript is usually dropped if the value of n is clear from the context.

2.1 John's position

Let \mathcal{J}_n be the family of convex bodies $K \in \mathcal{K}_n$ which are in John's position, i.e., B_2^n is the maximal volume ellipsoid of K (see, e.g., [9, Section 1.5.1]). Similarly, define \mathcal{J}_n^s to be the family of centrally symmetric convex bodies $K \in \mathcal{K}_n^s$ which are in John's position. For any $K \in \mathcal{K}_n$ (or $K \in \mathcal{K}_n^s$) there exists an affine image of K in \mathcal{J}_n (or in \mathcal{J}_n^s).

John's theorem (see e.g., [2, Theorem 2.1.3, Remark 2.1.17]) implies that

$$(2.1) \quad K \subseteq nB_2^n \quad \text{for any } K \in \mathcal{J}_n, \quad \text{and} \quad K \subseteq \sqrt{n}B_2^n \quad \text{for any } K \in \mathcal{J}_n^s.$$

2.2 Quermassintegrals

In order to bound H_n and H_n^s we will use the inequality (1.8), and so we start by discussing upper bounds on $|K + tB_2^n|$, where K is in John's position.

By Steiner's formula, for any $K \in \mathcal{K}_n$

$$(2.2) \quad |K + tB_2^n| = \sum_{j=0}^n \binom{n}{j} W_j(K) t^j,$$

where

$$W_j(K) = V(\underbrace{K, \dots, K}_{n-j \text{ times}}, \underbrace{B_2^n, \dots, B_2^n}_j)$$

is the j th quermassintegral of K and $V(\cdot)$ is the mixed volume, see, e.g., [2, Section 1.1.5], [9, Section 1.4.2] or [33, Section 4.2]. Additionally, $W_0(K) = |K|$, $W_1(K) = \partial(K)/n$, where $\partial(K)$ is the surface area of K , and $W_n(K) = |B_2^n|$. Also note that if $B_2^n \subseteq K$, then due to the monotonicity of mixed volumes we have $W_i(K) \geq W_j(K)$ for $i \leq j$.

Next, we will discuss upper bounds on $W_i(K)$ for $K \in \mathcal{J}_n$ and $K \in \mathcal{J}_n^s$ respectively.

Let T^n be a regular simplex in \mathbb{R}^n of unit edge length, and let Δ^n be a dilation of T^n for which B_2^n is the inscribed ball, then $\Delta^n = \sqrt{2(n+1)n} T^n$. Also, let $C^n := [-1, 1]^n$ be a cube circumscribed about B_2^n . Ball [3] proved that among all convex bodies in \mathbb{R}^n simplexes have maximal volume ratio (volume ratio measures how much of the volume of the whole body can be contained in the largest inscribed ellipsoid) while for the symmetric ones such a maximizer is the cube [4]. These results can be stated in our terms as follows:

$$(2.3) \quad W_0(K) \leq \begin{cases} W_0(\Delta^n), & \text{if } K \in \mathcal{J}_n, \\ W_0(C^n), & \text{if } K \in \mathcal{J}_n^s. \end{cases}$$

For $K \in \mathcal{K}_n$ and a direction $u \in \mathbb{S}^{n-1}$, the support function is defined as $h_K(u) = \sup\{\langle x, y \rangle : y \in K\}$. Let σ be the rotationally invariant probability measure on \mathbb{S}^{n-1} ,

then the mean width of K is defined by

$$w(K) = \int_{\mathbb{S}^{n-1}} h_K(u) d\sigma(u).^1$$

We have $W_{n-1}(K) = |B_2^n|w(K)$, which is a partial case of Kubota's formula [2, eq (1.1.1)]. Barthe [5, Theorem 3] proved² that among the bodies from \mathcal{J}_n , the mean width is maximized for Δ^n , while Schechtman and Schmuckenschläger [32] (the proof is also included in [5, Theorem 2]) remarked that C^n maximizes the mean width among the bodies from \mathcal{J}_n^s , i.e.,

$$(2.4) \quad W_{n-1}(K) \leq \begin{cases} W_{n-1}(\Delta^n), & \text{if } K \in \mathcal{J}_n, \\ W_{n-1}(C^n), & \text{if } K \in \mathcal{J}_n^s. \end{cases}$$

We estimate $w(\Delta^n)$ for the required values of n in the next section. It is known [15], [17, Section 13.2.3] (or can be obtained by a straightforward computation) that $w(C^n) = 2 \frac{|B_2^{n-1}|}{|B_2^n|}$ for $n \geq 2$, and so $W_{n-1}(C^n) = 2|B_2^{n-1}|$.

Remark 2.1 The results [3, Theorem 1, Theorem 1'] imply that $W_1(K) \leq W_1(\Delta^n)$ for any $K \in \mathcal{J}_n$ (recall that $W_1(K) = \partial(K)/n$, where $\partial(K)$ is the surface area of K). This would not lead to any improvements in our context as $W_1(\Delta^n) = W_0(\Delta^n)$ and we get the same upper bound on $W_1(K)$ as from $W_1(K) \leq W_0(K)$. A similar remark regarding the inequality $W_1(K) \leq W_1(C^n)$ also holds for $K \in \mathcal{J}_n^s$, and follows from [3, Theorem 2], [4, Theorem 3].

Remark 2.2 It is natural to conjecture that $W_j(K) \leq W_j(\Delta^n)$ (and that $W_j(K) \leq W_j(C^n)$) for any $K \in \mathcal{J}_n$ (respectively, $K \in \mathcal{J}_n^s$) and all $0 \leq j \leq n$. This is an obvious equality for $j = n$ and is valid for $j \in \{0, 1, n-1\}$ as described above ((2.3), Remark 2.1, (2.4)).

Finally, we need a Bonnesen-style inequality by Bokowski and Heil [7]. If $K \in \mathcal{K}_n$ satisfies $K \subseteq RB_2^n$, then for all $0 \leq i < j < k \leq n$

$$(2.5) \quad W_j(K) \leq \frac{(k-j)(i+1)R^i W_i(K) + (j-i)(k+1)R^k W_k(K)}{(k-i)(j+1)R^j} =: B_{R,i,j,k}(W_i(K), W_k(K)).$$

Recall that in our settings we can choose R according to (2.1).

2.3 Density of coverings by balls

One of the approaches to construct specific efficient coverings of the space by balls is to use lattices, see [11, Chapter 2]. Considering the A_n^* lattice yields the following

¹Note that sometimes the mean width is defined as $\int_{\mathbb{S}^{n-1}} h_K(u) + h_K(-u) d\sigma(u)$.

²The result of Barthe is stated in terms of ℓ -norm, and can be translated in the language of mean width using the formula [5, p. 685] for the ℓ -norm of the dual body.

Table 3: Least known lattice covering densities, as in [34].

n	Least known lattice covering density	n	Least known lattice covering density
2	1.209199	8	3.142202
3	1.463505	9	4.340185
4	1.765529	10	5.251713
5	2.124286	11	5.598338
6	2.464801	12	7.510113
7	2.900024	13	7.864060

estimate (valid for all n):

$$(2.6) \quad \theta(B_2^n) \leq |B_2^n| \sqrt{n+1} \left(\frac{n(n+2)}{12(n+1)} \right)^{n/2}.$$

It turns out that the above is optimal (smallest possible *lattice* covering density) for $2 \leq n \leq 5$ and provides the best known upper bound on $\theta(B_2^n)$ in most dimensions $10 \leq n \leq 21$. Better lattices were found by Schürmann and Vallentin [34] for $6 \leq n \leq 8$, which is important for our applications. To the best of our knowledge, there have been no improvements after the work [34], where an interested reader can find a brief survey of the topic. We list the corresponding lattice covering densities of \mathbb{R}^n by balls in Table 3.

3 Estimates on mean width of regular simplex

The values of $2w(T^n)$ for $2 \leq n \leq 6$ expressed as certain integrals and numerically evaluated with high precision can be found in [15]. We remark that in [15] expected values of widths of simplexes are considered, while the mean width as we defined here (and as commonly defined in the literature on geometry, see [2, 9]) is the expected value of the support function, this discrepancy results in the mean width of T^n from [15] (and [35]) being equal to $2w(T^n)$. We need upper estimates of $w(T^n)$ for $n = 7, 8$, which we were unable to find in the literature. We present a simple computational technique to obtain upper and lower estimates on $w(T^n)$ which will suffice for our purposes.

The starting point is the following representation that follows the work [15] by Finch and references therein, in particular, Sun [35].

Let $F(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-t^2/2} dt$ be the cumulative distribution function of the standard normal distribution, then $F(x) = \frac{1}{2} + \frac{1}{2} \operatorname{erf}\left(\frac{x}{\sqrt{2}}\right)$, where $\operatorname{erf}(z) = \frac{2}{\sqrt{\pi}} \int_0^z e^{-t^2} dt$ is the error function.

For $n \geq 0$ let

$$g_{n+1}(x) := 1 - F(x)^{n+1} - (1 - F(x))^{n+1} = 1 - \left(\frac{1 + \operatorname{erf}\left(\frac{x}{\sqrt{2}}\right)}{2} \right)^{n+1} - \left(\frac{1 - \operatorname{erf}\left(\frac{x}{\sqrt{2}}\right)}{2} \right)^{n+1}.$$

The following formula for mean width follows from [35] and was derived in [15]:

$$(3.1) \quad w(T^n) = \frac{\Gamma(\frac{n}{2})}{2\Gamma(\frac{n+1}{2})} \int_0^\infty g_{n+1}(x) dx.$$

Next, we estimate the integral in the formula.

Proposition 3.1 *For any $a > 2$ and any positive integers n, N , the following estimates hold:*

$$(3.2) \quad \frac{a}{N} \sum_{k=1}^N g_{n+1}\left(\frac{ak}{N}\right) \leq \int_0^\infty g_{n+1}(x) dx \leq \frac{a}{N} \sum_{k=0}^{N-1} g_{n+1}\left(\frac{ak}{N}\right) + \frac{n+1}{\sqrt{2\pi}} \exp(-a).$$

Proof We directly estimate the “tail” of the integral and use simple endpoint Riemann sums for the “main” part of the integral in (3.1).

Recalling that $\frac{d}{dx}F(x) = \frac{1}{\sqrt{2\pi}}e^{-x^2/2}$ it is straightforward to verify that g_{n+1} is a positive strictly decreasing function on $[0, \infty)$. Hence, considering the upper and the lower Riemann sums for $\int_0^a g_{n+1}(x) dx$ and the uniform partition of $[0, a]$ into N subintervals, we obtain

$$(3.3) \quad \frac{a}{N} \sum_{k=1}^N g_{n+1}\left(\frac{ak}{N}\right) \leq \int_0^a g_{n+1}(x) dx \leq \frac{a}{N} \sum_{k=0}^{N-1} g_{n+1}\left(\frac{ak}{N}\right).$$

For any $x > a \geq 2$

$$\begin{aligned} g_{n+1}(x) &\leq 1 - F(x)^{n+1} = 1 - (1 - F(-x))^{n+1} = 1 - \left(1 - \frac{1}{\sqrt{2\pi}} \int_x^\infty e^{-t^2/2} dt\right)^{n+1} \\ &\leq \frac{n+1}{\sqrt{2\pi}} \int_x^\infty e^{-t^2/2} dt \leq \frac{n+1}{\sqrt{2\pi}} \int_x^\infty e^{-t} dt = \frac{n+1}{\sqrt{2\pi}} \exp(-x), \end{aligned}$$

so

$$\int_a^\infty g_{n+1}(x) dx \leq \frac{n+1}{\sqrt{2\pi}} \exp(-a).$$

Taking this inequality and the evident $\int_a^\infty g_{n+1}(x) dx \geq 0$ into account, we deduce (3.2) from (3.1) and (3.3). ■

Since the error function can be computed numerically with any given precision, employing a simple SageMath [31] computation [1], we obtain the following corollary.

Corollary 3.2 *The following inequalities hold:*

$$0.4208 \leq w(T^5) \leq 0.4215,$$

$$0.4067 \leq w(T^6) \leq 0.407,$$

$$0.39425 \leq w(T^7) \leq 0.39427,$$

$$0.383 \leq w(T^8) \leq 0.38301.$$

These estimates for $n = 5, 6$ are consistent with the values obtained in [15]; we include them here for completeness. While our computational method allows to obtain a tighter gap in the estimates, we only derived what was necessary for our application of estimating H_n from above ensuring that no further improvement is possible (even if the value of the lower bound on $w(T^n)$ is used, the upper bound on the integer value H_n does not improve). The computations take less than two hours on a modern personal computer.

4 Proof of Theorem 1.1

Let us begin with the general (not necessarily symmetric) case. Since $C(K, \text{int}(K))$ is invariant under affine transforms, we can assume that $K \in \mathcal{J}^n$. Then using $B_2^n \subseteq K$, compactness and (1.6), we have

$$\begin{aligned} C(K, \text{int}(K)) &\leq C(K, \text{int}(B_2^n)) \leq \lim_{r \rightarrow 1^-} C(K, rB_2^n) \\ &\leq \lim_{r \rightarrow 1^-} \frac{|K - rB_2^n|}{|rB_2^n|} \theta(B_2^n) = \frac{|K + B_2^n|}{|B_2^n|} \theta(B_2^n). \end{aligned}$$

Hence, applying (2.2),

$$(4.1) \quad C(K, \text{int}(K)) \leq \frac{\theta(B_2^n)}{|B_2^n|} \sum_{j=0}^n \binom{n}{j} W_j(K).$$

The upper bounds for $\theta(B_2^n)$ come from Table 3, so it remains to estimate $W_i(K)$. As $\Delta^n = \sqrt{2n(n+1)}T^n$, by (2.3),

$$(4.2) \quad W_i(K) \leq W_0(K) \leq W_0(\Delta^n) = (2n(n+1))^{n/2} |T^n| = \frac{n^{n/2}(n+1)^{(n+1)/2}}{n!}$$

for any $0 \leq i \leq n$. Using (2.4),

$$(4.3) \quad W_{n-1}(K) \leq W_{n-1}(\Delta^n) = |B_2^n| w(\Delta^n) = |B_2^n| \sqrt{2n(n+1)} w(T^n).$$

Trivially, $W_n(K) = |B_2^n|$.

Next we combine the above, and use (2.5) (valid with $R = n$ due to (2.1)) for appropriate parameters. The calculations were performed in the script [1].

Bound on H_5 . We apply (4.1) and estimate the quermassintegrals as follows. For $0 \leq i \leq 2$, use (4.2); (4.3) and (2.5) give $W_4(K) \leq W_4(\Delta^5)$ and $W_3(K) \leq B_{5,2,3,4}(W_0(\Delta^5), W_4(\Delta^5))$; recall that $W_5(K) = |B_2^5|$. Combining the above, using Corollary 3.2 and calculating the actual value of the bound yields $H_5 \leq 933$.

Bound on H_6 . The arguments are similar to the previous case. The required application of (2.5) is $W_4(K) \leq B_{6,3,4,5}(W_0(\Delta^6), W_5(\Delta^6))$. The result is $H_6 \leq 6137$.

Bounds on H_n for $n = 7, 8$. We proceed similarly with the only difference that $W_j(K) \leq B_{n,n-4,j,n-1}(W_0(\Delta^n), W_{n-1}(\Delta^n))$ is used for $j \in \{n-3, n-2\}$ yielding $H_7 \leq 41377$ and $H_8 \leq 284096$.

When the body is centrally symmetric, we follow a similar route, with the following differences. In place of (4.2), we have

$$(4.4) \quad W_i(K) \leq W_0(K) \leq W_0(C^n) = 2^n$$

for any $0 \leq i \leq n$. Using (2.4) we get

$$(4.5) \quad W_{n-1}(K) \leq W_{n-1}(C^n) = 2|B_2^{n-1}|.$$

Finally, by (2.1), the inequality (2.5) can be used with $R = \sqrt{n}$.

Bound on H_4^s . Apply (4.1) and bound the quermassintegrals as follows. For $i = 0, 1$ use (4.4); use (4.5) to get $W_3(K) \leq 2|B_2^3|$; we have $W_4(K) = W_4(C^4) = |B_2^4|$; by (2.5) $W_2(K) \leq B_{2,1,2,3}(W_1(C^4), W_3(C^4))$. Combining the above and calculating the value of the bound yields $H_4^s \leq 72$.

Bounds on H_n^s for $n = 5, 6$. We use similar arguments: (4.4) for $i = 0, 1, 2$; $W_n(K) = W_n(C^n) = |B_2^n|$; (4.5) for $i = n - 1$; and $W_j(K) \leq B_{\sqrt{n}, 2, j, n-1}(W_0(C^n), W_{n-1}(C^n))$ for $3 \leq j \leq n - 2$, which imply $H_5^s \leq 305$ and $H_6^s \leq 1292$.

Remark 4.1 In the above computations, we used the exact value for the density arising from A_n^* lattice from the estimate (2.6) for $n = 4, 5$. For $n = 6, 7, 8$ we used the values of covering densities from [34] presented in Table 3 increased by $5 \cdot 10^{-6}$ as they were given to 6 decimal places, while we require an upper bound. Even if such an increase is not performed, the resulting integer valued upper bounds in Theorem 1.1 would not change, so the accuracy given in [34] is more than sufficient for our needs.

5 Estimates via optimized Rogers bound

Proposition 5.1 Suppose $n \geq 3$. If

$$(5.1) \quad r_n = \min_{x \in (0, 1/n)} f_n(x), \quad \text{where} \quad f_n(x) = (1+x)^n(1-n \ln(x)),$$

then

$$(5.2) \quad \theta(K) \leq r_n \quad \text{for any} \quad K \in \mathcal{K}_n, \quad H_n \leq \binom{2n}{n} r_n, \quad \text{and} \quad H_n^s \leq 2^n r_n.$$

Proof The first inequality in (5.2) is established in [28, p. 5], p. 5. (The bound (1.5) was obtained in [28] by taking $x = \frac{1}{n \ln n}$ in (5.1).) The other two inequalities in (5.2) follow from (1.6) in the same way as (1.1) and (1.4) do. ■

For our purposes, it suffices to use the straightforward $r_n \leq \min\{f_n(\frac{j}{Nn}) : 1 \leq j \leq N-1\}$ with $N = 1000$. The resulting bounds on r_n for $3 \leq n \leq 14$ are given in Table 4 (the computed values of $\min\{f_n(\frac{j}{Nn}) : 1 \leq j \leq N-1\}$ are rounded up in the sixth digit, so they represent actual upper bounds), and the estimates on H_n and H_n^s can be found in Tables 1 and 2. For the computations, see [1]. We remark that $\max\{\theta(K), K \in \mathcal{K}_2\} = \frac{3}{2}$ was established by Fáry [14].

Table 4: Upper bounds on $\max\{\theta(K), K \in \mathcal{K}_n\}$, for $3 \leq n \leq 14$.

n	$\max\{\theta(K), K \in \mathcal{K}_n\} \leq$	n	$\max\{\theta(K), K \in \mathcal{K}_n\} \leq$
3	10.064123	9	42.458503
4	14.916986	10	48.445515
5	20.024359	11	54.551530
6	25.362768	12	60.765566
7	30.898293	13	67.078451
8	36.603890	14	73.482436

Acknowledgments The authors are grateful to the referees for the numerous comments that improved the paper.

References

- [1] A. Arman, A. Bondarenko, and A. Prymak, *Sagemath script for computations on Hadwiger's covering problem in small dimensions*. available at <https://github.com/andriyprym/illum-small-dims/releases/tag/v1> (accessible as of May 2025).
- [2] S. Artstein-Avidan, A. Giannopoulos, and V. D. Milman, *Asymptotic geometric analysis. Part I*, Mathematical Surveys and Monographs, 202, American Mathematical Society, Providence, RI, 2015.
- [3] K. Ball, *Volume ratios and a reverse isoperimetric inequality*. J. London Math. Soc. (2) 44(1991), no. 2, 351–359.
- [4] K. Ball, *Volumes of sections of cubes and related problems*. In: Geometric aspects of functional analysis (1987–88), Lecture Notes in Mathematics, 1376, Springer, Berlin, 1989, pp. 251–260. <https://doi.org/10.1007/BFb0090058>. MR1008726
- [5] F. Barthe, *An extremal property of the mean width of the simplex*. Math. Ann. 310(1998), no. 4, 685–693.
- [6] K. Bezdek and M. A. Khan, *The geometry of homothetic covering and illumination*. In: Discrete geometry and symmetry, Springer Proceedings in Mathematics & Statistics, 234, Springer, Cham, 2018, pp. 1–30.
- [7] J. Bokowski and E. Heil, *Integral representations of quermassintegrals and Bonnesen-style inequalities*. Arch. Math. (Basel) 47(1986), no. 1, 79–89.
- [8] V. Boltyanski, *The problem of illuminating the boundary of a convex body*. Izv. Mold. Fil.ANSSSR 76(1960), 77–84.
- [9] S. Brazitikos, A. Giannopoulos, P. Valettas, and B.-H. Vritsiou, *Geometry of isotropic convex bodies*, Mathematical Surveys and Monographs, 196, American Mathematical Society, Providence, RI, 2014.
- [10] M. Campos, P. van Hintum, R. Morris, and M. Tiba, *Towards Hadwiger's conjecture via Bourgain slicing*. Int. Math. Res. Not. IMRN 10(2024), 8282–8295.
- [11] J. H. Conway and N. J. A. Sloane, *Sphere packings, lattices and groups*. 3rd ed., Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 290, Springer-Verlag, New York, NY, 1999. With additional contributions by E. Bannai, R. E. Borcherds, J. Leech, S. P. Norton, A. M. Odlyzko, R. A. Parker, L. Queen and B. B. Venkov.
- [12] M. Diao, *On Hadwiger covering problem in five- and six-dimensional Euclidean spaces*. M.Sc. thesis, University of Manitoba, 2022. available at <http://hdl.handle.net/1993/36378>.
- [13] P. Erdős and C. A. Rogers, *The star number of coverings of space with convex bodies*. Acta Arith. 9(1964), 41–45.
- [14] I. Fáry, *Sur la densité des réseaux de domaines convexes*. Bull. Soc. Math. France 78(1950), 152–161 (French).
- [15] S. R. Finch, *Mean width of a regular simplex*. Preprint. available at <https://arxiv.org/abs/1111.4976v2>.
- [16] I. C. Gohberg and A. S. Markus, *A certain problem about the covering of convex sets with homothetic ones*. Izv. Moldavskogo Filiala Akad. Nauk SSSR 10(1960), no. 76, 87–90 (Russian).

- [17] J. E. Goodman and J. O'Rourke (eds.), *Handbook of discrete and computational geometry*, CRC Press Series on Discrete Mathematics and its Applications, CRC Press, Boca Raton, FL, 1997. Edited by Jacob E. Goodman and Joseph O'Rourke. MR1730156
- [18] H. Hadwiger, *Ungelöstes problem Nr. 20*. Elem. Math. 12(1957), 121 (German).
- [19] H. Huang, B. A. Slomka, T. Tkocz, and B.-H. Vritsiou, *Improved bounds for Hadwiger's covering problem via thin-shell estimates*. J. Eur. Math. Soc. (JEMS) 24(2022), no. 4, 1431–1448.
- [20] F. John, *Extremum problems with inequalities as subsidiary conditions*. In: Studies and essays presented to R. Courant on his 60th birthday, Interscience Publishers, Inc., New York, NY, 1948, Jan. 8, 1948, pp. 187–204.
- [21] M. Lassak, *Solution of Hadwiger's covering problem for centrally symmetric convex bodies in E^3* . J. London Math. Soc. (2) 30(1984), no. 3, 501–511.
- [22] M. Lassak, *Covering the boundary of a convex set by tiles*. Proc. Amer. Math. Soc. 104(1988), no. 1, 269–272.
- [23] M. Lassak, *Covering a three-dimensional convex body by smaller homothetic copies*. Beitrage Algebra Geom. 39(1998), no. 2, 259–262.
- [24] F. W. Levi, *Überdeckung eines Eibereiches durch Parallelverschiebung seines offenen Kerns*. Arch. Math. (Basel) 6(1955), 369–370 (German).
- [25] I. Papadoperakis, *An estimate for the problem of illumination of the boundary of a convex body in E^3* . Geom. Dedicata 75(1999), no. 3, 275–285.
- [26] A. Prymak, *A new bound for Hadwiger's covering problem in E^3* . SIAM J. Discrete Math. 37(2023), no. 1, 17–24.
- [27] A. Prymak and V. Shepelska, *On the Hadwiger covering problem in low dimensions*. J. Geom. 111(2020), no. 3, Paper No. 42, 11.
- [28] C. A. Rogers, *A note on coverings*. Mathematika 4 (1957), 1–6.
- [29] C. A. Rogers and G. C. Shephard, *The difference body of a convex body*. Arch. Math. (Basel) 8(1957), 220–233.
- [30] C. A. Rogers and C. Zong, *Covering convex bodies by translates of convex bodies*. Mathematika 44(1997), no. 1, 215–218.
- [31] The Sage Developers, *Sagemath, the sage mathematics software system (version 9.2)*, 2020. <https://www.sagemath.org>.
- [32] G. Schechtman and M. Schmuckenschläger, *A concentration inequality for harmonic measures on the sphere*. In: Geometric aspects of functional analysis (Israel, 1992), Operator Theory: Advances and Applications, 77, Birkhäuser, Basel, 1995, pp. 255–273.
- [33] R. Schneider, *Convex bodies: The Brunn-Minkowski theory, expanded edition*, Encyclopedia of Mathematics and its Applications, 151, Cambridge University Press, Cambridge, 2014.
- [34] A. Schürmann and F. Vallentin, *Computational approaches to lattice packing and covering problems*. Discrete Comput. Geom. 35(2006), no. 1, 73–116.
- [35] Y. Sun, *Stochastic iterative algorithms for signal set design for Gaussian channels and optimality of the signal set*. IEEE Trans. Inform. Theory 43(1997), no. 5, 1574–1587.

Department of Mathematics, University of Manitoba, Winnipeg, MB, R3T 2N2, Canada

e-mail: andrew0arman@gmail.com prymak@gmail.com

Department of Mathematical Sciences, Norwegian University of Science and Technology, NO-7491

Trondheim, Norway

e-mail: andriybond@gmail.com